



## ABSTRACT

*Da spedire entro il 30 Settembre a*

*Prof. Paolo Quadrelli  
e-mail: paolo.quadrelli@unipv.it*

### **AUTORE**

<i>Valentina Pirovano</i>		
<i>Università degli Studi di Milano, Dipartimento di Scienze Farmaceutiche, Sezione di Chimica Generale e Organica A. Marchesini</i>		
<i>Via G. Venezian 21</i>		<i>Milano</i>
<i>+390250314477</i>	<i>+390250314476</i>	<i>valentina.pirovano@unimi.it</i>

*Catalytic transformations of indoles: recent achievements and new perspectives for the synthesis of complex indole derivatives*


*Valentina Pirovano, Elisa Brambilla, Elisabetta Rossi, Giorgio Abbiati*

# CATALYTIC TRANSFORMATIONS OF INDOLES: RECENT ACHIEVEMENTS AND NEW PERSPECTIVES FOR THE SYNTHESIS OF COMPLEX INDOLE DERIVATIVES

Valentina Pirovano, Elisa Brambilla, Elisabetta Rossi and Giorgio Abbiati

Dipartimento di Scienze Farmaceutiche, Sez. di Chimica Generale e Organica "A. Marchesini", Università degli Studi di Milano

Transformation of simple indoles into (polycyclic) complex scaffolds has become the object of intensive studies in synthetic organic chemistry due to ubiquitous occurrence of indole core in the structure of relevant molecules.<sup>1</sup> In particular, catalytic-promoted manipulation of indoles have become an incomparable tool to increase indole structural complexity working under exceedingly mild conditions and in a regio- and stereo-controlled fashion.<sup>2</sup> In this context we reported in the last years the synthesis of complex carbazole derivatives through gold-catalyzed (4+2) cycloaddition reactions of vinylindoles with different  $\pi$ -systems<sup>3</sup> and the functionalization of structurally simple indoles with challenging metal-activated electrophiles.<sup>4</sup> Taking into account these premises, this oral communication will deal with our recent achievements in the field of cycloaddition reactions involving vinylindoles as  $4\pi$  systems and in gold-catalyzed cascade reactions on indole-based scaffolds. In particular we were able to extend cycloaddition of vinylindoles besides (4+2) processes to synthesize cyclohepta[b]indoles. Moreover, we explored the reactivity of 4*H*-furo[3,2-*b*]indoles with gold-activated  $\pi$ -systems to synthesize 2-spiro-3-oxindoles and 2-alkenyliden-indolin-3-ones selectively.<sup>5</sup> Advantages of our methods as well as limitations and future perspectives will be discussed.



1. a) H. Plieninger, *Angew. Chem. Int. Ed.*, **10**, 361-631 (1971); b) P. Kumar et al., *J. Heterocycl. Chem.*, **47**, 491-502 (2010); c) S. L. Manju et al., *Eur. J. Pharm. Sci.*, **91**, 1-10 (2016) d) O. Silakari et al., *Eur. J. Med. Chem.*, **134**, 159-184 (2017)
2. a) M. Bandini et al., *Angew. Chem. Int. Ed.*, **48**, 9608-9644, (2009); b) Y.-X. Jia et al., *Org. Biomol. Chem.*, **15**, 3550-3567 (2017).
3. E. Rossi et al., *Eur. J. Org. Chem.*, 4512-4529 (2017).
4. a): V. Pirovano, *Eur. J. Org. Chem.*, 1925-1945 (2018); b) for a recent copper-catalyzed example see: V. Pirovano et al., *Org. Lett.*, **20**, 405-408 (2018).
5. a) E. Rossi et al, *J. Org. Chem.*, **84**, 5150-5166 (2019); b) E. Rossi, E. Brambilla et al., *Org. Chem. Front.*, **6**, 3078-3084 (2019).