

X-ray properties in massive galaxy clusters: XMM-Newton observations of the REFLEX-DXL sample (*Corrigendum*)

Y.-Y. Zhang¹, H. Böhringer¹, A. Finoguenov¹, Y. Ikebe^{1,2}, K. Matsushita^{1,3}, P. Schuecker¹,
L. Guzzo⁴, and C. A. Collins⁵

¹ Max-Planck-Institut für extraterrestrische Physik, Giessenbachstraße, 85748 Garching, Germany
e-mail: yyzhang@astro.uni-bonn.de

² National Museum of Emerging Science and Innovation, Tokyo, Japan

³ Tokyo University of Science, Tokyo, Japan

⁴ INAF – Osservatorio Astronomico di Brera, Merate/Milano, Italy

⁵ Liverpool John Moores University, Liverpool, UK

A&A 456, 55–74 (2006), DOI:[10.1051/0004-6361:20053650](https://doi.org/10.1051/0004-6361:20053650)

ABSTRACT

The article by Zhang et al. (2006, A&A, 456, 55) contains an error in Fig. 5 on page 64, in which we inadvertently re-called the mass distribution of RXCJ0528.9–3972. We note that this does not affect any results in Zhang et al. (2006). Nevertheless, we regret this error and provide an update of Fig. 5 here.

Key words. cosmology: observations – galaxies: clusters: general – X-rays: galaxies: clusters – dark matter – errata, addenda

References

Zhang, Y.-Y., Böhringer, H., Finoguenov, A., et al. 2006, A&A, 456, 55




Fig. 5. Measured mass profiles of the REFLEX-DXL clusters and their best fits by the extended-NFW models. Residuals scaled by the data uncertainties are plotted in the upper part of the diagrams.