



# The emerging predictive role of C-reactive protein after ventral hernia repair with mesh

A. Aiolfi<sup>1</sup> · G. Bonitta<sup>1</sup> · D. Bona<sup>1</sup>

Received: 17 February 2020 / Accepted: 22 February 2020  
© Springer-Verlag France SAS, part of Springer Nature 2020

Dear Editor,

We have read with interest the retrospective analysis by Pochhammer et al. [1]. The authors reported a series of 373 patients who underwent conventional ventral hernioplasty with mesh augmentation in a referral hernia center. Postoperative C-reactive protein (CRP) measurement was consecutively performed in all patients during the hospital course. Prevalence of postoperative infectious complications (ICs) was 13.7% ( $n = 51$ ). Authors reported positive predictive values (PPV) and negative predictive values (NPV) at different postoperative days (POD). PPVs and NPVs for POD 2/3 and POD were 0.29–0.46 and 1.00–0.93, respectively. The authors concluded that postoperative serum CRP allows for early prediction of the postoperative course with low CRP values being associated with reduced risk of ICs.

We congratulate the authors for the idea to investigate the postoperative CRP predictive value in the diagnosis of ICs after mesh-reinforced ventral hernioplasty. This paper strengthens and broadens this concept previously investigated in other general surgery settings [2–4]. After carefully reading, CRP seems to be a useful biomarker to rule out postoperative ICs. This is supported by the estimated low POD 2/3 and POD 5/6 PPVs (0.29–0.46, respectively) and high NPVs (1.00 and 0.93, respectively). It is true that PPV and NPV are influenced by the disease prevalence (prior probability) but having such high estimated NPVs ( $> 0.90$ ), results are less affected by disease prevalence.

We thank the authors for their remarkable work and for having added further empirical evidence that confirms the emerging role of postoperative CRP predictive values. Supported by previous published evidence and authors' data,

CRP may have a diagnostic role with a high level of confidence. In conclusion, CRP should not be interpreted as marker to diagnose but rather as a useful tool to rule out a diagnosis. However, postoperative CRP should not be considered a panacea, but in conjunction with reassuring clinical signs it may be an additional arrow in each surgeon's quiver.

**Conflict of interest** AA, GB, and DB declare no conflicts of interest.

**Ethical approval** Approval from the Institutional review board was not required for this study.

**Human and animal right** This article does not contain any studies with human participants or animals performed by any of the authors.

**Informed consent** For this type of study formal consent is not required.

## References

1. Pochhammer J, Scholtes B, Keuler J et al (2020) Serum C-reactive protein level after ventral hernia repair with mesh reinforcement can predict infectious complications: a retrospective cohort study. *Hernia* 24(1):41–48
2. Bona D, Micheletto G, Bonitta G et al (2019) Does C-reactive protein have a predictive role in the early diagnosis of postoperative complications after bariatric surgery? systematic review and Bayesian meta-analysis. *Obes Surg* 29(11):3448–3456. <https://doi.org/10.1007/s11695-019-04013-0>
3. Singh PP, Zeng IS, Srinivasa S et al (2014) Systematic review and meta-analysis of use of serum C-reactive protein levels to predict anastomotic leak after colorectal surgery. *Br J Surg* 101(4):339–346. <https://doi.org/10.1002/bjs.9354>
4. Aiolfi A, Asti E, Rausa E et al (2018) Use of C-reactive protein for the early prediction of anastomotic leak after esophagectomy: systematic review and Bayesian meta-analysis. *PLoS ONE* 13(12):e0209272

✉ A. Aiolfi  
alberto.aiolfi86@gmail.com

<sup>1</sup> Department of Biomedical Science for Health, Istituto Clinico Sant'Ambrogio, University of Milan, Via Luigi Giuseppe Faravelli, 16, 20149 Milan, Italy

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.