

1  
2  
3 **O<sub>2</sub>-requiring molecular reporters of gene**  
4 **expression for anaerobic microorganisms**  
5

6  
7 Simone Guglielmetti<sup>1</sup>, Ville Santala<sup>2</sup>, Rahul Mangayil<sup>2</sup>, Alessandro  
8 Ciranna<sup>1, 2</sup> and Matti T. Karp<sup>\*2</sup>  
9

10  
11 <sup>1</sup>Department of Food, Environmental and Nutritional Sciences (DeFENS),  
12 University of Milan, Italy and <sup>2</sup>Laboratory of Chemistry and Bioengineering,  
13 Tampere University of Technology, Finland  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33

34 \*Corresponding author: Matti Karp, Laboratory of Chemistry and  
35 Bioengineering, Tampere University of Technology, Korkeakoulunkatu 8, PO  
36 Box 541, 33101 Tampere Finland, Phone +358-40-5533189, e-mail  
37 matti.karp@tut.fi  
38  
39

40 **Key words:** bioluminescence, Gram (+), Gram (-), luciferase, arabinose induction  
41

42 **ABSTRACT**

43

44 Many genetic reporter systems require molecular oxygen; therefore, the use of  
45 reporter genes to study molecular mechanisms in anaerobic microorganisms has  
46 been hampered by the lack of convenient reporting systems. We describe reporter  
47 gene whole cell-based biosensor systems based on luciferase genes and the  
48 associated oxygen-requiring enzymes. By using two different oxygen-dependent  
49 reporters, insect and bacterial luciferases, and two bacterial hosts, Gram (+)  
50 *Bifidobacterium longum* and Gram (-) *Escherichia coli*, we show that the enzymes  
51 can be used in gene expression studies of anaerobic bacteria. *E. coli*, a facultative  
52 anaerobe, was grown both in aerobic and anaerobic conditions with an arabinose-  
53 inducible expression system. We show that a short treatment time of few minutes in  
54 ambient atmosphere is sufficient to detect light emission from living cells that is  
55 directly proportional to the number of cells and to the inducer concentration. The  
56 induction levels were the same in both the aerobically and anaerobically cultured  
57 cells. Similar results were obtained in the case of *B. longum* cultured in anaerobic  
58 conditions.

59

60

61

62

63

## 64 **1. INTRODUCTION**

65 Anaerobic microbes are an important class of organisms because they can be used in  
66 a vast number of biotechnological applications, and these microbes have been used since  
67 ancient times in the production of food and feed, with wine and beverages being the most  
68 well-known products. Upon the origin of life on Earth approximately 3.9 billion years ago  
69 (Rizzotti 2009), the conditions were anaerobic, and the first living organisms were  
70 heterotrophic. The abundance of sub-seafloor microbes is expected to be  $35.5 \cdot 10^{29}$  cells,  
71 comprising 55–86% of Earth's prokaryotic biomass and 27–33% of Earth's living biomass,  
72 and most of these cells are anaerobic, methanogenic bacteria or archaea, which dominate  
73 in deep sea sediments (Whitman et al. 1998); therefore, they present a vast, nearly unknown  
74 reservoir of organisms that have not yet been studied at the gene expression level.  
75 Furthermore, anaerobic microorganisms have an enormous market potential in various  
76 biotechnological applications, such as anaerobic digestion in sewage treatment systems,  
77 bioenergy production ( $H_2$ ), alcohols and methane bioprocesses, production of fermented  
78 food products and in metabolic engineering in cosmetics, prebiotics and medicines.  
79 Therefore, it is strongly justified to identify convenient and simple methods for studying gene  
80 expression in both Gram (+) and Gram (-) anaerobic microorganisms, and also fungi and  
81 yeasts.


82 Research with anaerobic organisms has lagged behind that with aerobic  
83 microorganisms, partly because there are too few powerful molecular tools for efficiently  
84 studying the intracellular mechanisms in physiology and biochemistry at the molecular level.  
85 Reporter genes are tools that are used in place of each target regulon in cases where gene  
86 expression and its regulation are studied (Daurert et al. 2000; Elad et al. 2008). The gene

87 encoding  $\beta$ -galactosidase has been the most widely used reporter in anaerobic research  
88 because its product can be easily measured by a simple colour reaction (Feustel et al. 2004).  
89 Green fluorescent protein (GFP) and its colour variants have been shown to be unsuitable  
90 reporters in anaerobic bacteria because their correct chromophore formation requires  
91 aerobic conditions (Hansen et al. 2001; Tsien 1998). Drepper et al. (Drepper et al. 2007)  
92 showed that blue-light photoreceptors from *Bacillus subtilis* and *Pseudomonas aeruginosa*  
93 contain light-oxygen-voltage-sensing domains that can be engineered to work as fluorescent  
94 reporters for gene expression analysis under both aerobic and anaerobic conditions.  
95 Bioluminescent proteins require oxygen as a substrate in their light-emitting reaction.  
96 Reporter genes supporting bioluminescent or fluorescent detection technologies have  
97 several benefits over other approaches whose activities are detected by spectrophotometry,  
98 such as  $\beta$ -galactosidase (Feustel et al. 2004) and alkaline phosphatase (Edwards et al.  
99 2015), or radioactivity (chloramphenicol acetyltransferase), namely, high sensitivity and real-  
100 time detection. Most of these approaches, with the exception of luciferases, require cell  
101 disruption as an extra step in activity measurements. In principle, luciferase reporters have  
102 the highest sensitivity because biological samples do not produce light of their own (except  
103 for light-emitting organisms), given that the measurement chamber is a black, light-tight box  
104 that blocks ambient light. The use fluorescent reporters suffers from cellular  
105 autofluorescence that is triggered upon excitation, thus lowering the sensitivity. Both  
106 bacterial and insect luciferases (Michelini et al. 2008) are widely used as molecular reporter  
107 systems, but their use in anaerobic applications is believed to be restricted by the  
108 requirement for oxygen in the light-emitting reaction.

109 There are few reports on the use of luciferase-based reporters in anaerobic  
110 microorganisms. However, there are no in-depth analyses of how they function. For


111 instance, the first paper by Phillips-Jones (Phillips-Jones 1993) described the use of the  
112 *Vibrio fischeri luxAB* genes for monitoring gene expression in anaerobically cultured  
113 *Clostridium perfringens*. Not until ten years later were the *luxAB* genes used to monitor the  
114 effects of antimicrobial agents against nonreplicating, anaerobically grown *Mycobacterium*  
115 *tuberculosis* (Cho et al. 2007). In a decade-old report, fusions of the mercury resistance  
116 operon (*mer*) with a promoterless *luxCDABE* operon from *Vibrio fischeri* (Selifonova et al.  
117 1993) were used to study the effect of intracellular pH on the accumulation of trace  
118 concentrations in *Escherichia coli* in anaerobic conditions (Golding et al. 2008). In that study,  
119 as in the previous examples, the measurement of light emission from the luciferase reporters  
120 was measured by shaking the samples in aerobic conditions after growth in anaerobic  
121 conditions, and the experimental setups were not shown in detail. It was found that only the  
122  $\beta$ -galactosidase reporter gene worked properly in a tumour hypoxia model when monitored  
123 with *in vivo* imaging (Cecic et al. 2007). This result shows that it is important to provide  
124 molecular oxygen to the cells expressing reporter gene products that require oxygen for  
125 catalysis.

126 The light emission reaction by bacterial luciferases has the following formula:



129 where RCHO is a long-chain aldehyde and RCOOH is the corresponding fatty acid. Light-  
130 emission can be obtained from cells containing the structural genes of the bacterial  
131 luciferase operon, *luxCDABE*, without any external additives.

132 The reaction catalysed by insect luciferases has the following formula:



135 Light-emission can be obtained from cells that are supplied with external D-luciferin.

136 Both types of luciferase reporter proteins require molecular oxygen for catalysis. We  
137 report here that by applying a short recovery step under aerobic atmosphere prior to  
138 bioluminescence measurements, bacterial and insect luciferases (biosensing elements) can  
139 be used as molecular reporters in anaerobically cultured *Escherichia coli* and  
140 *Bifidobacterium longum* model organisms. We also show that gene expression closely  
141 parallels the dose-response using induction of the arabinose-inducible promoter in both  
142 aerobic and anaerobic conditions.

143

144

## 145 **2. MATERIALS AND METHODS**

### 146 **2.1 Construction of Sak-Lux plasmid for *Escherichia coli* expression**

147

148 The construction of the arabinose-inducible, bacterial luciferase operon-containing plasmid  
149 is shown in the **Supplementary Material**.

150

### 151 **2.2 Construction and cultivation of bifidobacteria harbouring the shuttle plasmid** 152 **expressing the click beetle luciferase gene.**

153 Electrocompetent *Bifidobacterium longum* cells were prepared as described earlier  
154 (Guglielmetti et al. 2007). The human intestinal anaerobic bacterium *B. longum* NCC2705  
155 was transformed by electroporation with a shuttle vector (pGBL8b) containing the insect  
156 luciferase gene from a click beetle (*Pyrophorus plagiophthalmus*) under the control of a  
157 strong phage T5 promoter. The bioluminescence reporter vector pGBL8b was constructed  
158 earlier (Guglielmetti et al. 2008). Electroporation was carried out with a MicroPulser

159 Electroporator (Bio-Rad, Milano, Italy) set at 12.5 kV/cm and employing a 2-mm cuvette (the  
160 time constants obtained were between 3.9 and 4.2 ms). This recombinant bifidobacterial  
161 strain was cultivated in anaerobic conditions at 37°C in MRS medium (Difco Laboratories  
162 Inc., Detroit, MI) supplemented with 0.05% cysteine-HCl and 10 µg/ml of chloramphenicol.  
163 Anaerobic conditions were created by the use of an anaerobic glove box (80% N<sub>2</sub>, 10% CO<sub>2</sub>,  
164 10% H<sub>2</sub>). The resulting clones displayed a bioluminescent phenotype. Light emission was  
165 studied in relation to anaerobic conditions.

166

### 167 **2.3 Aerobic and anaerobic induction experiments for bacterial and insect luciferase** 168 **reporter systems**

169 pSak-Lux plasmid-containing *E. coli* XL1 was inoculated in 5 ml of L-broth with ampicillin  
170 (100 µg/ml) for overnight (O/N) incubation. O/N cells were inoculated into anaerobic tubes  
171 containing 10 ml LB at a 1% dilution and were grown to an optical density of 0.6. Further  
172 steps were carried out in an anaerobic glove box or in ambient oxygen concentrations.

173 500 µl aliquots of culture were transferred to Eppendorf tubes to which different  
174 concentrations of arabinose were added. One set of Eppendorf tubes was incubated at 37°C  
175 in a glove box under anaerobic conditions, and another set was incubated at 37°C,  
176 aerobically for 1 h. Thereafter, 200 µl of the samples was added to each well of the microtiter  
177 plate.

178 The recombinant *B. longum*/pGBL8b strain was grown and treated essentially the same  
179 as the *E. coli* strain containing the reporter construct. The measurement of the reporter  
180 activity in anaerobic conditions is described below.

181 Light emission was measured using Hidex Chameleon multilabel reader (Turku, Finland)  
182 from 96-well microtiter plates that had been kept either in ambient atmosphere (*E. coli*/Sak-  
183 Lux plasmid), or in an anaerobic chamber (*E. coli*/pSak-Lux plasmid and *B. longum*  
184 NCC2755/pGBL8b). For *B. longum*, 10 µl of 1 mM D-luciferin (Aboatox Oy, Turku, Finland) in  
185 0.1 M Na-citrate buffer, pH 5.0, was pipetted into the plate wells to start the light emission  
186 reaction. The bioluminescence emission from the cultures was measured after aerobic  
187 treatment to quantify the gene expression. A short incubation time (from 5 min on) under  
188 ambient atmosphere was adequate for the bioluminescence detection from anaerobic  
189 cultures, and this bioluminescence closely paralleled that of aerobically grown *E. coli*/pSak-  
190 Lux cells. *B. longum*/pGBL8b cells from anaerobically grown cultures were measured  
191 essentially in the same way as with *E. coli*. The measurements were performed directly from  
192 unbroken, living cells.

193

#### 194 **2.4 Dilution tests**

195 pSak-Lux plasmid-containing *E. coli* XL1 cells were inoculated into anaerobic tubes and  
196 incubated O/N at 150 rpm at 37°C. The next day, 1% of the O/N culture was inoculated to  
197 another anaerobic tube and incubated at 37°C/150rpm until it reached an O.D.<sub>600nm</sub> of 0.6.  
198 Serial dilutions were prepared from the 0.6 O.D. culture in different anaerobic tubes in the  
199 glove box (anaerobic chamber). The dilutions were as follows: 1; 0.1; 0.01; 0.001; 0.0001;  
200 0.00001). A stock solution of 1% arabinose was added to each Eppendorf tube (containing  
201 500 µl of the sample, 3 tubes each per sample) and the tubes were incubated at 37°C in the  
202 glove box for 1 h. Next, the tubes were removed and a 200 µl sample from each Eppendorf  
203 tube was added to microtiter plate wells (3 replicates for each sample) and 10 µg/ml of  
204 tetracycline was added. The plates were read for 20 minutes at 5-min intervals.



205 The *B. longum*/pGBL8b cells were diluted and measured essentially the same as the *E. coli*  
206 reporter cells.


207

### 208 3. RESULTS

209 *E. coli* XL-1 cells containing the plasmid pSak-Lux (map in Fig. S1), a derivative of  
210 plasmid pCGLS-11 (Frackman et al. 1990), which encodes a bacterial luciferase operon,  
211 was used as a model facultative anaerobic Gram (-) bacterial strain to study the reporter  
212 system in anaerobic conditions. As a Gram (+) model bacterium, we used *Bifidobacterium*  
213 *longum* NCC2705 that was transformed with the pGBL8b shuttle vector (Guglielmetti et al.  
214 2008) containing an insect luciferase gene from a click beetle (*P. plagiophthalmus*) under  
215 the control of a strong phage T5 promoter.

216 The experimental data collected from the measurements of the insect luciferase reporter  
217 system confirmed that O<sub>2</sub> is strictly necessary for the luciferase/luciferin reaction as shown  
218 in logarithmic phase *B. longum*/pGBL8b cells. Nevertheless, oxygen was shown to rapidly  
219 diffuse into the culture medium and into the reporter cells after removing the plate from the  
220 anaerobic hood, even when all of the reagents were prepared and mixed under stringent  
221 anaerobic conditions. Furthermore, the addition of 0.2 volumes of D-luciferin solution  
222 prepared in an oxygenic environment to the bacterial broth culture supplied sufficient oxygen  
223 to detect significant light production, which showed saturation after approximately 40 to 50  
224 minutes of incubation at room temperature (Fig. 1). Hereafter, the light emission  
225 measurements were performed 40 minutes after substrate addition.

226 Fig. 1




227

228 The light emission as a function of cell density was studied by diluting the cells and  
 229 measuring the bioluminescence of anaerobically grown *B. longum*/pGBL8b. Fig. 2 shows  
 230 that even as few as approximately  $2 \times 10^3$  log-phase-growing cells/ml can be detected via  
 231 their light emitting activity. This is due to the use of a very strong phage T5 promoter  
 232 (Reynolds et al. 1986) in front of the insect luciferase reporter gene. These results clearly  
 233 indicate the convenience of using these powerful, but never-used, reporters in anaerobic  
 234 organisms. The data show that the light emission is directly proportional to the cell count,  
 235 also highlighting the wide dynamic measurement range in cell densities over several orders  
 236 of magnitude. This sensitivity allows for the use of different types of light-emission  
 237 measurement devices with various light gathering efficiencies, including luminometers,  
 238 multilabel and scintillation counters, and even Polaroid film. The measurements were  
 239 performed directly from unbroken, living cells, making the approach even more robust than  
 240 the conventional  $\beta$ -galactosidase-based reporter systems, which require cell disruption  
 241 steps. The measurement of light emission from intact cells is possible due to the lowering of

242 the pH to acidic values, as shown by Wood and DeLuca (Wood and DeLuca 1987), under  
243 which the substrate for insect luciferase, D-luciferin, is in its protonated form and can easily  
244 cross through the *E. coli* membranes. We have previously shown that this technique is also  
245 useful for other bacterial strains, such as *Streptococcus mutans* (Loimaranta et al. 1998)  
246 and *Staphylococcus aureus* (Tenhami et al. 2001).

247 Fig. 2




248  
249 Similar experiments were performed with the laboratory bacterium *E. coli* grown in  
250 anaerobic and aerobic conditions. The *E. coli* XL-1 strain with the bacterial luciferase operon  
251 in the plasmid pSak-Lux (*luxCDABE* from *P. luminescens*) allows the monitoring of gene  
252 expression without any addition of exogenous substrates, since the products of the operon  
253 produce the long chain aldehyde needed for light emission. The *lux* operon, containing the  
254 *luxCDABE*-genes, was inserted under the control of the arabinose promoter and the light-  
255 emission was monitored after arabinose induction in both aerobic and anaerobic conditions.

256 As expected, the induction of protein expression is not affected by the absence of oxygen.  
257 Reporter proteins are expressed according to the concentration of the sugar used (in top  
258 right corner of Fig. 3). Five minutes of exposure to ambient oxygen, after the withdrawal of  
259 the plate from the anaerobic hood, is enough to get a saturated level of light emission for  
260 gene expression analysis in a microtiter plate-based measuring system. A dilution of several  
261 orders of magnitude was done to the cells induced with 1% arabinose and measured for  
262 RLU as a function of conventional plate counting. The light emitted is directly proportional to  
263 the amount of reporter cells (Fig. 4). Compared to the bifidobacterial strain used in this study,  
264 *E. coli* is much faster, presumably due to the different luciferase reporter. In *B. longum*, the  
265 insect luciferase substrate D-luciferin must first pass the cytoplasmic membrane to reach  
266 the active centre of the reporter protein to achieve light emission. This delay has also been  
267 noticed earlier in aerobically grown whole-cell bacterial reporter systems from several  
268 different types of microbial strains, from bacteria to yeasts (Lampinen et al. 1995; Leskinen  
269 et al. 2003).

270


Fig. 3



271

272

Fig. 4



273

274

275

In addition, the induction of gene expression is not affected by whether the *E. coli* cells with the arabinose-inducible system are grown aerobically or anaerobically (Fig. 5 and Fig.

276 6). The induction closely parallels the concentration of the inducer, arabinose, regardless of  
277 the incubation atmosphere. Here, the bacterial luciferase again shows a faster response to  
278 a change to anaerobic conditions compared to the insect luciferase.

279

280

281

282

283


284

285

286

287


Fig. 5



288

289

Fig. 6



290

## 291 4. DISCUSSION

292 All organisms capable of emitting light as a by-product of their metabolism,  
 293 bioluminescence, need oxygen for the reaction catalysed by the luciferase enzyme.  
 294 Bioluminescent organisms exist on all of the continents of the globe, and include planktonic  
 295 bacteria, algae, shrimps, fish and jelly fish in the sea, as well as terrestrial organisms, with  
 296 the *Photinus pyralis* beetle being the most well-known example. One bacterial species, *P.*  
 297 *luminescens*, is the only example of a terrestrial bacteria capable of bioluminescence.  
 298 Sometimes, fluorescence goes hand-in-hand with bioluminescence in nature. This  
 299 relationship was originally demonstrated by Osamu Shimomura in his studies with a  
 300 luminous hydromedusan, *Aequorea victoria*, from which he was able to extract and purify  
 301 two activities, a fluorescent protein that was excited by a bioluminescent enzyme  
 302 (Shimomura et al. 1962). More and more organisms capable of producing fluorescence upon



303 excitation have been found, and their fluorescence emission and mode of action have been  
304 characterized. Furthermore, the proteins responsible for fluorescence have been mutated  
305 to change, for instance, the emission wavelength, making it possible to normalize  
306 experiments by measuring two or more different spectra at the same time. Fluorescent  
307 proteins require oxygen for proper folding of the chromophore inside the can-like structure  
308 of the protein to produce fluorescence light; on the contrary, bioluminescent proteins  
309 (luciferases) are enzymes (Tsien 1998). The mCherry fluorescent protein was used as a  
310 bioreporter in a recent study, and it was shown to also work in anaerobic conditions (Muhr  
311 et al. 2015). Sixty minutes of incubation in ambient oxygen (air) was sufficient to ensure  
312 maturation for microscopic analysis. However, in this report no optimizations or quantitative  
313 analysis were shown. Newer fluorescent proteins that don't require molecular oxygen have  
314 recently been found for studies in anaerobic or microaerobic conditions (Landete et al.  
315 2015). It was found that the plasmid evoglow-Pp1 (Evocatal GmbH, Germany) allowed  
316 fluorescence generation in anaerobic conditions. However, fluorescence was affected by  
317 external pH in lactobacilli. Furthermore, a pitfall in the use of fluorescent proteins is that all  
318 cells, from prokaryotic to eukaryotic organisms, emit intrinsic fluorescence. This property  
319 causes a background fluorescence level that should be taken into account by subtracting  
320 the fluorescence of cells carrying the empty plasmid. This issue lowers the detection limit of  
321 the system to nanomolar levels and causes extra work (Hakkila et al. 2002). Recently, a  
322 flavin mononucleotide (FMN)-based fluorescent protein (FbFP) was described as reporter  
323 for anaerobic promoter screening in *Clostridium cellulolyticum* (Teng et al. 2015). The  
324 authors showed that is possible to measure promoter activity in situ from anaerobically  
325 growing clostridial strains. However, they did not mention the sensitivity of the system, and,  
326 according to the graphs, the dynamic measurement range was only a few hundred

327 fluorescence units, while luciferase-based reporters have a range of hundreds of thousands  
328 of relative light emission units (see Fig. 2 and Fig. 4).

329 Cronin et al. (Cronin et al. 2008) constructed a bacterial luciferase cassette to study  
330 *Bifidobacterium breve* persistence in living mice. They took into account the possible  
331 absence of oxygen needed for the light emission reaction in the mouse gut and found that  
332 sufficient aerobic conditions existed for the luciferase reaction to occur. They did not further  
333 study the role of oxygen in this living animal setup.

334 Bacterial luciferase reporter genes have also been used in several other applications. An  
335 *E. coli mer-luc* bioreporter was used in anaerobic conditions to monitor the intracellular  
336 accumulation of trace concentrations of Hg(II) ions (Golding et al. 2008). All of the  
337 incubations of the cells were performed in anaerobic conditions, and the actual reporter gene  
338 activity analysis was done by opening the cap of the incubation tube for three minutes to  
339 aerate the sample, followed by quick measurement in a liquid scintillation counter. They  
340 concluded that this approach would be unlikely to have an effect on the bioreporter  
341 response, because it was assumed to take 20 min for the bacteria to sufficiently express the  
342 bioluminescence reporter proteins to detectable levels. However, they did not further study  
343 or optimize the conditions for the bioluminescence reaction after anaerobic growth  
344 conditions.

345 Bioluminescent *Mycobacterium tuberculosis* was made by transferring the *luxAB*  
346 reporter genes into strain ATCC27294 (Cho et al. 2007). Light emission was obtained from  
347 cells cultured in low oxygen conditions by adding the substrate n-decanal, and the emission  
348 was correlated to the antibiotic dosage used to cure the pathogenic bacterium. As in the  
349 previous study, the oxygen-dependent reporter was successfully utilized in anaerobic

350 conditions; however, the conditions were not thoroughly optimized for the reporter's  
351 performance. The *luxAB* luciferase reporter genes from the bacterium *Aliivibrio fischeri* were  
352 used to monitor bioluminescence expression in the anaerobe *Clostridium perfringens*  
353 (Phillips-Jones 1993). The author further expanded the study to monitor rapid changes in  $\alpha$ -  
354 toxin gene expression in the same host to correlate it to reporter gene expression (Phillips-  
355 Jones 2000). She found that the reporter gene expression closely paralleled the  $\alpha$ -toxin  
356 expression and that measurement of *lux*-gene expression provided a rapid and real-time  
357 non-invasive measurement of gene expression. However, she had to add the bacterial  
358 luciferase substrate n-decanal similar to in the study of Cho et al. described above.

359 There is no robust and universal reporter gene system for studies of gene expression  
360 and regulation in anaerobic organisms. Nevertheless, anaerobic organisms represent the  
361 vast majority of all living creatures on Earth. Furthermore, these organisms allow man to  
362 manufacture high value chemicals, food and drink of various sorts by fermentation and to  
363 help purify water and municipal wastes. Here, we have shown that well-known light-emitting  
364 proteins, luciferases and their encoding genes, can be efficiently utilized as genetic reporters  
365 in Gram (+) and Gram (-) bacteria via short exposure to ambient oxygen and without extra  
366 cell disruption steps. Combined with sensitive measurement technologies with sophisticated  
367 electronics, this approach forms a bioreporter system that can be miniaturized to be suitable  
368 for field or in situ usage. The dynamic measurement range is several orders of magnitude  
369 due to the lack of background noise that disturbs other reporter systems utilized thus far.

370

## 371 **5. CONCLUSIONS**

372 Biosensors often utilize whole-cell, engineered cell systems in combination with sensitive  
373 devices to measure light emission. From the examples shown above, quite a few studies  
374 have used reporters that need oxygen for their activity measurements to successfully  
375 quantify gene expression in anaerobically grown bacterial cells. Common to all of these  
376 studies is that the phenomenon itself was not studied and characterized in detail. In our  
377 study, we compared two luciferase reporter systems in two different bacterial strains. Taken  
378 together, the examples using two aerobic reporter systems in two anaerobically grown  
379 microorganisms showed that the real-time monitoring of gene expression in anaerobic  
380 microorganisms is robust and simple, even though it was previously considered impossible.  
381 The reported methodology is universal for bacteria and probably also applies to yeasts and  
382 fungi. We have shown that the above approach can be used for the measurement of gene  
383 expression both in Gram (+) and Gram (-) bacterial strains using insect and bacterial  
384 luciferase reporters. Future gene expression studies of anaerobic microorganisms will prove  
385 the strength of our approach.

386

### 387 **Acknowledgements**

388 This research was funded by the Academy of Finland (HYDROGENE project no.  
389 107425) and The Nordic Energy Research (BIOHYDROGEN project no. 28-02).

**REFERENCES**

- 391 Cecic, I., Chan, D.A., Sutphin, P.D., Ray, P., Gambhir, S.S., Giaccia, A.J., Graves, E.E., 2007. *Mol*  
392 *Imaging* 6(4), 219-228.
- 393 Cho, S.H., Warit, S., Wan, B., Hwang, C.H., Pauli, G.F., Franzblau, S.G., 2007. *Antimicrob Agents*  
394 *Chemother* 51(4), 1380-1385.
- 395 Cronin, M., Sleator, R.D., Hill, C., Fitzgerald, G.F., van Sinderen, D., 2008. *BMC Microbiol* 8, 161.
- 396 Daunert, S., Barrett, G., Feliciano, J.S., Shetty, R.S., Shrestha, S., Smith-Spencer, W., 2000. *Chem*  
397 *Rev* 100(7), 2705-2738.
- 398 Drepper, T., Eggert, T., Circolone, F., Heck, A., Krauss, U., Guterl, J.K., Wendorff, M., Losi, A.,  
399 Gartner, W., Jaeger, K.E., 2007. *Nat Biotechnol* 25(4), 443-445.
- 400 Edwards, A.N., Pascual, R.A., Childress, K.O., Nawrocki, K.L., Woods, E.C., McBride, S.M., 2015.  
401 *Anaerobe* 32, 98-104.
- 402 Elad, T., Lee, J.H., Belkin, S., Gu, M.B., 2008. *Microb Biotechnol* 1(2), 137-148.
- 403 Feustel, L., Nakotte, S., Durre, P., 2004. *Appl Environ Microbiol* 70(2), 798-803.
- 404 Frackman, S., Anhalt, M., Nealson, K.H., 1990. *J Bacteriol* 172(10), 5767-5773.
- 405 Golding, G.R., Sparling, R., Kelly, C.A., 2008. *Appl Environ Microbiol* 74(3), 667-675.
- 406 Guglielmetti, S., Ciranna, A., Mora, D., Parini, C., Karp, M., 2008. *Int J Food Microbiol* 124(3), 285-  
407 290.
- 408 Guglielmetti, S., Karp, M., Mora, D., Tamagnini, I., Parini, C., 2007. *Appl Microbiol Biotechnol*  
409 74(5), 1053-1061.
- 410 Hakkila, K., Maksimow, M., Karp, M., Virta, M., 2002. *Anal Biochem* 301(2), 235-242.
- 411 Hansen, M.C., Palmer, R.J., Jr., Udsen, C., White, D.C., Molin, S., 2001. *Microbiology* 147(Pt 5),  
412 1383-1391.
- 413 Lampinen, J., Virta, M., Karp, M., 1995. *Appl Environ Microbiol* 61(8), 2981-2989.
- 414 Landete, J.M., Langa, S., Revilla, C., Margolles, A., Medina, M., Arques, J.L., 2015. *Appl Microbiol*  
415 *Biotechnol* 99(16), 6865-6877.
- 416 Leskinen, P., Virta, M., Karp, M., 2003. *Yeast* 20(13), 1109-1113.
- 417 Loimaranta, V., Tenovuo, J., Koivisto, L., Karp, M., 1998. *Antimicrob Agents Chemother* 42(8),  
418 1906-1910.
- 419 Michelini, E., Cevenini, L., Mezzanotte, L., Ablamsky, D., Southworth, T., Branchini, B., Roda, A.,  
420 2008. *Anal Chem* 80(1), 260-267.
- 421 Muhr, E., Leicht, O., Gonzalez Sierra, S., Thanbichler, M., Heider, J., 2015. *Front Microbiol* 6, 1561.
- 422 Phillips-Jones, M.K., 1993. *FEMS Microbiol Lett* 106(3), 265-270.
- 423 Phillips-Jones, M.K., 2000. *FEMS Microbiol Lett* 188(1), 29-33.
- 424 Reynolds, A.E., Mahadevan, S., LeGrice, S.F., Wright, A., 1986. *J Mol Biol* 191(1), 85-95.
- 425 Rizzotti, M., 2009. In: UNESCO (Ed.).
- 426 Selifonova, O., Burlage, R., Barkay, T., 1993. *Appl Environ Microbiol* 59(9), 3083-3090.
- 427 Shimomura, O., Johnson, F.H., Saiga, Y., 1962. *J Cell Comp Physiol* 59, 223-239.
- 428 Teng, L., Wang, K., Xu, J., Xu, C., 2015. *J Microbiol Methods* 119, 37-43.
- 429 Tenhami, M., Hakkila, K., Karp, M., 2001. *Antimicrob Agents Chemother* 45(12), 3456-3461.
- 430 Tsien, R.Y., 1998. *Annu Rev Biochem* 67, 509-544.
- 431 Whitman, W.B., Coleman, D.C., Wiebe, W.J., 1998. *Proc Natl Acad Sci U S A* 95(12), 6578-6583.
- 432 Wood, K.V., DeLuca, M., 1987. *Anal Biochem* 161(2), 501-507.
- 433

## 434 **Figure Legends**

435 **Fig. 1.** Time curve for anaerobically grown *Bifidobacterium longum*/pGBL8b  
436 after removal from the anaerobic hood. The substrate for the luciferase reporter  
437 protein, D-luciferin, was added aerobically immediately after the anaerobic  
438 incubation of the cells, and the light-emission was continuously monitored in the  
439 plate reader. The values are the means of two independent experiments conducted  
440 in triplicates. The variation between the replicates was less than 10%.

441 **Fig. 2.** Emission of light as a function of cell count in *B. longum*/pGBL8b. Vertical  
442 bars at each point represent the standard deviation of the data obtained from three  
443 independent measurements conducted in quadruplicate.

444 **Fig. 3.** Light emission by *E. coli* XL-1/pSak-lux from anaerobically grown cells  
445 induced with different concentrations of arabinose (% in the right corner box) as a  
446 function of time after aerobic incubation. The values are the means of two  
447 independent experiments conducted in triplicates. The variation between the  
448 replicates was less than 10%.

449 **Fig. 4.** Emission of light as a function of cell count in anaerobically grown *E. coli*  
450 XL-1/pSAK-Lux. The values are the means of two independent experiments  
451 conducted in triplicates. The variation between the replicates was less than 10%.

452 **Fig. 5.** Emission of light as a function of the concentration of the inducer,  
453 arabinose, after transferring the anaerobically grown cells to the ambient oxygen  
454 concentration. The values are the means of two independent experiments  
455 conducted in triplicates. The variation between the replicates was less than 10%.

456 **Fig. 6.** Emission of light as a function of the concentration of the inducer,  
457 arabinose, after growing the cells aerobically. The different time points represent the  
458 relatively light units (RLU) after placing the microtiter plate into the plate counter.  
459 The values are the means of two independent experiments conducted in triplicates.  
460 The variation between the replicates was less than 10%.