

A ROUTE TO BENZODITHIOPHENE SYSTEMS BY EXPLOITING A LIGAND-FREE SUZUKI-MIYaura COUPLING REACTION IN DEEP EUTECTIC SOLVENTS

Valentina Pelliccioli,^{a*} Silvia Cauteruccio,^a Giuseppe Dilauro,^b Filippo Perna,^b Vito Capriati,^b Emanuela Licandro^a

^aDipartimento di Chimica, Università degli Studi di Milano, via Golgi 19 I-20133, Milan, Italy;

^bDipartimento di Farmacia–Scienze del Farmaco, Università degli Studi di Bari "Aldo Moro", Consorzio C.I.N.M.P.I.S., I-70125, Bari, Italy

* e-mail: valentina.pelliccioli@unimi.it

Benzo[1,2-*b*:4,3-*b'*]dithiophene (**BDT**) and its derivatives belong to an interesting class of thiophene-based aromatic π -conjugated compounds that are widely studied as functional organic materials inserted, for instance, as units in mono and polydisperse oligomers [1], or as π -spacers in push-pull organic chromophores for photovoltaic applications [2]. Moreover, **BDTs** are key intermediates for the synthesis of inherent chiral tetrathia[7]helicenes, which are an attractive class of heterohelicenes with unique physicochemical and chiroptical properties due to their helix-like structure [3]. Thus, **BDT** is a key starting molecule which can allow access to more complex and interesting systems through a selective and judicious functionalization of the α and β -positions of the terminal thiophene rings. Building on our recent studies on the synthesis and functionalization of **BDTs** [4], we questioned whether a novel class of 2,7-diarylsubstituted **BDTs** **1** (Figure 1) could be synthesized *via* a palladium-catalysed Suzuki-Miyaura reaction between heteroaryl halides **2** and organoboron derivatives **3** in *Deep Eutectic Solvents* (**DESS**), which have proven to be effective as sustainable and environmentally responsible reaction media in several transition-metal-catalyzed reactions [5].




Figure 1. General scheme for the synthesis of 2,7-diarylsubstituted BDTs **1**

In this communication, we report our preliminary results on the preparation of diarylsubstituted **BDTs** **1**, and discuss the substrate scope of the proposed protocol. Some of the compounds so far obtained display interesting photophysical properties, which are currently under investigation.

[1] Y. Nishide, H. Osuga, M. Saito, T. Aiba, Y. Inagaki, Y. Doge, K. Tanaka, *J. Org. Chem.* **2007**, *72*, 9141–9151.

[2] E. Longhi, A. Bossi, G. Di Carlo, S. Maiorana, F. De Angelis, P. Salvatori, A. Petrozza, M. Binda, V. Roiati, P. R. Mussini, C. Baldoli, E. Licandro, *Eur. J. Org. Chem.* **2013**, 84–94.

[3] E. Licandro, S. Cauteruccio, D. Dova, *Adv. Het. Chem.* **2016**, *118*, 1–46.

[4] S. Cauteruccio, D. Dova, C. Graiff, C. Carrara, J. Doulctet, G. R. Stephenson, E. Licandro, *New J. Chem.* **2014**, *38*, 2241–2244.

[5] (a) F. Messa, S. Perrone, M. Capua, F. Tolomeo, L. Troisi, V. Capriati, A. Salomone, *Chem. Commun.* **2018**, *54*, 8100–8103; (b) G. Dilauro; S. Mata García, D. Tagarelli, P. Vitale, F. M. Perna, V. Capriati, *ChemSusChem*, **2018**, *11*, 3495–3501.