

1 **Impact of *Staphylococcus aureus* infection on the late lactation goat milk proteome:**
2 **new perspectives for monitoring and understanding mastitis in dairy goats**

3 Salvatore Pisanu^a, Carla Cacciotto^a, Daniela Pagnozzi^a, Sergio Uzzau^{a,b,c}, Claudia Pollera^d, Martina Penati^d,
4 Valerio Bronzo^d, Maria Filippa Addis^{c,d*}

5

6 ^aPorto Conte Ricerche, Alghero, Italy

7 ^bDipartimento di Scienze Biomediche, Università degli Studi di Sassari, Sassari, Italy

8 ^cMediterranean Center for Disease Control, Sassari, Italy

9 ^dDipartimento di Medicina Veterinaria, Università degli Studi di Milano, Milan, Italy

10

11 Short title: **Goat milk proteins and *S. aureus***

12

13

14 *Corresponding author at the Department of Veterinary Medicine, University of Milan, Via dell'Università 6,
15 26900 Lodi (LO). Tel. ++39 02 503 34515; E-mail: filippa.addis@unimi.it.

16 **Abstract**

17 The milk somatic cell count (SCC) is a standard parameter for monitoring intramammary infections (IMI) in
18 dairy ruminants. In goats, however, the physiological increase in SCC occurring in late lactation heavily
19 compromises its reliability. To identify and understand milk protein changes specifically related to IMI, we
20 carried out a shotgun proteomics study comparing high SCC late lactation milk from goats with subclinical
21 *Staphylococcus aureus* IMI and from healthy goats to low SCC mid-lactation milk from healthy goats. As a
22 result, we detected 52 and 19 differential proteins (DPs) in *S. aureus*-infected and uninfected late lactation
23 milk, respectively. Unexpectedly, one of the proteins higher in uninfected milk was serum amyloid A. On the
24 other hand, 38 DPs were increased only in *S. aureus*-infected milk and included haptoglobin and numerous
25 cytoskeletal proteins. Based on STRING analysis, the DPs unique to *S. aureus* infected milk were mainly
26 involved in defense response, cytoskeleton organization, cell-to-cell, and cell-to-matrix interactions. Being
27 tightly and specifically related to infectious/inflammatory processes, these proteins may hold promise as
28 more reliable markers of IMI than SCC in late lactation goats.

29

30 *Significance*

31 The biological relevance of our results lies in the increased understanding of the changes specifically related
32 to bacterial infection of the goat udder in late lactation. The DPs present only in *S. aureus* infected milk may
33 find application as markers for improving the specificity of subclinical mastitis monitoring and detection in
34 dairy goats in late lactation, when other widespread tools such as the SCC lose diagnostic value.

35

36 **Keywords:**

37 Goat mastitis; late lactation milk; *Staphylococcus aureus*; somatic cell count; shotgun proteomics;
38 haptoglobin.

39 **1. Introduction**

40 Intramammary infections (IMI) and mastitis cause milk production losses and reduce dairy goat product
41 quality. Subclinical mastitis due to chronic IMI can be especially problematic, and reliable monitoring and
42 detection tools are needed for maintaining good profitability of goat productions [1]. The somatic cell count
43 (SCC), that is, the number of cells per mL of milk, is largely considered a reliable IMI indicator in dairy
44 ruminants [2]. In goats, however, the SCC is subjected to physiological variations related to age, parity, stage
45 of lactation, estrus, and other factors [3–5], undermining specificity and limiting the diagnostic value of this
46 practical and cost-effective marker. Late lactation, in particular, is associated with SCC increase in cow, sheep
47 and goat milk [6], but the magnitude of this increase in goats is so high that SCC may not enable to distinguish
48 infected from uninfected udders in late lactation [3,7,8]. Consequently, the reliability of the most widespread
49 field tool, the California Mastitis Test (CMT), is severely affected [9]. The availability of a protein marker
50 appearing in the milk only upon infection would increase the specificity of subclinical mastitis detection and
51 support the screening of late lactation goats for IMI, enabling more meaningful management decisions
52 especially at the dry-off [10,11].

53 The widespread adoption of milk SCC as an indicator of IMI is based on the notion that the number of cells
54 in milk increases due to the active influx of neutrophils recalled into the milk as a result of the inflammation
55 elicited by a microbial insult. Being this accompanied by increased permeability of the blood-milk barrier,
56 with consequent leakage of serum contents into the milk, other ways to detect subclinical mastitis are based
57 on these “leaked” proteins and other molecules found in the milk as a result of active secretion, cellular lysis
58 or tissue rearrangements [11]. Investigating the proteome changes specifically associated with subclinical IMI
59 is a suitable way to identify marker proteins that may represent a reliable alternative when the SCC loses
60 specificity.

61 Gram-positive bacteria, and staphylococci in particular, are the most prevalent intramammary pathogens in
62 dairy goats [12–15]. Gram-positive bacteria cause mainly subclinical, chronic infections that persist along the
63 dry period [6,8] justifying the need for more sensitive and specific screening tools for monitoring mammary

64 gland health in dairy goats. Therefore, we selected *Staphylococcus aureus* subclinical IMI as the model
65 condition for this study.

66 In summary, we applied a shotgun proteomics pipeline to compare late lactation, high SCC, *S. aureus* infected
67 and uninfected milk with mid-lactation, low SCC uninfected milk to understand the changes induced by
68 infection and to identify differential proteins with potential as subclinical mastitis markers in late lactation.

69

70 **2. Materials and methods**

71

72 *2.1. Animals and milk samples*

73 Half-udder goat milk was retrieved from a frozen sample bank collected along the course of two entire
74 lactations in a herd of Alpine goats farmed in Lombardy, Italy. All goats were clinically healthy for the two
75 lactation years and showed no signs of mastitis. The detailed description of the herd and of experimental
76 procedures was reported in a previous work [3]. Briefly, bacteriological analysis was carried out bi-monthly
77 according to the National Mastitis Council standards [16] as described previously [17]. Ten μl of milk was
78 spread on blood agar plates and incubated aerobically at 37°C. After 24 h, plates were examined, and colonies
79 were provisionally identified based on Gram stain, morphology, and haemolysis patterns. Gram-positive cocci
80 were tested for catalase and coagulase production for identification as *Staphylococcus aureus*, and colonies
81 were re-isolated on Baird-Parker medium for further confirmation. Somatic cell count (SCC) was measured
82 with an automated somatic cell counter (Bentley Somacount 150, Bentley Instrument, USA) [3]. Nine samples
83 from multiparous goats were selected for the current study as follows: i) three mid-lactation samples (40 \pm 10
84 Days in milk - DIM) with very low SCC (19,000 \pm 7000) from half-udders producing a sterile milk bacterial
85 culture for two consecutive samplings (MLU, Mid-lactation, Low SCC, Uninfected); ii) three late lactation
86 samples (> 250 DIM) with SCC > 2,000,000 cells/mL (2,932,000 \pm 439,000) from half-udders producing a sterile
87 milk bacterial culture for the whole lactation (LHU, Late lactation, High SCC, Uninfected); and iii) three late
88 lactation samples (> 250 DIM) with SCC > 2,000,000 cells/mL (3,980,000 \pm 74,000) from goat half-udders with
89 a milk bacterial culture repeatedly positive for *S. aureus* in the previous lactation year (LHS, Late lactation,

90 High SCC, *Staphylococcus aureus* infected). *S. aureus* positive goats were culled at the end of lactation in the
91 first year. The whole herd tested negative to *S. aureus* in the second year, when MLU and LHU samples were
92 collected. The SCC > 2,000,000 cells/mL threshold was selected because the California Mastitis Test (CMT)
93 scores are the highest over this value.

94

95 2.2. Milk sample preparation for proteomic analysis

96 Milk sample preparation for proteomic analysis was carried out as described previously [17]. Briefly, milk was
97 allowed to thaw at room temperature and centrifuged at 800 x g at 4°C for 10 min, the fat ring was removed,
98 and skim milk was diluted 1:1 with lysis buffer, incubated at 95°C for 10 min and sonicated in a refrigerated
99 water bath for 10 min, after which the suspension was centrifuged at 10.000 x g for 10 min at 4°C. Then, 7 µl
100 of extract was subjected to filter-aided sample preparation (FASP) [18]. Protein samples were reduced,
101 alkylated, and digested with trypsin on 3 kDa cut-off Amicon Ultra-0.5 mL centrifugal filter units (Millipore,
102 Billerica, MA, USA). Peptide concentration was determined with a NanoDrop 2000 spectrophotometer
103 (Thermo Scientific, San Jose, CA, USA).

104

105 2.3. Tandem mass spectrometry analysis of peptides

106 All peptide mixtures were analysed on a Q-Exactive interfaced with an UltiMate 3000 RSLCnanoLC system
107 (Thermo Scientific, San Jose, CA, USA), as detailed previously [19], using 4 µg of peptide mixture. Protein
108 identification was carried out with Proteome Discoverer (version 1.4; Thermo Scientific) and Sequest-HT as
109 the search engine. MS/MS spectra were analyzed as follows. Database: custom, obtained by merging *Bos*
110 *taurus*, *Capra hircus* and *Staphylococcus* databases. These were downloaded from Swiss-Prot (*Bos taurus*)
111 and TrEMBL (*Capra hircus* and *Staphylococcus*) release2017_05 and 2016_11, respectively; enzyme: trypsin,
112 with two missed cleavages allowed; precursor mass tolerance: 10 ppm; MS/MS tolerance: 0.02 Da; charge
113 states: +2, +3, and +4; cysteine carbamidomethylation as static modification and methionine oxidation as
114 dynamic modifications. The percolator algorithm was used for protein significance and for peptide validation

115 (false discovery rate, FDR, < 0.01). Peptide and protein grouping according to the Proteome Discoverer's
116 algorithm were allowed, applying the strict maximum parsimony principle.

117

118 *2.4. Proteomic data analysis*

119 Protein abundance changes were assessed by the spectral counting (SpC) approach as described previously
120 (Pisanu et al., 2019). When proteins had more than one entry, only those with the highest number of unique
121 peptides and SpCs were considered. Only proteins identified in at least two biological replicates and having
122 ≥ 2 SpCs (Peptide Spectrum Matches, PSMs) in at least one sample of the group were considered for
123 differential analysis. Relative abundance of single proteins in all samples and abundance changes of proteins
124 between groups were calculated by considering the normalised spectral abundance factor (NSAF) and the R_{SC}
125 (the log₂ of the protein abundance ratio), respectively [20,21]. Statistical significance was assessed by the
126 beta-binomial test with FDR correction according to Benjamini-Hochberg [22]. Only proteins with $R_{SC} \geq 1.0$ or
127 ≤ -1.0 between the compared groups and having a p-value ≤ 0.05 were considered differential. The biological
128 processes and molecular functions reported by UniProtKB database were used for gene ontology (GO)
129 analysis of differential proteins (DPs), integrated with manual curation. Protein-protein interaction network
130 was assessed with the STRING database (Version 11, <http://string-db.org/>), after replacing all *Capra hircus*
131 UniProt IDs with the corresponding *Bos taurus* UniProt IDs using the Basic Local Alignment Search Tool
132 (BLAST) [23] and by taking into account only functional interactions with high confidence (combined score >
133 0.7) [24].

134

135 *2.5. Data Availability*

136 The data have been deposited to the ProteomeXchange with identifier PXD017243 [25]. A complete
137 description of the dataset is available in Pisanu et al., 2020 (Data in Brief, submitted).

138

139

140

141 **3. Results**

142

143 *3.1. Shotgun proteomics and differential analysis*

144 *S. aureus* infected and uninfected milk was subjected to a shotgun proteomics workflow combining FASP, RP-
145 HPLC, and high-resolution orbitrap MS. This led to the identification of 540 total unique proteins, of which
146 256 eligible for differential analysis. The complete description of the proteomic datasets is available in Pisanu
147 et al. 2020 (Data in Brief, submitted).

148 To identify the changes specifically induced by *S. aureus* as opposed to the physiological changes occurring
149 in late lactation, we compared late lactation, high SCC infected and uninfected milk with mid-lactation, low
150 SCC uninfected milk. As a result, late lactation infected milk showed 52 significant DPs, while late lactation
151 uninfected milk showed only 19 DPs. Results are summarised in Table 1 and are detailed in Supplementary
152 file (sheets 1 and 2, respectively). The higher number of DPs in *S. aureus* positive milk indicated that the
153 presence of *S. aureus* was more impacting on the milk proteome than the physiological late lactation changes
154 alone.

155 Table 2 lists all the DPs obtained in the two comparisons with the respective \log_2 ratio abundance values
156 (R_{SC}). Protein abundance changes were generally more intense in *S. aureus* infected milk, as most common
157 DPs showed higher R_{SC} values in this sample group. The top DPs in both *S. aureus* infected and uninfected
158 milk were lactotransferrin and cathelicidin-2. Vimentin, the third top DP, changed significantly only in *S.*
159 *aureus* infected milk. In uninfected milk, only complement C3, olfactomedin-like protein 3, and serum
160 amyloid A showed higher R_{SC} values, and only three DPs were unique: fatty acid synthase, calreticulin, and
161 lactoperoxidase. On the other hand, 38 proteins showed significant changes only in *S. aureus* infected milk
162 and are highlighted in bold in Table 2.

163

164 *3.2. Functional analysis*

165 The 38 DPs unique to *S. aureus* infected milk (Table 2, bold) were analyzed for their interactions and biological
166 functions by STRING. Several proteins were strongly connected, such as tubulins with 14-3-3 proteins and

167 heat-shock proteins, myosin light chains, other cytoskeletal proteins, and proteins involved in cell-to-cell and
168 cell-to-matrix interactions.

169 Supplementary file, sheets 4-7, report the list of significant GO terms and pathways enriched for the
170 categories Biological Process, KEGG Pathways, Cellular Component, INTERPRO Protein Domains and
171 Features, and Reactome Pathways, respectively. The most relevant Biological Process GO terms and KEGG
172 pathways are indicated in Figure 1A and in Figure 1B.

173

174 **4. Discussion**

175 Aim of this work was to detect and understand milk changes specifically related to IMI in dairy goats,
176 especially focusing on late lactation, by differential label-free shotgun proteomics. This was also the first
177 proteomic investigation of milk from goats with subclinical *S. aureus* mastitis.

178 When comparing the DPs observed in late lactation *S. aureus* infected and uninfected milk, the number,
179 identity, and abundance indicated that the presence of *S. aureus* had a specific and strong impact on the goat
180 milk proteome. Although some DPs, especially those with the highest R_{SC} values, were increased in both
181 infected and uninfected milk, most were detected only in infected milk and are the most relevant for
182 understanding the differences between the two conditions and for identifying useful mastitis markers.

183 Lactotransferrin, cathelicidins, serum amyloid A, and haptoglobin are long known to increase in cow milk
184 during mastitis [10,26–32] and have been evaluated as protein markers also in sheep and goats [3,33–35]. In
185 this study, lactotransferrin and cathelicidins increased in late lactation *S. aureus* infected milk but increased
186 also in late lactation uninfected milk, although at a slightly lower extent. These proteins might increase in the
187 milk as a result of neutrophil influx, and as opposed to other dairy species, in goats this occurs also
188 physiologically as lactation progresses [5]. This is in line with the observations recently made by our group
189 when comparing the value of cathelicidins in late lactation sheep and goats [3] and, in spite of the great value
190 in other dairy species, might reduce mastitis detection specificity in late lactation goats.

191 On the other hand, the unexpected inverse behavior of serum amyloid A and haptoglobin was of significant
192 interest. In fact, previous gel-based and gel-free proteomic studies carried out in goat milk are contrasting in

193 this respect. Olumee-Shabon et al. [36] observed a significant increase of both haptoglobin and serum
194 amyloid A in agreement with previous studies in cows, while Wang et al. detected serum amyloid A but not
195 haptoglobin [37]. However, both studies evaluated an experimentally induced lipopolysaccharide (LPS)
196 mastitis, and specificities in the host response to Gram-negative and Gram-positive microorganisms might
197 partly account for these differences [38,39]. This is especially relevant when considering that Gram-positive
198 bacteria are by far the leading intramammary pathogens in dairy goats, with *Staphylococcus* being the most
199 prevalent genus [12–15]. *Staphylococcus* spp. also cause mainly chronic, subclinical infections that persist
200 along the dry period [8] justifying the need for more sensitive and specific screening tools. In our study, serum
201 amyloid A was increased in high SCC milk, but such increase was not specific for the presence of an infection;
202 actually, the R_{SC} value was higher in uninfected (2.41) than in *S. aureus* infected (1.21) milk, raising the
203 question that it might be related to physiological rather than pathological processes. On the other hand,
204 haptoglobin increased significantly only in *S. aureus* infected milk ($R_{SC} = 1.70$). Therefore, haptoglobin might
205 have potential as a specific mastitis marker also in late lactation, high SCC goat milk. A dedicated study with
206 large sample numbers, different etiological agents, and thoroughly validated antibodies will be needed to
207 further investigate this finding.

208 The STRING protein network analysis provided interesting information on the biological processes and
209 pathways involving the 38 DPs detected only in *S. aureus* infected milk. Notably, tight junction, regulation of
210 the actin cytoskeleton and leukocyte transendothelial migration were among the most significant KEGG
211 pathways highlighted by STRING analysis. Tight junctions participate actively in regulating the passage of
212 blood-derived antimicrobial factors, cytokines, and neutrophils [40,41]. Accordingly, loss of tight junction
213 integrity has been linked to reduced milk secretion and increased paracellular mixture of serum and milk
214 components [42,43]. The specific function of haptoglobin in the context of mammary gland inflammation is
215 mainly attributed to hemoglobin scavenging to inhibit its oxidative activity [44], but the full-length precursor
216 of haptoglobin, zonulin, increases epithelial permeability by mediating intercellular tight junction
217 disassembly [45]. Altogether, this suggests that its increase only in late lactation infected milk might be a

218 specific consequence of the blood-milk barrier dynamics related to inflammation [42,43,46–48] and
219 highlights its potential as specific goat mastitis marker.

220 The highest R_{SC} value for proteins increased only in *S. aureus* infected milk was observed for vimentin, a highly
221 abundant intermediate filament protein [49] involved in the innate immune response to pathogens [50] by
222 regulating inflammasome activity [51]. Vimentin was one of the top 15 up-regulated proteins at 57, 81, and
223 312 hours after intramammary challenge of cows with *Streptococcus uberis* [52], and it was the first DP in
224 the milk of buffaloes with *S. aureus* IMI [17]. Several members of the annexin family were also significantly
225 higher in *S. aureus*-infected milk. Annexins are involved in vesicular trafficking and might be increased as a
226 result of cell degranulation, especially by neutrophils [59]. Interestingly, clumping factor A of *S. aureus*
227 interacts with annexin A2 on mammary epithelial cells mediating its entry into the host cell [60]. The increase
228 in apolipoprotein A4 only in infected milk is in line with the observations of Olumee-Shabon *et al.* in the milk
229 of goats challenged with LPS [36].

230 Other DPs increased also in late lactation, uninfected milk. Among other causes, these might be the result of
231 physiological processes involved in mammary gland tissue dynamics and recycling associated with the natural
232 involution of the mammary gland at the cessation of lactation [6,53].

233 In conclusion, this work provided the first characterization of *S. aureus* infected goat milk; identified the
234 differences between infected and uninfected late lactation, high SCC milk; identified several proteins that are
235 increased in milk only upon infection; provided insights on the mechanisms leading to the specific changes
236 found in the milk proteome when an IMI is present; and, most importantly, identified putative markers that
237 might improve specificity of subclinical mastitis detection and enable more meaningful management
238 decisions especially in late lactation, when the diagnostic value of SCC is reduced.

239

240 **Acknowledgements**

241 This work was funded by Sardegna Ricerche and the University of Milan.

242

243 **Author contributions**

244 Proteomic analysis and differential proteomics: SP, CC, DP. Microbiological analysis of milk: CP, MP. Animal
245 examination, sample collection: VB. Funding and contribution to study design: SU, VB. Study conception,
246 design and coordination, data analysis and interpretation, manuscript drafting: MFA. Data interpretation and
247 manuscript revision: All authors.

248

249 **Conflict of interest**

250 The authors declare no conflict of interest.

251

252 **Appendix A. Supplementary data**

253 Supplementary data to this article can be found online at

254 <https://www.sciencedirect.com/science/article/pii/S1874391920301317?via%3Dihub>.

255 **References**


256

- 257 [1] P.I. Menzies, S.Z. Ramanoon, Mastitis of Sheep and Goats, *Vet. Clin. North Am. Food Anim. Pract.* 17
 258 (2001) 333–358. [https://doi.org/10.1016/S0749-0720\(15\)30032-3](https://doi.org/10.1016/S0749-0720(15)30032-3).
- 259 [2] I.R. Dohoo, A.H. Meek, Somatic cell counts in bovine milk., *Can. Vet. Journal. La Rev. Vétérinaire Can.*
 260 23 (1982) 119–25. <http://www.ncbi.nlm.nih.gov/pubmed/17422127> (accessed June 17, 2016).
- 261 [3] V. Tedde, V. Bronzo, G.M.G. Puggioni, C. Pollera, A. Casula, G. Curone, P. Moroni, S. Uzzau, M.F.
 262 Addis, Milk cathelicidin and somatic cell counts in dairy goats along the course of lactation, *J. Dairy*
 263 *Res.* 86 (2019) 217–221.
- 264 [4] A.M. Rota, C. Gonzalo, P.L. Rodriguez, A.I. Rojas, L. Martín, J.J. Tovar, Effects of stage of lactation and
 265 parity on somatic cell counts in milk of Verata goats and algebraic models of their lactation curves,
 266 *Small Rumin. Res.* 12 (1993) 211–219. [https://doi.org/10.1016/0921-4488\(93\)90085-V](https://doi.org/10.1016/0921-4488(93)90085-V).
- 267 [5] R. Jimenez-Granado, M. Sanchez-Rodriguez, C. Arce, V. Rodriguez-Estevez, Factors affecting somatic
 268 cell count in dairy goats: A review, *Spanish J. Agric. Res.* 12 (2014) 133–150.
 269 <https://doi.org/10.5424/sjar/2014121-3803>.
- 270 [6] G. Leitner, U. Merin, O. Krifucks, S. Blum, A.L. Rivas, N. Silanikove, Effects of intra-mammary
 271 bacterial infection with coagulase negative staphylococci and stage of lactation on shedding of
 272 epithelial cells and infiltration of leukocytes into milk: comparison among cows, goats and sheep.,
 273 *Vet. Immunol. Immunopathol.* 147 (2012) 202–10. <https://doi.org/10.1016/j.vetimm.2012.04.019>.
- 274 [7] P. Moroni, G. Pisoni, G. Ruffo, P.J. Boettcher, Risk factors for intramammary infections and
 275 relationship with somatic-cell counts in Italian dairy goats, *Prev. Vet. Med.* 69 (2005) 163–173.
 276 <https://doi.org/10.1016/j.prevetmed.2004.10.013>.
- 277 [8] V. Bernier Gosselin, S. Dufour, M.J. Calcutt, P.R.F. Adkins, J.R. Middleton, Staphylococcal
 278 intramammary infection dynamics and the relationship with milk quality parameters in dairy goats
 279 over the dry period, *J. Dairy Sci.* 102 (2019) 4332–4340. <https://doi.org/10.3168/JDS.2018-16030>.
- 280 [9] F.N. Souza, M.G. Blagitz, C.F.A.M. Penna, A.M.M.P. Della Libera, M.B. Heinemann, M.M.O.P.
 281 Cerqueira, Somatic cell count in small ruminants: Friend or foe?, *Small Rumin. Res.* 107 (2012) 65–
 282 75. <https://doi.org/10.1016/j.smallrumres.2012.04.005>.
- 283 [10] C. Viguier, S. Arora, N. Gilmartin, K. Welbeck, R. O’Kennedy, Mastitis detection: current trends and
 284 future perspectives, *Trends Biotechnol.* 27 (2009) 486–493.
 285 <https://doi.org/10.1016/j.tibtech.2009.05.004>.
- 286 [11] T. Stuhr, K. Aulrich, Intramammary infections in dairy goats: Recent knowledge and indicators for
 287 detection of subclinical mastitis, *Landbauforsch. Volkenrode.* 60 (2010) 267–280.
 288 <https://www.scopus.com/inward/record.uri?eid=2-s2.0-78651506795&partnerID=40&md5=c66a7d7a4f179507f1df1ad04e819117>.
- 289 [12] S. Dore, M. Liciardi, S. Amatiste, S. Bergagna, G. Bolzoni, V. Caligiuri, A. Cerrone, G. Farina, C.O.
 290 Montagna, M.A. Saletti, M.L. Scatassa, G. Sotgiu, E.A. Cannas, Survey on small ruminant bacterial
 291 mastitis in Italy, 2013–2014, *Small Rumin. Res.* 141 (2016) 91–93.
 292 <https://doi.org/10.1016/j.smallrumres.2016.07.010>.
- 293 [13] A. Contreras, D. Sierra, A. Sanchez, J.C.C. Corrales, J.C.C. Marco, M.J.J. Paape, C. Gonzalo, A. Sánchez,
 294 J.C.C. Corrales, J.C.C. Marco, M.J.J. Paape, C. Gonzalo, Mastitis in small ruminants., *Small Rumin. Res.*
 295 68 (2007) 145–153. <https://doi.org/10.1016/j.smallrumres.2006.09.011>.
- 296 [14] M.G. Dominguez-Bello, E.K. Costello, M. Contreras, M. Magris, G. Hidalgo, N. Fierer, R. Knight,
 297 Delivery mode shapes the acquisition and structure of the initial microbiota across multiple body
 298 habitats in newborns., *Proc. Natl. Acad. Sci. U. S. A.* 107 (2010) 11971–5.
 299 <https://doi.org/10.1073/pnas.1002601107>.
- 300 [15] S. McDougall, D. Malcolm, C. Prosser, Prevalence and incidence of intramammary infections in
 301 lactating dairy goats, *N. Z. Vet. J.* 62 (2014) 136–145.
 302 <https://doi.org/10.1080/00480169.2013.865294>.
- 303 [16] J.R. Middleton, L.K. Fox, G. Pighetti, *Laboratory Handbook on Bovine Mastitis*, National Mastitis
 304 Council, Madison, WI, New Prague, MN, 2017. <http://nmconline.org/docs/lhb.htm> (accessed May 4,
 305 2015).
 306

- 307 [17] S. Pisanu, C. Cacciotto, D. Pagnozzi, S. Uzzau, P. Ciaramella, J. Guccione, M. Penati, C. Pollera, P.
308 Moroni, V. Bronzo, M.F. Addis, Proteomic changes in the milk of water buffaloes (*Bubalus bubalis*)
309 with subclinical mastitis due to intramammary infection by *Staphylococcus aureus* and by non-
310 aureus staphylococci, *Sci. Rep.* (2019) in press.
- 311 [18] S. Pisanu, T. Cubeddu, D. Pagnozzi, S. Rocca, C. Cacciotto, A. Alberti, G. Marogna, S. Uzzau, M.F.
312 Addis, Neutrophil extracellular traps in sheep mastitis, *Vet. Res.* 46 (2015) 59.
313 <https://doi.org/10.1186/s13567-015-0196-x>.
- 314 [19] S. Pisanu, G. Biosa, L. Carcangiu, S. Uzzau, D. Pagnozzi, Comparative evaluation of seven commercial
315 products for human serum enrichment/depletion by shotgun proteomics, *Talanta.* 185 (2018) 213–
316 220. <https://doi.org/10.1016/j.talanta.2018.03.086>.
- 317 [20] W.M. Old, K. Meyer-Arendt, L. Aveline-Wolf, K.G. Pierce, A. Mendoza, J.R. Sevinsky, K.A. Resing, N.G.
318 Ahn, Comparison of label-free methods for quantifying human proteins by shotgun proteomics.,
319 *Mol. Cell. Proteomics.* 4 (2005) 1487–502. <https://doi.org/10.1074/mcp.M500084-MCP200>.
- 320 [21] B. Zybailov, A.L. Mosley, M.E. Sardu, M.K. Coleman, L. Florens, M.P. Washburn, Statistical analysis of
321 membrane proteome expression changes in *Saccharomyces cerevisiae*., *J. Proteome Res.* 5 (2006)
322 2339–47. <https://doi.org/10.1021/pr060161n>.
- 323 [22] S. Pisanu, S. Ghisaura, D. Pagnozzi, G. Biosa, A. Tanca, T. Roggio, S. Uzzau, M.F. Addis, The sheep milk
324 fat globule membrane proteome., *J. Proteomics.* (2011) 350–358.
- 325 [23] D. Szklarczyk, A.L. Gable, D. Lyon, A. Junge, S. Wyder, J. Huerta-Cepas, M. Simonovic, N.T. Doncheva,
326 J.H. Morris, P. Bork, L.J. Jensen, C. von Mering, STRING v11: protein–protein association networks
327 with increased coverage, supporting functional discovery in genome-wide experimental datasets,
328 *Nucleic Acids Res.* 47 (2019) D607–D613. <https://doi.org/10.1093/nar/gky1131>.
- 329 [24] J. Campbell, C.J. Ryan, R. Brough, I. Bajrami, H.N. Pemberton, I.Y. Chong, S. Costa-Cabral, J. Frankum,
330 A. Gulati, H. Holme, R. Miller, S. Postel-Vinay, R. Rafiq, W. Wei, C.T. Williamson, D.A. Quigley, J. Tym,
331 B. Al-Lazikani, T. Fenton, R. Natrajan, S.J. Strauss, A. Ashworth, C.J. Lord, Large-Scale Profiling of
332 Kinase Dependencies in Cancer Cell Lines, *Cell Rep.* 14 (2016) 2490–2501.
333 <https://doi.org/10.1016/j.celrep.2016.02.023>.
- 334 [25] J.A. Vizcaíno, A. Csordas, N. Del-Toro, J.A. Dienes, J. Griss, I. Lavidas, G. Mayer, Y. Perez-Riverol, F.
335 Reisinger, T. Ternent, Q.W. Xu, R. Wang, H. Hermjakob, 2016 update of the PRIDE database and its
336 related tools, *Nucleic Acids Res.* 44 (2016) D447–D456. <https://doi.org/10.1093/nar/gkv1145>.
- 337 [26] P.D. Eckersall, F.J. Young, a M. Nolan, C.H. Knight, C. McComb, M.M. Waterston, C.J. Hogarth, E.M.
338 Scott, J.L. Fitzpatrick, Acute phase proteins in bovine milk in an experimental model of
339 *Staphylococcus aureus* subclinical mastitis., *J. Dairy Sci.* 89 (2006) 1488–501.
340 [https://doi.org/10.3168/jds.S0022-0302\(06\)72216-0](https://doi.org/10.3168/jds.S0022-0302(06)72216-0).
- 341 [27] M. Singh, A. Sharma, R. Sharma, D. Mittal, P. Yadav, G. Charaya, Estimation of acute phase proteins
342 as early biomarkers of buffalo subclinical mastitis, *Asian J. Anim. Vet. Adv.* 10 (2015) 894–902.
343 <https://doi.org/10.3923/ajava.2015.894.902>.
- 344 [28] F.C. Thomas, M. Waterston, P. Hastie, T. Parkin, H. Haining, P.D. Eckersall, The major acute phase
345 proteins of bovine milk in a commercial dairy herd, *BMC Vet. Res.* 11 (2015) 207.
346 <https://doi.org/10.1186/s12917-015-0533-3>.
- 347 [29] T.T. Wheeler, G.A. Smolenski, D.P. Harris, S.K. Gupta, B.J. Haigh, M.K. Broadhurst, A.J. Molenaar, K.
348 Stelwagen, Host-defence-related proteins in cows' milk, *Animal.* 6 (2012) 415–422.
349 <https://doi.org/10.1017/S1751731111002151>.
- 350 [30] M. Akerstedt, L. Björck, K. Persson Waller, A. Sternesjö, Biosensor assay for determination of
351 haptoglobin in bovine milk., *J. Dairy Res.* 73 (2006) 299–305.
352 <https://doi.org/10.1017/S0022029906001774>.
- 353 [31] M. Akerstedt, K. Persson Waller, A. Sternesjö, Haptoglobin and serum amyloid A in relation to the
354 somatic cell count in quarter, cow composite and bulk tank milk samples., *J. Dairy Res.* 74 (2007)
355 198–203. <https://doi.org/10.1017/S0022029906002305>.
- 356 [32] M. Åkerstedt, K.P. Waller, L.B. Larsen, L. Forsbäck, Å. Sternesjö, Relationship between haptoglobin
357 and serum amyloid A in milk and milk quality, *Int. Dairy J.* 18 (2008) 669–674.
358 <https://doi.org/10.1016/j.idairyj.2008.01.002>.

- 359 [33] A. Miglio, L. Moscati, G. Fruganti, M. Pela, E. Scoccia, A. Valiani, C. Maresca, Use of milk amyloid A in
360 the diagnosis of subclinical mastitis in dairy ewes., *J. Dairy Res.* 80 (2013) 496–502.
361 <https://doi.org/10.1017/S0022029913000484>.
- 362 [34] M.F. Addis, V. Tedde, S. Dore, S. Pisanu, G.M.G. Puggioni, A.M. Roggio, D. Pagnozzi, S. Lollai, E.A.
363 Cannas, S. Uzzau, Evaluation of milk cathelicidin for detection of dairy sheep mastitis, *J. Dairy Sci.*
364 (2016) 1–13.
- 365 [35] M.F. Addis, V. Tedde, G.M.G. Puggioni, S. Pisanu, A. Casula, C. Locatelli, N. Rota, V. Bronzo, P.
366 Moroni, S. Uzzau, Evaluation of milk cathelicidin for detection of bovine mastitis, *J. Dairy Sci.* 99
367 (2016) 8250–8258. <https://doi.org/10.3168/jds.2016-11407>.
- 368 [36] Z. Olumee-Shabon, T. Swain, E.A. Smith, E. Tall, J.L. Boehmer, Proteomic analysis of differentially
369 expressed proteins in caprine milk during experimentally induced endotoxin mastitis, *J. Dairy Sci.* 96
370 (2013) 2903–2912. <https://doi.org/10.3168/jds.2012-5956>.
- 371 [37] N. Li, Y. Wang, C. You, J. Ren, W. Chen, H. Zheng, Z. Liu, Variation in Raw Milk Microbiota Throughout
372 12 Months and the Impact of Weather Conditions, *Sci. Rep.* 8 (2018) 2371.
373 <https://doi.org/10.1038/s41598-018-20862-8>.
- 374 [38] E.M. Ibeagha-Awemu, A.E. Ibeagha, S. Messier, X. Zhao, Proteomics, genomics, and pathway
375 analyses of escherichia coli and staphylococcus aureus infected milk whey reveal molecular
376 pathways and networks involved in mastitis, *J. Proteome Res.* 9 (2010) 4604–4619.
377 <https://doi.org/10.1021/pr100336e>.
- 378 [39] O. Wellnitz, E.T. Arnold, R.M. Bruckmaier, Lipopolysaccharide and lipoteichoic acid induce different
379 immune responses in the bovine mammary gland, *J. Dairy Sci.* 94 (2011) 5405–5412.
380 <https://doi.org/10.3168/JDS.2010-3931>.
- 381 [40] T.T. Wheeler, A.J. Hodgkinson, C.G. Prosser, S.R. Davis, Immune components of colostrum and milk--
382 a historical perspective., *J. Mammary Gland Biol. Neoplasia.* 12 (2007) 237–47.
383 <https://doi.org/10.1007/s10911-007-9051-7>.
- 384 [41] Y. Lin, L. Xia, J.D. Turner, X. Zhao, Morphologic observation of neutrophil diapedesis across bovine
385 mammary gland epithelium in vitro., *Am. J. Vet. Res.* 56 (1995) 203–7.
386 <http://www.ncbi.nlm.nih.gov/pubmed/7717587> (accessed July 10, 2019).
- 387 [42] K. Stelwagen, Effect of Milking Frequency on Mammary Functioning and Shape of the Lactation
388 Curve, *J. Dairy Sci.* 84 (2001) E204–E211. [https://doi.org/10.3168/JDS.S0022-0302\(01\)70219-6](https://doi.org/10.3168/JDS.S0022-0302(01)70219-6).
- 389 [43] K. Stelwagen, K. Singh, The role of tight junctions in mammary gland function, *J. Mammary Gland
390 Biol. Neoplasia.* 19 (2014) 131–138. <https://doi.org/10.1007/s10911-013-9309-1>.
- 391 [44] I.-H. Lai, J.H. Tsao, Y.P. Lu, J.W. Lee, X. Zhao, F.L. Chien, S.J.T. Mao, Neutrophils as one of the major
392 haptoglobin sources in mastitis affected milk., *Vet. Res.* 40 (2009) 17.
393 <https://doi.org/10.1051/vetres:2008055>.
- 394 [45] A. Tripathi, K.M. Lammers, S. Goldblum, T. Shea-Donohue, S. Netzel-Arnett, M.S. Buzza, T.M. Antalis,
395 S.N. Vogel, A. Zhao, S. Yang, M.-C. Arrietta, J.B. Meddings, A. Fasano, Identification of human
396 zonulin, a physiological modulator of tight junctions, as prehaptoglobin-2., *Proc. Natl. Acad. Sci. U. S.
397 A.* 106 (2009) 16799–804. <https://doi.org/10.1073/pnas.0906773106>.
- 398 [46] D.A. Nguyen, M.C. Neville, Tight junction regulation in the mammary gland., *J. Mammary Gland Biol.
399 Neoplasia.* 3 (1998) 233–46. <http://www.ncbi.nlm.nih.gov/pubmed/10819511> (accessed July 1,
400 2019).
- 401 [47] G. Smolenski, S. Haines, F.Y.S. Kwan, J. Bond, V. Farr, S.R. Davis, K. Stelwagen, T.T. Wheeler,
402 Characterisation of host defense proteins in milk using a proteomic approach, *J. Proteome Res.* 6
403 (2007) 207–215.
- 404 [48] O. Wellnitz, E.T. Arnold, M. Lehmann, R.M. Bruckmaier, Short communication: Differential
405 immunoglobulin transfer during mastitis challenge by pathogen-specific components, *J. Dairy Sci.* 96
406 (2013) 1681–1684. <https://doi.org/10.3168/jds.2012-6150>.
- 407 [49] J.E. Eriksson, T. Dechat, B. Grin, B. Helfand, M. Mendez, H.-M. Pallari, R.D. Goldman, Introducing
408 intermediate filaments: from discovery to disease, *J. Clin. Invest.* 119 (2009) 1763–1771.
409 <https://doi.org/10.1172/JCI38339>.
- 410 [50] T.N. Mak, H. Brüggemann, Vimentin in Bacterial Infections., *Cells.* 5 (2016).

- 411 <https://doi.org/10.3390/cells5020018>.
- 412 [51] G. dos Santos, M.R. Rogel, M.A. Baker, J.R. Troken, D. Urich, L. Morales-Nebreda, J.A. Sennello, M.A.
413 Kutuzov, A. Sitikov, J.M. Davis, A.P. Lam, P. Cheresch, D. Kamp, D.K. Shumaker, G.R.S. Budinger, K.M.
414 Ridge, Vimentin regulates activation of the NLRP3 inflammasome, *Nat. Commun.* 6 (2015) 6574.
415 <https://doi.org/10.1038/ncomms7574>.
- 416 [52] M. Mudaliar, R. Tassi, F.C. Thomas, T.N. McNeilly, S.K. Weidt, M. McLaughlin, D. Wilson, R.
417 Burchmore, P. Herzyk, P.D. Eckersall, R.N. Zadoks, Mastitomics, the integrated omics of bovine milk
418 in an experimental model of *Streptococcus uberis* mastitis: 2. Label-free relative quantitative
419 proteomics, *Mol. BioSyst.* 12 (2016) 2748–2761. <https://doi.org/10.1039/C6MB00290K>.
- 420 [53] C.A. Ogden, A. deCathelineau, P.R. Hoffmann, D. Bratton, B. Ghebrehiwet, V.A. Fadok, P.M. Henson,
421 C1q and mannose binding lectin engagement of cell surface calreticulin and CD91 initiates
422 macropinocytosis and uptake of apoptotic cells., *J. Exp. Med.* 194 (2001) 781–95.
423 <https://doi.org/10.1084/jem.194.6.781>.
- 424



426

427 **Figure 1. STRING interaction networks of the proteins significantly changed only in *S. aureus* infected**
 428 **milk. Proteins associated with relevant statistically significant Biological Process GO terms (A) and KEGG**
 429 **Pathways (B) are marked with different colors as indicated. Gene names correspond to the proteins**
 430 **reported in Table 2.**

431 TABLES

432 Table 1. Summary of differential proteomic results.

	Eligible for comparison*	Changed**	Differential*** $R_{SC} \leq -1.0$ or ≥ 1.0	Increased*** $R_{SC} \geq 1.0$	Decreased*** $R_{SC} \leq -1.0$
Infected ^b	243	62	52	52	0
Uninfected ^b	152	20	19	18	1

433 ^aLate lactation *S. aureus* infected milk vs mid-lactation uninfected milk. ^bLate lactation uninfected milk
434 vs mid-lactation uninfected milk. *Proteins identified in at least two biological replicates and with ≥ 2
435 spectral counts in at least one sample of the experimental group. ** $p \leq 0.05$ by the beta-binomial test
436 with FDR correction according to Benjamini-Hochberg. *** $p \leq 0.05$ by the beta-binomial test with FDR
437 correction according to Benjamini-Hochberg and $R_{SC} \leq -1.0$ or ≥ 1.0 .
438

439 **Table 2. Significantly differential proteins observed in late lactation *S. aureus* infected and uninfected milk**
 440 **in comparison to mid-lactation uninfected milk.** The respective R_{SC} values are reported for all the differential
 441 proteins in the two comparisons. Bold: proteins significantly increased only in infected milk.

Accession	Gene name	Description	Infected ^{a*}	Uninfected ^{b*}
Q29477	LTF	Lactotransferrin	4.22	4.05
P82018	CATHL2	Cathelicidin-2	3.30	2.75
P48616	VIM	Vimentin	3.01	-
Q9XSJ4	ENO1	Alpha-enolase	2.98	1.90
P46193	ANXA1	Annexin A1	2.84	2.24
P62808	HIST1H2B	Histone H2B type 1	2.74	2.12
P60712	ACTA1	Actin, cytoplasmic 1	2.49	1.74
Q8SPQ0	CHI3L1	Chitinase-3-like protein 1	2.38	2.12
P62803	H4	Histone H4	2.37	1.71
Q2UVX4	C3	Complement C3	2.35	2.62
P07589	FN1	Fibronectin	2.33	-
Q28178	THBS1	Thrombospondin-1	2.33	-
A5D7D1	ACTN4	Alpha-actinin-4	2.24	1.10
P02584	PFN1	Profilin-1	2.15	1.10
P68138	ACTA1	Actin, alpha skeletal muscle	2.10	1.23
Q2KJD0	TUBB	Tubulin beta-5 chain	2.10	-
Q3MHM5	TUBB4B	Tubulin beta-4B chain	2.05	-
P62871	GNB1	Guanine nucleotide-binding protein G(I)/G(S)/G(T) sub β -1	2.00	-
Q3B7N2	ACTN1	Alpha-actinin-1	2.00	-
Q5VI41	ITGB2	Integrin beta-2	2.00	-
P10096	GAPDH	Glyceraldehyde-3-phosphate dehydrogenase	1.88	-
A7E3Q8	PLS3	Plastin-3	1.88	-
Q71SP7	FASN	Fatty acid synthase	-	1.85
Q3SWX7	ANXA3	Annexin A3	1.82	-
P31976	EZR	Ezrin	1.76	1.65
Q2TBU0	HP	Haptoglobin	1.70	-
P63103	YWHAZ	14-3-3 protein zeta/delta	1.70	-
P60661	MYL6	Myosin light polypeptide 6	1.63	-
P68250	YWHAB	14-3-3 protein beta/alpha	1.56	-
Q5E956	TPI1	Triosephosphate isomerase	1.70	1.56
P02253	HIST1H1C	Histone H1.2	1.48	-
Q3SZI4	YWHAQ	14-3-3 protein theta	1.48	-
O18739	CTGF	Connective tissue growth factor	1.48	-
P81947	TUBA1B	Tubulin alpha-1B chain	1.48	-
Q0VCP3	OLFML3	Olfactomedin-like protein 3	1.23	1.45
Q76LV2	HSP90AA1	Heat shock protein HSP 90-alpha	1.42	-
P81287	ANXA5	Annexin A5	1.40	-
Q3SX14	GSN	Gelsolin	1.38	-
Q32PJ2	APOA4	Apolipoprotein A-IV	1.38	-
Q3TOP6	PGK1	Phosphoglycerate kinase 1	1.32	-
Q92176	CORO1A	Coronin-1A	1.32	-
P21809	BGN	Biglycan	1.23	-
Q76LV1	HSP90AB1	Heat shock protein HSP 90-beta	1.23	-
P62157	CALM2	Calmodulin	1.23	-
Q03247	APOE	Apolipoprotein E	1.22	-
P35541	SAA2	Serum amyloid A protein	1.21	2.41
P62261	YWHAE	14-3-3 protein epsilon	1.14	-

Accession	Gene name	Description	Infected ^{a*}	Uninfected ^{b*}
P04272	ANXA2	Annexin A2	1.14	-
P0C0S9	HIST1H2AG	Histone H2A type 1	1.14	-
Q29443	TF	Serotransferrin	1.13	-
P52193	CALR	Calreticulin	-	1.10
A4IF97	MYL12B	Myosin regulatory light chain 12B	1.04	-
Q6B855	TKT	Transketolase	1.04	-
P01030	C4	Complement C4	1.04	-
Q2KIS7	CLEC3B	Tetranectin	1.04	-
Q5KR47	TPM1	Tropomyosin alpha-3 chain	1.04	-
P80025	LPO	Lactoperoxidase	-	-1.04

442 ^aLate lactation, high SCC, *S. aureus*-infected milk vs mid-lactation, low SCC, uninfected milk. ^bLate lactation, high SCC,
443 uninfected milk vs mid-lactation, low SCC, uninfected milk. *R_{SC} ≥ 1.0 or ≤ -1.0 and p-value ≤ 0.05 with FDR correction
444 according to Benjamini-Hochberg.
445

446