

Knowledge and Management of Aquatic Ecosystems

Is landscape of fear of macroinvertebrate communities a major determinant of mesopredator and prey activity?

--Manuscript Draft--

Manuscript Number:	kmae190137
Full Title:	Is landscape of fear of macroinvertebrate communities a major determinant of mesopredator and prey activity?
Article Type:	Short Communication
Keywords:	seepage; triclad; isopod; behaviour; predator.
Corresponding Author:	Raoul Manenti Universita degli Studi di Milano Milano, Milano ITALY
Corresponding Author E-Mail:	raoulmanenti@gmail.com
Order of Authors:	Raoul Manenti Benedetta BARZAGHI
Abstract:	Macroinvertebrate foragers play an important role on the trophic structures of freshwater environments, and multiple trophic levels occur among macroinvertebrate communities providing very interesting scenarios for testing scientific hypotheses. One of the most intriguing aspect to understand is the role played by the landscape of fear (LOF) on macrobenthos density and activity.
Additional Information:	
Question	Response
Suggested Reviewers:	Daniela Ghia Universita degli Studi di Pavia daniela.ghia@unipv.it Expert of invertebrates Ronald Sluys Naturalis Nationaal Natuurhistorisch Museum ronald.sluys@naturalis.nl Expert of behaviour and ecology of triclads Diego Fontaneto Istituto per lo Studio degli Ecosistemi Consiglio Nazionale delle Ricerche d.fontaneto@ise.cnr.it Studies freshwater invertebrates distribution and ecology.
Opposed Reviewers:	

Milano, 01/11/2019

Dear, Prof. Daniel Gerdeaux,

Thank you fro the quick answer.

We shortened consistently the paper to submit it as a short communication. As we are convinced that testing the landscape of fear (LOF) on freshwater macroinvertebrate communities is quite novel and very interesting we believe that a short and focused communication can be a good basis for further and more detailed studies.

We used multiple surveys and multiple season to assess if the LOF at the macroinvertebrate communities affected the density of three target invertebrate species, a detrtivor and two mesopredators.

The broad implication of the research is that the abundance of freshwater macroinvertebrates is mainly linked to some environmental than to the predation risk at the community level.

Please find enclosed the short communication entitled “Is landscape of fear of macroinvertebrate communities a major determinant of mesopredator and prey activity?”, to be considered for publication in Knowledge and Management of Aquatic Ecosystems.

We confirm that:

- The enclosed work was never submitted or published and to another journal;
- its submission for publication was approved by all relevant authors and institutions
- all persons entitled to authorship have been so named
- all authors have seen and agreed to submit this version of the manuscript.

Yours sincerely,

Raoul Manenti, Benedetta Barzaghi

1 Title page

2

3 **Is landscape of fear of macroinvertebrate communities a major determinant of mesopredator**
4 **and prey activity?**

5

6 Raoul MANENTI*, Benedetta BARZAGHI

7

8 Department of Environmental Science and Policy, università degli Studi di Milano, Via Celoria, 26,
9 20133 Milano (Italy)

10 *= corresponding author: raoulmanenti@gmail.com

11

12

13 Running head: landscape of fear of macroinvertebrates

14

15 **Abstract**

16 Macroinvertebrate foragers play an important role on the trophic structures of freshwater
17 environments, and multiple trophic levels occur among macroinvertebrate communities providing
18 very interesting scenarios for testing scientific hypotheses. One of the most intriguing aspect to
19 understand is the role played by the landscape of fear (LOF) on macrobenthos density and activity.

20 With this pilot study we wanted to test if LOF at the macrobenthos community levels play a role in
21 determining the density of both prey and mesopredators.

22 During two consecutive years, we evaluated, with both day and night surveys, the density of two
23 mesopredator triclad species and of one detritivore prey crustacean species comparing it to the number
24 of respective predators occurring in the macroinvertebrate community.

25 LOF levels at the macroinvertebrate community did not reduce the abundance of the target taxa. One
26 of the triclad species was instead positively related to the levels of LOF assessed for it on the basis of
27 the available knowledge.

28 The broad implication of the research is that the abundance of freshwater macroinvertebrates is not
29 mainly linked to the predation risk at the community level, suggesting that also for researches on
30 macrobenthos LOF analyses should take in consideration the role of top predators.

31

32 **Keywords:** seepage, triclad, isopod, behaviour, predator.

33

34 The communities of freshwater invertebrates are regarded as fundamental indicator of the status and
35 pollution of freshwater habitats. Several factors may determine differences in macroinvertebrate
36 activity and distribution; in general, all the aspects under the constraints of Darwinian natural
37 selection as: food availability, predation risk and other inter- and intra-specific interactions may
38 concur to determine differences in macroinvertebrate species density (Elliott, 2000, 2002; Kusano
39 and Kusano, 1991). While food availability is a well-recognised element regulating macrobenthos
40 abundance and diel activity (Elliott, 2002; Fiser et al., 2007), in freshwater habitats less attention is
41 paid to the predation risk that may involve different taxa. In particular, one of the most intriguing
42 aspect to understand is the role played by landscape of fear (LOF) on macrobenthos density and
43 activity: macroinvertebrate foragers play an important role on the trophic structures of freshwater
44 environments, and multiple trophic levels occur among macroinvertebrate communities furnishing
45 several opportunities to study LOF effects (Marino et al., 2016). A forager has to usefully adopt
46 strategies to forage based on the type of risk it is likely to face (Matassa and Trussell, 2011).
47 Generally, the activity patterns a forager must take to cope risk from habitats with high number of
48 predators, will differ greatly from those it will take to exploit safer habitats (Melotto et al., 2019).
49 Within predation risk both the diversity of the predator community and of the predator activity play
50 major roles in affecting the LOF (Gaynor et al., 2019); in particular, the predator diel activity levels
51 may strongly change the features of LOF ((Bleicher et al., 2019; Laundre, 2010) with consequent
52 reflections on prey activity itself. However, how much LOF levels may affect macroinvertebrate
53 species sampling and activity remains an intriguing aspect to be studied.

54 To assess if LOF affects the macrobenthos diel activity, we studied environments with similar aquatic
55 top predator presence, such as day active visual predators (fish) and night active wanderer predators
56 (the invasive crayfish *Procambarus clarkii*) and we focused only on the LOF at the macroinvertebrate
57 community level.

58 First of all, we tested if LOF varied between day and night conditions; second we evaluated the
59 relationship between LOF and density of both target mesopredator and prey invertebrate species. In
60 particular, we tested two hypotheses:

- 61 1) Fear hypothesis; LOF affects prey and/or mesopredator density activity.
- 62 2) No fear hypothesis; The density of predator and/or prey species varies with day/night conditions
63 notwithstanding to the levels of LOF.

64 In particular, during two consecutive years, we evaluated, with both day and night surveys, the density
65 of two mesopredator triclad species and of one detritivore prey crustacean species comparing it to the
66 predator occurring in the whole macroinvertebrate community.

67 We performed the study in Lombardy (NW Italy). We studied four “fontanili”; springs forming lentic
68 habitats fed by groundwater flow. Fontanili are springs anciently managed by humans pushing tubes
69 in the substrate to collect groundwaters and ease their flow toward the surface (Balderacchi et al.,
70 2016). Fontanili springs are generally characterised by a large head in which the tubes occur and a
71 straight section that allows water to flow out. We performed transects in the spring head along the
72 outflow tubes tracing 8 transects (2 for each site). The transects were all 1 m wide, but varied in length
73 depending on the site features (length average \pm SE = 4.3 ± 0.7 m).

74 During winter months, from December 2017 to February 2018 and from December 2018 to February
75 2019 we performed for each site 12 repeated surveys both during day and during night (6 surveys
76 during night and 6 during day). During surveys in each site we first assessed visually the occurrence
77 and the number of the target taxa such as crustaceans of the species *Asellus aquaticus* and planarians
78 of the species *Polycelis nigra* and *Dendrocoelum lacteum* along two transects per site.

79 During each survey, in each transect after 20 minutes of visual encounter numbering of the target
80 organisms we sampled the whole macrobenthos community using a dip-net. Net samplings lasted 10
81 minutes in each transect and were performed by intense movement of the substrate. All the collected
82 organisms were released in the transect of origin after having been numbered and recognised at
83 species, genus or family level according to the guidelines for the Italian Biotic Index assessment

84 (Ghetti 1997). We also we assessed the occurrence of wanderer top predator species like fish. From
85 each survey we kept at minimum 4 days of interval. During surveys we recorded also maximum
86 illuminance of the water surface (with a PCE EM882 luxmeter) and water temperature. LOF
87 assessment considered the taxa collected through the dip-netting of the substrate at each sampling
88 session. LOF was calculated using the number of potential predator taxa for each target species
89 occurring in the transects: we divided the number of occurring predators per the total number of
90 taxonomical units collected. Predator assessment was based on the information available in the
91 literature (Ghetti, 1997; Reynoldson and Young, 2000; Tachet, 2010).

92 To test if LOF was different between sites and day/night conditions, we developed a Linear Mixed
93 Model (LMM) using the log transformed levels of LOF as dependent variables and the transect
94 identity and the period (day/night) as fixed factors; we considered also the year of sampling as random
95 factor. Through a Wald F test we assessed the significance of the fixed factors composing the model.
96 We then used random-effect generalized mixed models (GLMMs) to assess the relationships between
97 the relative abundance of the target taxa and the LOF (Barker et al., 2017). In particular, we used a
98 negative binomial distribution to account for over dispersion as, especially for planarians we had
99 different 0 occurrences. As a dependent variable, we considered the number of active individuals of
100 the target taxa observed for each transect at each survey. We included the moment of observation
101 (day/night) and the sampling method (visual/net) as covariates. We included the year of survey, the
102 number of survey and the transect as random factors.

103 GLMMs and LMMs were run in R environment (R Development Core Team 2018) using a negative
104 binomial error, using the package glmmTMB, lmerTest and car (Brooks et al., 2017).

105 Considering the whole samplings, *Polycelis nigra* was the more abundant species (on average (\pm SE)
106 34.9 ± 8.5 individuals per sampling). Considering night samplings only the average number of *Asellus*
107 *aquaticus* observed overcame the average number of *P. nigra* (18 ± 5.2 *A. aquaticus* individuals' vs
108 16.9 ± 5.2 *P. nigra* individuals).

109 Water temperature was on average (\pm SE) $12,17\text{ }^{\circ}\text{C} \pm 0,19\text{ }^{\circ}\text{C}$; a significant difference, assessed
110 through ANOVA and post-hoc tukey test was observed only between two of the sites ($F= 3,2$; $P =$
111 $0,03$). Maximum illuminance in fontanili during sunny days was around 40000 lux and ranged
112 between 0,01 and 0,1 lux during night with no significant differences between sites. Wanderer top
113 predator taxa such as fish and the alien freshwater crayfish *Procambarus clarkii* were recorded in all
114 the sites with at least one observation in the proximity or inside the transects during each year of
115 monitoring. While fish were observed both during day and night, crayfish were detected mainly
116 during night. In the transects we recorded globally 13 macroinvertebrate predator taxa at which for
117 our crustacean target species must be added the two planarians target species. On average (\pm SE),
118 considering all the predator taxa, the number of the potential predator individual for our crustacean
119 target species was of $1,25 \pm 0,34$ individuals per net sampling.

120 LOF levels differed among sites (for both crustacean and planarians LOF levels: $F > 5,7$; $P < 0,001$),
121 but not between day/night conditions (for both crustacean and planarians LOF levels: $F < 0,22$; $P >$
122 $0,64$). *Dendrocoelum lacteum* was significantly more abundant during night (Table 1, Fig. 1). On the
123 other hand, the abundance of *Polycelis nigra* showed a weak, unexpected and significant relationship
124 with the LOF levels considered, being more abundant in transects with higher levels of LOF (Table
125 1, Fig. 1). *A. aquaticus* was more abundant when sampled with deep net (Table 1, and Fig. 2). No
126 significant effect was played by the LOF levels considered.

127 The broad implication of the present research is that LOF at the macroinvertebrate community level
128 does not seem to affect the activity of macroinvertebrate foragers. On the other side the abundance of
129 both macroinvertebrate predators and prey is strongly related to other factors irrespective to their
130 position on the food web and to LOF levels.

131 The ecological study of LOF is increasingly being recognised as central in understanding the patterns
132 driving predator-prey interactions (Gallagher et al., 2017). LOF can determine the population density
133 of a species, but interspecific competitive/predatory interactions in complex communities may
134 produce various combinations of impacts (Gallagher et al., 2017; Laundre et al., 2014). In freshwaters

135 communities, where food webs are often highly structured, the study of LOF effects can be intriguing
136 and reveal important insights in terms of management. As examples, assessing the role of predators
137 may increase the efficacy of restoration action in lotic environments, while understanding the role of
138 LOF in spring habitats may reveal important insights for understanding some of the evolutionary
139 pressures that drive groundwaters colonisation. However, our study suggests that further work is in
140 order. From one side mesopredator taxa considered in our study may not be the most important
141 determinants of the LOF in the system that hosts native fish and alien crayfish as top predators. Often
142 top predators feed both on mesopredators with which they share prey (Rodriguez-Lozano et al., 2015)
143 with likely LOF top-down control on both mesopredator and prey. Thus a finer scaled characterisation
144 of LOF levels based on the foraging activity of top wanderer predators like fish and crayfish may
145 reveal different patterns.

146 Moreover, some of our results suggest that the evaluation of LOF for planarians on the basis of the
147 available information could not be sufficiently reliable. The assessment of LOF level has been made
148 on the basis of the few information available for the genus *Polycelis* and its possible predator (Tachet,
149 2010). However some of the predator taxa included could not directly feed on *P. nigra* that it is itself
150 considered mainly a predator of living macroinvertebrates with a minor preference for also dead
151 invertebrates (Reynoldson and Young, 2000; Tachet, 2010). It is possible that *P. nigra* is an
152 opportunistic mesopredator feeding on already damaged/dead invertebrates and thus favoured by
153 other mesopredator occurrence.

154 A second argument of discussion originating from our results is the differential effect played by
155 day/night conditions on the invertebrate target species. While the abundance of individuals of both
156 crustacean and planarian pigmented species does not differ between day and night, the abundance of
157 individuals belonging to the unpigmented *Dendrocoelum lacteum* species is slightly higher during
158 night. Generally, nocturnal activity in both vertebrates and invertebrates is considered as an adaptive
159 strategy to minimize risk of predation (Huhta et al., 2000; Kotler et al., 2010) and is often supported
160 by heightened non-visual senses that allow detection of threatens in darkness conditions (Bleicher et

161 al., 2019; Vestheim et al., 2013). In our system that assessed LOF levels of invertebrate community
162 through substrate sampling, for both detritivor and mesopredator target species LOF levels did not
163 vary between day and night. *D. lacteum* is an unpigmented epigeal species for which some general
164 ecological study has been performed (Herrmann, 1986; Reynoldson and Young, 2000); however, no
165 detailed behavioural information exist; generally freshwater planarians are regarded as nocturnal
166 (Lombardo et al., 2011), but our results indicate that between different genera and species slight
167 differences in the diel activity may occur. As we have argued elsewhere the study of LOF in
168 freshwater environments may be considered a promising aspect to understand evolutionary and
169 ecological patterns shaping freshwater organisms' distribution. Our results suggest however that more
170 studies are necessary to increase the knowledge of species composing the microbenthic community
171 and that the potential role of top predators should be accounted at different habitat scales.

172

173 **Acknowledgments**

174 The research has been funded by The Mohamed Bin Zayed Species Conservation Fund, project n.
175 180520056. We are grateful to Enrico Pezzoli, Pietro Pozzoli, Giancarlo Ghioni, Tullio Muraro and
176 the Comitato Bevere ong for support during field observations. We thank Serena Ghioni for
177 comments on a previous version of the manuscript.

178 **References**

179

180 Balderacchi, M., Perego, A., Lazzari, G., Munoz-Carpena, R., Acutis, M., Laini, A., Giussani, A.,
181 Sanna, M., Kane, D. and Trevisan, M., 2016. Avoiding social traps in the ecosystem stewardship:
182 The Italian Fontanile lowland spring. *Science of the Total Environment* 539, 526-535.
183 Barker, R.J., Schofield, M.R., Link, W.A. and Sauer, J.R., 2017. On the reliability of N-Mixture
184 models for count data. *Biometrics* 74, 369-377.
185 Bleicher, S.S., Marko, H., Morin, D.J., Teemu, K. and Hannu, Y., 2019. Balancing food, activity
186 and the dangers of sunlit nights. *Behav Ecol Sociobiol* 73.

187 Brooks, M.E., Kristensen, K., van Benthem, K.J., Magnusson, A., Berg, C.W., Nielsen, A., Skaug,
188 H.J., Maechler, M. and Bolker, B., 2017. glmmTMB Balances Speed and Flexibility Among
189 Packages for Zero-inflated Generalized Linear Mixed Modeling. *The R Journal* 9, 378-400.

190 Elliott, J.M., 2000. Contrasting diel activity and feeding patterns of four species of carnivorous
191 stoneflies. *Ecol Entomol* 25, 26-34.

192 Elliott, J.M., 2002. A quantitative study of day-night changes in the spatial distribution of insects in
193 a stony stream. *J. Anim. Ecol.* 71, 112-122.

194 Fiser, C., Keber, R., Kerezi, V., Moskríc, A., Palandancic, A., Petkovska, V., Potocnik, H. and Sket,
195 B., 2007. Coexistence of species of two amphipod genera: *Niphargus timavi* (Niphargidae) and
196 *Gammarus fossarum* (Gammaridae). *J. Nat. Hist.* 41, 2641-2651.

197 Gallagher, A.J., Creel, S., Wilson, R.P. and Cooke, S.J., 2017. Energy Landscapes and the
198 Landscape of Fear. *Trends Ecol. Evol.* 32, 88-96.

199 Gaynor, K.M., Brown, J.S., Middleton, A.D., Power, M.E. and Brashares, J.S., 2019. Landscapes of
200 Fear: Spatial Patterns of Risk Perception and Response. *Trends Ecol. Evol.* 34, 355-368.

201 Ghetti, P.F., 1997. Indice Biotico Esteso (I.B.E.): Manuale di applicazione. Provincia Autonoma di
202 Trento, Trento. 222 p.

203 Herrmann, J., 1986. Reproductive Ecology of *Dendrocoelum-Lacteum* (Turbellaria) in a Rapid
204 Stream in Southern Sweden and Comparisons with a Lake Population. *Hydrobiologia* 132, 273-277.

205 Huhta, A., Muotka, T. and Tikkanen, P., 2000. Nocturnal drift of mayfly nymphs as a post-contact
206 antipredator mechanism. *Freshw. Biol.* 45, 33-42.

207 Kotler, B.P., Brown, J., Mukherjee, S., Berger-Tal, O. and Bouskila, A., 2010. Moonlight avoidance
208 in gerbils reveals a sophisticated interplay among time allocation, vigilance and state-dependent
209 foraging. *Proc R Soc Lond B Biol Sci* 277, 1469-1474.

210 Kusano, H. and Kusano, T., 1991. Diel Activity of Breeding Individuals of a Fresh-Water
211 Amphipod, *Jesogammarus-Spinopalpus*. *J Ethol* 9, 105-111.

212 Laundre, J.W., 2010. Behavioral response races, predator-prey shell games, ecology of fear, and
213 patch use of pumas and their ungulate prey. *Ecology* 91, 2995-3007.

214 Laundre, J.W., Hernandez, L., Medina, P.L., Campanella, A., Lopez-Portillo, J., Gonzalez-Romero,
215 A., Grajales-Tam, K.M., Burke, A.M., Gronemeyer, P. and Browning, D.M., 2014. The landscape
216 of fear: the missing link to understand top-down and bottom-up controls of prey abundance?
217 *Ecology* 95, 1141-1152.

218 Lombardo, P., Giustini, M., Miccoli, F.P. and Cicolani, B., 2011. Fine-scale differences in diel
219 activity among nocturnal freshwater planarias (Platyhelminthes: Tricladida). *Journal of Circadian*
220 *Rhythms* 9, 2.

221 Marino, N.A.C., Srivastava, D.S. and Farjalla, V.F., 2016. Predator kairomones change food web
222 structure and function, regardless of cues from consumed prey. *Oikos* 125, 1017-1026.

223 Matassa, C.M. and Trussell, G.C., 2011. Landscape of fear influences the relative importance of
224 consumptive and nonconsumptive predator effects. *Ecology* 92, 2258-2266.

225 Melotto, A., Ficetola, G.F. and Manenti, R., 2019. Safe as a cave? Intraspecific aggressiveness rises
226 in predator-devoid and resource-depleted environments. *Behav Ecol Sociobiol* 73.

227 Reynoldson, J.D. and Young, J.O., 2000. A key to the freshwater triclads of Britain and Ireland
228 with notes on their ecology. Freshwater Biological Association, Ambleside (Cumbria). 72 p.

229 Rodriguez-Lozano, P., Verkaik, I., Rieradevall, M. and Prat, N., 2015. Small but Powerful: Top
230 Predator Local Extinction Affects Ecosystem Structure and Function in an Intermittent Stream. *Plos*
231 *One* 10.

232 Tachet, H., 2010. Invertébrés d'eau douce : Systématique, biologie, écologie CNRS, Paris.

233 Vestheim, H., Brucet, S. and Kaartvedt, S., 2013. Vertical distribution, feeding and vulnerability to
234 tactile predation in *Metridia longa* (Copepoda, Calanoida). *Mar Biol Res* 9, 949-957.

235

236

237 **Table and figures legends**

238

	Variables	Estimate	z	P
<i>Asellus aquaticus</i>				
	Night	0.22	1.05	0.29
	LOF	0.85	0.31	0.75
	Deep netting	1.23	5.44	< 0.001
<i>Dendrocoelum lacteum</i>				
	Night	0.72	2.79	< 0.01
	LOF	0.09	0.05	0.95
	Deep netting	0.14	0.56	0.57
<i>Polycelis nigra</i>				
	Night	0.44	1.52	0.12
	LOF	7.02	1.96	0.04
	Deep netting	0.15	0.55	0.58

239 Table 1 Results of the GLMMs analysis. In bold the significant results. LOF represents the level of
 240 landscape of fear.

241

242

243

244 Figure 1. Plots and boxplots of the relationship between the number of planarians of the species
245 *Dendrocoelum lacteum* and *Polycelis nigra* and the parameters studied. A, B and C refer to
246 *Dendrocoelum lacteum*; D, E and F to *Polycelis nigra*. Fear_planarians indicates the level of landscape
247 of fear for the planarians.

248

249 Figure 2. Plots and boxplots of the relationship between the number of crustaceans of the species
250 *Asellus aquaticus*. Fear_crustaceans indicates the level of landscape of fear for the target taxon.



