



Polyphenols and Intestinal Permeability: Rationale and Future Perspectives

Stefano Bernardi,^{†,‡} Cristian Del Bo',^{†,‡} Mirko Marino,[‡] Giorgio Gargari,[‡] Antonio Cherubini,[§] Cristina Andrés-Lacueva,^{||,⊥} Nicole Hidalgo-Liberona,^{||,⊥} Gregorio Peron,^{||,⊥} Raúl González-Dominguez,^{||,⊥} Paul Kroon,[#] Benjamin Kirkup,[#] Marisa Porrini,[‡] Simone Guglielmetti,[‡] and Patrizia Riso^{*,‡}

[†]Department of Food, Environmental and Nutritional Sciences (DeFENS), Università degli Studi di Milano, 20122 Milan, Italy

[§]Geriatrics, Accettazione Geriatrica e Centro di Ricerca per l'Invecchiamento, Istituto di Ricovero e Cura a Carattere Scientifico (IRCCS)–l'Istituto Nazionale Ricovero e Cura Anziani (INRCA), 60127 Ancona, Italy

^{||}Biomarkers and Nutrimetabolomics Laboratory, Department of Nutrition, Food Sciences and Gastronomy, Food Technology Reference Net (XaRTA), Nutrition and Food Safety Research Institute (INSA), Faculty of Pharmacy and Food Sciences, University of Barcelona, 08028 Barcelona, Spain

[⊥]CIBER de Fragilidad y Envejecimiento Saludable (CIBERfes), Instituto de Salud Carlos III, 08028 Barcelona, Spain

[#]Quadram Institute Bioscience, Norwich Research Park, Norwich NR4 7UQ, United Kingdom

ABSTRACT: Increasing evidence links intestinal permeability (IP), a feature of the intestinal barrier, to several pathological or dysfunctional conditions. Several host and environmental factors, including dietary factors, can affect the maintenance of normal IP. In this regard, food bioactives, such as polyphenols, have been proposed as potential IP modulators, even if the mechanisms involved are not yet fully elucidated. The aim of the present paper is to provide a short overview of the main evidence from *in vitro* and *in vivo* studies supporting the role of polyphenols in modulating IP and briefly discuss future perspectives in this research area.

KEYWORDS: polyphenols, intestinal permeability, *in vitro* studies, animal studies, human studies

INTRODUCTION

Over the last 10 years, there has been significant research effort to investigate the central role of gut function and properties in the promotion of human health and/or the development of several pathological conditions.

The intestine is the main organ involved in the absorption of nutrients and water, and it is the largest area of contact with environmental factors. It contains a large number of specialized immune cells that can coordinate with defensive responses that prevent or counteract exposure of the host and its immune system to luminal antigens of different origins (e.g., microbial and dietary origin).¹

The definition and specific ontology related to the gut as a complex anatomical and functional system has been widely debated. Bischoff et al.² defined the intestinal barrier (IB) as a functional entity separating the gut lumen from the inner host and consisting of mechanical elements (mucus and epithelial layer), humoral elements [defensins and immunoglobulin A (IgA)], immunological elements (lymphocytes and innate immune cells), and muscular and neurological elements. Differently, intestinal permeability (IP), which contributes to the regulation of solute and fluid exchange between the lumen and tissues, should refer to a key feature of IB that is measurable as a whole or at a given site (e.g., evaluating specific molecule/factor flux rates). IP evaluation can be used to address a normal/stable or disturbed/compromised permeability related with IB function.² In this context, it is fundamental to underline that IB

integrity and functionality can be affected also by the characteristics of the intestinal microbial ecosystem and mucosal immune system.

From an anatomical point of view, a well-organized monolayer of epithelial cells is required to form a selective permeability system mainly controlled by the transcellular and paracellular pathways.³

While the absorption and/or transport of nutrients (i.e., sugars, amino acids, vitamins, fatty acids, and minerals) occur through specific transporters or membrane channels (transcellular path),³ a complex system of junctions crucial for the transport between adjacent cells [i.e., tight junction (TJ), gap junction (GJ), adherens junction (AJ), and desmosomes] constitutes the paracellular path.⁴

TJs have a composite molecular structure consisting of multiple protein complexes (with more than 50 proteins identified) that include a series of transmembrane tetraspan proteins, named occludin, claudins, and tricellulin, able to develop fibrils crossing the membranes and creating a connection with adjacent cell proteins. In addition, single-span transmembrane proteins are included and are mostly

Special Issue: Food Bioactives and Health

Received: April 16, 2019

Revised: June 21, 2019

Accepted: June 24, 2019

Published: June 24, 2019

represented by junctional adhesion molecules (JAMs, belonging to the immunoglobulin superfamily). The claudin proteins are considered to be the structural pillar of TJs.⁵ Specifically, TJ sealing, fundamental to avoid paracellular permeability, is provided by claudin-1, -3, -4, -5, and -8, while claudin-2 can form charge-selective pores. Less information is available for the specific activities of claudins-7, -12, and -15 and occludin.⁶

The transmembrane proteins strictly interact with the intracellular scaffold proteins, such as zonula occludens (ZO-1, ZO-2, and ZO-3) and cingulin tight fitting the actin cytoskeleton. In particular, increased paracellular permeability is activated by perijunctional actomyosin ring contraction induced by myosin light-chain kinase (MLCK). In addition, other signaling proteins, including protein kinase C (PKC) and mitogen-activated protein kinase (MAPK), together with phosphorylation are involved in the regulation pathways of assembly, disassembly, and maintenance of TJ-specific properties.⁷ Finally, AJs, together with desmosomes and GJs located beneath the TJs, are involved in the cell-to-cell adhesion and intracellular signaling but do not seem to contribute to paracellular permeability.⁸

By considering the complex interplay of functions and activities of TJ proteins and signals regulating the fluxes/exchanges of molecules between the lumen and the environment, it is clear that TJ barrier integrity is essential for human health and metabolic homeostasis.

In fact, an impairment or defect in IB function can lead to modest (i.e., subclinical) but chronic immune system activation that might contribute to the pathogenesis of intestinal diseases, such as inflammatory bowel disease,⁴ celiac disease,⁹ intestinal bowel syndrome,¹⁰ and colon cancer.¹¹ In addition, recent research showed a possible correlation of IB dysfunction with several clinical conditions, such as metabolic syndrome, obesity, non-alcoholic fatty liver disease (NAFLD),¹² diabetes,¹³ and inflammatory joint diseases,¹⁴ but also neurological conditions, such as major depression and degenerative disorders, such as Parkinson's disease¹⁵ and multiple sclerosis (MS), involving the central nervous system (CNS).¹⁶

It is noteworthy that emerging experimental evidence suggests that an alteration of IB function and/or increased IP can actually occur also during aging, thus potentially representing a further mechanism underpinning the activation of the low-grade systemic inflammation process (also named inflammaging) identified in older subjects.¹⁷ The alterations can take place at different levels of the IB, for example, induced by impairment of the epithelium (physical barrier) and/or the immune cell/function or by an alteration of the chemical barrier consisting of the thick mucus layer able to reduce the passage of bacteria through the epithelium (i.e., mucin secretion) or as a result of an inefficient/inadequate microbial barrier (represented by the commensal "protective" bacteria). In this regard, it has been demonstrated that age-associated microbial dysbiosis can increase gut microbiota lipopolysaccharide (LPS) production and promote IP with an increased risk of systemic endotoxemia and inflammation. In particular, bacteria LPS has been demonstrated to activate nuclear factor κ B (NF- κ B) and mitogen-activated protein kinase (MAPK) by triggering the toll-like receptor 4 (TLR4) inflammatory cascade in immune cells (e.g., macrophages and monocytes).¹⁸

In addition, dysbiosis is not only an age-associated characteristic but can also be found in different clinical conditions associated with inflammation (e.g., obesity, diabetes, and NAFLD).

Thus, intestinal microbiota can be considered a critical regulator of the IP. Gut microorganisms may act directly on IP by affecting TJ properties and activities and indirectly by modulating inflammation, which is a well-recognized factor promoting IP impairment.¹⁹ Consequently, the manipulation of the complex intestinal microbial ecosystem has been proposed as a novel strategy to restore IP.²

■ DIET AND IP

An adequate nutritional status is fundamental to maintain normal IB function (being able to affect all of the components of IB), and accordingly, malnutrition is associated with increased IP.²⁰ For example, Guerriero et al.²¹ showed that a depletion of glutamine, tryptophan, and zinc could lead to increased IP.

Overall, it has been demonstrated that dietary patterns are a dominant factor in shaping the intestinal microbiota.²² Hence, strategies to modify the relative abundance of specific bacterial groups by means of dietary interventions have been proposed with the aim also to modulate the concentrations of microbial metabolites in the gut, affecting inflammation.²³

It has been demonstrated that the Western diet, characterized by high-energy and high-fat intake or high-fructose consumption, can alter IP by affecting the gut microbiota composition.²¹ In addition, this dietary pattern often involves the consumption of food components, such as specific fatty acids, alcohol, additives, gliadin, and chitosan, and food-processing methods that are known to alter IB physical structure homeostasis and/or commensal microbial homeostasis. On the other hand, a healthy dietary pattern, such as the Mediterranean diet (MD), rich in fruit, vegetables, legumes, and unrefined cereals, has been suggested to positively affect IP and related conditions.²¹ This may be related to an increased production of short-chain fatty acids (SCFAs), including acetate, propionate, butyrate, and valerate,² by gut commensal bacteria following fiber degradation provided by the MD dietary pattern. These metabolites have been suggested to play an important role as substrates for a functional colonic epithelium and the maintenance of the IB. For example, butyrate was shown to affect TJ integrity but also inhibit tumor necrosis factor α (TNF- α) release and inflammation.²³ In addition, butyrate has shown to increase expression of claudin-1 and ZO-1, to reverse the aberrant expression of ZO-1 and decrease LPS translocation, leading to inhibition of macrophage activation and pro-inflammatory cytokine production.²⁴ Moreover, plant-based dietary patterns, including MD, are also commonly abundant in bioactive compounds, such as polyphenols (PPs), that have been recently in the spotlight of research for their potential modulatory properties with respect to IP.²⁵

■ RATIONALE FOR PP CONTRIBUTION TO A PROTECTIVE DIETARY PATTERN IN THE CONTEXT OF IP

PPs are secondary metabolites of plants, widely distributed in fruits, vegetables, and plant-derived foods. A diet rich in fruits, vegetables, and plant-based beverages has been estimated to provide about 1 g of PPs/day,²⁶ with significant variations depending also upon the extent of consumption of beverages rich in PPs (tea, wine, coffee, and fruit juices). The basic monomer in PPs is the phenolic ring. Phenols can be mainly classified into phenolic acids (hydroxycinnamic and hydroxybenzoic acids), flavonoids (flavons, flavanones, flavanols, flavonols, isoflavones, and anthocyanidins), stilbenes (i.e.,




Figure 1. Putative effects of PPs on IP at different physiological levels: (1) intraluminal level, modulation of microbiota composition, endotoxin and/or SCFA production, redox status, and dietary component absorption and/or activity; (2) intracellular level, regulation of expression of TJ, AJ, GJ, and desmosome proteins, upregulation of kinases and Nrf-2, and downregulation of NF- κ B and TLR4; and (3) systemic level, maintenance of the functional immune system and regulation of inflammatory processes (toward a reduced pro-inflammatory status).

resveratrol), and lignans. PPs are recognized to be poorly bioavailable, rapidly absorbed, and extensively metabolized by gut microbiota.²⁷ Additional biotransformation can occur in the liver and kidneys through methylation, glucuronidation, and sulfation reactions of phenolic hydroxyl groups;²⁸ for these reasons, the concentration of the native compounds in the blood is low compared to their metabolic derivatives (from nanomoles up to micromoles per liter).

PPs and their metabolites are widely studied for their numerous biological activities, including antimicrobial, anti-proliferative, antioxidant, and anti-inflammatory functions.²⁹ These effects are exerted at both intestinal and systemic levels. In particular, PPs may exert their effects by downregulating inflammatory genes (i.e., NF- κ B) and upregulating cytoprotective and antioxidant genes [i.e., nuclear factor erythroid 2-related factor 2 (Nrf-2)]. This modulation may bring a reduction of cytokine production [e.g., interleukin (IL)-8, IL-1 β , and TNF- α] and boost the bodies' own antioxidant status [heme oxygenase 1 (HO-1), superoxide dismutase (SOD), and glutathione peroxidase (GPx)].³⁰ Furthermore, recent reviews^{31,32} have shown that PPs may affect, in either a positive or negative way, pattern recognition receptors, such as toll-like receptors and nucleotide-binding oligomerization domain proteins, whose activation in epithelial cells may lead to intestinal inflammation. Moreover, PPs seem to be involved in the regulation of epigenetic factors through interaction with the enzymes responsible for DNA methylation and acetylation by reducing intestinal inflammation.³²

Several studies documented the effects of PPs in the modulation of the intestinal microbial ecosystem. However, the mechanisms by which these compounds modulate the gut microbiota remain unclear. Some studies report that the interaction between PPs and microbiota may involve interference with enzymatic expression and activity and modulation of

specific pathways related to antioxidant and anti-inflammatory activity.³³ In addition, PPs has been proposed to exert a prebiotic effect potentially inhibiting the pathogenic bacteria and stimulating the growth of beneficial microbes.^{34–36} In fact, the microbiota can extensively metabolize PPs in numerous derivatives that could affect not only the composition of microbiota but also specific signaling pathways.³³ Another important aspect regards the possible involvement of PPs in the metabolism of colonic products, such as SCFAs, sterols (cholesterol and bile acids), and microbial products of non-absorbed proteins, which may directly or indirectly counteract or suppress pro-oxidant and/or pro-inflammatory responses with an overall improvement of gut health.³⁷

To unravel the complex scenario related with PP–microbiota interaction *in vivo*, a combination of metabolomic, microbiome, and metagenomic approaches is strongly demanded.³³

Finally, in the last few decades, specific research has been devoted to the evaluation of PPs as promising protective factors and regulators of epithelial homeostasis and the IB function. In particular, a direct/indirect effect of the regulation of TJ proteins has been investigated.

MECHANISMS OF PP REGULATION OF IP

At present, the exact mechanisms linking PPs with the intestinal epithelial barrier function have not yet been established (Figure 1). Some studies hypothesized a direct/indirect involvement of NF- κ B signaling in the onset of IP. This pathway is recognized as one of the most important mediators of the inflammation; cytokines and interleukins have been shown to activate NF- κ B and impair the epithelial barrier function by TJ disassembly. Conversely, PPs have documented to block NF- κ B activation by inhibiting I κ B kinase (IKK) phosphorylation and/or preventing proteasomal degradation of I κ B.³⁸

Table 1. Summary of the Main *In Vitro* Studies Highlighting the Mechanisms of Action of PP Compounds in the Modulation of Barrier Integrity and Function

reference	cells	stimulation	PP source and dose	signaling pathway	response/marker	effect
Atkinson and Rao ⁴⁰	Caco-2	acetaldehyde	genistein (30–300 μ M)	\downarrow tyrosine kinase	TEER, ^a occludin, and ZO-1 ^b	\uparrow TEER \uparrow occludin \uparrow ZO-1 \uparrow TEER
Watson et al. ⁵⁹	T84	IFN- γ^c	epigallocatechin gallate (100 μ M)	\downarrow STAT-1 ^d \downarrow MAPK ^e	TEER	\uparrow TEER
Amasheh et al. ⁵¹	Caco-2		quercetin (0–200 μ M)	\downarrow MLCK ^f and PKC ^g	TEER, occludin, claudin-1, claudin-3, claudin-4, and claudin-7	\uparrow TEER \uparrow claudin-4 = claudin-1 = claudin-3 = claudin-7 = occludin \uparrow ZO-2
Suzuki and Hara ⁵²	Caco-2		quercetin (0–100 μ M)	\downarrow PKC δ	ZO-2, occludin, claudin-1, and claudin-4	\uparrow occludin \uparrow claudin-1 \uparrow claudin-4 \uparrow claudin-1 \downarrow claudin-2 \uparrow ZO-1 \uparrow occludin \uparrow TEER \uparrow ZO-1 \uparrow occludin \uparrow claudin-4
Amasheh et al. ⁶⁰	HT29/B6	TNF- α^h	berberine (50 μ M)	\downarrow NF- κ B, ⁱ PI3K/Akt, ^j and tyrosine kinase	claudin-1 and claudin-2	\uparrow ZO-1
Chuenkityanon et al. ⁶⁴	ECV304	H ₂ O ₂ ^k	quercetin (10 μ M)	\downarrow p38 ^l	ZO-1 and occludin	\uparrow TEER \uparrow ZO-1
Rogoll et al. ⁶¹	T84		(+)-catechin (10 μ M) (–)-epicatechin (10 μ M) quercetins (10 μ M) phlorretins (20 μ M) D-(–)-quinic acids (10–50 μ M) p-coumaric acids (10 μ M) caffeic acids (20 μ M)	\downarrow TJ permeability	TEER, ZO-1, occludin, and claudin-4	\uparrow TEER \uparrow ZO-1 \uparrow occludin \uparrow claudin-4
Shin et al. ⁶⁶	HCT-116		anthocyanin mixture (45 μ g/mL), delphinidin, cyanidin, petunidin, delphinidin, malvidin, peonidin-3,5-diglucoside, cyanidin, petunidin, peonidin, and malvidin-3-glucoside)	\uparrow p38	TEER, claudin-1, claudin-3, and claudin-4	\uparrow TEER \downarrow claudin-1 \downarrow claudin-3 \downarrow claudin-4
Suzuki et al. ⁵³	Caco-2		kaempferol (100 μ M)	\downarrow TJ permeability	TEER, ZO-1, ZO-2, occludin, claudin-1, claudin-3, and claudin-4	\uparrow TEER \uparrow occludin \uparrow claudin-1 \uparrow claudin-3 \uparrow claudin-4 \uparrow ZO-1 \uparrow ZO-2
Noda et al. ⁵⁴	Caco-2		chrysin, daidzein, genistein, hesperetin, luteolin, morin, and naringenin (100 μ M)	\downarrow TJ permeability	TEER, ZO-1, ZO-2, JAMI, claudin-1, claudin-3, and claudin-4	\uparrow TEER (negative effect for chrysin) effect on TJ proteins was compound-dependent
Amasheh et al. ⁶²	HT-29/B6	IFN- γ and TNF- α	quercetin (200 μ M)	\downarrow TJ permeability	TEER, claudin-1, claudin-2, claudin-3, claudin-4, claudin-7, and occludin	\uparrow TEER \downarrow claudin-2 \downarrow claudin-3 = claudin-1

Table 1. continued

reference	cells	stimulation	PP source and dose	signaling pathway	response/marker	effect
Noda et al. ⁵⁵	Caco-2		naringenin (100 μ M)	\uparrow Sp1 ^{WT} -dependent transcriptional regulation \downarrow TJ permeability	TEER, ZO-1, ZO-2, occludin, JAM-A, ⁷¹ claudin-1, claudin-3, and claudin-4	= claudin-4 = claudin-7 = occludin \uparrow TEER
Cao et al. ⁵⁶	Caco-2	IFN- γ and TNF- α	berberine (100 μ M)	\downarrow MLCK	occludin, claudin-1, ZO-1, and IP	\uparrow claudin-1 \uparrow claudin-4 \uparrow occludin = ZO-1 = JAM-A \uparrow occludin \uparrow claudin-1 \uparrow ZO-1 \downarrow IP \uparrow TEER (no effect with rutin)
Carasco-Pozo et al. ⁵⁷	Caco-2	indomethacin	mix of quercetin (33 μ M), resveratrol (438 μ M), rutin (164 μ M), and epigallocatechin gallate (218 μ M)	\uparrow epithelial barrier function	TEER, FD4, ⁶ ZO-1, and occludin	\downarrow FD4 (no effect with rutin) \uparrow ZO-1 after quercetin \uparrow occludin after quercetin \uparrow TEER
Priegholdt et al. ⁵⁸	Caco-2	TNF- α	biochanin A (50 μ M) and prunetin (50 μ M)	\downarrow NF- κ B, ERK, ⁷² and tyrosine kinase	TEER, claudin-1, occludin, ZO-1, and E-cadherin	= claudin-1 = ZO-1 = E-cadherin \uparrow occludin \uparrow claudin-1 \uparrow ZO-1 \uparrow ZO-1 = occludin
Park et al. ⁴¹	Caco-2		theaflavin-3'-O-gallate (20 μ M)	\downarrow MLCK	occludin, claudin-1, and ZO-1	= claudin-2 \uparrow TEER (only berberine)
Contreras et al. ⁴²	Caco-2	TNF- α	(-)-epicatechin (0.5–5 μ M)	\downarrow NF- κ B, p-IKK α , ⁴ p-IkB β , and MLCK	occludin, ZO-1, and claudin-2	quercetin (\uparrow claudin-2, claudin-4, and claudin-5 and \downarrow tricellulin) berberin (\downarrow claudin-2 and D-mannitol)
Valenzano et al. ⁴³	Caco-2		berberine (50–200 μ M) quercetin (100–400 μ M)	\uparrow epithelial barrier function	TEER, claudin-1, claudin-2, claudin-3, claudin-4, claudin-5, claudin-7, occludin, tricellulin, and D-mannitol	\uparrow TEER \uparrow occludin \uparrow claudin-3 \uparrow claudin-4 \downarrow FD4 = claudin-1
Ling et al. ⁶⁵	IPEC-J2	deoxynivalenol	resveratrol (0–200 μ M)	\downarrow p38, ERK, and p-JNK ⁷³	TEER, FD4, claudin-1, claudin-3, claudin-4, claudin-7, occludin, and ZO-1	\uparrow TEER \uparrow occludin \uparrow claudin-3 \uparrow claudin-4 \downarrow FD4 = claudin-1
Wang et al. ⁴⁴	Caco-2		PP-rich propolis extract (25 and 50 μ g/mL)	\uparrow AMPK- α , ⁵ ERK1/2, Akt, and p38	ZO-1 and occludin	\uparrow TEER \uparrow occludin \uparrow ZO-1

Table 1. continued

reference	cells	stimulation	PP source and dose	signaling pathway	response/marker	effect
Azzini et al. ⁴⁵	Caco-2		three different PP-rich extracts from chicory (0.2, 1.3, 10, 17, 34, and 70 μ M)	\uparrow epithelial barrier function	TEER	\uparrow TEER
Luescher et al. ³⁸	Caco-2	TNF- α	xanthohumol (chalcone, 10 μ M) and isoxanthohumol (prenylflavone, 10 μ M)	\downarrow TJ permeability	TEER	\uparrow TEER
Cremonini et al. ¹⁶	Caco-2	TNF- α	cyandin, delphinidin, malvidin, petunidin, or peonidin-3-O-glucoside (0.25–1 μ M)	\downarrow IKK and p65 phosphorylation	TEER	\uparrow TEER (only cyanidin, delphinidin, and ACN-rich plant extracts)
		IFN- γ	crowberry extract (1–10 μ g/mL)			
			anthocyanin-rich plant extracts [black chokeberry, black kernel rice, wild blueberry, bilberry, crowberry, domesticated blueberry, and red grape (5 μ g/mL)]			
Rybakovsky et al. ⁴⁷	Caco-2	¹⁴ C-D-mannitol	theaflavins (5–20 μ g/mL) quercetin (100–400 μ M)	\uparrow membrane permeability	claudin-1, claudin-2, claudin-4, and claudin-5	\uparrow TEER (quercetin) \downarrow transepithelial mannitol permeability (quercetin) \uparrow claudin-2 = claudin-1 = claudin-4 = claudin-5
			berberine (50–200 μ M)			\uparrow TEER
Van Buiten et al. ⁴⁸	Caco-2		decaffeinated green tea PPs (0–100 μ g/mL)	\downarrow paracellular permeability	TEER, IL-6, ⁱ and IL-8	\downarrow IL-6 \downarrow IL-8
Li et al. ⁶⁷	MODE-K	LPS ⁶⁴	naringin (50–200 μ M)	\downarrow NF- κ B and MLCK/MLC	TNF- α , IL-10, IL-6, MLCK, p-MLC/MLC, p-p65/p65, and p-I κ B α /I κ B α	\downarrow TNF- α \downarrow IL-10 \downarrow IL-6 \downarrow MLCK \downarrow p-MLC/MLC \downarrow p-p65/p65 \downarrow p-I κ B α /I κ B α
Cremonini et al. ¹⁹	Caco-2	TNF- α	(–)-epicatechin	\uparrow ERK1/2, and AMPK and \downarrow NF- κ B	NOX1/NOX4, ^v FITC ^w –dextran transport, and TEER	\uparrow TEER \downarrow FITC \downarrow NOX1/NOX4 \uparrow improvement of apparent membrane permeability
Vazquez-Olivo et al. ³⁰	Caco-2		four PP-rich mango extracts (100 μ g/mL) gallic acid (100 μ g/mL)	\uparrow membrane permeability	Papp	
Nunes et al. ⁶³	HT-29	TNF- α , IL-1, and IFN- γ	non-alcoholic polyphenolic red wine extract (catechin, oligomeric procyanidins, anthocyanin, phenolic acids, ethyl cinnamate, and condensed tannin), 200, 400, and 600 μ g/mL	\downarrow paracellular permeability	occludin, claudin-5, and ZO-1	\uparrow occludin \uparrow claudin-5 \uparrow ZO-1

^aTEER = transepithelial electrical resistance. ^bZO-1 = zonula occludens 1. ^cIFN- γ = interferon γ . ^dSTAT-1 = signal transducer and activator of transcription 1. ^eMAPK = mitogen-activated protein kinase. ^fMLCK/MLC = myosin light-chain kinase. ^gPKC = protein kinase C. ^hTNF- α = tumor necrosis factor α . ⁱNF- κ B = nuclear factor κ B. ^jPI3K/Akt = phosphoinositide 3-kinase. ^kH₂O₂ = hydrogen peroxide. ^lp38 = p38 pathway. ^mSP-1 = specific protein transcription factor 1. ⁿJAM-A = junctional adhesion molecule A. ^oFD4 = fluorescein isothiocyanate-labeled dextran. ^pERK1/2 = extracellular signal-regulated kinases 1 and 2. ^qp-IKK α = phosphorylated I κ B kinase α . ^rJNK = c-Jun N-terminal kinase. ^sIL-6, IL-8, and IL-10 = interleukins 6, 8, and 10. ^tAMPK = AMP-activated protein kinase. ^uLPS = lipopolysaccharide. ^vNOX = nicotinamide adenine dinucleotide oxidase. ^wFITC = fluorescein isothiocyanate.

Other important factors potentially involved in increasing IP are the multiple protein kinases, such as MAPK, phosphoinositide-3-kinase (PI3K)/Akt, PKC, tyrosine kinase, MLCK, and adenosine monophosphate (AMP)-activated protein kinase (AMPK). Most of them are regulators of fundamental biological processes in epithelial cells, including barrier function, primarily through regulating TJ expression. Some PPs (e.g., quercetin, curcumin, epigallocatechin 3-gallate, and myricetone) have shown to improve the epithelial barrier function through the inhibition of different kinases (PKC and MLCK) involved in phosphorylation of target proteins controlling IP.^{3,30,39}

To ascertain the availability of data supporting the role of PPs on IP, a literature search has been performed using the following terms “intestinal permeability” OR “intestinal barrier” AND “polyphenols” OR “bioactives” OR “phenolics” as keywords in PubMed. The use of the word “polyphenols” as a specific keyword consistently reduced the number of results. On the contrary, a more appropriate search with single PP subclasses AND “intestinal permeability” provided a larger number of *in vitro* and animal studies mainly summarized in Table 1 and 2 and an apparent lack of human intervention studies.

In Vitro Studies. The main lines of evidence on the *in vitro* effects of PPs in the modulation of the potential mediators and regulatory pathways involved in the IP are reported in Table 1. Most of the studies are performed on the Caco-2 cell line,^{38,40–58} as a model of the IB, followed by T84 and HT29/B6 cells (colonic adenocarcinoma cell line),^{59–63} IPEC-J2 cells (intestinal porcine enterocytes), and ECV304 cells (human endothelial cell line).^{64,65} The main evidence of protection is available for berberine, quercetin, and catechin tested in a range of concentrations between 10 and 200 μM (from physiological to pharmacological concentrations). Other PPs tested included genistein, anthocyanins, resveratrol, theaflavin, and a mix of PPs. Most the studies have shown an increase in transepithelial electrical resistance (TEER) across a cellular monolayer, confirming the integrity and functional permeability of the membranes.^{38,43–49,53–55,57,58,62,65,66} In addition, most the PPs tested have shown to increase the expression and/or production of numerous TJ proteins, including ZO-1, occludin, and the family of claudins, whose alteration may result in increased paracellular permeability.^{41,42,44,53,55–57,63,65} Finally, some studies have reported the capacity of PP to counteract the inflammatory process induced by TNF- α and interferon γ (IFN- γ) downregulating the expression of several interleukins, such as IL-8 and IL-6.^{48,67}

Animal Studies. Table 2 reported the effects of PPs and PP-rich extracts in the modulation of IP in animal models.^{44,49,67–74} Most of the studies were performed in healthy rat models (i.e., Wistar rats and Sprague Dawley rats), and IP was induced by stimuli, such high-fat diets, mannitol, inflammatory cytokines, or chemicals.^{44,72,74} Two studies used mice with IL-10 deficiency to test the effect on IP.^{69,70}

The main PPs used were obtained from grape seed extracts (1% GSE; grams of GSE per gram of dry food weight)^{69,70} and grape seed proanthocyanidin extracts (5–50 mg/kg).⁷⁴ Other studies included berberine (200 mg/kg),⁶⁸ (–)-epicatechin (2–20 mg/kg),⁴⁹ and epigallocatechin-3-gallate (about 3 mg/mL).⁷³ Some studies were performed by testing anthocyanin-rich raspberry extract, PP-rich propolis extract, and oregano essential oil.^{44,72} The doses administered ranged from nearly physiological (epicatechin) up to supraphysiological (i.e., berberine). The duration of the intervention varied from a few days (3–10 days) up to several weeks (15–16 weeks).

On the whole, the results obtained support an improvement of IP following the intervention with PPs and PP-rich extracts. In particular, the studies showed the capacity of PPs to upregulate some important genes, such as AMPK and ERK, and downregulate NF- κB as pathways involved in the inflammation process. In line with the observations reported in the *in vitro* studies, the compounds tested have shown to increase the expression of ZO-1, occludin, and several claudins involved in the functioning of TJs.

Human Studies. The number of human intervention studies with IP as primary or secondary outcome increased in the last few years, as also documented by the number of trials made available and reported in public registers (i.e., ISRCTN and ClinicalTrials.gov).

Most of these studies were performed or are ongoing using probiotics, prebiotic fibers, dietary supplements, and sugars. Only four studies seem to have explored the potential beneficial effects of PPs/PP-rich foods on IP in humans (Table 3).^{75–78} The studies differ in terms of population (overweight/obese, cyclists, and older subjects), foods administered (green tea, flavonoid-rich beverage, and mix of PP-rich foods), dose of bioactives (650 mg of flavonoids and 750 mg of PPs), duration of intervention (from 2 weeks up to 8 weeks), and marker of IP selected (endotoxin, lactulose/mannitol ratio, and zonulin levels). The trials are still ongoing, and the results will be useful to increase the understanding on the actual role of PPs and PP-rich foods in humans, where a large number of factors can interact, affecting IP. For example, it is well-recognized that PPs are poorly bioavailable and are biotransformed by gut microbiota into metabolites that can be absorbed in the colon. At the same time, PPs may modulate the composition of the gut microbial community, shaping toward protective symbionts and reducing pathobionts. The complex and not fully elucidated two-way interaction between PPs and gut microbiota is postulated to play a potential direct/indirect role on IP regulation.

In this context, the MaPLE project (Microbiome mAnipulation through Polyphenols for managing gut Leakiness in the Elderly) has been developed with the aim to test the hypothesis that changing the diet of older subjects with established enhanced IP by increasing their PP consumption can alter the intestinal microbial ecosystem (IME) in a way that is beneficial for IB function, resulting in reduced IP and decreased translocation of inflammogenic bacterial factors from the digestive tract into the bloodstream.⁷⁸ To test this hypothesis, a multidisciplinary approach has been used (i) to evaluate the impact of a PP-rich dietary pattern on IB, IP, and IME in a target group of older subjects and (ii) to investigate the possible mechanisms involved in the PP–microbiota–IP interactions through *in vitro* and animal models.

Findings obtained from our and other studies will be “pivotal” for the development of new and advanced hypothesis and experimental approaches in this complex area of research.

■ SOME CONSIDERATIONS ON IP ASSESSMENTS IN DIFFERENT CONTEXTS

IP can be evaluated through numerous methodologies, and consequently, data obtained can differ among studies. The techniques vary depending upon the setting (*in vitro*, *ex vivo*, or *in vivo* models), the models (cells, animals, and humans), and the markers (i.e., ions, macromolecules, bacteria, and bacterial products) but also the compartments (i.e., tissues, blood, and urine). The measurement of IP can be performed through *ex*

Table 2. Summary of the Main Evidence from Animal Models Reporting the Effects of PPs and PP-Rich Extracts in the Modulation of Barrier Integrity and Function

reference	animal model	diet	PP source and dose	signaling pathway	response/marker	main findings
Gu et al. ⁶⁸	male C57BL/6 mice	BBR versus C LPS stimulation	BBR, berberine (200 mg/kg) C, control diet 7 days	↓ MLCK ^{c1}	IP claudin-1 claudin-4 occludin ZO-1 ^b	↑ ZO-1 ↑ occludin ↑ claudin-1 ↑ claudin-4 ↓ IP
Yang et al. ⁶⁹	C57BL/6 (WT) and IL-10 ^{-/-} -deficient (IL-10 ^{-/-} , IL10KO) female mice	GSE versus C dextran sulfate sodium stimulation	GSE, grape seed extract (0 or 1% GSE) ^d C, standard rodent diet 16 weeks	↓ NF-κB ^e	claudin-1 claudin-2	↑ claudin-1 ↓ claudin-2
Wang et al. ⁷⁰	IL10-deficient mice (IL10KO)	GSE versus C dextran sulfate sodium stimulation	GSE, grape seed extract (0 or 1% GSE) ^d C, standard rodent diet 16 weeks	↓ AMPK ^f	claudin-1 claudin-2	↑ claudin-1 ↓ claudin-2
Li et al. ⁷¹	BALB/c mice	ARF versus C dextran sulfate sodium stimulation	ARF, anthocyanin-rich raspberry extract (20 mg/kg) C, saline solution as control treatment 10 days	↓ NF-κB ↓ MAPKs ^g	colonic histological architecture	↑ colonic histological architecture
Wei et al. ⁷²	male Wistar rats	OEO versus C diquat stimulation	OEO, oregano essential oil (5 or 20 mg/kg of BW) C, saline solution as control treatment 14 days	↓ SOD ^h ↓ GSH-Px ⁱ	ZO-1 occludin	↑ ZO-1 ↑ occludin
Wang et al. ⁴⁴	male Sprague Dawley rats	PPE versus C 2,4,6-trinitrobenzenesulfonic acid stimulation	PPE, PP-rich propolis extract (0.3%, w/w) ^d C, control diet 14 days	↑ AMPK ↑ ERK ^j	ZO-1 occludin	↑ ZO-1 ↑ occludin
Bitzer et al. ⁷³	male CF-1 mice	DSS ^k treatment + D (0.5% citric acid) DE (DDS + EGCG) + D (0.5% citric acid) C diet	EGCG, epigallocatechin-3-gallate (3.2 mg/mL) C, control diet 3 days		GLP-2 ^l LAC/RHA ^m	↓ GLP-2 ↓ LAC/RHA
Gil-Cardoso et al. ⁴⁹	female Wistar rats	CAF CAF + GSPE C group	CAF, cafeteria diet ^d CAF + GSPE, (cafeteria diet + grape seed proanthocyanidin extract, 5–50 mg/kg) C, control diet 15 weeks CAF 3 weeks CAF + GSPE		SUC/ERY ⁿ ZO-1 occludin claudin-1 JAM-A ^o	↓ SUC/ERY ↑ ZO-1 ↓ occludin claudin-1 JAM-A ^o
Cremonini et al. ⁴⁹	C57BL/6j mice	HF versus C HFE20 versus CE	CE, (-)-epicatechin (2–20 mg/kg) C, control diet HF, high-fat diet (60% total calories from fat); HFE20, high-fat diet + 20 mg/kg of epicatechin 15 weeks	↑ ERK1/2 ↑ NF-κB (p65) ↑ AMPK	p65 ^p GLP-2 NOXI/NOX4 ^q	↑ p65 (HF) ↑ GLP-2 (CE and HFE20) ↑ NOXI/NOX4 (HF)
Li et al. ⁶⁷	male Kunming mice	CLP ^r + vehicle CLP + NG (30) CLP + NG (60)	NG, naringin (30 and 60 mg/kg) C, no control diet 24–72 h		TEM ^s FITC ^t -dextrane D-lactate	↑ survival CLP + NG (30–60) ↑ IM impairment CLP + vehicle CLP ↑ FITC-dextrane and D-lactate CLP + NG ↓ FITC-dextrane (dose-dependent)

Table 2. continued

^aMLCK/MLC = myosin light-chain kinase. ^bZO-1 = zonula occludens 1. ^cIL = interleukin. ^dData on PP characterization were not provided. ^eNF- κ B = nuclear factor κ B. ^fAMPK = AMP-activated protein kinase. ^gMAPK = mitogen-activated protein kinase. ^hSOD = superoxide dismutase. ⁱGSH-Px = glutathione peroxidase. ^jERK1/2 = extracellular signal-regulated kinases 1 and 2. ^kDSS = dextran sulfate sodium. ^lGLP-2 = glucagon-like peptide 2. ^mLAC/RHA = lactulose/rhamnose ratio. ⁿSUC/ERY = sucralose/erythritol ratio. ^oJAM = junctional adhesion molecule. ^pp65 = transcription factor p65. ^qNOX1/NOX 4 = NADPH oxidases. ^rCLP = cecal ligation and puncture. ^sTEM = transmission electron microscopy. ^tFITC = fluorescein isothiocyanate.

vivo and *in vivo* approaches.⁷⁹ An example of an *ex vivo* approach includes the use of an Ussing chamber able to measure the transport of ions and molecules (i.e., nutrients and drugs) across various epithelial tissues using fresh intestinal tissue. *In vivo*, the assessment of IP can be performed through permeability assays (i.e., evaluation of the ratio of lactulose/mannitol, sucralose, sucrose, polyethylene glycols, or ⁵²Cr-EDTA in urine), analysis of bacterial-related markers (i.e., endotoxin test, EndoCAB, D-lactate, and butyrate production), markers of epithelial damage (i.e., citrullin, fatty-acid-binding protein, and claudin-3), and/or other related markers (i.e., fecal calprotectin). Finally, histological approaches measuring, for example, goblet cell analysis, shedding of the epithelium, or Paneth cell loss, can be performed.²

On the whole, on the basis of revised literature, it can be assumed that current *in vitro* permeability models are still far from reflecting an *in vivo* situation. This limits the relevance of data obtained within the cell culture and the possibility to transfer the results to humans. In fact, the comparison between *in vitro* and *in vivo* permeability data is difficult and dependent upon numerous factors, including the type of cells used, the molecule under study, the transport route evaluated, and the method used for the assessment of IB function and permeability [i.e., mainly transepithelial electrical resistance (TEER) and biomarkers of epithelial integrity], which can significantly affect the results obtained, making it difficult to identify the best approach.

A novel biomarker of IP *in vivo* is zonulin, a protein secreted by enterocytes but also from other type of cells (i.e., epithelial cells), known to be a physiological modulator and, thus, to control IP reversibly via intercellular TJs.⁸⁰ Increased zonulin serum levels have been observed in many gut-related diseases, and emerging evidence suggests an increased zonulin level in specific subjects (e.g., older persons)⁸¹ and in different diseases or conditions (e.g., diabetes and obesity).^{82,83} The reliability and accuracy of the different markers to assess IP is clearly a fundamental part of the recent discussion and a hot topic considering the increasing demand for non-invasive diagnosis tools.⁸⁴ In this regard, the concurrent evaluation of different markers of IP seems highly recommendable to improve reliability of findings on IB function.

CONCLUSIONS AND FUTURE PERSPECTIVES

There is increasing demand for non-invasive strategies able to modulate critical regulatory functions for human health, such as IP, which can play a role in the pathogenesis of intestinal and systemic diseases. The improvement or manipulation of the diet, for example, increasing or reducing specific nutrients and/or including food bioactives, such as PPs, is recognized as a potential powerful tool to be explored also in the context of IP. From data available, PP activity seems to be plausibly a consequence of multiple mechanisms, which may also depend upon the type and amount of compounds considered. The results from *in vitro* studies have shown the capacity of PPs to increase the expression and/or production of numerous TJ proteins and to reduce the release of several interleukins/cytokines. These results are partially in line with the findings obtained in the animal models, showing the capacity of PPs to up-/downregulate some important genes involved in the inflammatory process. With regard to human studies, recent literature suggests that PPs may modulate IP through a number of direct and indirect effects, including the impact on the intestinal ecosystem and immune system. This type of research

Table 3. Summary of the Ongoing Human Studies Evaluating the Effect of PPs and PP-Rich Foods on IP

title	source	subject number/characteristics and inclusion criteria	study design	intervention	duration of intervention	markers under study
dietary green tea confederation for resolving gut permeability-induced metabolic endotoxemia in obese adults	ClinicalTrials.gov NCT03413735	40 overweight/obese (BMI = 28–40 kg/m ²) fasting glucose of <126 mg/dL	randomized parallel design	test group, green tea extract (GTE)-rich confection placebo group, no GTE-rich confection	4 weeks	primary outcome: Endotoxin secondary outcome, gut permeability (lactulose/mannitol ratio and sucralose/erythritol ratio) microbiota Firmicutes/Bacteroidetes ratio)
effect of flavonoids on gut permeability in cyclists	ClinicalTrials.gov NCT03427879	normotensive (blood pressure of <140/90 mmHg) non-dietary supplement user non-smoker 22 male or female of any race or ethnicity between 18 and 49 years of age competed in a road race or triathlon in past 12 months	randomized crossover design	dose, daily (no information about the amount provided in terms of PPs) test group, a high-flavonoid, sports nutrition recovery beverage will be prepared from milk (78%), sugar (8.6%), maltodextrin (8.6%), blueberry powder (2.4%), cocoa powder (1.6%), green tea extract (0.1%), and whey protein isolate (0.6%) containing approximately 620 mg of flavonoids per serving placebo group, a low-flavonoid, sports nutrition recovery beverage will be prepared from milk (78%), sugar (8.6%), maltodextrin (8.6%), blueberry powder (2.4%), alkalized cocoa powder (1.6%), and whey protein isolate (0.6%), containing approximately 5 mg of flavonoids per serving	2 weeks	plasma intestinal fatty-acid-binding protein secondary outcome, fecal calprotectin urinary sucralose/mannitol ratio inflammatory markers (TNF- α and IL-10) ^c endotoxin other variables related to exercise performance
effect of dietary flavonoids on intestinal microbiota, intestinal inflammation and metabolic syndrome	ClinicalTrials.gov NCT02728570	free of chronic disease and gut inflammation conditions train at least 3 times per week, 1 h at a time on average willing to prepare and consume provided pre-workout beverage daily maintain weight (no more/less than 5 kg change) willing to avoid consumption of high flavonoid foods/supplements, large dose vitamin and mineral supplements, and NSAIDs ^d or other medications known to affect inflammation during the study period	randomized crossover design	test group, prepared diet with high levels of dietary flavonoids (340 mg of flavonoids/1000 kcal) with a macronutrient composition of 17% energy from protein, 30% energy from fat, and 53% energy from carbohydrate control group, prepared diet with high levels of dietary flavonoids (10 mg of flavonoids/1000 kcal) with a macronutrient composition of 17% energy from protein, 30% energy from fat, and 53% energy from carbohydrate	6 weeks	primary outcome, fecal calprotectin serum PCR ^e serum TNF- α serum insulin secondary outcome, fecal microbiome composition, SCFAs, eosinophil protein X, and myeloperoxidase IP by four sugar differential absorption tests serum endotoxin, IL-6, soluble TNF- α , and fasting glucose

Table 3. continued

title	source	subject number/characteristics and inclusion criteria	study design	intervention	duration of intervention	markers under study
effect of a polyphenol-rich diet on leaky gut in the elderly	ISRCTN registry ISRCTN10214981	60 healthy older subjects age of >60 years old	randomized crossover design	test group, habitual diet + PP-rich products (berries and derived products, blood oranges and derived products, pomegranate juice, Renetta apple and purée, green tea, and dark chocolate products) control group, comparable diet without the PP-rich products	8 weeks	calculated homeostatic model assessment for insulin resistance serum C peptide plasma lipid profile blood pressure other outcome measures, serum resistin, visfatin, adiponectin, and leptin body weight primary outcome, zonulin serum levels
	78	IP evaluated by zonulin serum level adequate nutritional status evaluated with a mini nutritional assessment (MNA) score of ≥ 24 good cognitive status tested with mini mental state evaluation (MMSE) score of ≥ 24 self-sufficiency assessed with validated tests (e.g., Barthel index for activities of daily living and Tinetti balance assessment)		dose, three portions of PP-rich food products daily (about 750 mg of PP _s)		secondary outcome, total blood bacterial load fecal microbiota composition and metabolism SCFAs and PP-derived metabolites inflammatory, oxidative stress, and related markers endotoxin LPS-BPs ^g metabolic markers metabolic and anthropometric markers

^aNSAID = non-steroidal anti-inflammatory drug. ^bTNF- α = tumor necrosis factor α . ^cIL-10 = interleukin 10. ^dBMI = body mass index. ^ePCR = C-reactive protein. ^fTNFR-2 = tumor necrosis factor receptor 2. ^gLPS-BP = lipopolysaccharide-binding protein.

is still in its infancy by considering the few human studies available. Future research should be targeted to identify the PPs and/or their metabolites eventually involved in the modulation of IP while also demonstrating their specific dose-dependent mechanisms of action. Meanwhile, *in vivo* studies should be performed to increase understanding of the diet–microbiota–IP axis possibly through the development of well-controlled dietary intervention studies. Finally, by considering the wide discussion in the literature on IP evaluation, further effort is needed to better define the reliability of the already available IP biomarkers and the potential exploitation of new and/or improved candidate biomarkers.

AUTHOR INFORMATION

Corresponding Author

*Telephone: +39-02-503-16726. Fax: +39-02-503-16721. E-mail: patrizia.riso@unimi.it.

ORCID

Cristina Andrés-Lacueva: 0000-0002-8494-4978

Gregorio Peron: 0000-0002-6007-6184

Paul Kroon: 0000-0002-9805-6947

Simone Guglielmetti: 0000-0002-8673-8190

Patrizia Riso: 0000-0002-9204-7257

Author Contributions

†Stefano Bernardi and Cristian Del Bo' contributed equally to this work.

Funding

This work was developed as part of the MAPLE project (Gut and Blood Microbiomics for Studying the Effect of a Polyphenol-Rich Dietary Pattern on Intestinal Permeability in the Elderly) supported within the European Joint Programming Initiative “A Healthy Diet for a Healthy Life” (JPI HDHL, <http://www.healthydietforhealthylife.eu/>) granted by Mipaaf (Italy, D.M. 8245/7303/2016), MINECO (Spain, PCIN-2015-238), and BBSRC (U.K., BB/R012512/1). Cristina Andres-Lacueva thanks 2017SGR1546 from the Generalitat de Catalunya's Agency AGAUR, funds from CIBERFES (co-funded by the FEDER Program from the European Union), and ICREA Academia Award 2018.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

European Cooperation for Science and Technology (COST Action) CA16112 “NutRedOx: Personalized Nutrition in Aging Society: Redox Control of Major Age-Related Diseases” is acknowledged.

ABBREVIATIONS USED

IP, intestinal permeability; IB, intestinal barrier; IME, intestinal microbial ecosystem; TJ, tight junction; GJ, gap junction; AJ, adherens junction; JAM, junctional adhesion molecule; ZO, zonula occludens; MLCK, myosin light-chain kinase; PKC, protein kinase C; MAPK, mitogen-activated protein kinase; TLR4, toll-like receptor 4; NAFLD, non-alcoholic fatty liver disease; MS, multiple sclerosis; CNS, central nervous system; TNF- α , tumor necrosis factor α ; MD, Mediterranean diet; SCFA, short-chain fatty acid; PP, polyphenol; NF- κ B, nuclear factor κ B; Nrf-2, nuclear factor erythroid 2-related factor 2; IL, interleukin; HO-1, heme oxygenase 1; SOD, superoxide dismutase; GPx, glutathione peroxidase; DNA, deoxyribonu-

cleic acid; IKK, I κ B kinase; PI3K, phosphoinositide-3-kinase; AMPK, AMP-activated protein kinase; TEER, transepithelial electrical resistance; INF- γ , interferon γ ; ERK, extracellular regulated kinase; MAPLE, microbiome manipulation through polyphenols for managing gut leakiness in the elderly

REFERENCES

- (1) Brandtzaeg, P. The Gut as Communicator between Environment and Host: Immunological Consequences. *Eur. J. Pharmacol.* **2011**, *668* (Suppl), S16–S32.
- (2) Bischoff, S. C.; Barbara, G.; Buurman, W.; Ockhuizen, T.; Schulzke, J. D.; Serino, M.; Tilg, H.; Watson, A.; Wells, J. M. Intestinal Permeability—A New Target for Disease Prevention and Therapy. *BMC Gastroenterol.* **2014**, *14* (1), 1–25.
- (3) Suzuki, T. Regulation of Intestinal Epithelial Permeability by Tight Junctions. *Cell. Mol. Life Sci.* **2013**, *70* (4), 631–659.
- (4) Lee, S. H. Intestinal Permeability Regulation by Tight Junction: Implication on Inflammatory Bowel Diseases. *Intest. Res.* **2015**, *13* (1), 11.
- (5) Furuse, M.; Fujita, K.; Hiiiragi, T.; Fujimoto, K.; Tsukita, S. Claudin-1 and -2: Novel Integral Membrane Proteins Localizing at Tight Junctions with No Sequence Similarity to Occludin. *J. Cell Biol.* **1998**, *141* (7), 1539–1550.
- (6) Furuse, M.; Hirase, T.; Itoh, M.; Nagafuchi, A.; Yonemura, S.; Tsukita, S.; Tsukita, S. Occludin: A Novel Integral Membrane Protein Localizing at Tight Junctions. *J. Cell Biol.* **1993**, *123* (6), 1777–1788.
- (7) Ulluwishewa, D.; Anderson, R. C.; McNabb, W. C.; Moughan, P. J.; Wells, J. M.; Roy, N. C. Regulation of Tight Junction Permeability by Intestinal Bacteria and Dietary Components. *J. Nutr.* **2011**, *141*, 769–776.
- (8) Oda, H.; Takeichi, M. Evolution: Structural and Functional Diversity of Cadherin at the Adherens Junction. *J. Cell Biol.* **2011**, *193* (7), 1137–1146.
- (9) Hollon, J.; Puppa, E. L.; Greenwald, B.; Goldberg, E.; Guerrero, A.; Fasano, A. Effect of Gliadin on Permeability of Intestinal Biopsy Explants from Celiac Disease Patients and Patients with Non-Celiac Gluten Sensitivity. *Nutrients* **2015**, *7* (3), 1565–1576.
- (10) Dunlop, S. P.; Hebden, J.; Campbell, E.; Naesdal, J.; Olbe, L.; Perkins, A. C.; Spiller, R. C. Abnormal Intestinal Permeability in Subgroups of Diarrhea-Predominant Irritable Bowel Syndromes. *Am. J. Gastroenterol.* **2006**, *101* (6), 1288–1294.
- (11) Melichar, B.; Zezulova, M. The Significance of Altered Gastrointestinal Permeability in Cancer Patients. *Curr. Opin. Support. Palliat. Care* **2011**, *5* (1), 47–54.
- (12) Vancamelbeke, M.; Vermeire, S. The Intestinal Barrier: A Fundamental Role in Health and Disease. *Expert Rev. Gastroenterol. Hepatol.* **2017**, *11* (9), 821–834.
- (13) Demir, E.; Ozkan, H.; Seckin, K. D.; Sahtiyanci, B.; Demir, B.; Tabak, O.; Kumbasar, A.; Uzun, H. Plasma Zonulin Levels as a Non-Invasive Biomarker of Intestinal Permeability in Women with Gestational Diabetes Mellitus. *Biomolecules* **2019**, *9* (1), 24.
- (14) Lerner, A.; Matthias, T. Rheumatoid Arthritis-Celiac Disease Relationship: Joints Get That Gut Feeling. *Autoimmun. Rev.* **2015**, *14* (11), 1038–1047.
- (15) Spielman, L. J.; Gibson, D. L.; Klegeris, A. Unhealthy Gut, Unhealthy Brain: The Role of the Intestinal Microbiota in Neurodegenerative Diseases. *Neurochem. Int.* **2018**, *120*, 149–163.
- (16) Buscarinu, M. C.; Romano, S.; Mechelli, R.; Pizzolato Umeton, R.; Ferraldeschi, M.; Fornasiero, A.; Renie, R.; Cerasoli, B.; Morena, E.; Romano, C.; Loizzo, N. D.; Umeton, R.; Salvetti, M.; Ristori, G. Intestinal Permeability in Relapsing-Remitting Multiple Sclerosis. *Neurotherapeutics* **2018**, *15* (1), 68–74.
- (17) Franceschi, C.; Bonafe, M.; Valensin, S.; Olivieri, F.; De Luca, M.; Ottaviani, E.; De Benedictis, G. Inflamm-Aging: An Evolutionary Perspective on Immunosenesescence. *Ann. N. Y. Acad. Sci.* **2000**, *908*, 244–254.

- (18) Manco, M.; Putignani, L.; Bottazzo, G. F. Gut Microbiota, Lipopolysaccharides, and Innate Immunity in the Pathogenesis of Obesity and Cardiovascular Risk. *Endocr. Rev.* **2010**, *31* (6), 817–844.
- (19) Malago, J. Contribution of Microbiota to the Intestinal Physicochemical Barrier. *Benefic. Microbes* **2015**, *6* (3), 295–311.
- (20) Norman, K.; Pirlich, M.; Schulzke, J.; Smoliner, C.; Lochs, H.; Valentini, L.; Bühner, S. Increased Intestinal Permeability in Malnourished Patients with Liver Cirrhosis. *Eur. J. Clin. Nutr.* **2012**, *66* (10), 1116.
- (21) Guerreiro, C. S.; Calado, Â.; Sousa, J.; Fonseca, J. E. Diet, Microbiota, and Gut Permeability—The Unknown Triad in Rheumatoid Arthritis. *Front. Med.* **2018**, *5*, 1–7.
- (22) De Filippo, C.; Cavalieri, D.; Di Paola, M.; Ramazzotti, M.; Poullet, J.; Massart, S.; Collini, S.; Pieraccini, G.; Lionetti, P. Impact of Diet in Shaping Gut Microbiota Revealed by a Comparative Study in Children from Europe and Rural Africa. *Proc. Natl. Acad. Sci. U. S. A.* **2010**, *107* (33), 14691–14696.
- (23) Plöger, S.; Stumpff, F.; Penner, G.; Schulzke, J.; Gäbel, G.; Martens, H.; Shen, Z.; Günzel, D.; Aschenbach, J. Microbial Butyrate and Its Role for Barrier Function in the Gastrointestinal Tract. *Ann. N. Y. Acad. Sci.* **2012**, *1258*, 52–59.
- (24) Wang, H. B.; Wang, P. Y.; Wang, X.; Wan, Y. L.; Liu, Y. C. Butyrate Enhances Intestinal Epithelial Barrier Function via Up-Regulation of Tight Junction Protein Claudin-1 Transcription. *Dig. Dis. Sci.* **2012**, *57* (12), 3126–3135.
- (25) Mrduljaš, N.; Krešić, G.; Bilušić, T. Polyphenols: Food Sources and Health Benefits. In *Functional Food—Improve Health through Adequate Food*; Hueda, M. C., Ed.; IntechOpen, Limited: London, U.K., 2017; Chapter 2, DOI: 10.5772/intechopen.68862.
- (26) Manach, C.; Scalbert, A.; Morand, C.; Rémésy, C.; Jiménez, L. Polyphenols: Food Sources and Bioavailability. *Am. J. Clin. Nutr.* **2004**, *79* (5), 727–747.
- (27) Cardona, F.; Andrés-Lacueva, C.; Tulipani, S.; Tinahones, F. J.; Queipo-Ortuño, M. I. Tinahones; María Isabel Queipo-Ortuño. Benefits of Polyphenols on Gut Microbiota and Implications in Human Health. *J. Nutr. Biochem.* **2013**, *24* (8), 1415–1422.
- (28) Monagas, M.; Urpi-Sarda, M.; Sánchez-Patán, F.; Llorach, R.; Garrido, I.; Gómez-Cordovés, C.; Andres-Lacueva, C.; Bartolomé, B. Insights into the Metabolism and Microbial Biotransformation of Dietary Flavan-3-ols and the Bioactivity of Their Metabolites. *Food Funct.* **2010**, *1* (3), 233–253.
- (29) Joseph, S. V.; Edirisinghe, I.; Burton-Freeman, B. M. Fruit Polyphenols: A Review of Anti-Inflammatory Effects in Humans. *Crit. Rev. Food Sci. Nutr.* **2016**, *56* (3), 419–444.
- (30) Kaulmann, A.; Bohn, T. Bioactivity of Polyphenols: Preventive and Adjuvant Strategies toward Reducing Inflammatory Bowel Diseases—Promises, Perspectives, and Pitfalls. *Oxid. Med. Cell. Longevity* **2016**, *2016* (c), 1–29.
- (31) Gil-Cardoso, K.; Ginés, I.; Pinent, M.; Ardévol, A.; Blay, M.; Terra, X. Effects of Flavonoids on Intestinal Inflammation, Barrier Integrity and Changes in Gut Microbiota during Diet-Induced Obesity. *Nutr. Res. Rev.* **2016**, *29* (2), 234–248.
- (32) Shimizu, M. Multifunctions of Dietary Polyphenols in the Regulation of Intestinal Inflammation. *J. Food Drug Anal.* **2017**, *25* (1), 93–99.
- (33) Mosele, J. I.; Macià, A.; Motilva, M. J. Metabolic and Microbial Modulation of the Large Intestine Ecosystem by Non-Absorbed Diet Phenolic Compounds: A Review. *Molecules* **2015**, *20* (9), 17429–17468.
- (34) Boto-Ordóñez, M.; Urpi-Sarda, M.; Queipo-Ortuno, M. I.; Tulipani, S.; Tinahones, F. J.; Andres-Lacueva, C. High Levels of Bifidobacteria Are Associated with Increased Levels of Anthocyanin Microbial Metabolites: A Randomized Clinical Trial. *Food Funct.* **2014**, *5* (8), 1932–1938.
- (35) Queipo-Ortuno, M. I.; Boto-Ordóñez, M.; Murri, M.; Gomez-Zumaquero, J. M.; Clemente-Postigo, M.; Estruch, R.; Cardona Diaz, F.; Andres-Lacueva, C.; Tinahones, F. J. Influence of Red Wine Polyphenols and Ethanol on the Gut Microbiota Ecology and Biochemical Biomarkers. *Am. J. Clin. Nutr.* **2012**, *95* (6), 1323–1334.
- (36) Guglielmetti, S.; Fracassetto, D.; Taverniti, V.; Del Bò, C.; Vendrame, S.; Klimis-Zacas, D.; Arioli, S.; Riso, P.; Porrini, M. Differential Modulation of Human Intestinal Bifidobacterium Populations after Consumption of a Wild Blueberry (*Vaccinium angustifolium*) Drink. *J. Agric. Food Chem.* **2013**, *61* (34), 8134–8140.
- (37) Edwards, C. A.; Havlik, J.; Cong, W.; Mullen, W.; Preston, T.; Morrison, D. J.; Combet, E. Polyphenols and Health: Interactions between Fibre, Plant Polyphenols and the Gut Microbiota. *Nutr. Bull.* **2017**, *42* (4), 356–360.
- (38) Luescher, S.; Urmann, C.; Butterweck, V. Effect of Hops Derived Prenylated Phenols on TNF- α Induced Barrier Dysfunction in Intestinal Epithelial Cells. *J. Nat. Prod.* **2017**, *80* (4), 925–931.
- (39) Das, J.; Ramani, R.; Suraju, M. O. Polyphenol Compounds and PKC Signaling. *Biochim. Biophys. Acta* **2016**, *1860* (10), 2107–2121.
- (40) Atkinson, K. J.; Rao, R. K. Role of Protein Tyrosine Phosphorylation in Acetaldehyde-Induced Disruption of Epithelial Tight Junctions. *Am. J. Physiol. Liver Physiol.* **2001**, *280* (6), G1280–G1288.
- (41) Park, H.-Y.; Kunitake, Y.; Matsui, T.; Hirasaki, N.; Tanaka, M. Theaflavins Enhance Intestinal Barrier of Caco-2 Cell Monolayers through the Expression of AMP-Activated Protein Kinase-Mediated Occludin, Claudin-1, and ZO-1. *Biosci., Biotechnol., Biochem.* **2015**, *79* (1), 130–137.
- (42) Contreras, T. C.; Ricciardi, E.; Cremonini, E.; Oteiza, P. I. (–)-Epicatechin in the Prevention of Tumor Necrosis Alpha-Induced Loss of Caco-2 Cell Barrier Integrity. *Arch. Biochem. Biophys.* **2015**, *573*, 84–91.
- (43) Valenzano, M. C.; DiGuilio, K.; Mercado, J.; Teter, M.; To, J.; Ferraro, B.; Mixson, B.; Manley, I.; Baker, V.; Moore, B. A.; Wertheimer, J.; Mullin, J. M. Remodeling of Tight Junctions and Enhancement of Barrier Integrity of the CACO-2 Intestinal Epithelial Cell Layer by Micronutrients. *PLoS One* **2015**, *10* (7), No. e0133926.
- (44) Wang, K.; Hu, F.; Conlon, M.; Song, Z.; Jin, X.; Jiang, X.; Chen, Y.; Topping, D. Polyphenol-Rich Propolis Extracts Strengthen Intestinal Barrier Function by Activating AMPK and ERK Signaling. *Nutrients* **2016**, *8* (5), 272.
- (45) Azzini, E.; Maiani, G.; Garaguso, I.; Polito, A.; Foddai, M. S.; Venneria, E.; Durazzo, A.; Intorre, F.; Palomba, L.; Rauseo, M. L.; Lombardi-Boccia, G.; Nobili, F. The Potential Health Benefits of Polyphenol-Rich Extracts from *Cichorium Intybus* L. Studied on Caco-2 Cells Model. *Oxid. Med. Cell. Longevity* **2016**, *2016*, 1–9.
- (46) Cremonini, E.; Mastaloudis, A.; Hester, S. N.; Verstraeten, S. V.; Anderson, M.; Wood, S. M.; Waterhouse, A. L.; Fraga, C. G.; Oteiza, P. I. Anthocyanins Inhibit Tumor Necrosis Alpha-Induced Loss of Caco-2 Cell Barrier Integrity. *Food Funct.* **2017**, *8* (8), 2915–2923.
- (47) Rybakovsky, E.; Valenzano, M. C.; Deis, R.; DiGuilio, K. M.; Thomas, S.; Mullin, J. M. Improvement of Human-Oral-Epithelial-Barrier Function and of Tight Junctions by Micronutrients. *J. Agric. Food Chem.* **2017**, *65* (50), 10950–10958.
- (48) Van Buiten, C. B.; Lambert, J. D.; Elias, R. J. Green Tea Polyphenols Mitigate Gliadin-Mediated Inflammation and Permeability in Vitro. *Mol. Nutr. Food Res.* **2018**, *62* (12), 1700879.
- (49) Cremonini, E.; Wang, Z.; Bettaieb, A.; Adamo, A. M.; Daveri, E.; Mills, D. A.; Kalanetra, K. M.; Haj, F. G.; Karakas, S.; Oteiza, P. I. (–)-Epicatechin Protects the Intestinal Barrier from High Fat Diet-Induced Permeabilization: Implications for Steatosis and Insulin Resistance. *Redox Biol.* **2018**, *14*, 588–599.
- (50) Vazquez-Olivo, G.; Antunes-Ricardo, M.; Gutiérrez-Urbe, J. A.; Osuna-Enciso, T.; León-Félix, J.; Heredia, J. B. Cellular Antioxidant Activity and in Vitro Intestinal Permeability of Phenolic Compounds from Four Varieties of Mango Bark (*Mangifera indica* L.). *J. Sci. Food Agric.* **2019**, *99* (7), 3481–3489.
- (51) Amasheh, M.; Schlichter, S.; Amasheh, S.; Mankertz, J.; Zeitz, M.; Fromm, M.; Schulzke, J. D. Quercetin Enhances Epithelial Barrier Function and Increases Claudin-4 Expression in Caco-2 Cells. *J. Nutr.* **2008**, *138* (6), 1067–1073.
- (52) Suzuki, T.; Hara, H. Quercetin Enhances Intestinal Barrier Function through the Assembly of Zonula Occludens-2, Occludin,

and Claudin-1 and the Expression of Claudin-4 in Caco-2 Cells. *J. Nutr.* **2009**, *139* (5), 965–974.

(53) Suzuki, T.; Tanabe, S.; Hara, H. Kaempferol Enhances Intestinal Barrier Function through the Cytoskeletal Association and Expression of Tight Junction Proteins in Caco-2 Cells. *J. Nutr.* **2011**, *141* (1), 87–94.

(54) Noda, S.; Tanabe, S.; Suzuki, T. Differential Effects of Flavonoids on Barrier Integrity in Human Intestinal Caco-2 Cells. *J. Agric. Food Chem.* **2012**, *60* (18), 4628–4633.

(55) Noda, S.; Tanabe, S.; Suzuki, T. Naringenin Enhances Intestinal Barrier Function through the Expression and Cytoskeletal Association of Tight Junction Proteins in Caco-2 Cells. *Mol. Nutr. Food Res.* **2013**, *57* (11), 2019–2028.

(56) Cao, M.; Sun, C.; He, W.; Wang, P.; Wang, F. Amelioration of IFN- γ and TNF- α -Induced Intestinal Epithelial Barrier Dysfunction by Berberine via Suppression of MLCK-MLC Phosphorylation Signaling Pathway. *PLoS One* **2013**, *8* (5), No. e61944.

(57) Carrasco-Pozo, C.; Morales, P.; Gotteland, M. Polyphenols Protect the Epithelial Barrier Function of Caco-2 Cells Exposed to Indomethacin through the Modulation of Occludin and Zonula Occludens-1 Expression. *J. Agric. Food Chem.* **2013**, *61* (22), S291–S297.

(58) Piegholdt, S.; Pallauf, K.; Esatbeyoglu, T.; Speck, N.; Reiss, K.; Ruddigkeit, L.; Stocker, A.; Huebbe, P.; Rimbach, G. Biochanin A and Prunetin Improve Epithelial Barrier Function in Intestinal CaCo-2 Cells via Downregulation of ERK, NF-KB, and Tyrosine Phosphorylation. *Free Radical Biol. Med.* **2014**, *70*, 255–264.

(59) Watson, J. L.; Wang, A.; Ansari, S.; Cameron, H.; McKay, D. M.; Akhtar, M. Green Tea Polyphenol (–)-Epigallocatechin Gallate Blocks Epithelial Barrier Dysfunction Provoked by IFN- γ but Not by IL-4. *Am. J. Physiol. Liver Physiol.* **2004**, *287* (5), G954–G961.

(60) Krug, S. M.; Fromm, M.; Zeitz, M.; Schulzke, J.-D.; Amasheh, S.; Fromm, A.; Amasheh, M.; Andres, S. TNF α -Induced and Berberine-Antagonized Tight Junction Barrier Impairment via Tyrosine Kinase, Akt and NF κ B Signaling. *J. Cell Sci.* **2010**, *123* (23), 4145–4155.

(61) Rogoll, D.; Bergmann, H.; Hellenschmidt, D.; Heinze, J.; Scheppach, W.; Melcher, R.; Richling, E. Influence of Apple Polyphenols on the Intestinal Barrier in a Colonic Cell Model. *J. Appl. Bot. Food Qual.* **2010**, *83*, 110–117.

(62) Amasheh, M.; Fromm, M.; Luettig, J.; Zeitz, M.; Schulzke, J.-D.; Amasheh, S. Effects of Quercetin Studied in Colonic HT-29/B6 Cells and Rat Intestine in Vitro. *Ann. N. Y. Acad. Sci.* **2012**, *1258* (1), 100–107.

(63) Nunes, C.; Freitas, V.; Almeida, L.; Laranjinha, J. Red Wine Extract Preserves Tight Junctions in Intestinal Epithelial Cells under Inflammatory Conditions: Implications for Intestinal Inflammation. *Food Funct.* **2019**, *10*, 1364–1374.

(64) Chuenkityanon, S.; Pengsuparp, T.; Jianmongkol, S. Protective Effect of Quercetin on Hydrogen Peroxide-Induced Tight Junction Disruption. *Int. J. Toxicol.* **2010**, *29* (4), 418–424.

(65) Ling, K. H.; Wan, M. L. Y.; El-Nezami, H.; Wang, M. Protective Capacity of Resveratrol, a Natural Polyphenolic Compound, against Deoxynivalenol-Induced Intestinal Barrier Dysfunction and Bacterial Translocation. *Chem. Res. Toxicol.* **2016**, *29* (5), 823–833.

(66) Shin, D. Y.; Lu, J. N.; Kim, G. Y.; Jung, J. M.; Kang, H. S.; Lee, W. S.; Choi, Y. H. Anti-Invasive Activities of Anthocyanins through Modulation of Tight Junctions and Suppression of Matrix Metalloproteinase Activities in HCT-116 Human Colon Carcinoma Cells. *Oncol. Rep.* **2011**, *25* (2), 567–572.

(67) Li, Z.; Gao, M.; Yang, B.; Zhang, H.; Wang, K.; Liu, Z.; Xiao, X.; Yang, M. Naringin Attenuates MLC Phosphorylation and NF-KB Activation to Protect Sepsis-Induced Intestinal Injury via RhoA/ROCK Pathway. *Biomed. Pharmacother.* **2018**, *103*, 50–58.

(68) Gu, L.; Li, N.; Gong, J.; Li, Q.; Zhu, W.; Li, J. Berberine Ameliorates Intestinal Epithelial Tight-Junction Damage and down-Regulates Myosin Light Chain Kinase Pathways in a Mouse Model of Endotoxemia. *J. Infect. Dis.* **2011**, *203* (11), 1602–1612.

(69) Yang, G.; Wang, H.; Kang, Y.; Zhu, M. J. Grape Seed Extract Improves Epithelial Structure and Suppresses Inflammation in Ileum of IL-10-Deficient Mice. *Food Funct.* **2014**, *5* (10), 2558–2563.

(70) Wang, H.; Xue, Y.; Zhang, H.; Huang, Y.; Yang, G.; Du, M.; Zhu, M.-J. Dietary Grape Seed Extract Ameliorates Symptoms of Inflammatory Bowel Disease in Interleukin-10 Deficient Mice. *Mol. Nutr. Food Res.* **2013**, *57* (12), 2253–2257.

(71) Li, L.; Wang, L.; Wu, Z.; Yao, L.; Wu, Y.; Huang, L.; Liu, K.; Zhou, X.; Gou, D. Anthocyanin-Rich Fractions from Red Raspberries Attenuate Inflammation in Both RAW264.7 Macrophages and a Mouse Model of Colitis. *Sci. Rep.* **2015**, *4*, 1–11.

(72) Wei, H.-k.; Chen, G.; Wang, R.-J.; Peng, J. Oregano Essential Oil Decreased Susceptibility to Oxidative Stress-Induced Dysfunction of Intestinal Epithelial Barrier in Rats. *J. Funct. Foods* **2015**, *18*, 1191–1199.

(73) Bitzer, Z. T.; Elias, R. J.; Vijay-Kumar, M.; Lambert, J. D. (–)-Epigallocatechin-3-Gallate Decreases Colonic Inflammation and Permeability in a Mouse Model of Colitis, but Reduces Macronutrient Digestion and Exacerbates Weight Loss. *Mol. Nutr. Food Res.* **2016**, *60* (10), 2267–2274.

(74) Gil-Cardoso, K.; Ginés, I.; Pinent, M.; Ardévol, A.; Arola, L.; Blay, M.; Terra, X. Chronic Supplementation with Dietary Proanthocyanidins Protects from Diet-Induced Intestinal Alterations in Obese Rats. *Mol. Nutr. Food Res.* **2017**, *61* (8), 1601039.

(75) United States National Library of Medicine (NLM). *Dietary Green Tea Confection for Resolving Gut Permeability-Induced Metabolic Endotoxemia in Obese Adults*; NLM: Bethesda, MD, 2018; <https://clinicaltrials.gov/ct2/show/NCT03413735>.

(76) United States National Library of Medicine (NLM). *Effect of Flavonoids on Gut Permeability in Cyclists*; NLM: Bethesda, MD, 2018; <https://clinicaltrials.gov/ct2/show/NCT03427879>.

(77) United States National Library of Medicine (NLM). *Effect of Dietary Flavonoids on Intestinal Microbiota, Intestinal Inflammation and Metabolic Syndrome*; NLM: Bethesda, MD, 2016; <https://clinicaltrials.gov/ct2/show/NCT02728570>.

(78) BioMed Central (BMC). *Effect of a Polyphenol-Rich Diet on Leaky Gut in the Elderly*; BMC: London, U.K., 2017; <http://www.isrctn.com/ISRCTN10214981>.

(79) Arrieta, M. C.; Bistriz, L.; Meddings, J. B. Alteration in Intestinal Permeability. *Gut* **2006**, *55*, 1512–1520.

(80) Wang, W.; Uzzau, S.; Goldblum, S. E.; Fasano, A. Human Zonulin, a Potential Modulator of Intestinal Tight Junctions. *J. Cell Sci.* **2000**, *113*, 4435–4440.

(81) Qi, Y.; Goel, R.; Kim, S.; Richards, E. M.; Carter, C. S.; Pepine, C. J.; Raizada, M. K.; Buford, T. W. Intestinal Permeability Biomarker Zonulin Is Elevated in Healthy Aging. *J. Am. Med. Dir. Assoc.* **2017**, *18* (9), 810.e1–810.e4.

(82) Sapone, A.; de Magistris, L.; Pietzak, M.; Clemente, M. G.; Tripathi, A.; Cucca, F.; Lampis, R.; Kryszak, D.; Carteni, M.; Generoso, M.; Iafusco, D.; Prisco, F.; Laghi, F.; Riegler, G.; Carratu, R.; Counts, D.; Fasano, A. Zonulin Upregulation Is Associated with Increased Gut Permeability in Subjects with Type 1 Diabetes and Their Relatives. *Diabetes* **2006**, *55* (5), 1443–1449.

(83) Zak-Golab, A.; Kocelak, P.; Aptekorz, M.; Zientara, M.; Juszczyk, L.; Martirosian, G.; Chudek, J.; Olszanecka-Glinianowicz, M. Gut Microbiota, Microinflammation, Metabolic Profile, and Zonulin Concentration in Obese and Normal Weight Subjects. *Int. J. Endocrinol.* **2013**, *2013*, 674106.

(84) Grootjans, J.; Thuijls, G.; Verdam, F.; Derikx, J. P.; Lenaerts, K.; A, W.; Joep, B. Non-Invasive Assessment of Barrier Integrity and Function of the Human Gut. *World J. Gastrointest. Surg.* **2010**, *2* (3), 61.