

## **The Z0011 trial reduced axillary dissections in early breast cancer: how to correctly stage the axilla in patients with 1-2 sentinel nodes retrieved only?**

Fabio Corsi, M.D.<sup>a,b\*</sup>, Luca Sorrentino, M.D.<sup>a</sup>, Sara Albasini, Ph.D.<sup>b</sup>, Daniela Bossi, M.D.<sup>b</sup>, Lea Regolo, M.D.<sup>b</sup>, Renzo Vanna, Ph.D.<sup>c</sup>, Carlo Morasso, Ph.D.<sup>c</sup>, Laura Villani, M.D.<sup>d</sup>, Marta Truffi, Ph.D.<sup>a</sup>

<sup>a</sup>Department of Biomedical and Clinical Sciences “Luigi Sacco”, University of Milan, via G. B. Grassi 74, 20157 Milan, Italy

<sup>b</sup>Surgery Department, Breast Unit, ICS Maugeri S.p.A. SB, via Maugeri 10, 27100 Pavia, Italy

<sup>c</sup>Nanomedicine and Molecular Imaging Lab, ICS Maugeri S.p.A. SB, via Maugeri 10, 27100 Pavia, Italy

<sup>d</sup>Service of Pathology, Breast Unit, ICS Maugeri S.p.A. SB, via Maugeri 10, 27100 Pavia, Italy

\*Corresponding Authors:

Prof. Fabio Corsi, MD

Department of Biomedical and Clinical Sciences “Luigi Sacco”, University of Milan, Via G. B. Grassi, 74, 20157 Milan, Italy

Surgery Department, Breast Unit, ICS Maugeri S.p.A. SB, via Maugeri 10, 27100 Pavia, Italy

E-mail: [fabio.corsi@unimi.it](mailto:fabio.corsi@unimi.it) - Phone: +39.02.5031.9850 - Fax: +39.02.5031.9846

**Declarations of interest:** None.

**Role of the funding source:** None.

**Funding:** This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

## Highlights

- Axillary dissection is not needed in early breast cancer with 1-2 positive nodes
- A mean of 1-2 SLNs per patients are retrieved in clinical practice
- It is not clear how to apply the Z0011 criteria in case of 1-2 SLNs only
- A nomogram to predict the risk of  $>2$  positive nodes was developed

## **Abstract**

*Background:* The ACOSOG Z0011 trial has decreased the rate of axillary dissections (ALND) after 1-2 positive sentinel lymph nodes (SLN) in breast cancer patients. Evidences showed that usually 1-2 SLNs only are retrieved: how can the surgeon know that only 1-2 nodes are positive, if 1-2 SLNs only are harvested? The aim of this study was to provide an innovative nomogram to specifically predict the risk of more than 2 positive nodes in patients who fulfill the Z0011 criteria.

*Methods:* From 2007 to 2017, at the Breast Unit of ICS Maugeri Hospital 271 patients with T1-T2 breast cancer, 1-2 macro-metastatic SLNs, were treated with breast-conserving surgery. In most cases 1-2 SLNs only were retrieved. Patients fulfilled the Z0011 inclusion criteria, but underwent ALND. A novel nomogram was developed based on preoperative variables.

*Results:* A mean of 1.5 ( $\pm 0.7$ ) SLNs per patient were identified and retrieved. 187 (69.0%) had 1-2 positive nodes, and 84 (31.0%) had 3 or more metastatic nodes. Independent predictors of axillary status were the ratio of positive SLNs on all retrieved SLNs (OR 10.95, 95%CI 2.50-47.93,  $p=0.001$ ), extranodal extension (OR 5.51, 95%CI 2.27-13.41,  $p=0.0002$ ), and multifocal disease (OR 2.9, 95%CI 1.44-5.84,  $p=0.003$ ). A nomogram based on these variables was constructed (AUC after bootstrap = 0.74).

*Conclusion:* The proposed nomogram might select those patients fulfilling the Z0011 criteria, with 1-2 SLNs only harvested, in whom ALND still could be necessary for a proper staging to guide subsequent adjuvant treatments.

**Keywords:** Breast Cancer; Axillary Nodal Status; Nomogram

## **Introduction**

Complete axillary lymph nodes dissection (ALND) after a positive sentinel lymph node (SLN) has become much less relevant in the era of multimodal treatment of breast cancer [1]. The ACOSOG Z0011 randomized clinical trial showed an excellent loco-regional disease control omitting ALND in patients undergoing breast-conserving surgery with 1-2 macro-metastatic SLNs, provided that whole-breast radiotherapy was planned [2]. Long-term outcomes of the Z0011 trial have been recently updated and corroborated previous results [3]. The Z0011 findings have been promptly translated into clinical practice in selected patients who fulfill all the requested criteria [4]. The application of the Z0011 in clinical practice resulted in a significant decrease of ALNDs [5, 6], but a major question arises: how can the surgeon know that only 1-2 nodes are positive, if 1-2 SLNs only are retrieved? This concern is not a collateral issue, because despite few authors report a mean of 3 identified SLNs per patient, the great majority of studies indicate that generally 1 or 2 SLNs are retrieved for each patient [7]. But at least 3 nodes should be removed and analyzed to decide if ALND can be safely omitted, and not surprisingly application of the Z0011 criteria has paradoxically increased the number of harvested SLNs per patient [8]. Moreover, keeping unknown the status of non-sentinel nodes after avoiding ALND in node-positive patients may lead to a “one size fits all” philosophy, widely offering chemotherapy and loco-regional irradiation to strengthen the loco-regional control [2, 9-11]. Therefore, if ALND is omitted after a positive SLN but no more than 2 nodes are removed, prediction of non-sentinel nodes status become fundamental to properly escalate or de-escalate systemic treatments if Z0011 is applied [12]. The aims of the present study were: 1) to assess the different clinical and prognostic features of patients with 1-2 vs. >2 positive nodes who would not have been discriminated avoiding ALND, according to the Z0011 inclusion criteria; 2) to evaluate the accuracy of currently validated nomograms to predict the non-SLN status; 3) to propose an innovative nomogram to specifically predict the risk of more than 2 positive axillary nodes in patients who fulfill the Z0011 criteria, if 1-2 SLNs only are retrieved.

## **Materials and Methods**

### *Study population*

From January 2007 to December 2017, 271 breast cancer patients treated at the Breast Unit of ICS Maugeri Hospital with breast-conserving surgery, clinically node-negative but with 1 or 2 SLNs positive for macrometastasis, were included in the study and received ALND. Inclusion criteria were: proven diagnosis of breast cancer by core-biopsy, stage cT1-cT2 breast cancer evaluated on preoperative imaging, stage cN0 evaluated clinically and by axillary ultrasound, indication for breast lumpectomy and SLN biopsy, and planned radiotherapy. Patients who underwent neoadjuvant chemotherapy or total mastectomy were excluded from the study, as well as patients with a clinically positive axillary status, more than 2 positive SLNs or micro-metastatic disease in the SLN, since pNmic status was not an indication for ALND. All selected patients were suitable not to receive ALND according to the ACOSOG Z0011 trial.

### *SLNs identification and pathological analysis procedures*

In all patients a peri-areolar subdermal injection of 0.2-0.5 mCi of colloidal human serum albumin labeled with <sup>99m</sup>Techneium was performed the day before surgery. During surgery, a gamma probe (Bluetooth Neoprobe Gamma Detection System) was used to localize the SLN. Further nodes with a radioactivity count over 10% of the excised node were considered as other SLNs and removed. All SLNs were sent for intraoperative pathological analysis with standard hematoxylin-eosin (HE), by embedding the sliced lymph node in paraffin wax, taking adjacent sections every 50 micron and staining them with HE.

### *Study design and endpoints*

First, mean number of retrieved SLNs per patient was reported. Based on axillary status evaluated on final histopathology, patients were divided into two groups: those who had only 1-2 positive lymph nodes (low axillary burden, IAB), and those who had >2 positive nodes (high axillary

burden, hAB). The proportions of lAB and hAB patients after 1 or 2 positive SLNs were assessed to evaluate the risk of axillary under-staging related to Z0011 inclusion criteria. Clinical and pathological variables were compared between the two groups, as well as the use of radiotherapy, hormone therapy and chemotherapy, to assess if lAB and hAB patients received different adjuvant treatments after surgery. Then, the crude rates of loco-regional recurrence (LRR) and distant metastases (DM) were compared between lAB and hAB patients, to verify a significant prognostic difference between the two groups of patients. LRR was defined as the occurrence of ipsilateral breast cancer or nodal disease at axillary, internal mammary or supraclavicular level, proven on core biopsy. DM was defined as the occurrence of distant lesions with computed tomography and positron emission tomography characteristics suggestive of malignancy.

Five validated nomograms to predict the non-SLN status in case of positive SLN were retrospectively applied to the present study population: Memorial-Sloan Kettering Cancer Center (MSKCC), Mayo, Tenon score, MD Anderson Cancer Center (MDA) score, and Cambridge formula [13-17]. In each patient all models were used to calculate the individual risk for non-SLN metastases, based on available variables. The accuracy of these nomograms or scoring systems to discriminate lAB and hAB patients was evaluated by each receiver operating curve (ROC) and the relative area under curve (AUC). For each model, the optimal probability cut-off to better predict >2 positive nodes was determined on ROC, and associated specificity and sensitivity were reported.

All relevant variables which emerged to be possibly related to hAB in descriptive analyses were included in a multivariate analysis with a binomial logistic model to find independent predictors of >2 positive nodes. Correlation of each variable with the outcome was assessed by calculating its odds ratio (OR) and  $\beta$  coefficient. Calibration was performed by graphical method (Supplementary Fig. S1). To construct the nomogram, the  $\beta$  coefficient with the higher absolute value selected the relative variable as the “driver variable” on which the model was based, independently from its

statistical significance [18]. A score of 0 points was attributed to the lowest value of that variable, and 100 points to the highest value; intermediate values were matched with corresponding points. Then, the other relevant variables possibly related to hAB were included in the model, also in the absence of a statistical significance on multivariate analysis. Each variable was matched with a score based on the ratio between its  $\beta$  coefficient and  $\beta$  coefficient of the driver variable. The maximum score was obtained with the subsequent formula:

$$\text{Maximum score} = (\beta_{\text{variable}}/\beta_{\text{driver variable}})*100$$

Once constructed the nomogram, it was applied to each patient of the present study to calculate the individual risk of hAB. A ROC with relative AUC was constructed for the proposed nomogram, based on the predicted probability to have >2 positive nodes, and the optimal cut-off value was determined to discriminate with the maximum accuracy IAB and hAB patients. Finally, an internal validation was performed with bootstrap method. Briefly, the original patient population was re-sampled 500 times and the optimism index (the mean of differences between AUC on bootstrap sample and AUC on original sample) was calculated. Optimism is the amount by which the AUC (or “the apparent prediction accuracy”) overestimates the true prediction accuracy of the model. Then, the corrected AUC after bootstrap was reported.

### *Statistical analysis*

Variables were reported as means  $\pm$  standard deviations or as absolute numbers and percentages. Categorical variables were compared using  $\chi^2$  test, while continuous variables were compared using Student’s T test or non-parametric Wilcoxon test in case of non-normal distribution of the variable. Statistical significance was set at  $p < 0.05$  (two tailed). Data analysis was performed using SAS software (v. 9.4, SAS Institute Inc., Cary, USA) and R software (v. 3.5.1, The R Foundation).

## Results

### *Baseline characteristics in lAB and hAB patients*

Mean lesion size on imaging was greater for hAB patients ( $17.6\pm 10.5$  mm vs.  $14.9\pm 7.1$  mm,  $p=0.014$ ). Multifocal disease was observed in 34.5% of hAB cases and in 16.0% of lAB patients ( $p=0.0007$ ). Lympho-vascular invasion was present in 67.9% of hAB lesions and in 54.0% of lAB tumors ( $p=0.034$ ). All the other baseline features were balanced between the two groups, as reported in Table 1.

### *SLN and axillary status in lAB and hAB patients*

187 patients (69.0%) had a maximum of 2 positive nodes, and the remaining 84 cases (31.0%) had 3 or more metastatic nodes. A mean of 1.5 ( $\pm 0.7$ ) SLNs per patient were identified and retrieved both in lAB and hAB cases ( $p=1.000$ ). lAB patients had a lower mean number of positive SLNs per patient compared to hAB cases ( $1.1\pm 0.3$  vs.  $1.3\pm 0.5$ ,  $p<0.0001$ ) and a higher mean number of negative SLNs (respectively  $0.4\pm 0.6$  vs.  $0.2\pm 0.5$ ,  $p=0.008$ ). At final pathology, all 187 lAB patients were staged as pN1, while 42.9% of hAB patients were staged as pN1, 42.9% as pN2 and 14.2% as pN3 ( $p<0.0001$ ). Extranodal extension (ENE) was observed in 6.4% of lAB patients vs. 26.2% of hAB cases ( $p<0.0001$ ). All these data are reported in Table 1.

### *Adjuvant treatments and long-term oncologic outcomes*

Hormone therapy was administered in 88.2% of lAB cases vs. 82.1% of hAB patients ( $p=0.18$ ). Locoregional irradiation was offered to 10.7% of lAB patients and to 53.6% of hAB cases, while remaining patients received whole-breast radiotherapy only ( $p<0.0001$ ). Chemotherapy was administered in 33.2% of lAB cases vs. 71.4% of hAB patients ( $p<0.0001$ , Table 2). In particular, an anthracyclines/taxanes-based regimen was proposed in 45.2% of lAB and 66.7% of hAB patients, trastuzumab was received respectively in 12.9% and 6.6% of cases, and other regimens were offered in 41.9% and 26.7% of patients ( $p=0.06$ ). Mean follow up was similar between

groups, being 52.3 months for lAB and 54.4 months for hAB patients ( $p=0.65$ ). No locoregional recurrence was observed among lAB patients, but it was reported in 6.0% of hAB cases (Log-rank test  $p=0.0005$ ). Distant metastases occurred respectively in 4.3% and 13.1% of patients (Log-rank test  $p=0.01$ ). No difference was reported in terms of cancer-related deaths ( $p=0.11$ , Table 2).

#### *Performance of validated nomograms to predict non-sentinel nodal disease*

ROC of each nomogram is showed in Fig. 1. The most predictive model applied to our series was the Mayo nomogram (AUC = 0.70), and a probability cut-off of 33% resulted in a sensitivity equal to 65% and a specificity of 71%. The MSKCC nomogram showed an AUC of 0.70 (cut-off 0.36, sensitivity 73%, specificity 55%). The Cambridge formula predicted hAB with an AUC of 0.66, while the Tenon score was associated with an AUC of 0.63. The MDA score showed the lowest prediction of >2 positive nodes, with an AUC of 0.59 (cut-off 0.34, sensitivity 70%, specificity 47%).

#### *Independent predictive factors of high axillary burden*

After multivariate analysis, the only independent predictive factors associated to >2 metastatic axillary lymph nodes were a higher ratio of positive SLNs on all retrieved SLNs (OR 10.95, 95%CI 2.50-47.93,  $p=0.001$ ), ENE (OR 5.51, 95%CI 2.27-13.41,  $p=0.0002$ ), and multifocal disease (OR 2.9, 95%CI 1.44-5.84,  $p=0.003$ ), as reported in Table 3.

#### *Development and internal validation of a nomogram to predict hAB*

Since the higher  $\beta$  coefficient was associated to the positive SLNs/retrieved SLNs ratio ( $\beta = 2.42$ ), it was considered the “driver variable” for constructing the nomogram. Based on their  $\beta$  coefficients, presence of ENE ( $\beta = 1.45$ ) was matched with a score of 60.2 and multifocality ( $\beta = 1.03$ ) with a score of 42.58. Once taken into account all the independent predictors for hAB, a set of non-significant variables clinically judged to be relevant for axillary status were included in the model:

estrogen receptor, HER2 status, histological type (invasive ductal vs. invasive lobular), lymphovascular invasion and grading (Fig. 2; Supplementary Table S1). Then, the constructed nomogram was applied to each patient: the minimum total score was 26.42 (predicted probability for hAB 4.0%) and the maximum total score was 276.48 points (predicted probability for hAB 90.0%). For each patient the predicted probability with relative confidence intervals was determined and a fitting plot for prediction related to score was constructed (Supplementary Fig. S2) with high goodness of model's fit (R-squared 0.92), then each interval of probability was associated to a score interval. A ROC based on predicted probability of axillary burden was designed (Fig. 3), and the calculated optimal probability cut-off was 0.31%, which corresponded to a score best cut-off equal to 144.79 points, associated to a sensitivity of 68% and a specificity of 72% for hAB. AUC was equal to 0.76. Finally, an internal validation of model prediction accuracy was performed by bootstrap technique. The optimism index was equal to 0.02, thus the calculated AUC after bootstrap was 0.74. Essentially, we have added a bias correction to the apparent original AUC.

## Discussion

Omission of ALND in the present study would not have correctly discriminated patients with 1-2 metastatic nodes only (69.0% of cases) from hAB patients. This distinction has a prognostic relevance, since no loco-regional recurrence was observed in the former group vs. 6.0% in the latter group ( $p=0.0005$ ), and distant metastases were 3-fold more frequent in hAB patients ( $p=0.01$ ), highlighting two distinct cohorts of patients from a prognostic point of view. Considering that cancer-related baseline variables were substantially balanced between the two groups, particularly biological subtype ( $p=0.98$ ), histological type ( $p=0.11$ ) and grading ( $p=0.12$ ), the extent of axillary nodal involvement confirmed to be a key prognostic determinant, as well established in literature [19].

A proper distinction of patients with low or high axillary burden is therefore mandatory to properly escalate or de-escalate adjuvant treatments, following what has been insistently recommended even in the 2017 St. Gallen consensus conference [20]. A personalized treatment of breast cancer should not simply translate into less ALNDs, but rather should address the most appropriate axillary surgery for each patient. Notably, chemotherapy was confidently avoided in 66.8% of lAB patients despite 1 or 2 macro-metastatic axillary nodes, and these patients experienced significantly less events compared to hAB (Log-rank  $p=0.0005$  for LRR and  $p=0.01$  for DM). Oppositely, up to 58% of patients accrued in the Z0011 trial were treated by upfront chemotherapy despite low-risk features (T1-T2 Luminal-type cancers) and despite a micro-metastatic SLN in 44.6% of cases, in the absence of precise information about their axillary status.<sup>2</sup> Is a less accurate staging of the axilla really a benefit for node-positive patients, if chemotherapy is then routinely used? Nodal involvement is still a fundamental predictive factor. As recently demonstrated by the MINDACT trial and recommended by the American Society of Clinical Oncology in 2017, the use of genomic biomarkers assays to guide decisions on adjuvant chemotherapy in these patients may be adopted,

but the correct knowledge of axillary status is indispensable [21]. Indeed, patients with more than 3 positive nodes are considered at higher risk and could take advantage from chemotherapy, while some patients with 1-3 positive nodes may safely avoid chemotherapy if classified as low-risk based on cancer biology [22].

Guidelines indicate the irradiation of supraclavicular and internal mammary nodes when 4 or more metastatic axillary nodes are encountered [23]. Accordingly, loco-regional radiotherapy was administered in 10.7% of lAB patients only vs. 53.6% of hAB cases ( $p < 0.0001$ ), but no loco-regional recurrence was observed in the formers. Therefore, distinguishing N2 from N1 patients is still relevant to properly tailor the adequate radiotherapy for each patient, because the majority of N1 patients undergoing breast-conserving surgery could safely receive only whole-breast irradiation [23-25]. Remarkably, the majority of Z0011 patients received whole-breast irradiation with high tangential field to cover at least axillary level 1 [2]. The AMAROS trial showed that axillary irradiation is not free from complications, despite fewer than ALND, and up to 15% of patients experienced lymphedema one year after treatment [9]. Thus, again the question is: always better to ignore the exact axillary nodal burden and propose routine axillary radiotherapy, also in patients with 1-2 positive nodes?

According to what emerged from the logistic regression in the present study, the strongest predictor is the ratio between the number of positive SLNs and the total of harvested SLNs (OR 10.95,  $p = 0.001$ ). In other words, also the number of negative SLNs is necessary to predict residual nodal disease [13-15], and a higher mean of negative SLNs per patient was associated to lAB ( $p = 0.008$ ). A patient with 1 retrieved SLN which resulted to be metastatic has a higher probability of residual nodal disease compared to another patient with 1 positive and 1 negative SLN. Another fundamental parameter to estimate the risk of hAB is the presence of ENE, as previously reported in other models [14, 26]. Indeed, ENE is strongly predictive of  $>2$  positive nodes, and better refines the risk prediction independently from the number of retrieved SLNs. Also multifocal disease was a

relevant predictor for hAB, but its usefulness is much less clear considering that it is observed in a low proportion of patients.

Among the tested nomograms to predict non-sentinel nodal disease, the Mayo and MSKCC models showed the best ROC curves, since these nomograms greatly rely on the number of negative SLNs, and a higher total score is assigned if zero negative SLNs are retrieved [13, 14]. An overall weak performance was observed with the other nomograms, with lack of specificity and classification of most patients as high-risk. The inaccuracy was due to the generally low number of negative SLNs and the fact that these nomograms were developed to globally predict non-sentinel nodal disease, and not to specifically discriminate lAB from hAB patients. Therefore, the use of currently validated nomograms may lead to over-estimation of axillary burden and subsequent increase in unnecessary ALND or use of adjuvant chemotherapy or loco-regional irradiation, thus losing the opportunity offered by the Z0011 findings if 1-2 SLNs only are retrieved.

## **Conclusions**

The present study proposes a novel nomogram specifically developed to predict the presence of >2 positive nodes in patients treated by breast-conserving surgery who fulfill the Z0011 inclusion criteria, in whom 1-2 SLNs only are retrieved. This nomogram, if externally validated, might safely select those patients in whom ALND still could be necessary for a proper staging to guide subsequent adjuvant treatments.

## **Conflict of Interest Statement**

None declared.

## References

1. Jatoi I, Benson JR, Toi M. De-escalation of axillary surgery in early breast cancer. *Lancet Oncol* 2016; 17(10):e430-e441.
2. Giuliano AE, McCall L, Beitsch P, et al. Locoregional recurrence after sentinel lymph node dissection with or without axillary dissection in patients with sentinel lymph node metastases: the American College of Surgeons Oncology Group Z0011 randomized trial. *Ann Surg* 2010; 252(3):426-32.
3. Giuliano AE, Ballman KV, McCall L, et al. Effect of Axillary Dissection vs No Axillary Dissection on 10-Year Overall Survival Among Women With Invasive Breast Cancer and Sentinel Node Metastasis: The ACOSOG Z0011 (Alliance) Randomized Clinical Trial. *JAMA* 2017; 318(10):918-926.
4. Gradishar WJ, Anderson BO, Balassanian R, et al. NCCN Guidelines Insights: Breast Cancer, Version 1.2017. *J Natl Compr Canc Netw* 2017; 15(4):433-451.
5. Gainer SM, Hunt KK, Beitsch P, Caudle AS, Mittendorf EA, Lucci A. Changing behavior in clinical practice in response to the ACOSOG Z0011 trial: a survey of the American Society of Breast Surgeons. *Ann Surg Oncol* 2012; 19(10):3152-8.
6. Caudle AS, Hunt KK, Tucker SL, et al. American College of Surgeons Oncology Group (ACOSOG) Z0011: impact on surgeon practice patterns. *Ann Surg Oncol* 2012; 19(10):3144-51.
7. Kim T, Giuliano AE, Lyman GH. Lymphatic mapping and sentinel lymph node biopsy in early-stage breast carcinoma: a metaanalysis. *Cancer* 2006; 106(1):4-16.

8. Subhedar P, Stempel M, Eaton A, Morrow M, Gemignani ML. Do the ACOSOG Z0011 Criteria Affect the Number of Sentinel Lymph Nodes Removed? *Ann Surg Oncol* 2015; 22 Suppl 3:S470-5.
9. Donker M, van Tienhoven G, Straver ME, et al. Radiotherapy or surgery of the axilla after a positive sentinel node in breast cancer (EORTC 10981-22023 AMAROS): a randomised, multicentre, open-label, phase 3 non-inferiority trial. *Lancet Oncol* 2014; 15(12):1303-10.
10. Kim H, Park W, Yu JI, et al. Prognostic Impact of Elective Supraclavicular Nodal Irradiation for Patients with N1 Breast Cancer after Lumpectomy and Anthracycline Plus Taxane-Based Chemotherapy (KROG 1418): A Multicenter Case-Controlled Study. *Cancer Res Treat* 2017; 49(4):970-980.
11. Haque W, Verma V, Hatch S, Klimberg VS, Butler EB, Teh BS. Omission of chemotherapy for low-grade, luminal A N1 breast cancer: Patterns of care and clinical outcomes. *Breast* 2018; 41:67-73.
12. Gruber I, Henzel M, Schönfish B, et al. Prediction of Non-sentinel Lymph Node Metastases After Positive Sentinel Lymph Nodes Using Nomograms. *Anticancer Res* 2018; 38(7):4047-4056.
13. Van Zee KJ, Manasseh DM, Bevilacqua JL, et al. A nomogram for predicting the likelihood of additional nodal metastases in breast cancer patients with a positive sentinel node biopsy. *Ann Surg Oncol* 2003; 10(10):1140-51.
14. Degnim AC, Reynolds C, Pantvaidya G, et al. Nonsentinel node metastasis in breast cancer patients: assessment of an existing and a new predictive nomogram. *Am J Surg* 2005; 190(4):543-50.
15. Barranger E, Coutant C, Flahault A, Delpech Y, Darai E, Uzan S. An axilla scoring system to predict non-sentinel lymph node status in breast

- cancer patients with sentinel lymph node involvement. *Breast Cancer Res Treat* 2005; 91(2):113-9.
16. Hwang RF, Krishnamurthy S, Hunt KK, et al. Clinicopathologic factors predicting involvement of nonsentinel axillary nodes in women with breast cancer. *Ann Surg Oncol* 2003; 10(3):248-54.
  17. Pal A, Provenzano E, Duffy SW, Pinder SE, Purushotham AD. A model for predicting non-sentinel lymph node metastatic disease when the sentinel lymph node is positive. *Br J Surg* 2008; 95(3):302-9.
  18. Iasonos A, Schrag D, Raj GV, Panageas KS. How to build and interpret a nomogram for cancer prognosis. *J Clin Oncol* 2008; 26(8):1364-70.
  19. Sivridis E, Giatromanolaki A, Galazios G, Koukourakis MI. Node-related factors and survival in node-positive breast carcinomas. *Breast* 2006; 15(3):382-9.
  20. Curigliano G, Burstein HJ, P Winer E, et al. De-escalating and escalating treatments for early-stage breast cancer: the St. Gallen International Expert Consensus Conference on the Primary Therapy of Early Breast Cancer 2017. *Ann Oncol* 2017; 28(8):1700-1712.
  21. Krop I, Ismaila N, Andre F, et al. Use of Biomarkers to Guide Decisions on Adjuvant Systemic Therapy for Women With Early-Stage Invasive Breast Cancer: American Society of Clinical Oncology Clinical Practice Guideline Focused Update. *J Clin Oncol* 2017; 35(24):2838-2847.
  22. Goldhirsch A, Winer EP, Coates AS, et al. Personalizing the treatment of women with early breast cancer: highlights of the St Gallen International Expert Consensus on the Primary Therapy of Early Breast Cancer 2013. *Ann Oncol* 2013; 24:2206-2223.
  23. Recht A, Comen EA, Fine RE, et al. Postmastectomy Radiotherapy: An American Society of Clinical Oncology, American Society for Radiation Oncology, and Society of Surgical Oncology Focused Guideline Update. *Ann Surg Oncol* 2017; 24:38–51.

24. Morrow M, Van Zee KJ, Patil S, et al. Axillary Dissection and Nodal Irradiation Can Be Avoided for Most Node-positive Z0011-eligible Breast Cancers: A Prospective Validation Study of 793 Patients. *Ann Surg* 2017; 266(3):457-462.
25. Siponen ET, Vaalavirta LA, Joensuu H, Leidenius MH. Axillary and supraclavicular recurrences are rare after axillary lymph node dissection in breast cancer. *World J Surg* 2012; 36(2):295-302.
26. Aziz S, Wik E, Knutsvik G, et al. Extra-nodal extension is a significant prognostic factor in lymph node positive breast cancer. *PLoS ONE* 2017; 12(2): e0171853.

## Tables

**Table 1.** Distribution of baseline features and axillary status between lAB and hAB patients.

	Low axillary burden (n = 187)	High axillary burden (n = 84)	p Value
<b>Age at diagnosis (years)</b>	58.5 ( $\pm$ 11.6)	58.6 ( $\pm$ 13.1)	0.95
<b>Mammography</b>			
BI-RADS 1-2	28 (15.0%)	12 (14.3%)	0.99
BI-RADS 3	14 (7.4%)	6 (7.0%)	
BI-RADS 4	126 (67.4%)	58 (69.0%)	
BI-RADS 5	19 (10.2%)	8 (9.5%)	
<b>Breast Ultrasound</b>			
U-RADS 1-2	16 (8.6%)	10 (11.9%)	0.55
U-RADS 3	12 (6.4%)	3 (3.5%)	
U-RADS 4	132 (70.6%)	56 (66.7%)	
U-RADS 5	27 (14.4%)	15 (17.9%)	
<b>Clinical findings:</b>			
Palpable lesion	12 (6.4%)	2 (2.4%)	0.38
Non-palpable lesion	137 (73.3%)	65 (77.4%)	
Others	38 (20.3%)	17 (20.2%)	
<b>Lesion size on imaging (mm)</b>	14.9 ( $\pm$ 7.1)	17.6 ( $\pm$ 10.5)	0.014
<b>Clinical T</b>			
T1a	10 (5.3%)	6 (7.1%)	0.37
T1b	45 (24.1%)	14 (16.7%)	
T1c	102 (54.5%)	45 (53.6%)	
T2	30 (16.0%)	19 (22.6%)	
<b>Lesion size on pathology (mm)</b>	15.7 ( $\pm$ 7.2)	16.8 ( $\pm$ 6.4)	0.23
<b>pT stage</b>			
pT1	157 (84.0%)	60 (71.4%)	0.05

pT2	28 (15.0%)	23 (27.4%)	
pT3-4	2 (1.0%)	1 (1.2%)	
<b>Multifocal disease</b>			
Yes	30 (16.0%)	29 (34.5%)	0.0007
No	157 (84.0%)	55 (65.5%)	
<b>Histological type</b>			
Invasive ductal	165 (88.2%)	68 (81.0%)	0.11
Invasive lobular	22 (11.8%)	16 (19.0%)	
<b>Grading</b>			
G1	19 (10.2%)	4 (4.8%)	0.12
G2	114 (61.0%)	47 (56.0%)	
G3	54 (28.9%)	33 (39.3%)	
<b>Biological portrait</b>			
Luminal A	98 (52.4%)	45 (53.5%)	0.98
Luminal B	73 (39.0%)	33 (39.3%)	
HER2-enriched	3 (1.6%)	1 (1.2%)	
Triple-negative	13 (7.0%)	5 (6.0%)	
<b>Lymphovascular invasion</b>			
Yes	101 (54.0%)	57 (67.9%)	0.034
No	86 (46.0%)	27 (32.1%)	
<b>Extensive intraductal component</b>			
Yes	86 (46.0%)	32 (38.1%)	0.23
No	101 (54.0%)	52 (61.9%)	
<b>Total identified SLNs per patient (mean)</b>			
	1.5 ( $\pm$ 0.7)	1.5 ( $\pm$ 0.7)	1.000
<b>Positive SLNs per patient (mean)</b>			
	1.1 ( $\pm$ 0.3)	1.3 ( $\pm$ 0.5)	<0.0001
<b>Negative SLNs per patient (mean)</b>			
	0.4 ( $\pm$ 0.6)	0.2 ( $\pm$ 0.5)	0.008
<b>Total removed nodes per patient (mean)</b>			
	19.3 ( $\pm$ 10.1)	22.6 ( $\pm$ 10.5)	0.015
<b>Total positive nodes per patient (mean)</b>			
	1.4 ( $\pm$ 0.5)	5.8 ( $\pm$ 3.7)	<0.0001
<b>pN stage</b>			
pN1	187 (100.0%)	36 (42.9%)	<0.0001
pN2	0 (0.0%)	36 (42.9%)	

pN3	0 (0.0%)	12 (14.2%)	
<b>Extranodal extension</b>			
Yes	12 (6.4%)	22 (26.2%)	<0.0001
No	175 (93.6%)	62 (73.8%)	

**Table 2.** Adjuvant treatments and long-term outcomes between lAB and hAB patients

	Low axillary burden (n = 187)	High axillary burden (n = 84)	p Value
<b>Hormone therapy</b>			
Yes	165 (88.2%)	69 (82.1%)	0.18
No	22 (11.8%)	15 (17.9%)	
<b>Radiation therapy</b>			
Whole breast	167 (89.3%)	39 (46.4%)	<0.0001
Loco-regional	20 (10.7%)	45 (53.6%)	
<b>Chemotherapy</b>			
Yes	62 (33.2%)	60 (71.4%)	<0.0001
No	125 (66.8%)	24 (28.6%)	
<b>Mean follow up (months)</b>	52.3 (±35.5)	54.4 (±34.0)	0.65
<b>Loco-regional recurrence</b>			
Yes	0 (0.0%)	5 (6.0%)	0.0005
No	187 (100.0%)	79 (94.0%)	
<b>Distant metastases</b>			
Yes	8 (4.3%)	11 (13.1%)	0.01
No	179 (95.7%)	73 (86.9%)	
<b>Cancer-related death</b>			
Yes	3 (1.6%)	5 (6.0%)	0.11
No	184 (98.4%)	79 (94.0%)	

**Table 3.** Multivariate analysis for preoperative predictive factors of high axillary burden

	Prediction of high axillary burden		
	OR	95% CI	p Value
<b>Age at diagnosis</b>	1.00	0.97-1.02	0.90
<b>Lesion size on imaging</b>	1.04	1.00-1.08	0.05
<b>Mammographic findings</b>			
BI-RADS 3 vs. 1	1.56	0.37-6.68	0.55
BI-RADS 4 vs. 1	0.98	0.39-2.49	0.97
BI-RADS 5 vs. 1	0.50	0.12-2.01	0.33
<b>Ultrasound findings</b>			
U-RADS 3 vs. 1	0.26	0.04-1.74	0.17
U-RADS 4 vs. 1	0.63	0.21-1.82	0.39
U-RADS 5 vs. 1	1.35	0.37-4.92	0.65
<b>Multifocal disease</b>			
Yes vs. no	2.90	1.44-5.84	0.003
<b>Histological type</b>			
Invasive lobular vs. ductal	1.35	0.56-3.26	0.51
<b>Grading</b>			
2 vs. 1	1.45	0.37-5.74	0.60
3 vs. 1	2.34	0.51-10.74	0.27
<b>Biological portrait</b>			
Luminal A vs. TNBC	1.25	0.31-5.09	0.76
Luminal B vs. TNBC	1.05	0.29-3.76	0.94
HER2-positive vs. TNBC	1.01	0.06-18.40	0.99
<b>Lympho-vascular invasion</b>			
Present vs. absent	1.49	0.78-2.86	0.23
<b>Extranodal extension</b>			

Present vs. absent	5.51	2.27-13.41	0.0002
<b>Positive SLNs/Total SLNs ratio</b>	10.95	2.50-47.93	0.001

---

### Figure legends

**Fig. 1** ROC curves of five validated nomograms to predict non-sentinel nodal disease in patients fulfilling the Z0011 inclusion criteria

**Fig. 2** Nomogram to predict more than 2 positive axillary nodes in patients fulfilling the Z0011 inclusion criteria

**Fig. 3** Performance of the proposed nomogram in predicting more than 2 positive axillary nodes evaluated by ROC

Figure 1

[Click here to download high resolution image](#)




Figure 2

[Click here to download high resolution image](#)




Figure 3

[Click here to download high resolution image](#)

**Logistic Regression ROC Curve: AUC = 0.759**

