

*Corresponding author: Emanuele Miraglia, Department of Dermatology and Venereology, Sapienza University, Viale del Policlinico 155, 00161 Rome, Italy. E-mail: emanuele.miraglia@hotmail.it

References

1. Bottillo I, Torrente I, Lanari V, Pinna V, Giustini S, Divona L, et al. Germline mosaicism in neurofibromatosis type 1 due to a paternally derived multi-exon deletion. *Am J Med Genet A* 2010;152A:1467–73.
2. Miraglia E, Pecorella I, Persechino F, Visconti B, Calvieri S, Giustini S. A rare association between neurofibromatosis type 1 and vulvar sarcoma. *G Ital Dermatol Venereol* 2014;149:374–6.
3. Miraglia E, Calvieri S, Giustini S. Cutaneous lipomatosis: diagnosing a rare condition. *G Ital Dermatol Venereol* 2017;152:678–80.
4. Ronan SJ, Broderick T. Minimally invasive approach to familial multiple lipomatosis. *Plast Reconstr Surg* 2000;106:878–80.
5. Alkindy A, Chuzhanova N, Kini U, Cooper DN, Upadhyaya M. Genotype-phenotype associations in neurofibromatosis type 1 (NF1): an increased risk of tumor complications in patients with NF1 splice-site mutations? *Hum Genomics* 2012;6:12.

Conflicts of interest.—The authors certify that there is no conflict of interest with any financial organization regarding the material discussed in the manuscript.

History.—Article first published online: October 22, 2018. - Manuscript accepted: October 9, 2018. - Manuscript revised: September 19, 2018. - Manuscript received: April 12, 2018.

(Cite this article as: Miraglia E, Calvieri S, Giustini S. Lipomas in neurofibromatosis type 1: a single-institution experience. *G Ital Dermatol Venereol* 2020;155:375-6. DOI: 10.23736/S0392-0488.18.06044-3)

© 2018 EDIZIONI MINERVA MEDICA
 Online version at <http://www.minervamedica.it>
 Giornale Italiano di Dermatologia e Venereologia 2020 June;155(3):376-7
 DOI: 10.23736/S0392-0488.18.06046-7

Lymphocytic variant of hypereosinophilic syndrome presenting with polymorphic cutaneous manifestations and nonspecific histopathological findings

Hypereosinophilic syndrome (HES) is a heterogeneous group of disorders defined by persistent peripheral blood hypereosinophilia (HE) >1.5 g/L on 2 examinations and/or tissue HE, HE-attributable organ damage and/or dysfunction, and exclusion of other explanations for organ dysfunction.¹ Different variants of HES have been distinguished, including idiopathic, primary, and secondary. Among the latter group, L-HES is due to the presence of an aberrant T cell clone determining an over-production of T helper-2 (Th2) cytokines, *i.e.* interleukin (IL)-5, IL-4, and IL-13, that leads to reactive expansion and activation of eosinophils.¹

An 80-year-old woman presented with a 3-year history of widespread pruritic erythematous-papular rash. This condition had been diagnosed as prurigo-like eczema and treated with systemic steroid, with good response but relapsing to discontinuation. Her medical history revealed multiple superficial thrombophlebitides, a thrombosis of the external jugular vein and a 10-year history of hypereosinophilia. Hypereosinophilia and itch started briefly after hip joint replacement. The patient was not taking any drug. Dermatological examination showed erythematous maculo-papular and nodular lesions involving the trunk, (Figure 1A) gluteal area, lower and upper limbs (Figure 1B), dyshidrosiform vesicobullous lesions on the palmo-plantar areas (Figure 1C), facial flushing, and diffuse excoriated lesions (Figure 1A, B, C). Histopathological examination resulted nonspecific, showing acanthosis with spongiosis, focal parakeratosis with neutrophils, and dermal infiltrate consisting of lymphocytes, histiocytes and numerous eosinophils. Full blood count was normal except for a high eosinophil count (5 · 10³/mmc, 30% of total white blood cells). A blood film confirmed eosinophilia, without atypical cells. Serum immunoglobulin E (IgE) levels were 244 kUA/L. Tryptase levels were within normal ranges. Erythrocyte sedimentation rate was mildly increased (35 mm). Serology testing for parasites, human T-cell lymphotropic virus, and hepatitis B and C gave negative results. Autoimmunity panel was also negative. Plasma levels of tissue factor were high. Memory Lymphocyte Immuno Stimulation Assay (MELISA) for cobalt was positive, with an increase of blasts proliferation of 40%, at a cobalt concentration of 5 mg/mL. Bone marrow aspirate and biopsy exhibited normal findings except for marked eosinophilic hyperplasia. Medullary blood karyotype was normal. Molecular analysis on peripheral blood was negative for BCR/ABL t(9;22)-p210, JAK2 V617F, and FIP1L1/PDGFRα. Flow cytometry immunophenotyping of peripheral blood found an aberrant population of T cells, which showed a CD4-positive, CD7-negative immunophenotype. The same T-cell clone was identified by T-cell gene rearrangement studies on two skin biopsies and peripheral blood. Computed tomography imaging of the entire body found no abnormalities. Lymphocytic variant of hypereosinophilic syndrome (L-HES) was diagnosed. During hospitalization, the patient was administered oral prednisone 25 mg/day, at progressively tapering doses, and intramuscular chlorpheniramine maleate 10 mg/day. At a 2-month follow-up, with a maintenance corticosteroid dose of 10 mg/day, the rash was almost cleared and itch was partially controlled.




Figure 1.—Dermatological examination

This document is protected by international copyright laws. No additional reproduction is authorized. It is permitted for personal use to download and save only one file and print only one copy of this Article. It is not permitted to make additional copies (either sporadically or systematically, either printed or electronic) of the Article for any purpose. It is not permitted to distribute the electronic copy of the article through online internet and/or intranet file sharing systems, electronic mailing or any other means which may allow access to the Article. The use of all or any part of the Article for any Commercial Use is not permitted. The production of derivative works from the Article is not permitted. It is not permitted to remove, cover, overlay, obscure, block, or change any copyright notices or terms of use which the Publisher may post on the Article. It is not permitted to frame or use framing techniques to enclose any trademark, logo, or other proprietary information of the Publisher.

In the current case, our patient met the criteria of HES, and given the presence of clonal T-cell populations and cells with aberrant immunophenotype in the skin and peripheral blood, was classified as L-HES. L-HES is characterized by eosinophilia and eosinophil-infiltrating lesions, especially of the skin, subcutaneous tissue, and — less often — internal organs.¹ Lefèvre *et al.* demonstrated in a study involving 21 patients with L-HES that dermatologic manifestations were the most common, being observed in 81% of the patients.¹ L-HES is responsible for a wide spectrum of protean mucocutaneous manifestations that could also be combined in a single patient¹ (as observed in our case) and include diffused isolated pruritus,¹ recurrent attacks of facial angioedema,^{1,2} diffuse eczematous lesions,¹ macular and maculopapular rash,¹ urticarial plaques,¹ papular lesions,¹ ulcerated plaques,³ necrosis of the oral cavity,⁴ subcutaneous nodules⁵ and widespread papulo-nodular lesions.^{1, 6} Besides, cutaneous manifestations are usually the presenting sign of the disease.¹ To date, little is known about the molecular mechanisms underlying L-HES pathogenesis. The immunophenotype of the T-cell clone is most frequently represented by CD3+CD4-CD8- or CD3-CD4+ subsets of lymphocytes. Other immunophenotypic abnormalities consist of elevated CD5 expression on CD3-CD4+ cells, loss of surface CD7 (as in our patient) and/or expression of CD27.¹ L-HES usually has an indolent course, even though the prognosis is variable and rare cases of malignant evolution towards T-cell lymphoma or Sézary Syndrome have been described, usually after several years of stable disease.¹ The aim of treatment is to inhibit eosinophilopoietic cytokine production (IL-5 and IL-3, GM-CSF) by aberrant T cells and to control their expansion and prevent end-organ damage, thromboembolic events and malignant transformation. Cugno *et al.* reported an increased risk of thrombosis due to a higher tissue factor (TF) expression in eosinophil-mediated disorders. In fact, eosinophils have been described as source of TF, the main initiator of blood coagulation.⁷ As proof of this, our patient's medical history comprised a thrombophilic diathesis. First-line treatment in patients affected by L-HES is represented by systemic steroids, which are characterized by high response rates but need to be maintained at low dosages for long periods.¹ Considering the possible side effect of prolonged corticosteroid therapies, the use of mepolizumab, a monoclonal antibody that acts blocking IL-5,⁸ cyclosporine¹ or interferon- α ¹ as corticosteroid-sparing agents may have also a rationale.

In conclusion, an important point of discussion is the possible role of the prosthetic material in the induction of the T-cell aberrant clone. Indeed, in our patient there is a temporal link between the eosinophilia and prosthetic surgery. Moreover, by means of MELISA we demonstrated a type IV hypersensitivity to cobalt that was contained in the prosthesis of our patient.

Marco CUSINI¹ *, Alessio PIZZATI¹,
Giovanni GENOVESE¹, Simona MURATORI¹,
Giuseppina SCHINCO², Emilio BERTI¹

¹UO Dermatologia, Dipartimento di Fisiopatologia Medico-Chirurgica e dei Trapianti, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy; ²Department of Internal Medicine, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy

*Corresponding author: Marco Cusini, Department of Dermatology, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Via Pace 9, Milan, Italy. E-mail: marco.cusini@policlinico.mi.it

References

1. Lefèvre G, Copin MC, Staumont-Sallé D, Avenel-Audran M, Aubert H, Taieb A, *et al.*; French Eosinophil Network. The lymphoid variant of hypereosinophilic syndrome: study of 21 patients with CD3-CD4+ aberrant T-cell phenotype. *Medicine (Baltimore)* 2014;93:255–66.
2. González Delgado P, de la Sen Fernández ML, Soriano Gomis V, Pérez Crespo M, Muñoz Ruiz C, Hernández Niveiro E. Cyclical hypereosinophilia with skin manifestations and a clonal T cell population. *J Investig Allergol Clin Immunol* 2008;18:401–3.
3. Klion AD, Mejia R, Cowen EW, Dowdell KC, Dunleavy K, Fahle GA, *et al.* Chronic active Epstein-Barr virus infection: a novel cause of lymphocytic variant hypereosinophilic syndrome. *Blood* 2013;121:2364–6.
4. d'Elbée JM, Parnes M, Mercié P, Longy Boursier M, Dieval C, de Mascarel A, *et al.* Hypereosinophilic syndrome - lymphocytic variant transforming into peripheral T-cell lymphoma with severe oral manifestations. *Oral Surg Oral Med Oral Pathol Oral Radiol* 2013;116:e185–90.
5. Hanbali A, Shaheen M, Alfraih F, Al-Otaibi W, El Fakih R, Owaidah T, *et al.* A case of T-cell lymphoproliferative disorder associated with hypereosinophilia with excellent response to mycophenolate mofetil. *Hematol Oncol Stem Cell Ther* 2018;11:241–4.
6. Kempf W, Kazakov DV, Szep Z, Vanecek T. CD30+ clonal T-cell lymphoid proliferation of the skin in a patient with hypereosinophilic syndrome. *J Cutan Pathol* 2015;42:130–5.
7. Cugno M, Marzano AV, Lorini M, Carbonelli V, Tedeschi A. Enhanced tissue factor expression by blood eosinophils from patients with hypereosinophilia: a possible link with thrombosis. *PLoS One* 2014;9:e111862.
8. Roufosse FE, Kahn JE, Gleich GJ, Schwartz LB, Singh AD, Rosenwasser LJ, *et al.* Long-term safety of mepolizumab for the treatment of hypereosinophilic syndromes. *J Allergy Clin Immunol* 2013;131:461–7.e1, 5.

Conflicts of interest.—The authors certify that there is no conflict of interest with any financial organization regarding the material discussed in the manuscript.

History.—Article first published online: May 16, 2018. - Manuscript accepted: May 10, 2018. - Manuscript received: April 18, 2018.

(Cite this article as: Cusini M, Pizzati A, Genovese G, Muratori S, Schinco G, Berti E. Lymphocytic variant of hypereosinophilic syndrome presenting with polymorphic cutaneous manifestations and nonspecific histopathological findings. *G Ital Dermatol Venereol* 2020;155:376-7. DOI: 10.23736/S0392-0488.18.06046-7)

© 2018 EDIZIONI MINERVA MEDICA
Online version at <http://www.minervamedica.it>
Giornale Italiano di Dermatologia e Venereologia 2020 June;155(3):377-9
DOI: 10.23736/S0392-0488.18.06054-6

Anti-TNF α -induced psoriasis under treatment of hidradenitis suppurativa: report of 2 cases

Tumor necrosis factor (TNF)- α inhibitors are biologic drugs widely used for the treatment of inflammatory and autoimmune diseases, including psoriasis. Infliximab, adalimumab and etanercept are among the most popular ones in dermatology and may paradoxically be responsible for the onset of psoriasis or the wors-