

# Travel-associated chikungunya acquired in Myanmar in 2019

Marta Díaz-Menéndez<sup>1,2</sup>, Elena Trigo Esteban<sup>1,2</sup>, Mugen Ujiie<sup>3</sup>, Guido Calleri<sup>4</sup>, Camilla Rothe<sup>5</sup>, Denis Malvy<sup>6</sup>, Emanuele Nicastrì<sup>7</sup>, Alfred L Bissinger<sup>8</sup>, Marc Grandadam<sup>9</sup>, Jonathan D Alpern<sup>10</sup>, Federico Gobbi<sup>11</sup>, Patricia Schlagenhauf<sup>12</sup>, Alexandre Duvignaud<sup>13</sup>, Emilie Javelle<sup>14</sup>, Takato Nakamoto<sup>15</sup>, Spinello Antinori<sup>16</sup>, Davidson H Hamer<sup>17</sup>

1. National Referral Unit for Imported Tropical Diseases. Department of Internal Medicine. Hospital Universitario La Paz-Carlos III, IdiPAZ, Madrid, Spain
2. These authors contributed equally to the work
3. Vaccination Support Center, Disease Control and Prevention Center. National Center for Global Health and Medicine, Tokyo, Japan
4. Travel Medicine Unit. Amedeo di Savoia Hospital. ASL Città di Torino, Torino, Italy
5. LMU Hospital Centre, Division of Infectious Diseases and Tropical Medicine, Munich, Germany
6. Department of Infectious Diseases and Tropical Medicine, CHU de Bordeaux & INSERM 1219, University of Bordeaux, Bordeaux, France
7. National Institute for Infectious Diseases Lazzaro Spallanzani, IRCCS, Rome, Italy
8. Institute for Tropical Medicine, Department of Internal Medicine, University Hospital Tübingen, Germany
9. Arbovirus & Emerging viral diseases laboratory. Institut Pasteur du Laos, Vientiane, Laos
10. Department of Travel and Tropical Medicine, HealthPartners; Department of Medicine, University of Minnesota, Minneapolis, United States
11. Department of Infectious /Tropical Diseases and Microbiology, IRCCS Sacro Cuore Don Calabria Hospital, Negrar, Verona, Italy
12. University of Zürich Centre for Travel Medicine, WHO Collaborating Centre for Travellers' Health, Zürich, Switzerland
13. Department of Infectious Diseases and Tropical Medicine, CHU de Bordeaux & INSERM 1219, University of Bordeaux, Bordeaux, France
14. Laveran Military Teaching Hospital, Aix Marseille Univ, IRD, AP-HM, SSA, VITROME, Marseille, France
15. Disease Control and Prevention Center, National Center for Global health and Medicine, Tokyo, Japan
16. Department of Biomedical and Clinical Sciences "L Sacco", University of Milan. Tropical Medicine Unit, ASST Fatebenefratelli Sacco, Milan, Italy
17. Department of Global Health and Boston University School of Public Health and Section of Infectious Diseases, Department of Medicine, Boston University School of Medicine, Boston, MA, United States

**Correspondence:** Marta Díaz-Menéndez (marta.diazmenendez@gmail.com)

## Citation style for this article:

Díaz-Menéndez Marta, Esteban Elena Trigo, Ujiie Mugen, Calleri Guido, Rothe Camilla, Malvy Denis, Nicastrì Emanuele, Bissinger Alfred L, Grandadam Marc, Alpern Jonathan D, Gobbi Federico, Schlagenhauf Patricia, Duvignaud Alexandre, Javelle Emilie, Nakamoto Takato, Antinori Spinello, Hamer Davidson H. Travel-associated chikungunya acquired in Myanmar in 2019. *Euro Surveill.* 2020;25(1):pii=1900721. <https://doi.org/10.2807/1560-7917.ES.2020.25.1.1900721>

Article submitted on 24 Nov 2019 / accepted on 09 Jan 2020 / published on 09 Jan 2020

**Eighteen cases of chikungunya virus infection in travellers returning from Myanmar were reported to the GeoSentinel Surveillance Network, its subnetwork EuroTravNet and TropNet in 2019, reflecting an ongoing local outbreak. This report reinforces the importance of travellers as sentinels of emerging arboviral outbreaks and highlights the importance of vigilance for imported cases, due to the potential for dissemination of the virus into areas with competent local vectors and conducive environmental conditions.**

In early October 2019, a GeoSentinel Surveillance Network site in Madrid, Spain, identified two patients with chikungunya virus (CHIKV) infection who had recently visited Myanmar. Rapid outreach to GeoSentinel sites (including EuroTravNet), TropNet and Laos external collaborators identified 16 additional infected travellers who acquired CHIKV infection in Myanmar during 2019.

According to a newspaper report, the Myanmar Department of Public Health identified an outbreak of

CHIKV infection in 2019, mainly in Nay Pyi Taw, Kachin State, and Tanintharyi Region, Myanmar [1]. In the previous 8 years however, no cases of CHIKV infection had been officially reported in the country [2,3].

We describe travellers with imported CHIKV infection from Myanmar who were diagnosed in 2019 (January to November) and identified at GeoSentinel, EuroTravNet and TropNet sites. Some epidemiological and clinical information as well as places visited by the travellers are presented.

## Case finding

GeoSentinel is a global surveillance network for emerging infectious diseases that has 68 sites in 28 countries; EuroTravNet is its European subnetwork; TropNet represents a separate European surveillance entity represented by 75 specialised tropical medicine centres in Europe.

For chikungunya surveillance, GeoSentinel follows case definitions proposed by the World Health Organization

**TABLE 1**

Epidemiological and travel characteristics of confirmed and probable chikungunya cases among travellers returning from Myanmar, 2019 (n = 18)

Case	Reporting country	Places visited in Myanmar	Pre-travel consultation	Period of stay in Myanmar (length of exposure period)	Approximate age in years <sup>a</sup>	Underlying medical condition(s)	Month of symptom onset
1	Japan	Yangon	Yes	Apr 2017 to Jul 2019 (NA)	60	Hypertension	Jul 2019
2	Japan	Yangon, Naypyidaw	No	Aug 2019 (6 days)	25	Atopic dermatitis	Aug 2019
3	France	Shan state, Bagan, Yangon	Yes	Jul 2019 (12 days)	45	No	Aug 2019
4	Italy	Nyaungshwe	Yes	Lived in Myanmar until Aug 2019 (NA)	60	No	Aug 2019
5	Japan	Yangon	Yes	Aug 2018 to Aug 2019 (NA)	70	Hyperlipidaemia, hypertension, diabetes mellitus	Aug 2019
6	France	Yangon, Hpa-An, Moulmein, Ye, Mandalay, Bagan, Inle lake, Pindaya	Yes	Jul to Aug 2019 (23 days)	45	No	Aug 2019
7	United States	Ye, (Mon State), Yangon	Unknown	May to Aug 2019 (65 days)	20	No	Jul 2019
8	Spain	Yangon, Mandalay, Inle Lake	No	Aug 2019 (7 days)	30	No	Aug 2019
9	Spain	Yangon Mandalay, Inle Lake, Bagan	No	Aug 2019 (14 days)	55	No	Aug 2019
10	Italy	Yangon, Mandalay, Inle Lake,	No	Aug to Sep 2019 (12 days)	50	No	Sep 2019
11	Italy	Yangon, Mandalay, Inle Lake	No	Aug to Sep 2019 (12 days)	50	No	Sep 2019
12	Laos	Mandalay	No	Sep 2019 (12 days)	50	No	Sep 2019
13	Germany	Yangon, Inle Lake, Bagan, Mandalay	Yes	Jan 2019 (15 days)	65	Mitral valve disease	Feb 2019
14	Germany	Yangon, Inle Lake, Bagan, Mandalay	Yes	Jan 2019 (15 days)	30	No	Feb 2019
15	Italy	Yangon, Bagan, Mandalay	No	Jul to Aug 2019 (12 days)	65	Hypertension	Jul 2019
16	Italy	Rangoon, Bagan	No	Aug 2019 (10 days)	55	No	Aug 2019
17	Germany	Yangon, Inle Lake, Mandalay, Bagan, Thandwe/ West Coast	No	Oct 2019 (16 days)	25	No	Oct 2019
18	France	Yangon, Mandalay, Bagan, Inle Lake	Unknown	Oct to Nov 2019 (15 days)	55	No	Oct 2019

F: female; Ig: immunoglobulin; M: male; NA: not assessed; VFR: visiting friends and relatives.

<sup>a</sup> Age rounded by five.

## FIGURE

Possible places of exposure: destinations visited by travellers before chikungunya virus infection, Myanmar, 2019 (n = 18)



Asterisks represent the places where travellers in the current study were possibly exposed to chikungunya virus. Some travellers visited the same destinations.

Source: Map adapted from Wikimedia Commons, the free media repository: [https://commons.wikimedia.org/wiki/File:Myanmar\\_location\\_map.svg](https://commons.wikimedia.org/wiki/File:Myanmar_location_map.svg)

(WHO) Regional Office for Southeast Asia [4]. Such definitions were used in the current report. A possible case was a patient with acute onset of fever  $>38.5^{\circ}\text{C}$  and severe arthralgia/arthritis not explained by other medical conditions. A probable case was a patient meeting both the previously mentioned clinical criteria and the following epidemiological criteria: residing or having visited epidemic areas, having reported transmission within 15 days prior to the onset of symptoms. A confirmed case had to meet one or more of the following laboratory criteria, irrespective of the clinical presentation: (i) virus isolation in cell culture or animal inoculations from acute phase sera, or (ii) presence of viral RNA in acute phase sera as demonstrated by RT-PCR, or (iii) presence of virus-specific IgM antibodies in single serum sample in acute or convalescent stage, or (iv) fourfold increase in virus-specific IgG antibody titre in samples collected at least 3 weeks apart.

For this study, cases were excluded if they were not in Myanmar at the likely time of exposure, which was inferred from their date of symptom onset and the typical incubation period (defined as 3 to 7 days) [5] or if the travellers had more than one potential travel destination exposure based on the incubation period.

## Description of cases

Epidemiological and clinical details for each patient were collected from reporting sites.

The 18 cases reported in the current study had a median age of 51 years (range: 19–68 years) and 10 were male (Table 1). Among the overall cases, one had visited friends and relatives in Myanmar at the likely time of exposure, two were there on a business trip and three were expatriates. The rest of the cases, which represented the majority (n=12), were tourists who had experienced a median length of stay of 13 days in Myanmar (range: 6–65 days).

The majority of cases were imported to Europe (n=13) and Asia (n=4), with one case in the Americas. Fewer than half (7/16) of travellers with available information, received pre-travel advice. Destinations visited in Myanmar are shown in the Figure.

Among the 18 cases, acute symptoms reported included fever (n=18), arthralgia (n=16) and rash (n=14) (Table 2). A median of 9.5 days (range: 1–240 days) elapsed between symptom onset and diagnosis. Most travellers (n=13) were identified at reporting sites between August and October 2019. Seven cases were confirmed by PCR, including five who were also found IgM positive, and nine cases were confirmed by a positive IgM only; the remaining two patients had probable diagnoses based on a single IgG positive serology. No acute coinfections with other arboviruses were described in the 16 patients tested for additional pathogens.

**TABLE 2A**

Clinical and diagnostic characteristics of confirmed and probable chikungunya cases among returning travellers from Myanmar, 2019 (n = 18)

Case	Clinical acute symptoms	Outcome	Need of hospitalisation (country)	Clinical status as at 20 Oct 2019	Chikungunya diagnostic method	Day of diagnosis in 2019 (time from onset of illness)	Other arboviral diagnostic test results
1	Fever, rash, arthralgias (fingers, wrists, ankles, knees)	Persistent arthralgias needing NSAIDs	No	Unknown	IgM+ (NHealth Myanmar Co., Ltd, Yangon, Myanmar)	08 Jul (6 days)	Dengue: IgM- / IgG-; NS1- Zika: NS1-
2	Fever, rash, arthralgia (wrists, ankles), retro-orbital pain, conjunctival hyperaemia	Persistent arthralgias needing NSAIDs	No	Improvement of arthralgias	IgM+; PCR+ (NovaLisa, NovaTec, Dietzenbach, Germany)	13 Aug (6 days)	Dengue: PCR-; IgM- Zika: PCR-; IgM-
3	Fever, rash, arthralgia, myalgia, conjunctivitis	Persistent incapacitating arthralgias needing NSAIDs	No	Mild and improving arthralgia, no more NSAIDs needed	RT-PCR+ (RealStar chikungunya RT-PCR, Altona diagnostics, Hamburg, Germany)	14 Aug (1 day)	Dengue: PCR- Zika: PCR-
4	Fever, arthralgia, deep asthenia, profuse sweating	Persistent arthralgias needing NSAIDs and CT (prednisone)	Yes (Italy)	Unknown	IgM+ / IgG+ (Euroimmun, Luebeck, Germany)	14 Aug (9 days)	Unknown
5	Fever, rash, arthralgia (fingers, wrists, knees), ankle swelling, conjunctival hyperaemia	Persistent arthralgias needing NSAIDs	Yes (United States)	Improving existing arthritis and arthralgia	IgM+ / IgG+ (NovaLisa, NovaTec)	21 Aug (20 days)	Dengue: IgM- / IgG- Zika: IgM-
6	Fever, ankles oedema	Persistent and recurrent arthralgias needing NSAIDs; fatigue, headache	No	Persistent moderate arthralgia, still needing NSAIDs	IgM+ / IgG+ (Euroimmun)	03 Sep (12 days)	Dengue: IgM- Zika: IgM-
7	Fever, rash, chills, myalgias, symmetric arthralgias (knees, ankles, wrists, hands)	Improving with NSAIDs	No	Improving	IgM+ / IgG+ (Arup Laboratories, Utah, United States)	04 Sep (41 days)	Not tested
8	Fever, rash, arthralgias (ankles, wrists, feet, interphalangeal joints in hands, bilateral)	Persistent arthralgias needing NSAIDs and CT	No	Improved	IgM+ (Euroimmun)	09 Sep (10 days)	Dengue: IgM- / IgG- Zika: IgM ambiguous / IgG-
9	Fever, rash, arthralgia	Persistent arthralgias needing NSAIDs and CT	No	Improved	IgM+ (Chemiluminescence Virclia UNILABS, Madrid, Spain)	13 Sep (21 days)	Dengue: IgM- / IgG- Zika: IgM- / IgG-
10	Fever, rash, arthralgia, diarrhoea, paraesthesia lower limbs, swelling upper and lower limbs, lymphadenopathy	Persistent arthralgias needing NSAIDs	No	Persistent arthralgia	IgM+ / IgG-; PCR+ (ChLIA Lifax, Polverada, Italy)	13 Sep (7 days)	Dengue: RDT- Zika: PCR-; IgM- / IgG-

**TABLE 2B**

Clinical and diagnostic characteristics of confirmed and probable chikungunya cases among returning travellers from Myanmar, 2019 (n = 18)

Case	Clinical acute symptoms	Outcome	Need of hospitalisation (country)	Clinical status at 20 Oct 2019	Chikungunya diagnostic method	Day of diagnosis in 2019 (time from onset of illness)	Other arboviral diagnostic test results
11	Fever, rash, arthralgia, diarrhoea	Persistent arthralgias needing NSAID	No	Persistent arthralgia	IgM+/IgG-; PCR+ (ChLIAAlifax)	13 Sep (6 days)	Dengue: RDT- Zika: PCR-, IgM-/IgG-
12	Fever, pruritus, rash, head and limbs felt swollen	Persistent arthralgias needing paracetamol	No	Little light headed when exercise	RT-PCR+ (in house RT-PCR: [24])	23 Sep (4 days)	Dengue NS1-; RT-PCR- Zika: not tested
13	Fever, nausea, arthralgias, ankle oedema	Persistent incapacitating arthralgias and ankle oedema needing NSAIDs and CT	No	Persistent incapacitating arthralgias and ankle oedema needing NSAIDs and corticosteroids	IgG+ (Euroimmun)	07 Oct (8 months)	Dengue: NS1-; IgM-, IgG+ (multiple exposures before, IIFT+ with low titre) Zika: not tested
14	Fever, rash, arthralgia	Not needing any treatment	No	Mild ankles arthralgias	IgG+ (Euroimmun)	07 Oct (8 months)	Dengue: NS1-; IgM-/IgG- Zika: not tested
15	Fever, rash, arthralgias (wrists, right elbow, metacarpophalangeal joints), general malaise and nausea	Persistent arthralgias needing NSAIDs	Yes (Myanmar)	Improve	IgM+/IgG+ (NovaLisa, NovaTec)	14 Oct (74 days)	Dengue: NS1- Zika: not tested
16	Fever, rash, arthralgia	Persistent arthralgias needing NSAIDs	No	Persistent arthralgias	IgM+/IgG+ (Euroimmun)	16 Oct (60 days)	Not tested
17	Fever, rash, arthralgias (hands, ankles)	No drugs needed	No but presented to local outpatient clinic where no testing for CHIK was done.	Mild arthralgias in fingers and ankles	RT-PCR+; IgM+/IgG- (Fast Track Diagnostics. Ltd, Esch-sur-Alzette, Luxembourg)	21 Oct (7 days)	Zika: PCR-; IgM-/IgG- Dengue: PCR-; NS1-
18	Fever, arthralgia	Improving arthralgia, NSAIDs no more needed	Yes (France)	Moderate persisting arthralgias	RT-PCR+ (RealStar chikungunya RT-PCR, Altona diagnostics); IgM+/IgG- (Euroimmun)	02 Nov (4 days)	Dengue: PCR-; IgM- Zika: PCR-; IgM-

CT: corticosteroids; IIFT: indirect immunofluorescence test; Ig: immunoglobulin; NSAID: non-steroidal anti-inflammatory drug; NS1: nonstructural 1 protein antigen; RDT: rapid diagnostic test; RT-PCR: real time PCR.

Four patients were hospitalised, and 15 needed non-steroidal anti-inflammatory drugs; four had corticosteroids added to manage symptoms (Table 2). As at 20 October 2019, fourteen patients still had persistent symptoms, predominantly arthralgia.

## Discussion

Chikungunya is often a mild self-limited illness, but severe and life-threatening complications have been described [6]. Persistent polyarthralgia can affect up to 40% of infected individuals and may last for months or years [7,8]. A high proportion of cases in our case series (14 of 18) had persistent sequelae (mainly arthralgia) that interfered with their daily life. Given the potential for long-term morbidity, the absence of a curative treatment or a preventive vaccine, detailed pre-travel counselling should focus on mosquito-bite prevention, particularly for those at high risk of incapacitating complications such as women older than 40 years, people with underlying rheumatic diseases, and professional athletes [9].

Considering the short duration of stay in different parts of Myanmar of many of the currently reported travellers, as well as the usual incubation period of chikungunya, the specific place where they were infected cannot be determined with certainty. Most patients in the case series acquired chikungunya from August to October, likely due to enhanced vector activity at the end or after the monsoon season, which occurs between May and August.

CHIKV has been detected in South and South East Asia since the 1950s [3], but its current epidemiology is poorly understood [10]. In Myanmar, CHIKV infection was first described in 1973 [11]. Since this time, the virus has caused outbreaks in 1998, 2006 and 2008 [3]. The number of reported cases subsequently declined until 2010, when local surveillance identified the last reported case [3]. Thereafter, from 2011 to 2018 no CHIKV cases were officially reported in the country, however a review of travellers who acquired chikungunya in Myanmar in the GeoSentinel database from 2000 to October 2019 revealed two cases in 2015 and 2016, one probable and one confirmed.

In 2019, Myanmar's National Health Laboratory detected chikungunya cases again [1]. During the same year, other national surveillance systems also identified travellers who acquired chikungunya in Myanmar. For example, the Japanese National Institute of Infectious Diseases, which systematically tracks travellers, detected 28 imported cases of chikungunya from Myanmar to Japan [12]. The Italian Arbovirology Surveillance System also detected four imported cases in Italy [13]. Our report highlights additional cases of exported CHIKV infection from Myanmar.

Little is known about factors contributing to the multiple detections of CHIKV infections in or from Myanmar in 2019. While aside from this year, no cases within the

country had been officially observed since the end of 2010, the detection of exported cases in 2015 and 2016 could suggest that CHIKV might have continued to circulate in Myanmar after 2010. Myanmar's disease surveillance system includes a Central Epidemiology Unit (CEU) and several vertical control programmes [14]. Certain areas of the country are, however, difficult to access and there could have been underreporting due to limitations of surveillance capacity.

Myanmar has geographical and epidemiological characteristics that put it at risk for CHIKV epidemics: a highly susceptible population [15], a long border with two large neighbouring endemic countries (India and Thailand) with cross-border population movement for trade and travel purposes [16], and competent vectors (both *Aedes aegypti* and *Ae. albopictus*) [17]. According to the Thai Ministry of Public Health, CHIKV is currently circulating in Thailand with an increasing incidence: as at 30 October 2019, the country has reported 8,744 cases this year [18]. Coincidentally, GeoSentinel recently documented a rise in the number of travel-associated CHIKV infections from Thailand [19]. Without further epidemiological and phylogenetic evidence, it remains difficult to establish whether any links exist between cases in Thailand and those in Myanmar.

It should be noted that Myanmar is an increasingly popular tourist destination, with more than 3.5 million visitors in 2017 [20]. Hence, although we report more cases exported from Myanmar in 2019 than in previous years, a surge in tourism to this country may have led to an apparent increase of the number of exported cases. GeoSentinel data are moreover not population-based so rates or risk estimates cannot be derived. Also, diagnostics depend on local site interpretation and reporting.

Imported CHIKV infection by viraemic travellers returning to their home countries raises the possibility of virus spread to these countries if competent vectors are present, mainly in hot seasons, as European *Ae. albopictus* is affected by seasonal temperature and undergoes a winter diapause [21]. There is also a potential risk of transmission by other routes, such as blood donation [22]. In Europe, some countries (Italy in 2007 and 2017, France in 2010, 2014 and 2017) have experienced autochthonous transmission of CHIKV through viraemic travellers [23]. Surveillance and early detection of both imported and autochthonous CHIKV infections is therefore relevant in areas with competent vectors, as well as close vector monitoring and rapid public health response.

This report of imported CHIKV infections reinforces the importance of travellers as sentinels of local outbreaks, particularly in settings with limited public health surveillance and reporting infrastructure.

## Acknowledgements

We would like to thank Dr. Chang-Kweng Lim in the Department of Virology, National Institute of Infectious Diseases of Japan, for his kind assistance with chikungunya diagnostics. We thank Pauline Perreau of the Department of Infectious Diseases and Tropical Medicine, CHU de Bordeaux, for her kind assistance with patient data extraction. We would finally like to thank Kristina M. Angelo of the CDC/DDID/NCEZID/DGMQ for her editorial assistance.

## Conflict of interest

None declared.

## Authors' contributions

Marta Díaz-Menendez, Elena Trigo Esteban, Mugen Ujiie, Guido Calleri, Camilla Rothe, Denis Malvy, Emanuele Nicastrì, Alfred L. Bissinger, Marc Grandadam, Jonathan D. Alpern, Federico Gobbi, Alexandre Duvignaud, Takato Nakamoto, Spinello Antinori provided travel history as well as clinical and biological information on the travellers. Marta Díaz-Menendez, Elena Trigo Esteban and Davidson H. Hamer created the first draft of the manuscript and modified it according to the editorial team's comments. Emilie Javelle, Patricia Schlagenhauf and Davidson H. Hamer reviewed and edited the manuscript. All authors provided input and reviewed the final version of the manuscript.

## References

1. Myanmar Times. Chikungunya reappears after 10 years. [Accessed 20 Oct 2019]. Available from: <https://www.mmtimes.com/news/chikungunya-reappears-after-10-years.html>
2. Tun MM, Thant KZ, Inoue S, Nabeshima T, Aoki K, Kyaw AK, et al. Detection of east/central/south African genotype of chikungunya virus in Myanmar, 2010. *Emerg Infect Dis*. 2014;20(8):1378-81. <https://doi.org/10.3201/eid2008.131431> PMID: 25062511
3. Wimalasiri-Yapa BMCR, Stassen L, Huang X, Hafner LM, Hu W, Devine GJ, et al. Chikungunya virus in Asia - Pacific: a systematic review. *Emerg Microbes Infect*. 2019;8(1):70-9. <https://doi.org/10.1080/22221751.2018.1559708> PMID: 30866761
4. World Health Organization South-East Asia Regional Office (WHO SEARO). Proposed case definition of Chikungunya Fever (WHO, SEARO). New Delhi: WHO SEARO. [Accessed 19 Dec 2019]. Available from: [http://origin.searo.who.int/entity/emerging\\_diseases/topics/Def\\_Chikungunya\\_Fever.pdf](http://origin.searo.who.int/entity/emerging_diseases/topics/Def_Chikungunya_Fever.pdf)
5. Silva JVJ Jr, Ludwig-Begall LF, Oliveira-Filho EF, Oliveira RAS, Durães-Carvalho R, Lopes TRR, et al. A scoping review of Chikungunya virus infection: epidemiology, clinical characteristics, viral co-circulation complications, and control. *Acta Trop*. 2018;188:213-24. <https://doi.org/10.1016/j.actatropica.2018.09.003> PMID: 30195666
6. Economopoulou A, Dominguez M, Helynck B, Sissoko D, Wichmann O, Quenel P, et al. Atypical Chikungunya virus infections: clinical manifestations, mortality and risk factors for severe disease during the 2005-2006 outbreak on Réunion. *Epidemiol Infect*. 2009;137(4):534-41. <https://doi.org/10.1017/S0950268808001167> PMID: 18694529
7. Schwartz O, Albert ML. Biology and pathogenesis of chikungunya virus. *Nat Rev Microbiol*. 2010;8(7):491-500. <https://doi.org/10.1038/nrmicro2368> PMID: 20551973
8. Consuegra-Rodríguez MP, Hidalgo-Zambrano DM, Vásquez-Serna H, Jimenez-Canizales CE, Parra-Valencia E, Rodríguez-Morales AJ. Post-chikungunya chronic inflammatory rheumatism: Follow-up of cases after 1 year of infection in Tolima, Colombia. *Travel Med Infect Dis*. 2018;21:62-8. <https://doi.org/10.1016/j.tmaid.2017.11.013> PMID: 29203304
9. van Aalst M, Nelen CM, Goorhuis A, Stijns C, Grobusch MP. Long-term sequelae of chikungunya virus disease: A systematic review. *Travel Med Infect Dis*. 2017;15:8-22. <https://doi.org/10.1016/j.tmaid.2017.01.004> PMID: 28163198
10. Pulmanusahakul R, Roytrakul S, Auewarakul P, Smith DR. Chikungunya in Southeast Asia: understanding the emergence

- and finding solutions. *Int J Infect Dis*. 2011;15(10):e671-6. <https://doi.org/10.1016/j.ijid.2011.06.002> PMID: 21775183
11. Thaug U, Ming CK, Swe T, Thein S. Epidemiological features of dengue and chikungunya infections in Burma. *Southeast Asian J Trop Med Public Health*. 1975;6(2):276-83. PMID: 126493
  12. National Institute of Infectious Diseases (NIID) Japan; Infectious Diseases Control Division (IDCD), Ministry of Health, Labor and Welfare (MHLW), Japan. IDWR (Infectious Diseases Weekly Report), week 42, as of 6 Oct 2019: NIID; 2019. [Accessed 29 Oct 2019]. Available from: <https://www.who.nih.go.jp/niid/idsc/idwr/IDWR2019/idwr2019-42.pdf>
  13. Sistema di Sorveglianza delle Arbovirosi Rapporto n.6, 2019. [Accessed 19 Dec 2019]. Available from: [https://www.epicentro.iss.it/arbovirosi/pdf/Chik\\_2019.pdf](https://www.epicentro.iss.it/arbovirosi/pdf/Chik_2019.pdf)
  14. Latt NN, Myat Cho S, Htun NM, Myint MN, Aoki F, Reyer JA, et al. Yu Mon Saw. Healthcare in Myanmar. *Nagoya J Med Sci*. 2016;78(2):123-34. PMID: 27303099
  15. Ngwe Tun MM, Inoue S, Thant KZ, Talemaitoga N, Aryati A, Dimaano EM, et al. Retrospective seroepidemiological study of chikungunya infection in South Asia, Southeast Asia and the Pacific region. *Epidemiol Infect*. 2016;144(11):2268-75. <https://doi.org/10.1017/S095026881600056X> PMID: 27018566
  16. Pulmanusahakul R, Roytrakul S, Auewarakul P, Smith DR. Chikungunya in Southeast Asia: understanding the emergence and finding solutions. *Int J Infect Dis*. 2011;15(10):e671-6. <https://doi.org/10.1016/j.ijid.2011.06.002> PMID: 21775183
  17. World Health Organization (WHO). Prevent Dengue & Chikungunya, WHO Myanmar newsletter special; 9 September 2019. Available from: [https://www.who.int/docs/default-source/searo/myanmar/prevent-dengue-and-chikungunya.pdf?sfvrsn=9b1b3069\\_0](https://www.who.int/docs/default-source/searo/myanmar/prevent-dengue-and-chikungunya.pdf?sfvrsn=9b1b3069_0)
  18. Department of Disease Control. Weekly Disease Forecast No.234\_Chikungunya (3-9 November 2019). Department of Disease Control, Ministry of Public Health. [Accessed 08 Jan 2020]. Available from: [https://ddc.moph.go.th/uploads/files\\_en/28420191219085018.pdf](https://ddc.moph.go.th/uploads/files_en/28420191219085018.pdf)
  19. Javelle E, Florescu S-A, Asgeirsson H, Jmor S, Eperon G, Leshem E, et al. Increased risk of chikungunya infection in travellers to Thailand during ongoing outbreak in tourist areas: cases imported to Europe and the Middle East, early 2019. *Euro Surveill*. 2019;24(10):1900146. <https://doi.org/10.2807/1560-7917.ES.2019.24.10.1900146> PMID: 30862335
  20. World Tourism Organization. Myanmar: Country-specific: Basic indicators (Compendium) 2013-2017 (11.2018). [Accessed 24 Oct 2019]. Available from: <https://www.e-unwto.org/doi/abs/10.5555/unwtotfb0104010020142018201909>
  21. Medlock JM, Avenell D, Barrass I, Leach S. Analysis of the potential for survival and seasonal activity of *Aedes albopictus* (Diptera: Culicidae) in the United Kingdom. *J Vector Ecol*. 2006;31(2):292-304. [https://doi.org/10.3376/1081-1710\(2006\)31\[292:AOTPFS\]2.0.CO;2](https://doi.org/10.3376/1081-1710(2006)31[292:AOTPFS]2.0.CO;2) PMID: 17249347
  22. Appassakij H, Khuntikij P, Silpapojakul K, Promwong C, Rujirojindakul P, Suddeaugrai O, et al. Risk of chikungunya virus transmission associated with European travelers returning from southern Thailand (2008-2015). *Transfusion*. 2019;59(8):2612-21. <https://doi.org/10.1111/trf.15401> PMID: 31228360
  23. Díaz-Menéndez M, Crespillo-Andújar C. Literature review of mosquito-borne viral infections in non-tropical European Union territories: A cause of concern? *Enferm Infecc Microbiol Clin*. 2019;37(9):619-20. <https://doi.org/10.1016/j.eimc.2018.12.002> PMID: 30594320
  24. Pastorino B, Bessaud M, Grandadam M, Murri S, Tolou HJ, Peyrefitte CN. Development of a TaqMan RT-PCR assay without RNA extraction step for the detection and quantification of African Chikungunya viruses. *J Virol Methods*. 2005;124(1-2):65-71. <https://doi.org/10.1016/j.jviromet.2004.11.002> PMID: 15664052

## License, supplementary material and copyright

This is an open-access article distributed under the terms of the Creative Commons Attribution (CC BY 4.0) Licence. You may share and adapt the material, but must give appropriate credit to the source, provide a link to the licence and indicate if changes were made.

Any supplementary material referenced in the article can be found in the online version.

This article is copyright of the authors or their affiliated institutions, 2020.