



Correction to: Point estimate and reference normality interval of MRI-derived myocardial extracellular volume in healthy subjects: a systematic review and meta-analysis

Francesco Sardanelli^{1,2} · Simone Schiaffino¹ · Moreno Zanardo³ · Francesco Secchi¹ · Paola Maria Cannò¹ · Federico Ambrogi⁴ · Giovanni Di Leo¹

Published online: 5 July 2019
© European Society of Radiology 2019

Correction to: European Radiology

<https://doi.org/10.1007/s00330-019-06185-w>

The original version of this article, published on 02 May 2019, unfortunately contained a mistake. The following correction has therefore been made in the original: The presentation of Fig. 2 was incorrect. The corrected figure is given below.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The online version of the original article can be found at <https://doi.org/10.1007/s00330-019-06185-w>

✉ Moreno Zanardo
moreno.zanardo@unimi.it

¹ Radiology Unit, IRCCS Policlinico San Donato, via Morandi 30, 20097 San Donato Milanese, Milan, Italy

² Department of Biomedical Sciences for Health, Università degli Studi di Milano, Via Morandi 30, 20097 San Donato Milanese, Milan, Italy

³ Integrative Biomedical Research, Università degli Studi di Milano, Via Mangiagalli 31, 20133 Milan, Italy

⁴ Department of Clinical Sciences and Community Health, Università degli Studi di Milano, Via Vanzetti 5, 20133 Milan, Italy

Fig. 2 Forest plot of the 56 analyzed studies, for a total of 64 independent study parts. Heterogeneity among studies was very high ($I^2 = 92\%$). The last row shows the pooled extracellular volume obtained using the random effects model

