

1           Reconsidering the compound effect of geomorphology, vegetation, and climate  
2           change on paleopedogenesis in sensitive environments (Northern Apennines, Italy)

3

4           Masseroli A.<sup>1</sup>, Villa S.<sup>2</sup>, Mariani G.S.<sup>3</sup>, Bollati I.M.<sup>1</sup>, Pelfini M.<sup>1</sup>, Sebag D.<sup>4,5,6</sup>, Verrecchia E.P.<sup>4</sup>, Trombino

5

L.<sup>1</sup>

6

7           1 Department of Earth Sciences “A. Desio”, Università degli Studi di Milano, Milano, Italy;

8           2 Dipartimento di Scienze Agrarie e Ambientali – Produzione, Territorio, Agroenergia

9           (DiSAA), Università degli Studi di Milano, Milano, Italy;

10          3 Dipartimento di Scienze Chimiche e Geologiche, Università degli Studi di Cagliari, Cagliari, Italy;

11          4 Institute of Earth Surface Dynamics, Faculty of Geosciences and the Environment, Université de Lausanne,

12          Switzerland;

13          5 Normandie University, UNIROUEN, UNICAEN, CNRS, M2C, 76000 Rouen, France.

14          6 IFP Energies Nouvelles (IFPEN), Direction Géosciences, Rueil-Malmaison, France

15

16          Corresponding author:

17          [anna.masseroli@unimi.it](mailto:anna.masseroli@unimi.it)

18          <http://orcid.org/0000-0002-9845-2608>

19          Earth Science Department "A. Desio", Università degli Studi di Milano

20          Via Mangiagalli, 34; 20133 Milan - ITALY

21

22          **Abstract**

23          Complex sequences of paleosols are often formed by the interaction between pedogenesis and  
24          geomorphological evolution. Their study, particularly in mountain areas, is useful to reconstruct past  
25          environmental conditions as well as climate shifts, and to gather information on the morphodynamical  
26          processes affecting the landscape through time.

27          Since the combined role that all different factors can play in the soil formation and evolution through time  
28          and space influences the formation and evolution of those complex paleosol sequences, a multidisciplinary  
29          study was conducted at the NW slope of Mt. Cusna (Northern Apennines, Italy). This work aims to  
30          reconstruct and to evaluate how the interactions between the geomorphological context, the Holocene  
31          climate variations, and the modification of the vegetation cover and composition influence the soil  
32          development of this area.

33          A combination of routine soil analyses (i.e., grain-size distributions, total organic carbon, total nitrogen, pH,  
34          and Fe/Al extractions), soil micromorphology and the Rock-Eval<sup>®</sup> pyrolysis allowed to characterize and to  
35          correlate the different soil units constituting a toposequence of six soil profiles.

36          The presence of different pedological units that can be correlated along the slope underlines the occurrence  
37          of separate events of pedogenesis, spatio-temporally linked to recognizable stability phases at slope scale.  
38          These phases of biostasy, characterized by vegetation cover and soil development, alternate to phases of  
39          rhexistasy, characterized mainly by slope instability (i.e., aggradation/degradation).

40          In detail, in the Mt. Cusna toposequence three different soil units, linked to three different stability phases,  
41          have been identified: the earliest stability phase, characterized by the presence of well-developed Luvisols,

42 the subsequent stability phase typified by less expressed Luvisols, and the ongoing stability phase with  
43 Leptosols. This latter pedogenetic phase, in some cases, is superimposed to the previous one, so affecting the  
44 exhumed paleosols.

45 In this light, the Mt. Cusna toposequence characterization allowed to enlighten the complexity of soil  
46 polygenesis in higher detail than the previous studies, not only reconstructing the past environmental  
47 conditions but also inferring the succession of phases of slope stability and phases characterized by erosion  
48 and deposition processes.

49

## 50 **Keywords**

51 Complex paleosols, paleosols sequences, Rock-Eval<sup>®</sup> pyrolysis, soil micromorphology, slope dynamic,  
52 Northern Apennines

53

## 54 **1. Introduction**

55

56 Soil evolution in active landscapes, such as mountain environments, is mainly influenced and controlled by  
57 topography (Zanini et al., 2015). Slope dynamics and instability influence soil formation, development, and  
58 preservation: conditions of slope instability can dramatically impact on soil formation and conservation in  
59 both short and long terms (Bollati et al., 2019; Coltorti et al., 2019). Areas characterized by steep slopes are  
60 affected by frequent and, often, rapid mass movements related to gravity and water-driven processes, which  
61 are able to substantially disrupt the relief and modify the old surfaces and dismantling previous soils and  
62 paleosols (Dewolf and Bourrié, 2008). Slope processes can vary in frequency and intensity under changing  
63 climate, environmental conditions and anthropogenic influence, resulting in a succession of rhexistasy and  
64 biostasy phases (Erhart, 1967). Consequently, pedogenesis can be variously impaired over time, whether  
65 interrupted, and soils can be buried by deposition phases, or get partially or completely eroded. Moreover,  
66 during the successive stability phases, soil formation processes can restart under new environmental  
67 conditions. In this light, the result of this tight interaction between pedogenesis and geomorphological  
68 evolution is the formation of complex sequences of paleosols, which formed in different morphoclimatic  
69 environments associated to distinct paleosurfaces (Fedoroff et al., 1990; Coltorti and Pieruccini, 2006;  
70 Vittori Antisari et al., 2016). Therefore, an exhaustive investigation of such sequences of paleosols is useful  
71 to reconstruct past environmental conditions as well as climate shifts, and to gather information on the  
72 morphodynamic processes affecting the landscape through time (Ruellan, 1971; Magliulo et al., 2006;  
73 Sheldon and Tabor, 2009). This proxy, evidence of environmental changes recorded in soils and paleosols, is  
74 often used as a paleoenvironmental tool in the mountain areas (Kaiser et al., 2007; D'Amico et al., 2016),  
75 and specifically in the Apennines (Giraudi, 2005; Coltorti and Pieruccini, 2006; Magliulo et al., 2006).

76

77 In the Northern Italian Apennines, the area of Mt. Cusna has been investigated with an array of studies based  
78 on various approaches in order to reconstruct its climate and environmental history through the Holocene.  
79 Multidisciplinary paleoenvironmental studies carried out at the treeline (Compostella et al., 2013; 2014)  
80 helped in characterizing the climate history of the soils in the area. In addition, geoarchaeological  
81 investigations of Mesolithic sites allowed the past environmental conditions of the area to be described using  
82 soil data, archaeological evidences, and palynological studies (Biagi et al., 1980; Cremaschi et al., 1984).  
83 Two geomorphological maps, within a time distance of 25 years (Panizza et al., 1982; Mariani et al., 2018),  
84 were made with the aim of reconstructing the geomorphological evolution of the area through the  
85 representation of landforms and paleosurfaces and their reciprocal distribution. However, in the Mt. Cusna  
86 area, no studies have been focused yet on the combined role that all different soil formation factors (Jenny,  
87 1941) could have played in soil formation and evolution through time and space.

88

89 Therefore, this work aims at the characterization and the correlation of different soil units constituting a  
90 toposequence (Milne, 1936) of six soil profiles at the NW slope of Mt. Cusna, by means of a combination of  
91 routine soil analyses (i.e., grain-size distributions, total organic carbon, total nitrogen, pH, and Fe/Al  
92 extractions), soil micromorphology and a non-conventional approach to interpret soil organic matter  
93 dynamics: the Rock-Eval<sup>®</sup> pyrolysis. Moreover, we focused on the information recorded in soils to try to  
94 reconstruct and evaluate how the interactions between the geomorphological context, the Holocene climate  
95 variations, the vegetation change and human impact influence the formation and evolution of the studied  
96 complex paleosol sequences.

97

## 98 2. Materials and methods

### 99 2.1. Geological, geomorphological and soilscape settings of the study area

100 The study area is located on the NW slope of Mt. Cusna (2120 m a.s.l.; Fig. 1a), the second highest peak of  
101 the Northern Italian Apennines. Mt. Cusna is located in the territory of Febbio (Emilia Romagna region),  
102 inside the “*Parco Nazionale dell’Appennino Tosco-Emiliano*” (Tuscan-Emilian Apennine National Park).

103 The climate is sub-Mediterranean with abundant and well distributed precipitation (2000 mm/y), with a  
104 summer minimum (Compostella et al., 2014). Mean annual temperatures range from 8.8 °C (Ligonchio, 928  
105 m a.s.l., 44°31’N-10°35’E) to 2.2 °C (Mt. Cimone, 2165 m a.s.l., 44°21’N-10°70’E; observation period for  
106 both stations 1961-1990). The study area, located between 1600-1700 m a.s.l., is slightly below the current  
107 treeline position (1750 m a.s.l., Compostella et al., 2013), and it is characterized by an open deciduous forest  
108 dominated by beech (*Fagus sylvatica*). Sparse shrubs and grassland species are also present, mainly  
109 *Vaccinium myrtillus*, *Juniperus nana*, *Thymus* sp, and *Laburnum alpinum*.

110 The bedrock consists mainly of turbiditic sandstones (locally marlstones) with intercalated sequences of  
111 claystones and siltstones (Panizza et al., 1982; Bortolotti, 1992). This area was diffusely subject to glacial  
112 and periglacial processes during the last glacial phases as testified by the presence of cirques and till deposits

113 in the surroundings and by the general rounded and hilly aspect of the slopes (Losacco, 1949, 1982; Mariani  
114 et al., 2018). During the Holocene, the most widespread processes are due to gravity and water runoff  
115 (Panizza et al., 1982; Mariani et al., 2018). The Mt. Cusna area is affected by extremely active slope  
116 morphodynamics (Bertolini and Pellegrini, 2001) as demonstrated by the presence of rock and debris slides  
117 on the slopes of the main ridges, with varying dimensions and positions (Mariani et al., 2018). Moreover, the  
118 areas of Mt. Cusna, where slopes are more stable and flat, are covered by wider colluvium deposits (Mariani  
119 et al., 2018).

120

121 In the study area, processes related to surface running water play also an important role in shaping the  
122 landforms, in different ways, according to the substrate. Runoff and wash out phenomena have low intensity  
123 on sandstone outcroppings, due to their semipermeable property. On the contrary, where claystones and  
124 marlstones outcrop, water runoff often exposes surfaces due to their mostly impermeable property, and large  
125 washout areas are characterized by the presence of pseudo-gullies (Mariani et al., 2018).

126

127 Given the widespread presence of degradation processes that have deposited substantial amounts of  
128 reworked sedimentary material and have eroded surfaces, the soil landscape is directly affected by the  
129 morphological conditions and evolution of the area. Consequently, around the study area, Entisols are found  
130 on active morphologies and mainly on claystones, Inceptisols developed on more stable surfaces, mainly on  
131 sandstones, whereas Spodosols are located at higher altitudes (Panizza et al., 1982). A detailed pedological  
132 map of the area is missing, but the 1:250000 soil map of Italy (Carta Ecopedologica d'Italia 1:250000,  
133 Servizi WMS, Geoportale Nazionale, <http://www.pcn.minambiente.it/mattm/servizio-wms/>) emphasizes the  
134 presence of Regosols or Cambisols (IUSS Working Group WRB, 2015).

135

136 Moreover, in the Mt. Cusna area traces of older soil formation, in the form of relict or buried paleosols, are  
137 also found below colluvial deposits. In particular, the most important paleosols associated to the Mt. Cusna  
138 paleosurface are located on the northern slope of Mt. Cusna (Panizza et al., 1982). These paleosols have been  
139 described as Taptho-Luvisols (Compostella et al., 2014 according to Krasilnikov and Calderón, 2006): these  
140 are mature soils, mainly subject to clay illuviation and with well differentiated horizons.

141

142 The first traces of human settlements in the area belong to Mesolithic hunters, between the Early and the  
143 Mid-Holocene (Mt. Bagioletto site, 1.6 km N far from the summit of Mt. Cusna; Panizza et al., 1982).  
144 Sporadic occupation is recorded from the Late Holocene to the Roman Age (Biagi et al., 1980; Panizza et al.,  
145 1982; Cremaschi et al., 1984). Later on, historical sources show a progressive colonization of the higher  
146 northern Apennines since the Late Middle Ages (Mariani et al., 2019a), with communities surviving on  
147 livestock and forest exploitation. Agriculture played a minor role and was limited to small patches nearest to  
148 settled villages (Panizza et al., 1982). During the last few centuries, farming has been mostly relegated below

149 1000-1300 m a.s.l., while higher altitudes were used for extensive utilization of the forests for wood and  
150 charcoal production; pastures occupied the highest altitudes, up to the Mt. Cusna peak.

151

## 152 2.2. Soil sampling

153 Six soil profiles were dug and described according to Jahn et al. (2006; Table 1) along an altitudinal transect  
154 on the NW slope of Mt. Cusna (Fig. 1b). Five soil profiles were chosen in an area as they were affected by  
155 running water erosion, whereas one soil profile (04) was excavated in a stable area with a forest plant cover.  
156 The locations of the five profiles were chosen on high topographic position, currently preserved from erosion  
157 processes. The coordinates of each profile were recorded using a GPS device. Between 0.5 to 2 kg of  
158 material were sampled from all identified soil horizons for laboratory analyses (Avery and Bascomb, 1982;  
159 Gales and Hoare, 1991; Cremaschi and Rodolfi, 1991). Seven undisturbed samples were collected using  
160 Kubiěna boxes (Kubiěna, 1953) from selected soil horizons to obtain thin sections for micromorphological  
161 investigations.

162

## 163 2.3 Soil analyses

164 Air-dried soil samples were treated by wet sieving in order to separate skeleton particles (> 2mm) from the  
165 fine earth. The pH (in 1:2.5 soil: distilled water), total Organic Carbon (Org. C) content (Walkley and Black,  
166 1934), and total nitrogen content (Kjeldahl, 1883) were measured for each soil sample. Particle size  
167 distributions were obtained on samples pretreated with H<sub>2</sub>O<sub>2</sub> (130 volumes); sand fractions (from 2000 to 63  
168 μm) were collected by sieving, while silt and clay particles (< 63 μm) were measured by aerometry with the  
169 Casagrande aerometer (Casagrande, 1934).

170

171 Different iron and aluminium forms were quantified (Ministero delle Risorse Agricole Alimentari e  
172 Forestali, 1994). Non-silicate forms (“free” iron, Fe<sub>d</sub> and Al<sub>d</sub>) were extracted with a bicarbonate-dithionite-  
173 citrate buffer, iron and aluminium in amorphous oxides and hydroxides (“active” forms, Fe<sub>o</sub> and Al<sub>o</sub>) were  
174 extracted with acid ammonium oxalate and, iron and aluminium bound to organic matter with covalent or  
175 partially polar bond (Fe<sub>p</sub> and Al<sub>p</sub>) were extracted in a solution of sodium pyrophosphate. For all forms, the  
176 amount of solubilized iron and aluminium in the supernatant was determined by means of a 4100 MP-AES  
177 (Agilent), after appropriate dilutions. Data with a %RSD (Relative Standard Deviation) of concentration >  
178 3.5 and/or with a not detectable clear peak, were considered invalid (n.d. in Table 2), while the data close to  
179 the detection limit of the instrument were approximated to the minor concentration detectable (<n in Table  
180 2).

181

182 In order to compare the results of iron and aluminium extractions to soil characteristics, both iron activity  
183 index (Fe<sub>o</sub>/Fe<sub>d</sub>) (Rhodes and Sutton, 1978) and illuviation (podzolization) index (Al<sub>o</sub>+1/2Fe<sub>o</sub>) were calculated  
184 (IUSS Working Group WRB, 2015). Moreover, the amount of crystalline iron oxides (Fe<sub>cry</sub>) was calculated

185 as the difference between the dithionite- and the oxalate-extractable Fe ( $Fe_{cy} = Fe_d - Fe_o$ ) (Bascomb, 1968;  
186 Pawluk 1972; Cremaschi and Rodolfi, 1991; Zanelli et al., 2007).

187

188 Organic Matter (OM) analysis was performed using a Rock-Eval<sup>®</sup> 6 pyrolyser (Vinci Technologies, France).  
189 About 60 mg of crushed material, previously sieved (<2mm), was analyzed for each horizon. Total Organic  
190 Carbon (TOC), Mineral Carbon (MINC), Hydrogen (HI) and Oxygen (OI) Indices were calculated by  
191 integrating the amounts of Hydrocarbon Compounds (HC), CO, and CO<sub>2</sub> produced during thermal cracking  
192 of OM and oxidative decomposition of carbonate, between defined temperature limits (Lafargue et al., 1998;  
193 Behar et al., 2001). The I-index and R-index were computed according to Sebag et al. (2016). By  
194 construction, the R-index relates to the most thermally resistant and refractory pools of organic matter, while  
195 the I-index is related to the ratio between thermally labile and thermally resistant pools (see details in Sebag  
196 et al., 2016). As they arise from a mathematical composition, these two indexes may be inversely correlated  
197 to each other when OM stabilization results from progressive decomposition of organic components  
198 according to their biogeochemical stability. Previous studies show that this property is verified with both  
199 indices highly correlated along a constant line ("humic trend") in composts and undisturbed soil profiles  
200 (Albrecht et al., 2015; Matteodo et al., 2018; Schomburg et al., 2018, 2019; Sebag et al., 2016). For  
201 comparison, we used the Matteodo's dataset composed of 46 soil profiles selected across various ecounits in  
202 Swiss Alps (Matteodo et al., 2018). The "humic trend" equations in the I-index/R-index plot was calculated  
203 starting from both Matteodo's dataset and study area dataset (colored dots in Fig. 3b).

204 Finally, as shown in previous studies (Malou et al. 2020; Thoumazeau et al., 2020), a Delta I-index can be  
205 calculated: it refers to the difference between the I-index value of each sample and the I-index value  
206 calculated with the "humic trend" equation (in bold in Fig. 3b), calculated starting from study area sample  
207 data, at the R-index value of each sample.

208

## 209 2.4 Micropedology

210 Uncovered soil thin sections were prepared from undisturbed samples through impregnation with  
211 polystyrene. Thin sections were then observed by means of a petrographic microscope (Leica Laborlux 18  
212 POL), in parallel (PPL), cross-polarized (XPL), and oblique incident light (OIL), using different objectives  
213 (1.6x, 4x, 10x and 25x). Thin sections were described according to Stoops (2003). The interpretation of thin  
214 sections was performed according to Stoops et al. (2018).

215

## 216 3. Results

### 217 3.1. Soil profiles description and analyses

218 All the soil profiles are located between 1600 and 1700 m a.s.l. (Fig. 1b); five soil profiles (01; 02; 03; 05;  
219 06) are located in areas affected by running water erosion (Table 1). The vegetation cover is composed of  
220 semi-deciduous shrubs in all the area of soil profiles, except for profile 04 characterized by beech forest.

221

222 Total depths for all profiles range from around 50 to about 200 cm. The thickest profiles are characterized by  
223 the presence of two (01) or three (02, 06) distinct soil units, identifiable in the field by the presence of grain-  
224 size discontinuities or buried organic horizons, and/or a color change. Soil structure is moderately expressed  
225 by granular or subangular blocky aggregates. Sometimes, surface horizons (i.e., 05 A, 06 O and 06 OA)  
226 exhibit only a single-grained structural condition. On the contrary, well separated angular blocky pedes can be  
227 found in buried horizons. Colors range between 10YR and 2.5Y in their hue, with a tendency for chroma to  
228 increase with depth inside each single soil unit (see Appendix A for detailed data). All the profiles are  
229 characterized by acidic conditions ranging from pH 4 to 5.6, usually increasing with depth, except for surface  
230 O horizons, which are often less acid than the underlying horizon (Fig. 2).

231 Particle size distributions of the investigated soil profiles display some common traits. Silt is always the  
232 predominant fraction, ranging from 44% to about 65%, with variable amounts of clay (from 21% to 52%).  
233 On the contrary, gravels never exceed 5%, while sands are only rarely > 20% (Fig. 2). Moreover, the grain  
234 size distributions allow the presence of different soil units to be confirmed. Indeed, the 01 and 06 soil  
235 profiles show an increase in clays at the top of the buried units (e.g., in profile 01: clay increases by 13.2%  
236 between horizons BC and 2AB1; in profile 06: clay increases by 13.7% between horizons OA and 2AB1); on  
237 the contrary, the 02 soil profile displays an increasing clay content by 2.6% between the horizons 2AB and  
238 2Btg. 03 and 05 profiles are characterized by a clay increase in the B horizons (i.e., 03OB, 05ABt1 and  
239 ABt2), whereas an increase of coarse materials clearly appears in correspondence of the topographic surface  
240 (03 OA and 05 A) and the deepest (03 BC) horizons (see Appendix B for detailed data and Appendix C for  
241 cumulative particle size distribution curves). Lastly, the 04 soil profile, under a stable forested area, shows a  
242 constant increase of the coarse fractions (gravel and sand) with depth (Fig. 2).

243 The identified discontinuities are also marked by the results from chemical analyses. Indeed, a peak of total  
244 organic C content is found at the top horizon of each buried unit in correspondence of the above-mentioned  
245 discontinuities: 01 2AB1, 06 3AB, 02 2AB, and in the horizon 02 3AB, while in the horizon 06 2AB1 a total  
246 N peak is found (Fig. 2). Conversely, in profile 04, total organic C decreases from the surface horizon with  
247 depth, as expected in conventional soil profiles. The same trend can be observed in profiles 03 and 05, with  
248 the exception of the OB horizon of 03, which shows an increase in total organic C, and the A horizon of 05,  
249 which has a more marked decrease in total organic C than the deeper horizons (Fig. 2).

250 In the analyzed soil profiles, the total N content follows roughly the same trend as the total organic C (Fig.  
251 2), but with significantly lower values. In surface horizons, total organic C ranges between 14.7 g/kg (02)  
252 and over 100 g/kg (04 and 05), while total N content never exceeds 7.0 g/kg. When focusing on superficial  
253 OM-rich horizons, the highest total N contents (about 7 g/kg) are found in 01, 04 and 06, while the lowest  
254 values belong to 02 (about 1.4 g/kg), following the same trends as total organic C.

255 The total content of free iron oxides ( $Fe_d$ ) in four horizons (01 2AB2, 03 OB, 05 ABt2 and 06 2AB1)  
256 exceeds 30 g/kg but this parameter generally ranges between 10 and 20 g/kg (Table 2). The values of  
257 amorphous iron oxides ( $Fe_o$ ) tend to be lower (<0.90 - 15.73 g/kg), as well as iron bound with organic matter

258 ( $Fe_p$ ), which ranges between <3.00 and 15.98 g/kg. Regarding the aluminium content, its values are lower  
259 and less variable than those of iron, with the exception of aluminium bound with the organic matter ( $Al_p$ )  
260 (Table 2). Free aluminium oxides ( $Al_d$ ) reaches abundance from 2.64 to 5.83 g/kg, amorphous aluminium  
261 oxides ( $Al_o$ ) from 0.85 to 5.59 g/kg, and the aluminium bound with the organic matter ( $Al_p$ ) from 2.98 to  
262 12.09 g/kg. The crystalline iron oxides ( $Fe_{crv}$ ) are mainly concentrated in the B horizons (Fig. 2), and the  
263 highest value is found in the 05 ABt2 (21.08 g/kg). Along the profiles, the amorphous iron oxides ( $Fe_o$ )  
264 exhibit an opposite changing compared to crystalline iron oxides (Fig. 2), except for some superficial  
265 horizons (e.g., 06 OA) and in a few buried organic horizons (e.g., 01 2AB2). Indeed, in most of the profiles,  
266 the amorphous iron oxides show the lowest concentrations in Bt, Btg and partially in Bw horizons (Table 2;  
267 Fig. 2).

268 The iron activity index ranges from 0.18 to 0.66. In more detail, its lowest values are found in the Bt, Btg  
269 and Bw horizons (Table 2), in particular, in the B horizons of buried soils (e.g., 01 2Bt, 06 2Bw2 and 06  
270 2Bw3). Instead, the BC horizons show the highest values of the iron activity index (e.g., 02 BC, 03 BC).

271 Finally, the results of the podzolization index ( $Al_o + \frac{1}{2} Fe_o$ ) partially meet the conditions for the presence of  
272 some podzolization processes only in the profile 02 (ABt2 and BC horizons; Table 2).

273 Rock-Eval<sup>®</sup> indices and parameters (Fig. 3; see Appendix B for detailed data) show that all the superficial  
274 organic horizons are plotted at the top left of the HI/OI diagram (Fig. 3a), whereas the buried soils are  
275 located at the bottom right of the diagram. Moreover, in the I/R diagram (Fig. 3b) the superficial organic  
276 horizons correspond to low R-index values and high I-index values and are located at the top left of the I/R  
277 diagram, whereas the organo-mineral and mineral horizons have a R-index values > 0.65 and are located in  
278 the right portion of the diagram. Taking into consideration the organo-mineral and mineral horizons in the  
279 I/R diagram, two different trends can be recognized. A first trend groups the horizons belonging to the  
280 superficial units of the soil profiles, which have I-index values varying between -0.3 and 0 (colored dots in  
281 Fig. 3b), whereas a second trend groups the horizons belonging to the buried units, which have higher I-  
282 index values, ranging from -0.2 to 0.5 (colored diamonds in Fig. 3b), except for horizons 01 2AB1, 2AB2  
283 and 06 2AB1, 2AB2. On the other hand, some horizons from the superficial units are characterized by high I-  
284 index values (02 BC, ABt1 and ABt2; 04 Bt, Btg, AB; 03 BC). Moreover, when comparing R-index with  
285 depth, buried soils evidence a different trend; indeed, the 01 buried soil display R index values higher than  
286 the 02 buried soil, at the same depth and with similar TOC contents (Fig. 4a,b). The presence of two different  
287 trends can also be observed when taking into consideration the I-index/TOC diagram: the horizons belonging  
288 to the first trend show an expected increase of the I-index with TOC and a decrease with depth, whereas the  
289 horizons belonging to the second trend have (i) high I-index values even with low TOC and (ii) I-index  
290 values which are not decreasing with depth (Fig. 4c, d). Finally, horizons belonging to the second trend have  
291 high delta-I values (Fig. 4e), which means that the I-index values are high compared to those expected for  
292 the corresponding R-index values, when referring to the humic trend.

293

### 294 3.2. Soil thin sections micromorphology



295 The micromorphological observations were carried out on four out of the six examined profiles: 01, 02, 04  
296 and 05 (see Appendix D for detailed thin sections descriptions). Horizons 02 ABt1, 01 2Bt and 04 Bt show  
297 some similar features: they all are characterized by granular aggregates of fine material and angular-  
298 subangular blocky aggregates, together with Fe-Mn nodules and well-developed clay coatings. Moreover, in  
299 the 02 ABt1, reworked soil fragments of subangular aggregates of fine material (i.e., pedorelicts *sensu*  
300 Brewer, 1976) with a high degree clay illuviation, and few allochthonous weathered rock fragments are  
301 present (i.e., lithorelicts *sensu* Brewer, 1976). Furthermore, regarding the coarse mineral fraction, there are  
302 similar proportions of sandstone and claystone fragments in 02 ABt1 and 04 Bt, while in the 01 2Bt  
303 sandstone constitutes the prevalent lithology of fragments.  
304 Similarly, 01 2AB1 and 2AB2, 05 ABt1 and ABt2 and 04 Btg horizons show analogous characteristics, such  
305 as their complex microstructure including granules and reddish clayey subangular aggregates, which contain  
306 a high proportion of Fe-Mn nodules (Fig. 5a, b). Moreover, the 01 2AB1 horizon differs from the 2AB2  
307 horizon by the predominant sub-angular aggregates in the latter. Finally, in the 05 ABt1 and ABt2 and 04  
308 Btg horizons show the presence of clay illuviation features (coatings); in the latter horizon, hydromorphic  
309 features in the form of depletion pedofeatures and intercalations are also found.  
310 The micromorphological approach further underlines the presence of peculiar characteristics in 01 A2, BC  
311 and 2AB1 and 02 2Btg horizons. For example, the 02 2Btg horizon exhibits a compact, vughy  
312 microstructure, and, well-developed diversified pedofeatures, such as dense digitate Fe-Mn nodules, typic  
313 clay coatings, typic clay infillings, and Fe-Mn hypoc coatings (Fig. 5c, d).  
314 Otherwise, the 01 A2, BC and 2AB1 horizons show a twofold distribution of crumbs and reworked  
315 subangular aggregates reddish in color, weakly striated and characterized by a high degree of pedogenesis  
316 (i.e., pedorelicts *sensu* Brewer, 1976). These pedofeatures are more common in the BC horizon (Fig. 5e, f).  
317 Finally, 01 2AB1 and 2AB2 contain identifiable pedofeatures only as Fe-Mn nodules, which are more  
318 concentrated inside the subangular reworked soil fragments.

319

## 320 4. Discussion

### 321 4.1. Complex paleosol sequences and the characterization of buried units

322 Sedimentological and chemical data help in identifying the presence and defining the boundaries of the soil  
323 units recognized in the field. Trend anomalies detected in analytical values are found in profiles 01 (between  
324 horizons BC-2AB1), 02 (between horizons BC-2AB and 2Btg-3AB), and 06 (between horizons OA-2AB1  
325 and 2Bw3-3AB; Fig. 2). In horizons 2AB1 of 01, 2AB1 and 3AB of 06, high values of total organic C or  
326 total N contents as well as of fine material, underline the presence of paleosurfaces, subsequently buried by  
327 coarse colluvial deposits, which disconnected the soils from surface pedogenetic processes (see below for  
328 micromorphological evidence). On the contrary, it is not possible to identify precisely in profile 02 the  
329 paleosurface location at the top of the 2AB horizon from the colluvial material above it. Even if the organic  
330 matter content peaks in 2AB horizon, both 2AB and BC horizons possess a high percentage of sand and

331 gravel (Fig. 2). Moreover, both horizons in the field have a homogeneous aspect: it is possible that the  
332 distinction between these two units may be difficult to identify because of a higher energy of deposition of  
333 the colluvium, which mixed part of the materials during the process. In the same profile, the total organic C  
334 content allows a discontinuity at the top of the 3AB horizon to be identified (Fig. 2). The variations in  
335 particle size distributions within profiles 03, 04 and 05 are not attributed to the presence of paleosurfaces.  
336 Micromorphological observations of thin sections from 01 and 02 profiles provide further information  
337 confirming the presence of different soil units and their respective characteristics. In 01, the presence of two  
338 distinct units is highlighted by the nature of the coarse material: in the surface unit (unit I) fragments of  
339 claystones are more abundant while in the deeper one (unit II), fragments of sandstones are more common.  
340 This difference in the coarse fraction lithology likely indicates the occurrence of two different parent  
341 materials, separating clearly two soil units. In unit II, weathering processes occurred in oxidative conditions  
342 with water infiltration, emphasized by the yellowish color of the groundmass indicating the presence of iron  
343 hydroxides (Sauro et al., 2009; Stoops et al., 2010, 2018; Compostella et al., 2014). The presence of clay  
344 illuviation features, indicated by clay coatings found in the 2Bt horizon, requires alternating phases of water  
345 infiltration into the soil, together with conditions of strong vegetative protection of the soil surface, in order  
346 to permit clay translocation into the deep horizons (McCarthy et al., 1998; Stoops et al., 2010, 2018).  
347 Moreover, the presence of Fe-Mn nodules indicates temporary waterlogging conditions inside the soil  
348 (McCarthy et al., 1998; Stoops et al., 2010, 2018), probably amplified by the presence of the underlying 2Bt  
349 horizon. The diffuse and irregular boundary of the nodules witnesses their *in situ* formation, without  
350 evidence of transport from other locations (Fedoroff and Goldberg, 1982). In unit I, frequent blocky peds  
351 show a generally reddish micromass color associated to Fe-Mn nodules, which indicate a degree of  
352 weathering greater than in the surrounding soil groundmass (e.g., reddish micromass, typic nodules of Fe-Mn  
353 and moderately weathered mineral fragments): therefore, these blocky peds can be regarded as pedorelicts  
354 (*sensu* Brewer, 1976), i.e., reworked fragments of an older soil (Fig. 5e, f) redeposited within more recent  
355 horizons (Kemp, 1998; Nicosia, 2006; Rellini et al., 2007; Sauro et al., 2009; Cremaschi et al., 2018).  
356 Moreover, these reworked fragments of paleosol are similar, in terms of fabric, to the 2AB2 horizon; thus, it  
357 is reasonable to state that they were eroded from higher portions of the slope and deposited within the  
358 presently BC horizon, which implies a highly erosive phase.

359 Regarding 02 profile, the micromorphological analysis indicates that the unit II (e.g., 2Btg horizon) is  
360 characterized by a stronger degree of pedogenesis than the unit I (e.g., ABt1 horizon). This is probably due  
361 to a greater intensity and/or duration of a pedogenetic phase. In unit II, clay coatings are clearly visible (Fig.  
362 5c, d) and it is often possible to identify an orientation in the deposition of fine material due to the  
363 development of crescent internal fabric. This corroborates the hypothesis of transport of clay materials from  
364 upper horizons (McCarthy et al., 1998). Moreover, the 2Btg horizon is the only one showing the  
365 development of Fe-Mn hypocoatings, due to longer periods of waterlogging (McCarthy et al., 1998; Stoops  
366 et al., 2010). Finally, only rare reworked paleosol fragments (pedorelicts *sensu* Brewer, 1976), composed of  
367 clayey blocky peds, are found in the ABt1 horizon.

368

## 369 4.2. Correlation of soil units

370 As already described in section 4.1 of the Discussion, the studied toposequence is constituted by six soil  
371 profiles located at different altitudes along the NW slope of Mt. Cusna, chosen according to their  
372 topographic position. Among the analyzed soil profiles, three are composed of different units, whereas only  
373 one soil unit is present in the other three profiles (03; 04 and 05).

374 In the following section, the most widespread unit set (i.e., unit II of 01; unit I of 02; 03; 04; 05; unit II of 06)  
375 is first characterized in order to better correlate the various soil profiles (Fig. 6). Indeed, the unit II of 01  
376 (e.g., 2Bt) and the unit I of 02 (e.g., ABt1), although characterized by a different mineral component, are  
377 associated with the presence of clay illuviation and a very similar microstructure, together with the presence  
378 of moderately impregnated amorphous nodules. Therefore, these units could be formed by the same  
379 pedogenetic event on different parent materials, the latter deposited by distinct colluvial events.

380 Unit II of 01 can also be correlated with 05 profile (e.g., ABt1 and ABt2), which shows similar aggregates  
381 (both subangular blocky and granular) and microstructure with regard to the horizons 2AB1 and 2AB2 (01  
382 profile), as well as clay illuviation features in the horizon 2Bt (01 profile). Moreover, moderate impregnative  
383 amorphous nodules of Fe-Mn are present through the referred horizons. The same consideration regarding  
384 microstructure, clay illuviation features, and amorphous nodules can be extended to the 04 profile, even if, in  
385 the horizon 04 Btg, the presence of hydromorphic features (i.e., depletion pedofeatures and intercalation,  
386 *sensu* Fedoroff and Courty, 2012) underlines a further pedogenetic phase induced by water logging, probably  
387 due to the topographic position.

388 The sedimentological and chemical data allow the above described correlation based on micromorphological  
389 features to be extended (Fig. 6): the unit II of the 01 (e.g., 2AB1 and 2AB2) and 06 (e.g., 2AB1 and 2AB2)  
390 profiles show the same clay high and sand low contents, and similar values of total organic C and total N  
391 contents. On the other hand, 03 profile can be correlated to the unit I of 02, not only because both units are  
392 characterized by a very low total organic C and total N contents, but also for their relative position along the  
393 slope.

394 In two profiles, 01 and 06, the above correlated unit is overlain by soil horizons, pointing to another soil unit  
395 (Fig. 6). This surficial unit, even if it is thin, shows uniform soil characteristics (particle size distributions,  
396 total organic C and total N contents), and could therefore be related to the same pedogenetic phase.

397 Focusing on soil horizons underlying the above correlated unit, the micromorphological observations  
398 highlight a general greater weathering impact. The 02 2Btg horizon (unit II) has very few analogies with the  
399 horizons observed in other profiles: the presence of well-developed pedofeatures (i.e., dense digitate nodules  
400 of Fe-Mn, typic clay coatings, typic clay infillings; and Fe-Mn hypocoatings) testifies a more intense  
401 pedogenetic phase, followed by a secondary hydromorphic phase induced by waterlogging. Moreover, the  
402 peculiarity of this unit II is also underlined by the organic matter thermal signature: in the R-index/depth  
403 plot, horizons of unit II from 02 have a lower R-index compared with the horizons belonging to unit II from  
404 01, which are located at the same depth (Fig. 4b), due to a presence of different proportion of the most

405 refractory organic pool (related to pedogenetic and inherited contributions) in the two units. A similar trend,  
406 though not so marked, is also visible in the R-index/TOC plot (Fig. 4a), where the horizons belonging to unit  
407 II of O2 are separated from the others. Furthermore, unit II of O2 may be correlated to unit III of O6 (Fig. 6).  
408 This deepest unit of O6 is only represented by a single horizon (3AB), but its stratigraphic continuity, in  
409 addition to Rock-Eval<sup>®</sup> parameters, suggest a possible correspondence between this unit and unit II of O2.  
410 Horizon O6 3AB is always located close to the horizons belonging to unit II of O2 in both R-index/TOC (Fig.  
411 4a) and R-index/depth plots (Fig. 4b), underlining similar organic matter dynamics.

412 Regarding Rock-Eval<sup>®</sup> parameters, as discussed above, the HI/OI diagram emphasizes the way horizons  
413 belonging to unit II of O1 (red dots, 2AB1, 2AB2 in Fig. 3a) and unit II of O6 (violet dots, 2AB1, 2AB2 in  
414 Fig. 3a) group in the same set as the present-day surface horizons of all the profiles, while units II of O2 and  
415 III of O6 are located at the bottom right of the plot, probably due to the presence of an OM refractory pool.  
416 These horizon groupings are clearly distinguishable observing the results of cluster analysis (Fig. 7) carried  
417 out on the Rock-Eval<sup>®</sup> indices take into consideration (HI, OI, R-Index, I-Index): in the dendrogram it is  
418 possible to observe how the organic O horizons clearly differ from the organo-mineral and mineral horizons.  
419 Among the organo-mineral and mineral horizons, the horizons belonging to unit II of O1 and unit II of O6  
420 group are grouped in the same set as the present-day surface horizons of all the other profiles (O2,O3,O4 and  
421 O5), while units II of O2 and III of O6 are all grouped together, close to the deepest horizons of the other units  
422 (O1 2BC2; O2 BC, ABt2; O4 Bt, Btg; O6 2Bw3) (Fig. 7). Moreover, the cluster analysis grouped the organo-  
423 mineral and mineral horizons in two different great groups, which can be attributable to a different evolution  
424 trend of organic matter. Indeed, in the I/R diagram (Fig. 3b), the #\*B\* horizons belong to the first set (i.e., O1  
425 2Bw, 2Bt, 2BC1, 2BC2; O2 BC, 2AB, 2Btg, 3AB; O3 BC; O4 AB, Bt, Btg; O6 2Bw1, 2Bw2, 2Bw3, 3AB;  
426 colored diamonds in Fig. 3b; these horizons are not belonging solely to buried units) show a new trend,  
427 parallel to the “*Inherited Organic Matter Trend*” proposed by Sebag et al. (2016). On the other hand,  
428 horizons richer in organic matter (i.e., O1 A1, A2, BC, 2AB1, 2AB2; O2 OB; O3 OA, OB; O4 OA; O5 A,  
429 ABt1, ABt2; O6 OA, 2AB1, 2AB2; colored dots in Fig. 3b; these horizons belong to surface or buried units)  
430 fit the “*Humic Trend*” (Sebag et al., 2016; Matteodo et al., 2018).

431 This new identified trend, grouping #\*B\* horizons with a high degree of weathering observed in both surface  
432 and buried units, suggests the presence of a relationship between these different soil horizons. In these  
433 horizons, as already specified, the presence of features related to pedogenetic processes not in equilibrium  
434 with the present-day environmental conditions, clearly defines these soil horizons as parts of paleosols (i.e.,  
435 relict soils *sensu* Ruellan, 1971). Moreover, this new trend clearly differs from the “*Humic trend*” (Sebag et  
436 al., 2016; Matteodo et al., 2018), mainly for what concerns the I-index values (Fig. 4c,e). Indeed, the  
437 horizons belonging to this trend show high I-index values (even if the TOC content is low; Fig. 4c,e),  
438 emphasizing the relatively low contribution of resistant pool compared to the most labile pools. Similar  
439 results have been obtained when advanced decomposition of OM affects both thermally labile and resistant  
440 pools, as in Arenosols for example (Malou et al. 2020; Romanens et al., 2019). In the context of the present  
441 study, the decomposition of thermally resistant pools can be explained by an earlier phase of pedogenesis

442 and this peculiarity of the trend supports the hypothesis that these soils located along the trend are paleosols.  
443 The presence in most of these horizons of high I-index values allows these horizons to be identified as  
444 former superficial ones, probably influenced in the past by the pedological surface processes involving  
445 organic matter. Moreover, their present day low TOC contents is explained when considering the  
446 geomorphological context of the area. In addition to advanced decomposition of OM, the contribution of  
447 mineral material, especially in these horizons, is due to colluvium deposition episodes affecting the slope.  
448 The presence of these truncated and buried units can be attributed to the interaction of erosive and  
449 depositional processes happening at different times and ultimately led by colluvial dynamics. The remix and  
450 redeposition of this loose material contributed to organic matter dilution, and, at the same time, buried soil  
451 units, isolating the soil by preventing the modification and external contributions to the buried organic  
452 matter.

453 Two type of paleosols can be observed: (i) buried soils (i.e., 01 2Bw, 2Bt, 2BC1, 2BC2; 02 2AB, 2Btg, 3AB;  
454 06 2Bw1, 2Bw2, 2Bw3, 3AB), as considered soil units are buried, and (ii) exhumed paleosols (i.e., buried  
455 soils that have been brought to surface by erosion, *sensu* Cremaschi et al., 2018) (i.e., 02 ABt1, ABt2, BC;  
456 03 BC; 04 AB, Bt, Btg), when considered soil units outcrop at the surface. Therefore, the new observed trend  
457 in the I/R diagram identifies some horizons of maximum weathering in the paleosols, regardless their  
458 morphological position, within the given limits of the study area. This specific paleosols trend in the I/R  
459 diagram represents a potential approach for further investigations regarding new criteria for paleosols  
460 identification.

461 Finally, regarding the iron and aluminium content, all analyzed Bt, Btg and Bw horizons show an increase in  
462 crystalline iron oxides ( $Fe_{cry}$ ), underlining the expression of some pedogenesis at work (Table 2; Fig. 2).  
463 Furthermore, the iron activity index ( $Fe_o/Fe_d$ ) shows a decrement within the most mature Bt, Btg and Bw  
464 horizons, likely emphasizing a stronger weathering (Table 2). Unfortunately, this ratio does not change  
465 significantly when comparing recent soils and paleosols; consequently, it is not wise to use it as a proxy for  
466 soil age (Arduino et al., 1986).

467

### 468 4.3. The role of geomorphological processes in the development of complex 469 pedosequences

470 The presence of different pedological units, and their correlation along the slope, underlines the occurrence  
471 of separate events of pedogenesis, spatio-temporally related to recognizable stability phases at the slope  
472 scale. These phases of biostasy (Erhart, 1967) are characterized by the absence of erosion and/or deposition  
473 on the slope, a vegetation cover, and a soil development. They alternated with phases of rhexistasy (Erhart,  
474 1967), characterized by slope instability. The study area has been affected by various colluvial events, as  
475 testified by trends in coarse materials content and the presence of pedorelicts and lithorelicts (*sensu* Brewer,  
476 1976) in the surface units.

477 Moreover, the phases of rhexistasy strongly control the conditional conservation of soils along the slopes:  
478 higher energy conditions would produce excessive disruption along the slopes in form of increased colluvial

479 and mass wasting events. These events would destroy the soil cover and cause the disappearance of  
480 evidences of previous pedogenetic phases in the area and at the same time inhibit the formation of new soils.  
481 However, even if these multiple phases of slope activity may have locally eroded part of the pre-existing soil  
482 cover (e.g., unit I of 02), at the same time, they locally buried parts of it, preserving some paleosols over time  
483 (e.g., unit II of 01). In addition, the geomorphological evolution of the slope through time may also brought  
484 previously buried paleosols to the surface again as a consequence of erosion, superimposing later a new  
485 pedogenetic phase on pre-altered materials, as attested by unit I of 02 and probably 03 and 05. In this sense,  
486 slope morphodynamics do not seem to influence directly the intensity of *soil formation processes*, but they  
487 act as a key factor controlling the distribution and occurrence of soil units and paleosols. Therefore, the  
488 development of the studied complex pedosequences is not the exclusive result of vertical top-down processes  
489 but of a number of complex processes including near-surface processes such as material sedimentation and  
490 erosion.  
491 On account of this, in the study area the evidence of different colluvial deposits in these pedosequences  
492 shares similarities with cover-beds successions (Kleber and Terhorst, 2013), thought at a much smaller scale  
493 and magnitude.

494

#### 495 4.4. Reconstruction of the environmental changes during the Holocene along the Mt. 496 Cusna NW slope

497 From the pure pedogenetic point of view, three separate phases have been recorded inside the soil units all  
498 along the slope (Fig. 6). These three pedogenetic phases are preceded by another pedogenetic phase, which  
499 will not be discussed because, since it is represented by an only one horizon (02 3AB) and it is the first time  
500 that it is found, there is not enough information to carry out a detailed reconstruction of the pedogenetic  
501 processes that have affected it. From the oldest to the most recent, these three separate phases are identified  
502 as:

503 -  *$\alpha$  pedogenesis*: it is observed exclusively in unit II of 02 (and partially in unit III of 06), and displays the  
504 characteristics of a well-developed brunification with clay illuviation (Duchaufour, 1983), high Fe<sub>d</sub> content  
505 and clay increase in Btg horizon. This strongly expressed pedogenetic phase led to the formation of a Luvisol  
506 (IUSS Working Group WRB, 2015), developed under a forest vegetation cover, as evidenced by the  
507 yellowish brown matrix with speckled striated b-fabric (Douglas and Thompson, 1985) and by the  
508 pedofeatures (i.e., frequent typic clay coatings; rare crescent typic clay coatings; very few typic clay  
509 infillings) observed in thin sections. This phase was interrupted by a sudden deposition of colluvial material,  
510 mobilized upslope by water runoff. During this phase, the soil surface of the evolving soil was buried,  
511 interrupting its pedogenesis. During this runoff period, the slope was not likely covered by forests, as they  
512 would have effectively prevented such mass movements. Consequently, this colluvial event provided a new  
513 parent material for further soil development during the following phase;

514 -  *$\beta$  pedogenesis*: it is characterized by a moderately developed brunification with some clay illuviation (i.e.,  
515 unit I of 02, unit II of 01, 05, and probably 03 and unit II of 06) in different parent materials. The presence of

516 different parent materials is probably the results not only of various colluvial events, but also of a  
517 differentiated material deposition, both in quantity and composition, due to variable distance from the source  
518 and slope characteristic (e.g., steepness). The pedological characteristics of this second phase, comparatively  
519 to the first phase but in a lesser extent, point to the formation of Luvisols (IUSS Working Group WRB,  
520 2015) under a stable forest cover. Similarly, a second colluvial event interrupted this pedogenetic phase.  
521 However, in this case, the deposited material is more homogeneous, as it is mainly composed of claystone  
522 fragments (including lithorelicts *sensu* Brewer, 1976) and pedorelicts (*sensu* Brewer, 1976), coming from the  
523 B horizons of Luvisols developed during the former pedogenetic phase, then eroded and reworked by  
524 geomorphological processes.

525 - *γ pedogenesis*: the characteristics of the present-day pedogenetic phase suggest a slight change in  
526 conditions compared to the previous two phases. The values of the  $Al_o + \frac{1}{2} Fe_o$  index calculated for ABt2  
527 and BC horizons of 02 profile seem to evidence a weak podzolization (Do Nascimento et al., 2008;  
528 Waroszewski et al., 2013; IUSS Working Group WRB, 2015). However, this process is not recognizable in  
529 the field, and is therefore qualified as cryptopodzolization, forming a “ranker cryptopodzolique” soil (*sensu*  
530 Duchaufour, 1983) or a Leptosol (protosodic) (IUSS Working Group WRB, 2015), as already observed in  
531 other soils of the Mt. Cusna (Mariani, 2016). This process could be favored by the development of low shrub  
532 vegetation dominated by acidophilic species, e.g. *Vaccinium myrtillus* (Duchaufour, 1983; Chersich et al.,  
533 2007; Compostella et al., 2013; Mariani, 2016). As far as the other profiles are concerned, under the same  
534 vegetation cover, an even less conspicuous process than cryptopodzolization can act, inducing the formation  
535 of a “ranker subalpin” soil (Duchaufour, 1983) or an Umbric Leptosol (IUSS Working Group WRB, 2015),  
536 except for the 04 profile, located downslope in a forest context, which can be regarded as a “ranker brunifié”  
537 soil (Duchaufour, 1983) or a Brunic Leptosol (IUSS Working Group WRB, 2015).

538 Finally, in some cases, the present-day pedogenesis is superimposed to the previous one (see section 4.2 of  
539 the Discussion), the latter being identifiable by relict and textural pedofeatures (see section 4.2 of the  
540 Discussion), emphasizing a greater inertia (Duchaufour, 1983) than the ongoing pedofeatures development.

541

#### 542 4.5. Interactions between the geomorphological context, the Holocene climate 543 variations, and the vegetation changes as influencing factors on paleosols

544 The study area is characterized during the Holocene by alternating phases of rhexistasy and biostasy (Erhart,  
545 1967). The pedogenesis has been mainly affected by three factors throughout time (i.e., climate, vegetation,  
546 slope features), which strongly influenced the environmental evolution of the area. In Fig. 8, a sketch on the  
547 role of the main factors influencing the soil development through time in the study area is proposed and  
548 herein discussed. The most ancient phase, recorded by the paleosols, was a biostasy phase (*α* pedogenesis)  
549 during which the climate was probably warm with a stable forest vegetation cover at the Mt. Cusna NW  
550 slope. Soil processes were mainly governed by a well-developed brunification, together with clay illuviation.  
551 Even if the time control of the pedogenetic phases at Mt. Cusna area remains speculative, it is likely that this  
552 pedogenetic phase took place in the Early Holocene (see below). Subsequent climate changes and their

553 associated vegetation shift downward the slope caused a phase of strong instability, marked by the action of  
554 colluvial processes and the interruption of pedogenetic processes. At a regional scale, in the Italian  
555 Apennines, a relationship can be drawn between cold periods and an increase in mass wasting events  
556 (Bertolini, 2007). During such a phase, slope processes eroded soils and deposited un-weathered mineral  
557 material at the topographic surface. The change to warmer temperatures promoted a new stable phase ( $\beta$ ),  
558 characterized by similar environmental conditions observed during the previous warm period. According to  
559 Cremaschi et al., (1984) and Compostella et al. (2013), this phase could correspond to the Early/Middle  
560 Holocene, since in the neighboring area, the charcoals coming from the upper part of soils developed during  
561 the  $\beta$  pedogenesis were radiocarbon dated (3920-3700 cal yr BP: Compostella et al., 2013) after the  
562 Middle/Late Holocene boundary (Mariani et al., 2019b). The difference in soil development grade, in respect  
563 to the previous stable phase, can be attributed to the different duration of pedogenesis or to a different  
564 vegetation cover, as testified by the variation in organic material characteristics highlighted by Rock-Eval<sup>®</sup>  
565 parameters. Such change in vegetation was likely driven by human modifications, as in many other parts of  
566 Italy and the Mediterranean during this period (Cremaschi and Nicosia, 2010; Regattieri et al., 2019; Sevink  
567 et al., 2019). This stability phase was interrupted by slope instability, due to a likely climate deterioration  
568 (probably during the LIA: Mariani et al., 2019b; Zerboni et al., 2019) associated to a loss of the vegetation  
569 cover, again due to the increased pastoral impact at higher elevations since the Middle Ages (Panizza et al.,  
570 1982), coupled with the exploitation of the forest below for charcoal production. However, colluvial  
571 processes during this latter phase of rhexistasy seem to have been less intense than the previous one, with  
572 soils partially eroded, and in some case, with previously buried paleosols exhumed. Moreover, transported  
573 and deposited material was often pre-altered, as it was originating from paleosols developed in higher  
574 topographic positions. Finally, the present-day pedogenesis starts from these colluvial deposits and/or from  
575 the exhumed paleosols. The present-day phase of biostasy ( $\gamma$ ) is characterized by distinctive environmental  
576 conditions with a different vegetation cover (i.e., shrubs): present-day soils are apparently less developed,  
577 possibly because of the vegetation type (i.e., cryptopodzolization induced by *Vaccinium myrtillus*), colder  
578 conditions, or a short duration of pedogenesis (Duchaufour, 1983; Compostella et al., 2014). The sparse  
579 vegetation cover, mainly composed of shrubs and apparently not subject to reforestation, does not protect the  
580 soil enough from water driven erosion acting today, triggering a soil cover erosion along the edge of rills.

581

## 582 5. Conclusions

583 This paper presents the assessment of the different environmental conditions that have affected the NW slope  
584 of Mt. Cusna, based on the analysis of soils and paleosols. In the Mt. Cusna toposequence, three different  
585 soil units have been identified: (i) a first (and the most ancient) soil unit, characterized by a well-developed  
586 brunification with clay illuviation, (ii) a second unit, characterized by the same processes of the first unit, but  
587 less intense, and (iii) a third recent unit, presenting a weak pedogenesis. This latter, in some case,  
588 superimposed on an older truncated soil (paleosol), affecting these exhumed paleosols. Based on the study of  
589 these soil units, it has been possible to reconstruct environmental changes that affected the Mt. Cusna slope



590 during the Holocene, to identify the succession of phases of slope stability, during which soils developed,  
591 and phases characterized by erosion and deposition processes.

592 In this light, soils proved to be a useful archive, not only to reconstruct the past environmental conditions,  
593 but also to trace the geomorphological processes that affected the area. In addition, it has been possible to  
594 identify the role of the geomorphological processes that have affected the evolution of complex paleosol  
595 sequences. This work, based on different lab techniques, demonstrated that a multi-analytical approach is  
596 necessary to properly characterize soils and their genetic pedological processes; for example, the use of  
597 Rock-Eval<sup>®</sup> pyrolysis improved substantially our knowledge about the relationship between paleosols and  
598 organic matter and, as such, opens new research avenues in paleopedogenesis.

599

## 600 Acknowledgments

601 Rock-Eval<sup>®</sup> is a trademark registered by IFP Energies Nouvelles. The authors thank the staff at the  
602 University of Lausanne (Switzerland) for completing the Rock-Eval<sup>®</sup> analyses, and they are particularly  
603 grateful to Thierry Adatte (Institute of Earth Sciences) and Stéphanie Grand (Institute of Earth Surface  
604 Dynamics) for their technical and scientific supports. The authors are also grateful to Chiara Compostella  
605 and Elena Silvia Ferrari for their assistance during the analyses at the Sediments and Soils Laboratory at  
606 Earth Science Department of Università degli Studi di Milano. The authors want to thank two anonymous  
607 reviewers for their constructive remarks that substantially improved the first version of the manuscript.

608

609

610

611

612

613

614

615

616

617

618

619

620

621

622

623

624

625

626

627

628

629

630

631

632 **References**

- 633 Albrecht, R., Sebag, D., Verrecchia, E.P., 2015. Organic matter decomposition: bridging the gap between  
634 Rock-Eval pyrolysis and chemical characterization (CPMAS <sup>13</sup>C NMR). *Biogeochemistry* 122, 101–  
635 111.
- 636 Arduino, E., Barberis, E., Ajmone Marsan, F., Zanini, E., Franchini, M., 1986. Iron oxides and clay minerals  
637 within profiles as indicators of soil age in northern Italy. *Geoderma*, 37, 45–55.
- 638 Avery, B. W., Bascomb, C. L., (Eds.). 1982. *Soil survey laboratory methods*. Lawes agricultural trust.
- 639 Bascomb, C. L., 1968. Distribution of pyrophosphate-extractable iron and organic carbon in soils of various  
640 groups. *European Journal of Soil Science*, 19(2), 251-268.
- 641 Behar, F., Beaumont, V., De B. Penteado, H.L., 2001. Rock-Eval® 6 technology: performances and  
642 developments. *Oil & Gas Science and Technology* 56 (2), 111–134.
- 643 Biagi, P., Castelletti, L., Cremaschi, M., Sala, B., Tozzi, C., 1980. Popolazione e territorio nell'Appennino  
644 ToscoEmiliano e nel tratto centrale del bacino del Po e nelle Prealpi bresciane tra il IX ed il V  
645 millennio. *Emilia Preromana*, 8, 13–36.
- 646 Bertolini, G., 2007. Radiocarbon dating on landslides in the Northern Apennines (Italy). In: McInnes R.,  
647 Jakeways J., Fairbank H., Mathie E., (eds), *Landslides and Climate Change*. Taylor & Francis Group,  
648 London.
- 649 Bertolini, G., Pellegrini, M., 2001. The landslides of Emilia Apennines (Northern Italy) with reference to  
650 those which resumed activity in the 1994–1999 period and required civil protection interventions.  
651 *Quaderni di Geologia*, 8(1), 27–74.
- 652 Bollati, I. M., Masseroli, A., Mortara, G., Pelfini, M., Trombino, L., 2019. Alpine gullies system evolution:  
653 erosion drivers and control factors. Two examples from the western Italian Alps. *Geomorphology*,  
654 327, 248-263.
- 655 Bortolotti, V., 1992. *Guide Geologiche Regionali: Appennino Tosco-Emiliano*. Milano: BE-MA Editrice.
- 656 Brewer, R., 1976. *Fabric and Mineral Analysis of Soils*. Huntington, NY: Krieger.
- 657 Casagrande, A., 1934. *Die Aräometer-Methode zur Bestimmung der Kornverteilung von Böden und anderer*  
658 *Materialien*. – Springer, Berlin, 56 p.
- 659 Chersich, S., Galvan, P., Frizzera, L., Scattolin, L., 2007. Variabilità delle forme di humus in due siti  
660 campione di pecceta altimontana trentina. *Forest@-Journal of Silviculture and Forest Ecology*, 4 (2),  
661 220-226.
- 662 Coltorti, M., Pieruccini, P., 2006. The last interglacial pedocomplexes in the litho-and morpho-stratigraphical  
663 framework of the central-northern Apennines (Central Italy). *Quaternary International*, 156, 118-132.
- 664 Coltorti, M., Pieruccini, P., Arthur, K. J., Arthur, J., Curtis, M., 2019. Geomorphology, soils and palaeosols  
665 of the Chench area (Gamo Gofa, south western Ethiopian Highlands). *Journal of African Earth*  
666 *Sciences*, 151, 225-240.
- 667 Compostella, C., Mariani, G.S., Trombino, L., 2014. Holocene environmental history at the treeline in the  
668 Northern Apennines, Italy: A micromorphological approach. *The Holocene*, 24(4), 393–404.
- 669 Compostella, C., Trombino, L., Caccianiga, M., 2013. Late Holocene soil evolution and treeline fluctuations  
670 in the Northern Apennines. *Quaternary International*, 289, 46–59.
- 671 Cremaschi, M., Nicosia, C., 2010. Corso di Porta Reno, Ferrara (Northern Italy): a study in the formation  
672 processes of Urban Deposits. *Il Quat. Ital. J. Quat. Sci.*, 23, 395–408.
- 673 Cremaschi M., Rodolfi, G., 1991. *Il suolo - Pedologia nelle scienze della Terra e nella valutazione del*  
674 *territorio*. La Nuova Italia Scientifica, Roma.
- 675 Cremaschi, M., Trombino, L., Zerboni, A., 2018. Palaeosols and relict soils: a systematic review. In  
676 *Interpretation of Micromorphological Features of Soils and Regoliths* (pp. 863-894). Elsevier.
- 677 Cremaschi, M., Biagi, P., Accorsi, C.A., Bandini Mazzanti, M., Rodolfi, G., Castelletti, L., Leoni, L., 1984.  
678 *Il sito mesolitico di Monte Bagioletto (Appennino Reggiano) nel quadro delle variazioni ambientali*  
679 *oloceniche dell'Appennino Tosco-Emiliano*. *Emilia Preromana*, 9/10, 11-46. D'Amico, M. E., Catoni,  
680 M., Terribile, F., Zanini, E., Bonifacio, E., 2016. Contrasting environmental memories in relict soils  
681 on different parent rocks in the south-western Italian Alps. *Quaternary International*, 418, 61-74.
- 682 Dewolf, Y., Bourrié, G., 2008. *Les formations superficielles: genèse, typologie, classification, paysages et*  
683 *environnements, ressources et risques*. Ellipses.
- 684 Do Nascimento, N. R., Fritsch, E., Bueno, G. T., Bardy, M., Grimaldi, C., Melfi, A. J., 2008. Podzolization  
685 as a deferralitization process: dynamics and chemistry of ground and surface waters in an Acrisol–  
686 Podzol sequence of the upper Amazon Basin. *European Journal of Soil Science*, 59(5), 911-924.

- 687 Douglas, L. A., Thompson, M. L., 1985. Soil micromorphology and soil classification. Madison, WI: Soil  
688 Science Society of America.
- 689 Duchaufour, P., 1983. Pédologie. 1. Pédogenèse et classification. Masson, Paris.
- 690 Erhart, H., 1967. La Genèse des sols en tant que phénomène géologique: esquisse d'une théorie géologique et  
691 géochimique, biostasie et rhexistasie: exmaples d'application. Masson.
- 692 Fedoroff, N., Courty, M., 2012. Textural features and microfacies expressing temporary and permanent soil  
693 water saturation. In Poch-Claret, R., Casamitjana, M., and Francis, M., editors, Proceedings of the  
694 14th Intern. Working Meet. On Soil Micromporphology, page 1.1.K. Session I. Editions i Publications  
695 de la Universitat de Lleida, Spain.
- 696 Fedoroff, N., Goldberg, P., 1982. Comparative micromorphology of two late Pleistocene paleosols (in the  
697 Paris Basin). *Catena*, 9, 227-251.
- 698 Fedoroff, N., Courty, M. A., Thompson, M. L., 1990. Micromorphological evidence of paleoenvironmental  
699 change in Pleistocene and Holocene paleosols. In *Developments in soil science* (Vol. 19, pp. 653-665).  
700 Elsevier.
- 701 Gales, S.J., Hoare, P.G., 1991. *Quaternary Sediments: Petrographic Methods for the Study of Unlithified*  
702 *Rocks* Belhaven, London, p. 323
- 703 Giraudi, C., 2005. Middle to Late Holocene glacial variations, periglacial processes and alluvial  
704 sedimentation on the higher Apennine massifs (Italy). *Quaternary Research*, 64(2), 176-184.
- 705 IUSS Working Group WRB, 2015. World Reference Base for Soil Resources 2014, update 2015  
706 International soil classification system for naming soils and creating legends for soil maps. World Soil  
707 Resources Reports No. 106. FAO, Rome.
- 708 Jenny, H., 1941. *Factors of soil formation: a system of quantitative pedology*. McGraw-Hill book company  
709 inc., New York.
- 710 Jahn, R., Blume, H. P., Asio, V. B., Spaargaren, O., Schad, P., 2006. *Guidelines for soil description*. FAO.
- 711 Kaiser, K., Schoch, W. H., Miehe, G., 2007. Holocene paleosols and colluvial sediments in Northeast Tibet  
712 (Qinghai Province, China): properties, dating and paleoenvironmental implications. *Catena*, 69(2), 91-  
713 102.
- 714 Kemp, R.A., 1998. Role of Micromorphology in paleopedological research. *Quaternary International* 51-52,  
715 133-141.
- 716 Kjeldahl, J., 1883. Neue Methode zur Bestimmung des Stickstoffs in organischen Körpern. *J. Anal. Chem.*  
717 22, 366-382.
- 718 Kleber, A., Terhorst, B., 2013. *Mid-latitude slope deposits (cover beds)* (Vol. 66). Newnes.
- 719 Krasilnikov, P., Calderón, N.E.G., 2006. A WRB-based buried paleosol classification. *Quaternary*  
720 *International*, 156, 176-188.
- 721 Kubiëna, W.L., 1953. *Bestimmungsbuch und Systematik der Böden Europas*. F. Enke Verlag, Stuttgart.
- 722 Lafargue, E., Marquis, F., Pillot, D., 1998. Rock-Eval® 6 applications in hydrocarbon exploration,  
723 production, and soil contamination studies. *Oil & Gas Science and Technology* 53 (4), 421-437.
- 724 Losacco, U., 1949. La glaciazione quaternaria dell'Appennino Settentrionale. *Rivista Geografica Italiana*,  
725 56(2), 90-152.
- 726 Losacco, U., 1982. Gli antichi ghiacciai dell'Appennino settentrionale. Studio morfologico e  
727 paleogeografico. *Atti della Societa dei Naturalisti e Matematici di Modena*, 113, 1-24.
- 728 Magliulo, P., Terribile, F., Colombo, C., Russo, F., 2006. A pedostratigraphic marker in the  
729 geomorphological evolution of the Campanian Apennines (Southern Italy): The Paleosol of  
730 Eboli. *Quaternary international*, 156, 97-117.
- 731 Malou O.P., Sebag D., Moulin P., Chevallier T., Badiane-Ndour N.Y., Thiam A., Chapuis-Lardy L. 2020.  
732 The Rock-Eval signature of soil organic carbon in arenosols of the Senegalese groundnut basin. How  
733 do agricultural practices matter? *Agriculture, Ecosystems & Environment* 301,107030.
- 734 Mariani G.S., Brandolini F., Pelfini M., Zerboni A., 2019a. Mapping Matilda's castles in the northern  
735 Apennines: geological and geomorphological constrains. *Journal of Maps* 15/2, 521-529.
- 736 Mariani, G. S., Compostella, C., Trombino, L., 2019b. Complex climate-induced changes in soil  
737 development as markers for the Little Ice Age in the Northern Apennines (Italy). *Catena*, 181, 104074.

- 738 Mariani, G. S., Cremaschi, M., Zerboni, A., Zuccoli, L., Trombino, L., 2018. Geomorphology of the Mt.  
739 Cusna Ridge (Northern Apennines, Italy): evolution of a Holocene landscape. *Journal of Maps*, 14(2),  
740 392-401.
- 741 Mariani, G.S., 2016. The role of paleosols in paleoenvironmental studies: genesis and development of  
742 Apennine mountain soils during the Holocene. Phd Thesis, University of Milan.
- 743 Matteodo, M., Grand, S., Sebag, D., Rowley, M. C., Vittoz, P., Verrecchia, E. P., 2018. Decoupling of  
744 topsoil and subsoil controls on organic matter dynamics in the Swiss Alps. *Geoderma*, 330, 41-51.
- 745 McCarthy P.J., Martini, I.P., Leckie, D.A., 1998. Use of micromorphology for palaeoenvironmental  
746 interpretation of complex alluvial palaeosols: an example from the Mill Creek Formation (Albian),  
747 southwestern Alberta, Canada. *Palaeogeography, Palaeoclimatology, Palaeoecology* 143, 87-110.
- 748 Milne, G., 1936. Normal erosion as a factor in soil profile development. *Nature*, 138(3491), 548.
- 749 Ministero delle Risorse Agricole Alimentari e Forestali, 1994. *Metodi ufficiali di 1036 analisi chimica del*  
750 *suolo, con commenti ed interpretazioni*. ISMEA, Roma, 207 pp.
- 751 Nicosia, C., 2006. Indicatori micromorfologici di erosione dei suoli nel settore settentrionale delle Valli  
752 Grandi Veronesi durante l'età del Ferro. *Padusa*, 62, 108-112.
- 753 Panizza, M., Bettelli, G., Bollettinari, G., Carton, A., Castaldini, D., Piacente, S., Bernini, M., Clerici, A.,  
754 Tellini, C., Vittorini, S., Canuti, P., Moisello, U., Tenti, G., Dramis, F., Gentili, B., Pambianchi, G.,  
755 Bidini, D., Lulli, L., Rodolfi, G., Busoni, E., Ferrari, G., Cremaschi, M., Marchesini, A., Accorsi,  
756 C.A., Mazzanti, M., Francavilla, F., Marchetti, G., Vercesi, P.L., Di Gregorio, F., Marini, A., (Gruppo  
757 Ricerca Geomorfologia CNR) 1982. Geomorfologia del territorio di Febbio tra il M.Cusna e il  
758 F.Secchia (Appennino Emiliano). *Geografia Fisica Dinamica Quaternaria*, 5, 285-360.
- 759 Pawluk, S., 1972. Measurement of crystalline and amorphous iron removal in soils. *Can. J. Soil Sci.* 52: 119-  
760 123.
- 761 Regattieri, E., Zanchetta, G., Isola, I., Zanella, E., Drysdale, R.N., Hellstrom, J.C., Zerboni, A., Dallai, L.,  
762 Tema, E., Lanci, L., et al., 2019. Holocene Critical Zone dynamics in an Alpine catchment inferred  
763 from a speleothem multiproxy record: disentangling climate and human influences. *Sci. Rep.*, 9, 1–9.
- 764 Rellini, I., Trombino, L., Firpo, M., Piccazzo, M., 2007. Geomorphological context of “plinthitic paleosols”  
765 in the Mediterranean region: examples from the coast of western Liguria (northern Italy). *Rev. C. & G.*,  
766 21 (1-2), 27-40.
- 767 Rhodes, E. R., Sutton, P. M., 1978. Active iron ratio of some soils from three physiographic units in Sierra  
768 Leone. *Soil Science*, 125(5), 326-328.
- 769 Romanens, R., Pellacani, F., Mainga, A., Fynn, R., Vittoz, P., Verrecchia, E.P., 2019. Soil diversity and  
770 major soil processes in the Kalahari basin. *Botswana. Geoderma Reg.* 19, e00236.  
771 <https://doi.org/10.1016/j.geodrs.2019.e00236>.
- 772 Ruellan, A., 1971. The history of soils. Some problems of definition and interpretation. In: Yaalon, D.H.  
773 (Ed.), *Paleopedology. Origin, Nature and Dating of Paleosols*. Israel University Press, Jerusalem, 350  
774 pp.
- 775 Sauro, U., Ferrarese, F., Francese, R., Miola, A., Mozzi, P., Rondo, G.Q., Trombino, L., Valentini, G., 2009.  
776 Doline fills- case study of the Faverghera Plateau (Venetian Pre-Alps, Italy). *Acta Carsologica*, 38 (1),  
777 51-63.
- 778 Schomburg, A., Verrecchia, E. P., Guenat, C., Brunner, P., Sebag, D., Le Bayon, R. C., 2018. Rock-Eval@  
779 pyrolysis discriminates soil macro-aggregates formed by plants and earthworms. *Soil Biology and*  
780 *Biochemistry*, 117, 117-124.
- 781 Schomburg, A., Sebag, D., Turberg, P., Verrecchia, E. P., Guenat, C., Brunner, P., Adatte T, Schlaepfer R.,  
782 Le Bayon, R. C., 2019. Composition and superposition of alluvial deposits drive macro-biological soil  
783 engineering and organic matter dynamics in floodplains. *Geoderma*, 355, 113899.
- 784 Schwertmann, U., 1973. Use of oxalate for Fe extraction from soils. *Canadian Journal of Soil Science*, 53(2),  
785 244-246.
- 786 Sebag, D., Verrecchia, E. P., Cécillon, L., Adatte, T., Albrecht, R., Aubert, M., Bureau, F., Cailleau, G.,  
787 Copard, Y., Decaens, T., Disnar, J. R., Hetényi, M., Nyilas, T., Trombino, L., 2016. Dynamics of soil  
788 organic matter based on new Rock-Eval® indices. *Geoderma*, 284, 185-203.

789 Sevink, J., Bakels, C.C., Attema, P.A., Di Vito, M.A., Arienzo, I., 2019. Holocene vegetation record of  
790 upland northern Calabria, Italy: Environmental change and human impact. *The Holocene*, 29, 633–  
791 647.

792 Sheldon, N.D., Tabor, N.J., 2009. Quantitative paleoenvironmental and paleoclimatic reconstruction using  
793 paleosols. *Earth-Science Reviews*, 95(1-2), 1-52.

794 Stoops, G., 2003. Guidelines for analysis and description of soil and regolith thin sections. Soil Science  
795 Society of America, Inc., Madison, Wisconsin, USA.

796 Stoops, G., Marcelino, V., Mees, F., 2018. Interpretation of micromorphological features of soils and  
797 regoliths. Elsevier.

798 Stoops, G., Marcelino, V., Mees, F., 2010. Interpretation of micromorphological features of soils and  
799 regoliths. Elsevier.

800 Thoumazeau A., Chevallier T., Baron V., Rakotondrazafy N., Panklang P., Marichal R., Kibblewhite M.,  
801 Sebag D., Tivet F., Bessou C., Gay F., Brauman A., 2020. A new in-field indicator to assess the  
802 impact of land management on soil carbon dynamics. *Geoderma* 375, 114496.

803 Vittori Antisari, L., Bianchini, G., Cremonini, S., Di Giuseppe, D., Falsone, G., Marchesini, M., Marvelli, S.,  
804 Vianello, G., 2016. Multidisciplinary study of a late glacial-Holocene sedimentary sequence near  
805 Bologna (Italy): insights on natural and anthropogenic impacts on the landscape dynamics. *J. Soils*  
806 *Sediments* 16, 645-662.

807 Walkley, A., Black, I.A., 1934. An examination of the Degtjareff method for determining soil organic  
808 matter, and proposed modification of the chromic acid titration method. *Soil Sci*, 37(1), 29-38.

809 Waroszewski, J., Kalinski, K., Malkiewicz, M., Mazurek, R., Kozłowski, G., Kabala, C., 2013. Pleistocene–  
810 Holocene cover-beds on granite regolith as parent material for Podzols—an example from the Sudeten  
811 Mountains. *Catena*, 104, 161-173.

812 Zanelli, R., Egli, M., Mirabella, A., Giaccai, D., Abdelmoula, M., 2007. Vegetation effects on pedogenetic  
813 forms of Fe, Al and Si and on clay minerals in soils in southern Switzerland and northern  
814 Italy. *Geoderma*, 141(1-2), 119-129.

815 Zanini, E., Freppaz, M., Stanchi, S., Bonifacio, E., Egli, M., 2015. Soil variability in mountain areas. In:  
816 Romeo, R; Vita, A; Manuelli, S; Zanini, E; Freppaz, Michele; Stanchi, Silvia. *Understanding*  
817 *Mountain Soils: A Contribution from mountain areas to the International Year of Soils 2015*. Rome:  
818 FAO, 60-62.

819 Zerboni A., Mariani G.S., Castelletti L., Amit R., Ferrari E.S., Tremari M., Livio F., 2019. Was the Little Ice  
820 Age the coolest Holocene climatic period in the Italian central Alps? *Progress in Physical Geography*.  
821 DOI: 10.1177/0309133319881105  
822

823 **Tables**

## 824 Table 1.

Profile	Elevation (m a.s.l.)	Slope gradient (°)	Slope exposure	Profile exposure	Parent Material	Geomor- phological context	Vegetation
01	1680	10	NE	SW	Colluvium deposits composed of claystones	Middle slope, flank of residual hills preserved from water runoff	Semi-deciduous shrub mainly composed of <i>Vaccinium myrtillus</i> and <i>Juniperus nana</i> . Sparse arboreal vegetation is characterized by <i>Fagus sylvatica</i> and rare <i>Laburnum alpinum</i>
02	1669	2	N-NW	SW	Colluvium deposits composed of claystones	Middle slope, flank of residual hills preserved from water runoff	Semi-deciduous shrub mainly composed of <i>Vaccinium myrtillus</i> and <i>Juniperus nana</i> . Rare <i>Fagus sylvatica</i>
03	1665	4	NW	N	Colluvium deposits composed of claystones	Middle slope, portion of residual hills preserved from water runoff	Semi-deciduous shrub mainly composed of <i>Vaccinium myrtillus</i> and <i>Juniperus nana</i> . Rare <i>Fagus sylvatica</i>
04	1659	11	NW	N-NE	Colluvium deposits composed of claystones	Middle slope	Deciduous woodland composed of <i>Fagus sylvatica</i> with <i>Vaccinium myrtillus</i> undergrowth
05	1663	10	NW	N-NE	Colluvium deposits composed of claystones	Middle slope	Semi-deciduous shrub mainly composed of <i>Vaccinium myrtillus</i>
06	1661	22	S-SE	SW	Colluvium deposits composed of claystones	Middle slope, portion of residual hills preserved from water runoff	Semi-deciduous shrub mainly composed of <i>Vaccinium myrtillus</i>

825

826 Table 1. Site description of investigated soil profiles.

827

828

829 Table 2.  
830


Profile	Horizon	Depth (cm)	Fe <sub>o</sub> (g/Kg)	Al <sub>o</sub> (g/kg)	Fe <sub>d</sub> (g/kg)	Al <sub>d</sub> (g/kg)	Fe <sub>p</sub> (g/kg)	Al <sub>p</sub> (g/kg)	Fe <sub>cry</sub> (Fe <sub>d</sub> -Fe <sub>o</sub> ) g/kg	Fe <sub>o</sub> /Fe <sub>d</sub>	Al <sub>o</sub> +1/2Fe <sub>o</sub> %
01	O	0-4	6.24	2.87	15.16	3.65	4.70	11.41	8.91	0.41	0.60
	A1	4-10	6.33	3.46	16.39	4.09	5.81	8.67	10.06	0.39	0.66
	A2	10-31	6.84	3.29	21.99	4.88	6.24	10.90	15.15	0.31	0.67
	BC	31-35	9.08	4.40	17.78	4.13	5.54	6.62	8.70	0.51	0.89
	2AB1	35-44	9.52	2.74	25.98	4.37	9.70	8.50	16.46	0.37	0.75
	2AB2	44-65	15.73	4.44	31.47	5.71	15.98	10.37	15.74	0.50	1.23
	2Bw	65-80	9.84	5.06	17.39	5.05	7.44	7.46	7.55	0.57	1.00
	2Bt	80-100	3.85	2.88	16.20	4.97	4.10	6.09	12.35	0.24	0.48
	2BC1	100-120	5.80	4.30	12.17	4.17	<3.00	4.86	6.37	0.48	0.72
	2BC2	120-140+	4.19	3.45	11.82	4.00	n.d.	3.99	7.63	0.35	0.55
02	OB	0-20	11.49	4.02	22.76	4.44	8.68	7.46	11.27	0.50	0.98
	ABt1	20-60	5.73	3.45	17.01	5.16	n.d.	6.92	11.29	0.34	0.63
	ABt2	60-75	2.48	3.16	12.12	3.17	n.d.	3.84	9.64	0.20	0.44
	BC	75-95	8.82	5.40	13.37	3.62	n.d.	4.57	4.55	0.66	0.98
	2AB	95-121	<0.90	1.69	12.70	3.07	n.d.	3.32	n.d.	n.d.	n.d.
	2Btg	121-151	n.d.	0.85	14.35	3.27	n.d.	3.64	n.d.	n.d.	n.d.
	3AB	151-176+	3.48	2.40	12.14	2.84	n.d.	2.98	8.66	0.29	0.41
03	OA	0-8	7.73	3.39	20.43	4.10	6.21	7.08	12.70	0.38	0.73
	OB	8-22	10.11	2.74	30.57	4.25	12.26	9.42	20.47	0.33	0.78
	BC	22-44+	12.92	5.59	22.82	5.53	8.53	6.30	9.89	0.57	1.21
04	O	0-6	8.61	3.13	17.21	3.38	4.58	4.34	8.60	0.50	0.74
	OA	6-12	6.48	3.17	22.61	4.85	8.23	8.52	16.14	0.29	0.64
	AB	12-27	8.19	4.93	15.70	4.62	5.17	7.58	7.51	0.52	0.90
	Btg	27-54	2.82	2.64	15.57	4.53	<3.00	6.13	12.75	0.18	0.40
	Bt	54-80+	n.d.	n.d.	14.37	3.68	<3.00	4.52	n.d.	n.d.	n.d.
05	O	0-3	3.52	2.14	12.02	2.64	n.d.	3.42	8.49	0.29	0.39
	A	3-12	n.d.	n.d.	17.52	4.85	<3.00	6.53	n.d.	n.d.	n.d.
	ABt1	12-28	6.24	2.71	18.21	3.45	5.63	6.38	11.97	0.34	0.58
	ABt2	28-56+	13.06	3.78	34.13	5.83	13.27	12.09	21.08	0.38	1.03
06	O	0-6	3.52	2.12	13.44	4.02	<3.00	4.58	9.92	0.26	0.39
	OA	6-12	10.21	4.53	22.84	4.53	7.15	7.69	12.64	0.45	0.96
	2AB1	12-19	12.34	3.28	31.07	5.60	12.72	9.31	18.73	0.40	0.94
	2AB2	19-36	10.66	3.73	25.79	5.49	12.63	7.95	15.13	0.41	0.91
	2Bw1	36-54	7.28	4.39	19.03	5.72	7.54	9.21	11.76	0.38	0.80
	2Bw2	54-63	2.82	2.29	14.45	4.43	<3.00	7.30	11.63	0.20	0.37
	2Bw3	63-71	2.55	2.75	14.35	4.25	n.d.	3.92	11.80	0.18	0.40
	3AB	71-89+	5.60	3.77	13.94	3.66	n.d.	4.52	8.33	0.40	0.66

831  
832 Table 2. Ammonium oxalate (Fe<sub>o</sub>, Al<sub>o</sub>), dithionite-citrate-bicarbonate (Fe<sub>d</sub>, Al<sub>d</sub>) and sodium pyrophosphate  
833 (Fe<sub>p</sub>, Al<sub>p</sub>) extractable Fe and Al in the studied profiles and derivate indices of crystalline iron oxides (Fe<sub>cry</sub>),  
834 activity iron index (Fe<sub>o</sub>/Fe<sub>d</sub>) and podzolization index (Al<sub>o</sub>+1/2Fe<sub>o</sub>).  
835 <: low values approximate to the minor concentration detectable; n.d.: no data.  
836


837



Figure 1  
[Click here to download high resolution image](#)



**Figure 2**  
[Click here to download high resolution image](#)



**Figure 3**  
[Click here to download high resolution image](#)





Figure 4  
[Click here to download high resolution image](#)








Figure 5  
[Click here to download high resolution image](#)




Figure 6

[Click here to download high resolution image](#)




Figure 7  
[Click here to download high resolution image](#)




Figure 8  
[Click here to download high resolution image](#)

## Imprints of factors

		Climate	Vegetation	Relief
Biorhexistay phases	$\gamma$ Biostasy phase	↑ ⊖	↕ ⊖ <sup>*</sup>	↕ ⊖
	Rhexistasy phase	↓ ⊖	↓	↓ ⊖
	$\beta$ Biostasy phase	↑ ⊕	↑ ⊖	↑
	Rhexistasy phase	↓ ⊕	↓	↓ ⊕
	$\alpha$ Biostasy phase	↑ ⊕	↑ ⊕	↑



Figure 1. A) Study area and location of the sample site at Mt. Cusna on altitude contour map. B) Locations of profiles on the orthophotograph (2006). The digital sources are courtesy made available by the Geoportale Nazionale (<http://www.pcn.minambiente.it/GN/>, WMS service).

Figure 2. Particle size distribution (dark grey: sand\*; grey: silt; light grey: clay), organic C content, total N content, pH, contents in crystalline iron oxides ( $Fe_{cry}$ ) and ammonium oxalate extractable Fe ( $Fe_o$ ) in the studied profiles. The  $Fe_o$  is recognized as a measurement of the “activity” of iron oxides (Schwertmann, 1973). \* Since the gravel content is very low, it has been added to the sand fraction.

Figure 3. a) HI (mg HC/g TOC)/OI (mg  $CO_2$ /g TOC) diagram; the horizon 01 2BC1 is not plotted due to its out of range value, b) I-index/R-index diagram of the studied horizons. Colored dots are used to calculate the “humic trend” equation written in bold. In background, the Matteodo's dataset, composed of 46 soil profiles selected across various ecounits in Swiss Alps (Matteodo et al., 2018) and the relating “humic trend” equation is depicted, as comparison.

Figure 4. a) R-Index/TOC (%) diagram; b) R-index/depth (cm) diagram; c) I-Index/TOC (%) diagram; d) I-index/depth (cm) diagram; e) Delta I/TOC (%) diagram of the studied horizons.

Figure 5. Photomicrographs of some micromorphological features observed in soils and palaeosols. a,b) altered, reddish fine-material forming subangular aggregates, with Fe/Mn nodule concentrations in 05 ABt1 and ABt2 horizons (1.6x, PPL e XPL; field length: 8 mm); c,d) vughy structure with clay illuviation and redoximorphic features in a 02 2Btg thin section (1.6x, PPL and XPL; field length: 8 mm); e,f) crumb aggregates and reddish, subangular pedorelicts in the horizon BC of the 01 profile (1.6x, PPL e XPL; field length: 8 mm).

Figure 6. Correlation scheme between the investigated soil profiles. The lines show the correlation among the different pedological units (black lines for the correlations found observing soil thin sections; dotted lines for the correlations supposed based on soil physical and chemical properties).

Figure 7. Dendrogram output for clustering of studied horizons, using Rock-Eval indices (HI, OI, I-Index and R-Index). The horizon 01 2BC1 is not plotted due to its out of range values.

Figure 8. Role of the main factors influencing the soil development through time in the study area. The upward arrow represents favorable conditions to soil development, whereas the downward arrow represents unfavorable conditions regarding soil development. The presence of plus or minus signs underlines the active role of a factor, which can be more (plus) or less (minus) intense, in its influence on the pedogenesis.

\*: human impact influence in vegetation cover.

## Appendix A

[Click here to download Supplementary material for on-line publication only: Appendix A.pdf](#)

## Appendix B

[Click here to download Supplementary material for on-line publication only: Appendix B\\_rev.pdf](#)

## Appendix C

[Click here to download Supplementary material for on-line publication only: Appendix C\\_rev.pdf](#)

## Appendix D

[Click here to download Supplementary material for on-line publication only: Appendix D\\_rev.pdf](#)

**Declaration of interests**

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

*Anna Passoli*  
*Isabelle* *Laura* *15A* *Demofeleon*  
*Eui* *De* *Giuseppe S. Maini*