

Table 2. Random fluctuations in healthy controls

Random fluctuation for one day								
	$\geq 8 \text{ kPa}$		$\geq 10 \text{ kPa}$		$\geq 12 \text{ kPa}$		$\geq 14 \text{ kPa}$	
	<i>mean</i>	<i>maximum</i>	<i>mean</i>	<i>maximum</i>	<i>mean</i>	<i>maximum</i>	<i>mean</i>	<i>maximum</i>
n (%) meeting criterion	22 (73%)	26 (87%)	18 (60%)	23 (77%)	14 (47%)	20 (66%)	10 (33%)	12 (40%)
% of days with random fluctuations; mean; (range)	28% (3-73)	29% (3-73)	16% (3-68)	24% (3-73)	18% (3-60)	18% (3-73)	19% (8-40)	19% (3-45)
Maximum random fluctuations (kPa); mean (range)	15 (9-24)	14 (8-22)	15 (10-24)	15 (10-22)	17 (12-24)	15 (12-22)	19 (14-24)	17 (14-22)
Random fluctuation for at least four consecutive days								
	$\geq 8 \text{ kPa}$		$\geq 10 \text{ kPa}$		$\geq 12 \text{ kPa}$		$\geq 14 \text{ kPa}$	
	<i>mean</i>	<i>maximum</i>	<i>mean</i>	<i>maximum</i>	<i>mean</i>	<i>maximum</i>	<i>mean</i>	<i>maximum</i>
n (%) meeting criterion	7 (23%)	11 (37%)	6 (20%)	7 (23%)	4 (13%)	5 (17%)	3 (10%)	3 (10%)
% of days with random fluctuations; mean; (range)	33% (21-57)	29% (13-57)	25% (13-37)	31% (16-57)	23% (13-37)	38% (20-57)	28% (20-37)	19% (13-30)