

Journal Pre-proof

Cerebral regional oxygen saturation during cardiopulmonary resuscitation and return of spontaneous circulation: a systematic review and meta-analysis

Filippo Sanfilippo, Paolo Murabito, Antonio Messina, Veronica Dezio, Diana Busalacchi, Giuseppe Ristagno, Maurizio Cecconi, Marinella Astuto



PII: S0300-9572(20)30596-7

DOI: <https://doi.org/10.1016/j.resuscitation.2020.12.002>

Reference: RESUS 8816

To appear in: *Resuscitation*

Received Date: 20 September 2020

Revised Date: 21 November 2020

Accepted Date: 1 December 2020

Please cite this article as: Sanfilippo F, Murabito P, Messina A, Dezio V, Busalacchi D, Ristagno G, Cecconi M, Astuto M, Cerebral regional oxygen saturation during cardiopulmonary resuscitation and return of spontaneous circulation: a systematic review and meta-analysis, *Resuscitation* (2020), doi: <https://doi.org/10.1016/j.resuscitation.2020.12.002>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier.

Cerebral regional oxygen saturation during cardiopulmonary resuscitation and return of spontaneous circulation: a systematic review and meta-analysis.

Filippo Sanfilippo^a, Paolo Murabito^{a,b,c}, Antonio Messina^{d,e}, Veronica Dezio^c, Diana Busalacchi^c, Giuseppe Ristagno^{f,g}, Maurizio Cecconi^{d,e}, Marinella Astuto^{a,b,c}.

^aDepartment of Anaesthesia and Intensive Care, A.O.U. "Policlinico-San Marco", Catania, Italy

^bDepartment of General Surgery and Medical-Surgical Specialties, Section of Anesthesia and Intensive Care, University of Catania, Catania, Italy

^cSchool of Anaesthesia and Intensive Care, University Hospital "G. Rodolico", University of Catania, Catania, Italy

^dHumanitas Clinical and Research Center – IRCCS, Milano, Italy

^eDepartment of Biomedical Sciences, Humanitas University, Pieve Emanuele, MI, Italy

^fDepartment of Pathophysiology and Transplantation, University of Milan, Milan, Italy

^gDepartment of Anesthesiology, Intensive Care and Emergency, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy

Corresponding Author: Filippo Sanfilippo, MD, PhD, EDIC

Department of Anaesthesia and Intensive Care, A.O.U. "Policlinico-San Marco", via S. Sofia, Post Code 95100, Catania, Italy.

Tel: 00393289178766 email: filipposanfi@yahoo.it

Abstract

Aim: Predicting the return of spontaneous circulation (ROSC) during cardiopulmonary resuscitation in victims of cardiac arrest (CA) remains challenging. Cerebral regional oxygen saturation (rSO₂) measured during resuscitation is feasible, and higher initial and overall values seem associated with ROSC. However, these observations were limited to the analysis of few small single-centre studies. There is a growing number of studies evaluating the role of cerebral rSO₂ in the prediction of ROSC.

Methods: We conducted an updated meta-analysis aimed at investigating the association of initial and overall values of cerebral rSO₂ with ROSC after CA. We performed subgroups analyses

according to the location of CA and conducted a secondary analysis according to the country where the study was conducted (resuscitation practice varies greatly for out-of-hospital CA).

Results: We included 17 studies. Higher initial rSO₂ values (11 studies, n=2870, 16.6% achieved ROSC) were associated with ROSC: Mean Difference (MD) -11.54 [95% Confidence Interval (CI)-20.96, -2.12]; p=0.02 (I²=97%). The secondary analysis confirmed this finding when pooling together European and USA studies, but did not for Japanese studies (p=0.06). One multi-centre Japanese study was an outlier with large influence on 95% CI. Higher overall rSO₂ values during resuscitation (9 studies, n=894, 33.7% achieving ROSC) were associated with ROSC: MD-10.38; [-13.73, -7.03]; p<0.00001 (I²=77%). All studies were conducted in Europe/USA.

Conclusions: This updated meta-analysis confirmed the association between higher initial and overall values of cerebral rSO₂ and ROSC after CA. However, we found geographical differences, since this association was not present when Japanese studies were analysed separately. **Keywords:** near infrared spectrometry; advanced cardiac life support; resuscitation order; cardiopulmonary resuscitation.

Introduction

Predicting the return of spontaneous circulation (ROSC) during cardiopulmonary resuscitation (CPR) represents a clinical challenge. Despite improvements in CPR and in the post-resuscitation care, mortality in patients suffering from cardiac arrest (CA) remains high. The rate of ROSC is very low, especially among out of hospital (OH)CA; furthermore, less than 50% of patients admitted to hospital after CA are alive at 6-month follow-up.[1-4] To date, there is no formally validated tool to help clinicians in deciding how long to prolong their efforts in CPR, and such decision is often made on

clinical grounds only. Importantly, when ROSC is achieved, the principal cause of death is attributable to neurological damage resulting from the ischemia-reperfusion injury.[5, 6]

Oximetry is a non-invasive monitoring technology based on near-infrared spectrometry (NIRS), and it has been approved by the Food and Drug Administration. When applied to the patient's forehead, it allows the evaluation of cerebral regional oxygen saturation (rSO_2) from superficial brain cortex regions, which are amongst the most susceptible areas to the ischemic-hypoxic injury.[7] The feasibility of using cerebral rSO_2 monitoring in patients suffering from CA has been repeatedly shown, both by studies investigating NIRS ability in predicting ROSC[8, 9] and/or focused on neurological outcome after CA,[10, 11]. Moreover, NIRS is under evaluation as tool for the assessment of the CPR quality.[12]

An initial meta-analysis performed in 2015 showed a significant association between the initial (five studies included) or the overall (five studies included) values of cerebral rSO_2 measured during CPR with ROSC in CA patients[13]. This meta-analysis included studies performed on both in-hospital (IH)CA, OHCA or mixed CA populations; moreover, at that time, there was a major contribution from two research groups, possibly decreasing the external validity of the meta-analysis findings. Although the introduction of NIRS during CA is still limited in clinical practice, additional research studies have been published after the previous meta-analysis. Therefore, considering the challenges in assessing ROSC probabilities in CA patients, we undertook an updated systematic review and meta-analysis with the aim of confirming the role of rSO_2 values in predicting ROSC. We hypothesized that both initial and overall cerebral rSO_2 values are positively associated with ROSC in CA patients undergoing CPR, in both subgroups of patients suffering from IHCA and OHCA.

Methods

We performed this systematic review using an automated literature search performed with the *NHS Library Evidence* tool. A further manual search was conducted independently by four authors (FS,

VD, DB, AM), exploring also the list of references of the findings of the systematic search. We followed the approach suggested by the PRISMA statement for reporting systematic reviews and meta-analyses (checklist provided in Supplemental Digital Content – Appendix 1).[14] We registered the protocol of our systematic review and meta-analysis in the PROSPERO database (CRD42020202993).

We included prospective and retrospective studies conducted in adult patients suffering from CA where NIRS sensors were applied for the evaluation of cerebral rSO₂ values. We included both studies performed on IHCA, OHCA and mixed populations. As primary outcome we evaluated the association between the initial and/or the overall cerebral rSO₂ values and ROSC in the selected population (see PICOS approach, Supplemental Digital Content – Appendix 2). We excluded articles referring to the paediatric population. Language restrictions were applied: only articles published in English were considered. We considered case series if reporting at least 10 patients; series 5 to 9 patients were included for sensitivity analyses only. Smaller series and case reports were excluded.

Searches, screening and data collection

The computerized search included MEDLINE, from inception until 15.06.2020. We also performed a search on EMBASE limited to the findings from 2017 in order to retrieve the newest conference abstracts not yet published, to allow a reasonable time for peer-review process. In both searches we combined terms from two groups, under the condition that at least one term per group was included in the article title and/or abstract. The first group included the terms “cerebral oximetry”, “near infrared”, “oxygen consumption”, “spectrophotometry” and “spectroscopy”; the second one consisted of the terms “cardiac arrest”, “heart arrest”, “resuscitation” and “return of spontaneous circulation”.

Two authors (VD, DB) independently searched the databases and selected studies for the inclusion in the systematic review. Discordances were resolved by involving two other reviewer (FS, MA). Two

authors (VD, DB) extracted data from the included studies in an Excel spreadsheet and two authors (FS, PM) performed a double-check. When needed, we contacted via email the corresponding authors for retrieving more data. Meta-analysis was performed by one experienced authors (FS) and checked by other two authors with experience (AM, MC).

Analyses of outcomes

We planned to analyse the association between cerebral rSO₂ values and ROSC dividing studies in subgroups according to the site of CA (IH, OH, mixed).

We also planned a secondary analysis dividing studies according to the country where these were conducted. In particular, we separated studies conducted in Europe and USA from those performed in Japan. We justify this choice as in the Japanese emergency medical personnel are not allowed to terminate CPR on site, and all OHCA are brought to the hospital with ongoing CPR.

Three sensitivity analyses were planned: the first one excluding one study at time (*“leave-one-out at time approach”*), the second including case series with less than 10 patients, and the third excluding studies with high risk of bias.

Risk of bias assessment

Five authors (FS, PM, AM, GR, MA) independently assessed the methodological quality of the included studies using the Newcastle-Ottawa Assessment Scale (NOS) which is recommended for the assessment of the quality of non-randomized studies.[15] The scale has three main domains and assigns one point for each of nine items. Studies are then classified as at high (1-3 points), intermediate (4-5 points) or low (6-9 points) risk of bias. **Statistical analysis**

Continuous outcome differences were analysed using an inverse variance model with a 95% Confidence Interval (CI). If data were reported only as median and interquartile range or CI, we followed current recommendations to approximate the values of mean and standard deviation

(SD)[16-18]. Values are reported as mean difference (MD), p values were two-tailed and considered significant if <0.05 . A random model was used. The presence of statistical heterogeneity was assessed using the X^2 (Cochran Q) test. Heterogeneity was likely if $Q > df$ (degrees of freedom) suggested and confirmed if $P \leq 0.10$. Quantification of heterogeneity performed using I^2 statistic. Values of 0-24.9%, 25-49.9%, 50-74.9% and $>75\%$ were considered as none, low, moderate and high heterogeneity respectively.[19]

Results

The literature search with the above mentioned criteria produced 894 findings on MEDLINE and 292 findings on EMBASE. The PRISMA flow diagram for screening and selection is shown in Figure 1. Overall, 1081 were judged not pertinent, and other 81 were excluded with reasons, leaving 24 findings. Of these, after verification of the period of inclusion/publication, we further excluded 3 studies since they included patients subsequently reported in updated versions of the same study by the same group of authors. This avoided to count twice or three times some patients. In particular, Meex et al. 2013[20] and Genbrugge et al. 2015[21] included patients reported subsequently in Genbrugge et al. 2018[22]. Similarly, Singer et al. 2014[9] included patients reported subsequently in Singer et al. 2018[23]. Of the remaining 21 studies, 3 were conference abstract on EMBASE and added for sensitivity analyses. Table 1 shows the characteristics of the remaining 18 studies (published in peer-reviewed journals) included in the primary analysis. Among these, one was considered only for sensitivity analysis as represented a small case series of 9 patients[24]. Studies are divided according to the location of CA and for each study we report the design, the country where it was performed, the baseline data on demographics and confounders, the NIRS device(s) used and the values of cerebral rSO_2 investigated by the authors.

All the studies included in this meta-analysis were non-randomized studied with observational design, and only one was a retrospective study. In particular, five studies included only patients presenting

with IHCA[12, 25-28], one study included a mixed population of both OHCA (28%) and IHCA (72%)[8] and 11 studies were conducted on OHCA patients[22, 23, 29-37]. For practicality, we considered in the population of OHCA studies one article where few patients (11%) experienced CA within the Emergency Department (on arrival to hospital)[29]. One case series was included for sensitivity analysis[24]. Among the included studies, one multicentre study was by far larger than the others (data on 1921 patients conducted in 15 hospitals in Japan)[32]. Another study was a multicentre collaboration between USA and United Kingdom[28].

Regarding the primary outcomes, initial cerebral rSO₂ values were reported by 11 studies (1 IHCA[26], 9 OHCA[22, 29-33, 35-37] and 1 mixed population[8]; 5 of these studies were conducted in Japan[30, 32, 36-38], 3 in Europe[22, 33, 35] and 3 in USA[8, 26, 29]); the values of overall cerebral rSO₂ during CPR were provided by nine studies (5 IHCA[12, 25-28] and 4 OHCA[22, 23, 29, 34]; one multicentre study across Europe and USA[28], 3 single center studies conducted in Europe[22, 25, 34] and 5 in USA[12, 23, 26, 27, 29] – none in Japan). No studies clearly reported the exact time between the CA and the first detected cerebral rSO₂ value.

Initial rSO₂ values

A total of 11 studies including 2870 patients reported data on initial cerebral rSO₂ values[8, 22, 26, 29-33, 35-37]. Of these, 476 achieved ROSC (16.6%), but excluding the largest study conducted in Japan, ROSC was achieved in 34.6% (n=328/949). ROSC was associated with significantly higher values of initial rSO₂ compared to the population that did not achieve ROSC (MD -11.54; 95%CI -20.96, -2.12; p=0.02; Figure 2), with high heterogeneity ($I^2=97%$) and no subgroup differences (p=0.52).

We performed a secondary analysis with geographic criteria, differentiating studies conducted in Japan vs Europe-USA. The analysis of five Japanese studies[30-32, 36, 37] (all of them in OHCA; 2212 patients; ROSC 11.2%) showed a trend towards an association between higher initial cerebral

rSO₂ values and ROSC (MD -16.16; 95% CI -32.76, 0.44; p=0.06; Figure 3), with high heterogeneity ($I^2=98\%$). The analysis conducted on six European-American studies[8, 22, 26, 29, 33, 35] (1 IHCA, 4 OHCA, 1 mixed population; 658 patients; ROSC 34.7%) showed a significant association between higher initial cerebral rSO₂ values and ROSC (MD -6.68; 95% CI -10.91, -2.45; p=0.002; Figure 4), with moderate heterogeneity ($I^2=64\%$) and no subgroup differences (p=0.56).

Overall rSO₂ values

Nine studies including 894 patients reported data on overall rSO₂ values during CPR, and 301 of them achieved ROSC (33.7%)[12, 22, 23, 25-29, 34]. ROSC was associated with significantly higher values of overall cerebral rSO₂ compared to the population that did not achieve ROSC (MD -10.38; 95% CI -13.73, -7.03; p<0.00001; Figure 5) with high heterogeneity ($I^2=77\%$) and subgroup differences (p=0.03). All the studies reporting the overall cerebral rSO₂ values during CPR and ROSC were conducted in Europe or USA, thus the secondary analysis with geographical criteria was not meaningful.

Sensitivity analyses

An initially unplanned sensitivity analysis was performed excluding the study of Nishihama et al.[32] from the analysis of initial cerebral rSO₂ values as this large multicentre study represented a clear outlier. Its exclusion did not change the results of the primary analysis (MD -8.30; 95% CI -12.07, -4.54; p=0.008; high heterogeneity $I^2=80\%$; no subgroup differences p=0.58). However, its exclusion from the secondary analysis of initial rSO₂ values in the Japanese studies changed the trend (p=0.06) towards a significant association between higher initial cerebral rSO₂ values and ROSC (MD -10.51; 95% CI -20.52, -0.50; p=0.04; high heterogeneity $I^2=91\%$). Readers should keep in mind that exclusions of outliers are not associated with worse study design or quality and indeed in this case the outlier was a high-quality multicentre study.[32]

The first group of planned sensitivity analyses was conducted with *leave-one-out at time* approach; none of these changed the results of the primary analyses regarding the initial or the overall values of cerebral rSO₂ and ROSC.

The sensitivity analyses performed including the feasibility study (case series of 9 patients) conducted by Tajima et al.[24] did not change any of the results. Similarly, the inclusion of the data from the 3 conference abstracts (Supplemental Digital Content – Appendix 3) did not change the results. Regarding the risk of bias assessment, the criteria for assigning (or not) the points are reported; as all the included studies were at low risk of bias scoring between 7 and 9 points of the NOS (Supplemental Digital Content – Appendix 4); therefore, no sensitivity analyses were conducted in this regard.

We conducted a post-hoc calculation on the mean initial and overall values of rSO₂, calculating it from the values of each study weighted for their sample size. For the mean initial rSO₂ value, we found that patients achieving ROSC had a mean of 43.8% vs 19% in those not achieving ROSC (absolute difference 24.8%). Similarly, for the mean overall value during resuscitation, we calculated that patients who achieved ROSC had a mean cerebral rSO₂ value of 40.8% vs 30.9% in those not achieving ROSC (absolute difference 9.9%).

Discussion

This updated meta-analysis confirmed a significant association between both higher initial and overall values of cerebral rSO₂ and ROSC in CA patients undergoing CPR. This finding is reinforcing the results of the precedent meta-analysis[13], with the advantage of a much larger sample size. Moreover, the previous meta-analysis included 9 studies but 4 of them were conducted by one research group, and 2 by another group, thus limiting the external validity of previous findings. In the present study, the inclusion of 17 studies and the removal of studies from the same research group

with overlaps in the recruitment period (and thus included patients) increases the external validity of our findings.

Importantly, we also conducted analyses based on “geographical” criteria, separating the studies reporting initial rSO₂ values conducted in Europe-USA from those performed in Japan. Our idea was based on the significant difference in the management of OHCA in Japan, where the emergency medical personnel is not allowed to terminate CPR in the OH setting (unless rare occasions), and consequently all OHCA patients are brought to ED. These analyses confirmed the results of the primary analysis, though the association between the initial cerebral rSO₂ values and ROSC was less clear when analysing Japanese studies only (p=0.06, though become significant excluding one large multicentre study behaving as outlier[32]).

The association of rSO₂ values with ROSC is expected for several reasons. Values of cerebral rSO₂ are correlated with oxygenation measured at the jugular venous bulb[39, 40] and, more importantly, with coronary and cerebral perfusion pressure.[41-43] Thus, higher values of cerebral oxygenation may theoretically indicate better coronary perfusion pressure and a greater efficacy of CPR. Indeed, a coronary perfusion pressure above 15 mmHg should be maintained during CPR in order to obtain a successful resuscitation.[44] As there is no available tool for measuring coronary perfusion pressure during CPR, cerebral oxygenation may represent a reasonable surrogate, considering also that application of NIRS sensors on the forehead is easy and feasible without causing delays in CPR. However, one study on quality controlled manual chest compression found that high quality CPR was not significantly reflected in the cerebral rSO₂ quantified by NIRS.[45] Studies evaluating the value of rSO₂ according to the CPR quality are needed. Nonetheless, it is important to consider that cerebral monitoring has been consistently used in patients undergoing cardiac surgery during the non-pulsatile perfusion with cardiopulmonary bypass.[46]

Considering the mean values of each study and accounting for each study’s size we calculated the weighted mean for the initial and the overall values of cerebral rSO₂. We found that the difference

between weighted means for ROSC and no-ROSC group were larger for the initial NIRS values (almost 25% greater in successfully resuscitated patients) as compared to overall NIRS values (almost 10% higher in those achieving ROSC). Whether the initial value may be of greater help in prognostication remains to be established.

The end-tidal CO₂ (etCO₂) is the most extensively studied indicator of CPR quality and predictor of ROSC.[47] It is interesting that Engel et al.[29] showed that both rSO₂ and etCO₂ are good predictors of ROSC, but rSO₂ performed better than etCO₂. This may be partially explained by the dependency of etCO₂ on both effective minute ventilation and administration of vasoactive drugs.[48-50] The etCO₂ has been investigated and it is recommended as a tool indicating the efficacy of chest compressions;[51] also rSO₂ is currently investigated for this scope but more studies are needed before drawing conclusions. The rSO₂ has the advantage of being theoretically applicable to all patients, whilst the use of etCO₂ requires advanced airway management.

Values of cerebral rSO₂ have been also considered for prognostication of neurological outcome after CA. Two studies[10, 11] found significantly higher rSO₂ values measured at hospital arrival in patients with favourable neurological outcome (56% and 68%, respectively) as compared with values in patients with poor outcome (20% and 58%, respectively). However, Bougle et al. reported no difference in the mean values rSO₂ during the first 48 hours after CA between favourable (62%) and poor neurological outcome (58%)[52]. Of note, all these data should be interpreted considering that several variables may influence cerebral oximetry, starting from hemodynamic and respiratory management, but taking into account also the influence of target temperature management.[53]

Limitations

This meta-analysis has several limitations. First, our meta-analysis investigates only the associations between NIRS values and ROSC, but the lack of individual patient data does not allow to correct for confounders performing analysis accounting for age, cause and location of CA, first detected rhythm,

witnessed CA and presence of bystander CPR, etc. This should not be confused with the risk of bias assigned to each study according to the quality in its design. Indeed, the risk of bias in individual studies is subjective and open to interpretation, and the low risk of bias in the included studies does not mean low risk of bias in the meta-analysis findings. In practice, our meta-analysis investigated the association between NIRS values and ROSC, but it cannot comment on correlation and causation. Secondly, the delay between CA and first application of the cerebral oximetry sensors was not clearly specified by the included studies and it greatly influences the first recorded cerebral oximetry value. For this reason we conducted a “geographical” analysing separately the studies conducted in Japan where OHCA patients are transported to the nearest hospital (not permitted to terminate resuscitation on the scene) and the application of the NIRS sensors in these patients may happen long time after the CA. Third, study size varied greatly from small single center to large multicentre (international) studies. In this context, it should be kept in mind that the largest muticenter study conducted in Japan[32] did not have a great influence on the results but it significantly skewed the 95% CI. Fourth, as shown in Table 1, different devices have been used to measure rSO₂. The multiple NIRS devices commercially available use different algorithms to adjust for confounding factors such as the interference of extra-cerebral tissue and spatial resolution between and superficial brain tissue,[54] the differences in the cerebral arterial/venous blood partitioning[54] (commonly considered as a ratio of 30%/70%),[55] the effect of blood- and tissue- derived chromophores (i.e. melanin, haemoglobin, myoglobin, conjugated bilirubin).[56] Myoglobin absorbency signal overlaps with hemoglobin. Chromophore concentration cannot be calculated by the Beer-Lambert law equations which is a limitation of tissue NIRS. Thus, the use of different devices potentially introduces a monitoring bias due to differences in calibration and algorithms of each device.[57] Moreover, there is a significant risk of contamination of the NIRS signal by extracerebral sources as strict probe placement on frontal lobe skin avoiding hair follicles and attachment of the probes may be difficult if the skin is wet. Probes may also be influenced by ambient light.[58]

Finally, the value of cerebral oxygen saturation for the prognostication on neurological outcome after CA is at early stage of research, and it could not be properly analysed. More studies are warranted to understand the relationship between initial and overall values of cerebral oxygen saturation with neurological outcome after CA.

Conclusions

Greater initial and overall values of cerebral oxygen saturation monitored with near infrared spectrometry are both associated with more possibilities of achieving return of spontaneous circulation in patients with cardiac arrest. This finding seems consistent for both in-hospital and out-hospital cardiac arrest settings. Moreover, we showed consistency with a geographical analysis conducted differentiating Japan from other countries.

Conflict of interest statement: The authors declare no conflict of interest in regards of this work.

On behalf of all co-authors,

F Sanfilippo

Conflict of interest statement

All the authors declare no conflict of interest with regards of this topic.

Fundings

No authors received funding for undertaking this project.

References


- [1] Gräsner JT, Lefering R, Koster RW, Masterson S, Böttiger BW, Herlitz J, et al. EuReCa ONE-27 Nations, ONE Europe, ONE Registry: A prospective one month analysis of out-of-hospital cardiac arrest outcomes in 27 countries in Europe. *Resuscitation*. 2016;105:188-95.
- [2] Okubo M, Schmicker RH, Wallace DJ, Idris AH, Nichol G, Austin MA, et al. Variation in Survival After Out-of-Hospital Cardiac Arrest Between Emergency Medical Services Agencies. *JAMA cardiology*. 2018;3:989-99.
- [3] Perkins GD, Ji C, Deakin CD, Quinn T, Nolan JP, Scomparin C, et al. A Randomized Trial of Epinephrine in Out-of-Hospital Cardiac Arrest. *The New England journal of medicine*. 2018;379:711-21.
- [4] Drennan IR, Lin S, Sidalak DE, Morrison LJ. Survival rates in out-of-hospital cardiac arrest patients transported without prehospital return of spontaneous circulation: an observational cohort study. *Resuscitation*. 2014;85:1488-93.
- [5] Laver S, Farrow C, Turner D, Nolan J. Mode of death after admission to an intensive care unit following cardiac arrest. *Intensive care medicine*. 2004;30:2126-8.
- [6] Sanfilippo F, Li Volti G, Ristagno G, Murabito P, Pellis T, Astuto M, et al. Clinical biomarkers in brain injury: a lesson from cardiac arrest. *Frontiers in bioscience (Scholar edition)*. 2010;2:623-40.
- [7] Skhirtladze-Dworschak K, Dworschak M. Cerebral oximetry and cardiac arrest. *Seminars in cardiothoracic and vascular anesthesia*. 2013;17:269-75.
- [8] Ahn A, Nasir A, Malik H, D'Orazi F, Parnia S. A pilot study examining the role of regional cerebral oxygen saturation monitoring as a marker of return of spontaneous circulation in shockable (VF/VT) and non-shockable (PEA/Asystole) causes of cardiac arrest. *Resuscitation*. 2013;84:1713-6.
- [9] Singer AJ, Ahn A, Inigo-Santiago LA, Thode HC, Jr., Henry MC, Parnia S. Cerebral oximetry levels during CPR are associated with return of spontaneous circulation following cardiac arrest: an observational study. *Emergency medicine journal : EMJ*. 2015;32:353-6.
- [10] Ito N, Nishiyama K, Callaway CW, Orita T, Hayashida K, Arimoto H, et al. Noninvasive regional cerebral oxygen saturation for neurological prognostication of patients with out-of-hospital cardiac arrest: a prospective multicenter observational study. *Resuscitation*. 2014;85:778-84.
- [11] Storm C, Leithner C, Krannich A, Wutzler A, Ploner CJ, Trenkmann L, et al. Regional cerebral oxygen saturation after cardiac arrest in 60 patients--a prospective outcome study. *Resuscitation*. 2014;85:1037-41.
- [12] Parnia S, Nasir A, Ahn A, Malik H, Yang J, Zhu J, et al. A feasibility study of cerebral oximetry during in-hospital mechanical and manual cardiopulmonary resuscitation*. *Critical care medicine*. 2014;42:930-3.
- [13] Sanfilippo F, Serena G, Corredor C, Benedetto U, Maybauer MO, Al-Subaie N, et al. Cerebral oximetry and return of spontaneous circulation after cardiac arrest: A systematic review and meta-analysis. *Resuscitation*. 2015;94:67-72.
- [14] Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gotzsche PC, Ioannidis JP, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration. *BMJ (Clinical research ed)*. 2009;339:b2700.
- [15] Wells G, Shea B, O'Connell D, Peterson J, Welch V, Losos M, et al. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomized studies in meta-analyses. http://www.ohri.ca/programs/clinical_epidemiology/oxford.htm.
- [16] Wan X, Wang W, Liu J, Tong T. Estimating the sample mean and standard deviation from the sample size, median, range and/or interquartile range. *BMC medical research methodology*. 2014;14:135.
- [17] Luo D, Wan X, Liu J, Tong T. Optimally estimating the sample mean from the sample size, median, mid-range, and/or mid-quartile range. *Statistical methods in medical research*. 2018;27:1785-805.
- [18] Hozo SP, Djulbegovic B, Hozo I. Estimating the mean and variance from the median, range, and the size of a sample. *BMC medical research methodology*. 2005;5:13.
- [19] Higgins J, Thompson S, Deeks J, Altman D. Measuring inconsistency in meta-analyses. *BMJ*. 2003;327:557-60.
- [20] Meex I, De Deyne C, Dens J, Scheyltjens S, Lathouwers K, Boer W, et al. Feasibility of absolute cerebral tissue oxygen saturation during cardiopulmonary resuscitation. *Critical care*. 2013;17:R36.
- [21] Genbrugge C, Meex I, Boer W, Jans F, Heylen R, Ferdinande B, et al. Increase in cerebral oxygenation during advanced life support in out-of-hospital patients is associated with return of spontaneous circulation. *Critical care*. 2015;19:837.

- [22] Genbrugge C, De Deyne C, Eertmans W, Anseeuw K, Voet D, Mertens I, et al. Cerebral saturation in cardiac arrest patients measured with near-infrared technology during pre-hospital advanced life support. Results from Copernicus I cohort study. *Resuscitation*. 2018;129:107-13.
- [23] Singer AJ, Nguyen RT, Ravishankar ST, Schoenfeld ER, Thode HC, Jr., Henry MC, et al. Cerebral oximetry versus end tidal CO₂ in predicting ROSC after cardiac arrest. *The American journal of emergency medicine*. 2018;36:403-7.
- [24] Tajima G, Shiozaki T, Izumino H, Yamano S, Hirao T, Inokuma T, et al. Portable system for monitoring of regional cerebral oxygen saturation during prehospital cardiopulmonary resuscitation: a pilot study. *Acute medicine & surgery*. 2015;2:48-52.
- [25] Yazar MA, Açıkgöz MB, Bayram A. Does chest compression during cardiopulmonary resuscitation provide sufficient cerebral oxygenation? *Turkish journal of medical sciences*. 2019;49:311-7.
- [26] Ibrahim AW, Trammell AR, Austin H, Barbour K, Onuorah E, House D, et al. Cerebral Oximetry as a Real-Time Monitoring Tool to Assess Quality of In-Hospital Cardiopulmonary Resuscitation and Post Cardiac Arrest Care. *Journal of the American Heart Association*. 2015;4:e001859.
- [27] Parnia S, Nasir A, Shah C, Patel R, Mani A, Richman P. A feasibility study evaluating the role of cerebral oximetry in predicting return of spontaneous circulation in cardiac arrest. *Resuscitation*. 2012;83:982-5.
- [28] Parnia S, Yang J, Nguyen R, Ahn A, Zhu J, Inigo-Santiago L, et al. Cerebral Oximetry During Cardiac Arrest: A Multicenter Study of Neurologic Outcomes and Survival. *Critical care medicine*. 2016;44:1663-74.
- [29] Engel TW, 2nd, Thomas C, Medado P, Bastani A, Reed B, Millis S, et al. End tidal CO₂ and cerebral oximetry for the prediction of return of spontaneous circulation during cardiopulmonary resuscitation. *Resuscitation*. 2019;139:174-81.
- [30] Fukuda T, Ohashi N, Nishida M, Gunshin M, Doi K, Matsubara T, et al. Application of cerebral oxygen saturation to prediction of the futility of resuscitation for out-of-hospital cardiopulmonary arrest patients: a single-center, prospective, observational study: can cerebral regional oxygen saturation predict the futility of CPR? *The American journal of emergency medicine*. 2014;32:747-51.
- [31] Koyama Y, Wada T, Lohman BD, Takamatsu Y, Matsumoto J, Fujitani S, et al. A new method to detect cerebral blood flow waveform in synchrony with chest compression by near-infrared spectroscopy during CPR. *The American journal of emergency medicine*. 2013;31:1504-8.
- [32] Nishiyama K, Ito N, Orita T, Hayashida K, Arimoto H, Abe M, et al. Characteristics of regional cerebral oxygen saturation levels in patients with out-of-hospital cardiac arrest with or without return of spontaneous circulation: A prospective observational multicentre study. *Resuscitation*. 2015;96:16-22.
- [33] Prosen G, Strnad M, Doniger SJ, Markota A, Stožer A, Borovnik-Lesjak V, et al. Cerebral tissue oximetry levels during prehospital management of cardiac arrest - A prospective observational study. *Resuscitation*. 2018;129:141-5.
- [34] Schewe JC, Thudium MO, Kappler J, Steinhagen F, Eichhorn L, Erdfelder F, et al. Monitoring of cerebral oxygen saturation during resuscitation in out-of-hospital cardiac arrest: a feasibility study in a physician staffed emergency medical system. *Scandinavian journal of trauma, resuscitation and emergency medicine*. 2014;22:58.
- [35] Storm C, Wutzler A, Trenkmann L, Krannich A, von Rheinbarben S, Luckenbach F, et al. Good neurological outcome despite very low regional cerebral oxygen saturation during resuscitation--a prospective preclinical trial in 29 patients. *Scandinavian journal of trauma, resuscitation and emergency medicine*. 2016;24:43.
- [36] Takegawa R, Shiozaki T, Ogawa Y, Hirose T, Mori N, Ohnishi M, et al. Usefulness of cerebral rSO₂ monitoring during CPR to predict the probability of return of spontaneous circulation. *Resuscitation*. 2019;139:201-7.
- [37] Tsukuda J, Fujitani S, Morisawa K, Shimosawa N, Lohman BD, Okamoto K, et al. Near-infrared spectroscopy monitoring during out-of-hospital cardiac arrest: can the initial cerebral tissue oxygenation index predict ROSC? *Emergency medicine journal : EMJ*. 2019;36:33-8.
- [38] Koyama Y, Wada T, Takamatsu Y, Takita M, Lohman BD, Matsumoto J, et al. Real-time Cerebral Hemoglobin Concentration Changes From NIRS; a Valid Evaluation Method for Quality CPR Chest Compression in Out-of-Hospital Cardiac Arrest. *Circulation*. 2013;128:A279.
- [39] Ikeda K, MacLeod DB, Grocott HP, Moretti EW, Ames W, Vacchiano C. The Accuracy of a Near-Infrared Spectroscopy Cerebral Oximetry Device and Its Potential Value for Estimating Jugular Venous Oxygen Saturation. *Anesthesia and analgesia*. 2014.

- [40] Daubeney PE, Pilkington SN, Janke E, Charlton GA, Smith DC, Webber SA. Cerebral oxygenation measured by near-infrared spectroscopy: comparison with jugular bulb oximetry. *The Annals of thoracic surgery*. 1996;61:930-4.
- [41] Friess SH, Sutton RM, French B, Bhalala U, Maltese MR, Naim MY, et al. Hemodynamic directed CPR improves cerebral perfusion pressure and brain tissue oxygenation. *Resuscitation*. 2014;85:1298-303.
- [42] Debaty G, Moore J, Duhem H, Rojas-Salvador C, Salverda B, Lick M, et al. Relationship between hemodynamic parameters and cerebral blood flow during cardiopulmonary resuscitation. *Resuscitation*. 2020;153:20-7.
- [43] Yannopoulos D, McKnite S, Aufderheide TP, Sigurdsson G, Pirralo RG, Benditt D, et al. Effects of incomplete chest wall decompression during cardiopulmonary resuscitation on coronary and cerebral perfusion pressures in a porcine model of cardiac arrest. *Resuscitation*. 2005;64:363-72.
- [44] Paradis NA, Martin GB, Rivers EP, Goetting MG, Appleton TJ, Feingold M, et al. Coronary perfusion pressure and the return of spontaneous circulation in human cardiopulmonary resuscitation. *JAMA : the journal of the American Medical Association*. 1990;263:1106-13.
- [45] Kamarainen A, Sainio M, Olkkola KT, Huhtala H, Tenhunen J, Hoppu S. Quality controlled manual chest compressions and cerebral oxygenation during in-hospital cardiac arrest. *Resuscitation*. 2012;83:138-42.
- [46] Douds M, Straub E, Kent A, Bistrick C, Sistino J. A systematic review of cerebral oxygenation-monitoring devices in cardiac surgery. *Perfusion*. 2014;29:545-52.
- [47] Hamrick JL, Hamrick JT, Lee JK, Lee BH, Koehler RC, Shaffner DH. Efficacy of chest compressions directed by end-tidal CO₂ feedback in a pediatric resuscitation model of basic life support. *Journal of the American Heart Association*. 2014;3:e000450.
- [48] Xu J, Yu X, Zhang L, Fu Y, Jin K, Yin L, et al. Modified volumetric capnography-derived parameter: A potentially stable indicator in monitoring cardiopulmonary resuscitation efficacy in a porcine model. *Resuscitation*. 2020;150:94-101.
- [49] Lindberg L, Liao Q, Steen S. The effects of epinephrine/norepinephrine on end-tidal carbon dioxide concentration, coronary perfusion pressure and pulmonary arterial blood flow during cardiopulmonary resuscitation. *Resuscitation*. 2000;43:129-40.
- [50] Callaham M, Barton C, Matthay M. Effect of epinephrine on the ability of end-tidal carbon dioxide readings to predict initial resuscitation from cardiac arrest. *Critical care medicine*. 1992;20:337-43.
- [51] Soar J, Nolan JP, Böttiger BW, Perkins GD, Lott C, Carli P, et al. European Resuscitation Council Guidelines for Resuscitation 2015: Section 3. Adult advanced life support. *Resuscitation*. 2015;95:100-47.
- [52] Bouglé A, Daviaud F, Bougouin W, Rodrigues A, Geri G, Morichau-Beauchant T, et al. Determinants and significance of cerebral oximetry after cardiac arrest: A prospective cohort study. *Resuscitation*. 2016;99:1-6.
- [53] Pellis T, Sanfilippo F, Ristagno G. The optimal hemodynamics management of post-cardiac arrest shock. *Best practice & research Clinical anaesthesiology*. 2015;29:485-95.
- [54] Ohmae E, Ouchi Y, Oda M, Suzuki T, Nobesawa S, Kanno T, et al. Cerebral hemodynamics evaluation by near-infrared time-resolved spectroscopy: correlation with simultaneous positron emission tomography measurements. *NeuroImage*. 2006;29:697-705.
- [55] Ito H, Kanno I, Fukuda H. Human cerebral circulation: positron emission tomography studies. *Annals of nuclear medicine*. 2005;19:65-74.
- [56] Madsen PL, Skak C, Rasmussen A, Secher NH. Interference of cerebral near-infrared oximetry in patients with icterus. *Anesthesia and analgesia*. 2000;90:489-93.
- [57] Murkin JM, Arango M. Near-infrared spectroscopy as an index of brain and tissue oxygenation. *British journal of anaesthesia*. 2009;103 Suppl 1:i3-13.
- [58] Wik L. Near-infrared spectroscopy during cardiopulmonary resuscitation and after restoration of spontaneous circulation: a valid technology? *Current opinion in critical care*. 2016;22:191-8.



Modified PRISMA 2009 Flow Diagram



From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(6): e1000097. doi:10.1371/journal.pmed1000097

For more information, visit www.prisma-statement.org.

Figure 1. PRISMA flow diagram for screening and selection of studies found by the systematic search.




Figure 2. Forest plot investigating the association between initial values of cerebral regional oxygen saturation (rSO₂) and return of spontaneous circulation (ROSC). Subgroup analysis is conducted according to location of cardiac arrest (CA). CI: Confidence interval; IV=inverse variance; IH: In Hospital; OH: Out of Hospital; SD: standard deviation.




Figure 3. Forest plot investigating the association between initial values of cerebral regional oxygen saturation (rSO₂) and return of spontaneous circulation (ROSC), with analysis conducted including Japanese studies only. CA: cardiac arrest; CI: Confidence interval; IV=inverse variance; IH: In Hospital; OH: Out of Hospital; SD: standard deviation.




Figure 4. Forest plot investigating the association between initial values of cerebral regional oxygen saturation (rSO₂) and return of spontaneous circulation (ROSC), with analysis conducted including studies from Europe and USA only (excluding Japanese studies). CA: cardiac arrest; CI: Confidence interval; IV=inverse variance; IH: In Hospital; OH: Out of Hospital; SD: standard deviation.




Figure 5. Forest plot investigating the association between overall values of cerebral regional oxygen saturation (rSO_2) during resuscitation and return of spontaneous circulation (ROSC). Subgroup analysis is conducted according to location of cardiac arrest (CA). CI: Confidence interval; IV=inverse variance; IH: In Hospital; OH: Out of Hospital; SD: standard deviation.

Table 1. Characteristics of the studies included in the meta-analysis. Studies are divided according to the site of cardiac arrest (IH: in-hospital; OH: out-of-hospital). The summary of characteristics also includes the country where the study was performed and its design, the number of patients with data available with the data on population characteristics, the near-infrared spectroscopy (NIRS) device used for monitoring cerebral oximetry, and the type of data available for outcome analysis. The last column indicates the reference number.

*Ahn et al. 2013: we used only the values of overall rSaO₂. Values of initial rSaO₂ are presented in subsequent studies from the same group of authors, thus have not been used to avoid duplicates.

** Engle et al 2019 was included amongst OHCA but a small amount of patients suffered from CA in the Emergency Department (n=19/173, 11%).

***Nishihama 2015: authors provided both values from left and right hemisphere. We arbitrarily used values from left hemisphere (dominant). However, values from right hemisphere were very similar and using these data did not change results.

#Parnia 2012: The data on age and first detected rhythm refers to a population of 19 patients (of these 15 were included in the study on cerebral oximetry).

##Tajima 2015: the study is in italic font as it was used only for sensitivity analysis,. Indeed, the study was an initial experience (case series) on nine patients with OHCA and monitored with a new NIRS portable device.

Study	Country, Design	N	Age -Males	NIRS device used	Oxymetry value	Ref
			ROSC rate			
		CPR duration	Shockable rhythm			
MIXED POPULATION						
Ahn 2013	USA	36 IH, 14 OH	65 y - 72%	Equanox 7600,Nonin Medical, Inc, Plymouth, MN,USA	Initial and overall*	8
	Prospective	52%	NR			
		34 min	24%			
OUT OF HOSPITAL CARDIAC ARREST						
Engel 2019	USA	173**	62 y - 65%	CerOx INVOS	Initial and overall	29
	Prospective	20%	67%			
		NR	26%			

Takegawa 2019	Japan	90	75 y - 66%	TOS-OR, Fujita Medical Instruments Co., LTD, Tokyo, Japan	Initial	36
	Retrospective	39%	44%			
		NR	8%			
Genbrugge 2018	Belgium	329	69 y - 70%	Equanox 7600 or SenSmart Model X-100, Nonin Medical Inc Plymouth, MN, USA	Initial and overall	22
	Prospective	33%	58%			
		26 min	20%			
Singer 2018	USA	100	69 y - 73%	Equanox 7600, Nonin Medical, Plymouth, MN, USA	Overall	23
	Prospective	33%	67%			
		NR	14%			
Prosen 2018	Slovenia	53	68 y - 85%	Invos Oximeter, Somanetics Corporation, Troy, MI, USA	Initial	33
	Prospective	42%	91%			
		37 min	28%			
Tsukuda 2018	Japan	117	70 y - 57%	NIRO-200 NX, Hamamatsu Photonics, Hamamatsu-City, Shizuoka, Japan	Initial	37
	Prospective	38%	51%			
		NR	11%			
Storm 2016	Germany	23	62 y - 78%	Invos 5100 C, Covidien, Mansfield, USA	Initial	11
	Prospective	22%	NR			
		NR	22%			
Tajima 2015^{##}	Japan	9	74 y - 44%	HAND ai TOS; TOSTEC Co., Ltd, Tokyo, Japan	Initial	24
	Prospective	33%	55%			
		NR	11%			
Nishihama 2015 (multicenter=15)	All in Japan	1921	76 y - 61%	INVOS™ 5100C, Covidien, Boulder, CO, USA	Initial***	32
	Prospective	8%	50%			
		NR	11%			
Shewe 2014	Germany	10	73 y - 80%	Equanox 7600, Nonin Medical, Plymouth, Minnesota, USA	Overall	34
	Prospective	30%	NR			
		NR	30%			
Fukuda 2014	Japan	69	66 y - 70%	Invos 5100C, Covidien, Boulder, CO, USA	Initial	30
	Prospective	23%	43%			
		NR	17%			

Koyama 2013	Japan	15	80 y - 67%	NIRO Hamamatsu Photonics, Hamamatsu-shi, Shizuoka, Japan	Initial	31
	Prospective	33%	NR			
		84 min	7%			
IN HOSPITAL CARDIAC ARREST						
Yazar 2019	Turkey	20	75 y - 40%	Invos 5100c, Somanetics, Troy, MI, USA	Overall	25
	Prospective	40%	NR			
		24 min	NR			
Parnia 2016 (multicenter=5)	1 USA, 4 UK	183	69 y - 61%	Equanox 7600, Nonin Medical, Plymouth, MN, USA	Overall	28
	Prospective	34%	NR			
		29 min	10%			
Ibrahim 2015	USA	27	66 y - 96%	Invos 5100c, Somanetics, Troy, USA	Initial and overall	26
	Prospective	70%	NR			
		20 min	15%			
Parnia 2014	USA	34	71 y - 65%	Equanox 7600; Nonin, Plymouth, MI, and Invos; Somanetics, Troy, MI	Overall	12
	Prospective	44%	NR			
		19 min	15%			
Parnia 2012	USA	15	74 y [#] - NR	Invos Somanetics, Troy, USA	Overall	27
	Prospective	33%	NR			
		16 min	5% [#]			

Table 2. Characteristics of the studies included in the meta-analysis.

Studies are divided according to the site of cardiac arrest (IH: in-hospital; OH: out-of-hospital). The summary of characteristics includes country where the study was performed **and its design**, the number of patients with data available **with the data on population characteristics**, the near-infrared spectroscopy (NIRS) device used for monitoring cerebral oximetry, and the type of data available for outcome analysis. **The last column indicates the reference number.**

*Ahn et al. 2013: we used only the values of overall rSaO₂. Values of initial rSaO₂ are presented in subsequent studies from the same group of authors, thus have not been used to avoid duplicates.

** Engle et al 2019 was included amongst OHCA but a small amount of patients suffered from CA in the Emergency Department (n=19/173, 11%).

***Nishihama 2015: authors provided both values from left and right hemisphere. We arbitrarily used values from left hemisphere (dominant). However, values from right hemisphere were very similar and using these data did not change results.

Parnia 2012: The data on age and first detected rhythm refers to a population of 19 patients (of these 15 were included in the study on cerebral oximetry).

Tajima 2015: the study is in italic font as it was used only for sensitivity analysis,. Indeed, the study was an initial experience (case series) on nine patients with OHCA and monitored with a new NIRS portable device.

Journal Pre-proof