Low influenza vaccination coverage among hospitalized COVID-19 patients in Milan: a gap to be urgently filled.

Anna Lisa Ridolfo, Laura Milazzo, Andrea Giacomelli, Letizia Oreni, Catia Borriello, Spinello Antinori

 PII:
 S0953-6205(20)30471-4

 DOI:
 https://doi.org/10.1016/j.ejim.2020.12.019

 Reference:
 EJINME 4701

To appear in: European Journal of Internal Medicine

Received date:20 October 2020Accepted date:24 December 2020

Please cite this article as: Anna Lisa Ridolfo, Laura Milazzo, Andrea Giacomelli, Letizia Oreni, Catia Borriello, Spinello Antinori, Low influenza vaccination coverage among hospitalized COVID-19 patients in Milan: a gap to be urgently filled., *European Journal of Internal Medicine* (2020), doi: https://doi.org/10.1016/j.ejim.2020.12.019

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier B.V. on behalf of European Federation of Internal Medicine.



Low influenza vaccination coverage among hospitalized COVID-19 patients in Milan: a gap to be urgently filled.

Anna Lisa Ridolfo<sup>1</sup>, Laura Milazzo<sup>1</sup>, Andrea Giacomelli<sup>1,2</sup>, Letizia Oreni<sup>1</sup>, Catia Borriello<sup>3</sup>, Spinello Antinori

<sup>1</sup>Department of Infectious Diseases, ASST Fatebenefratelli Sacco, Milan, Italy

<sup>2</sup> Luigi Sacco Department of Biomedical and Clinical Sciences, University of Milan, Italy

<sup>3</sup> Vaccination Unit, ASST Fatebenefratelli Sacco, Milan, Italy

The first wave of the COVID-19 epidemic in Italy during March and April 2020 put extreme pressure on the national healthcare system, particularly on the hospitals in the most effected regions of northern Italy. Although the drastic measures taken to contain SARS-CoV-2 transmission (including a nationwide lockdown) led to a marked decrease in the number of new COVID-19 cases and deaths during the summer, the number of new cases is currently rising again at an alarming rate [1], and the coming winter raises further concerns about the additional effects the seasonal influenza epidemic will have on already stressed healthcare services.

Both influenza and COVID-19 can present with non-specific symptoms (fever, myalgia, headache, nonproductive cough and shortness of breath), most of which are self-limiting but may also progress to severe conditions. Moreover, like COVID-19, influenza puts older people and those with chronic medical conditions at higher risk of morbidity and mortality [2-4]. This is why extended influenza vaccination programmes aimed particularly at such risk groups are more than ever necessary to avoid the overload of health services and hospitals due to the concomitant diffusion of influenza and COVID-19 infections with the subsequent challenge of differential diagnosis. However, this means overcoming the existing barriers to vaccinations that are responsible for the historical sub-optimal use of influenza vaccine in many countries (including Italy, where it is offered free of charge to people at high risk and healthcare providers) [5].

In a bid to contextualise this issue, we have analysed the patients living in the province of Milan who were admitted to our referral COVID-19 centre between 21 February and 31 May 2020 in order to calculate the proportion who received the influenza vaccination for the 2019-2020 season. The data regarding seasonal influenza vaccination uptake were extracted from the electronic immunisation register of Milan's Health Protection Agency.

The study population consisted of 428 subjects (146 females and 282 males with a median age of 60.6 years, range 48.9-72.7), of whom 335 (78.3%) belonged to one or more of the categories for which influenza vaccination is recommended by the Italian national immunisation guidelines: 174 (40.7%) were aged  $\geq$ 65 years, 291 (68.0%) had underlying chronic diseases, and 57 (13.1%) were healthcare workers. However, only 109 of the patients who should have been vaccinated actually received the vaccine: 44.2% of those aged  $\geq$ 65 years, 33.0% of those with underlying chronic diseases, and 31.6% of healthcare workers.

Our data show disappointingly low influenza vaccination coverage among the subjects at risk and healthcare workers admitted to our COVID-19 centre during the first wave of the epidemic. Particularly in the elderly the coverage was far below the minimum *recommended threshold of 75%* and consistent with the worrying decline of influenza vaccination registered in Lombardy over the last decade [6]. This clearly calls for the prompt reinforcement of the seasonal influenza vaccination information campaign and a greater supply of vaccine in order to make the most of the benefits of influenza vaccination for individual well-being and the functioning of the healthcare system at large over such a critical time. Moreover, emerging data suggest that influenza vaccinations may have a protective effect against the risk of experiencing severe COVID-19, especially among high-risk groups [7-9], which may be of paramount importance given the paucity of effective COVID-19 treatments.

#### References

1. Covid-19 - Situazione in Italia. http://opendatadpc.maps.arcgis.com/apps/opsdashboard/

index.html#/b0c68bce2cce478eaac82fe38d4138b.

- Clark A, Jit M, Warren-Gash C, Guthrie B, Wang HHX, Mercer SW, et al. Global, regional, and national estimates of the population at increased risk of severe COVID-19 due to underlying health conditions in 2020: a modelling study. Lancet Glob Health 2020;8:e1003-e1017. DOI: 10.1016/S2214-109X(20)30264-3.
- 3. Nicoll A, Ciancio B, Tsolova S, Blank P, Yilmaz C. The scientific basis for offering seasonal influenza immunisation to risk groups in Europe. Euro Surveill 2008;13:19018. DOI: 10.2807/ese.13.43.19018-en.
- 4. Demicheli V, Jefferson T, Di Pietrantonj C, Ferroni E, Thorning S, Thomas RE, et al. Vaccines for preventing influenza in the elderly. Cochrane Database Syst Rev 2018;2:CD004876.
- European Centre for Disease Prevention and Control. Seasonal influenza vaccination and antiviral use in EU/EEA Member States – Overview of vaccine recommendations for 2017–2018 and vaccination coverage rates for 2015–2016 and 2016–2017 influenza seasons. Stockholm: ECDC; 2018.
- Rossi D, Croci R, Affanni P, Odone A, Signorelli C. Influenza vaccination coverage in Lombardy Region: a twenty-year trend analysis (1999-2019). Acta Biomed 2020;91 (Supplement 3):141-5 DOI: 10.23750/abm.v91i3-S.9455.
- Amato M, Werba JP, Frigerio B, Coggi D, Sansaro D, Ravani A, et al. Relationship between Influenza Vaccination Coverage Rate and COVID-19 Outbreak: An Italian Ecological Study. Vaccines 2020;8:E535; doi: 10.3390/vaccines8030535.
- Zanettini C, Omar M, Dinalankara W, Imada EL, Colantuoni E, Parmigiani G, et al. Influenza Vaccination and COVID19 Mortality in the USA. Preprint. medRxiv. 2020;2020.06.24.20129817. Published 2020 Jun 26. doi:10.1101/2020.06.24.20129817

9. Fink G, Orlova-Fink N, Schindler T, Grisi S, Ferrer AP, Daubenberger C, et al. Inactivated trivalent influenza vaccine is associated with lower mortality among Covid-19 patients in Brazil. medRxiv 2020.06.29.20142505; DOI: https://doi.org/10.1101/2020.06.29.20142505.

#### Contributors

ALR, LM, and AG conceived the study, collected data and drafted the manuscript. LO conducted data analysis. SA and CB critically revised the manuscript. All of the authors contributed to writing the manuscript and approved the final version.

## **Declaration of interests**

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

### **Declarations of interest**

The authors declare they have no conflict of interest

#### Fundings

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-

profit sectors.

# Table 1. Vaccination coverage rate for influenza season 20019/2020 among patients hospitalized with

# COVID-19 in Milan, Italy.

Characteristic	Total	Vaccinated	Not vaccinated
	(n=428)	(109)	(319)
Sex, n (%)			
Females	146 (34.1)	36 (24.7)	110 (75.3)
Males	282 (65.9)	73 (25.9)	209 (74.1)
Age, n (%)		O	
< 65	254 (59.3)	32 (12.6)	222 (87.4)
≥ 65	174 (40.7)	77 (44.3)	97 (55.7)
Heath care workers	57 (13.3)	18 (31.6)	39 (68.4)
At least one underlying chronic disease, n (%)	291 (68.0)	96 (33.0)	195 (67.0)
Type of underlying comorbidities, n (%)			
Chronic lung disease	68 (15.9)	25 (36.8)	43 (63.2)
Heart disease	202 (47.2)	75 (37.1)	127 (62.9)
Diabetes mellitus	53 (12.4)	21 (39.6)	32 (60.4)
Chronic renal disease	36 (8.4)	14 (38.9)	22 (61.1)
Cancer	41 (9.6)	14 (34.1)	27 (65.9)
Immune disorders	32 (7.5)	10 (31.3)	22 (68.8)
Chronic liver disease	10 (2.3)	3 (30.0)	7 (70.0)