

# Oesophageal manometry and gas exchange in patients with COVID-19 acute respiratory distress syndrome

Silvia Coppola<sup>1</sup>, Tommaso Pozzi<sup>2</sup>, Mattia Busana<sup>3</sup>, Francesca Bichi<sup>2</sup>, Virginia Camponetti<sup>2</sup> and Davide Chiumello<sup>1,2,4,\*</sup>

<sup>1</sup>Department of Anesthesia and Intensive Care, ASST Santi Paolo e Carlo, San Paolo University Hospital, Milan, Italy, <sup>2</sup>Department of Health Sciences, University of Milan, Milan, Italy, <sup>3</sup>Medical University of Göttingen, Göttingen, Germany and <sup>4</sup>Coordinated Research Center on Respiratory Failure, University of Milan, Milan, Italy

\*Corresponding author. E-mail: [davide.chiumello@unimi.it](mailto:davide.chiumello@unimi.it)

**Keywords:** acute respiratory distress syndrome; COVID-19; hypoxaemic respiratory failure; mechanical ventilation; oesophageal manometry; positive end-expiratory pressure

Editor—Since the first description of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) pneumonia, the associated coronavirus disease 2019 (COVID-19) has rapidly spread worldwide.<sup>1</sup> Up to 35% of patients with COVID-19 have required ICU admission as a result of hypoxaemic respiratory failure.<sup>2</sup> The majority of patients meet current criteria for acute respiratory distress syndrome (ARDS). Invasive mechanical ventilation has been applied in 29–91% of cases.<sup>1</sup>

In one of the largest studies of ICU patients, it was found that the median PEEP in mechanically ventilated patients was 14 cm H<sub>2</sub>O.<sup>3</sup> The recent Surviving Sepsis Campaign guidelines for management of critically ill adults with COVID-19 suggest, as a weak recommendation with low quality of evidence, the use of a higher PEEP strategy compared with a lower PEEP strategy (>10 cm H<sub>2</sub>O vs <10 cm H<sub>2</sub>O, respectively).<sup>4</sup> Several strategies have been proposed to select optimal PEEP during ARDS, including PEEP/FiO<sub>2</sub> titration according to the ARDS Network protocol, use of electrical impedance tomography, the decremental PEEP test, the volume–pressure curve, and oesophageal manometry. Oesophageal manometry, by measuring oesophageal pressure swings during mechanical ventilation, allows measurement of the mechanical characteristics of the lung and chest wall and computation of transpulmonary pressure, the real distending force of the lung.<sup>5</sup> Lung and chest-wall mechanics in COVID-19 are still unknown. Reports have described the heterogeneity of the disease characterised by a

dissociation between relatively well-preserved lung mechanics and severe hypoxaemia<sup>6</sup> with variable degrees of lung recruitability<sup>7</sup> and perfusion.<sup>8–11</sup>

We studied 23 consecutive mechanically ventilated Covid-19–ARDS (C-ARDS) patients admitted to the ICU of ASST Santi Paolo e Carlo Hospital in Milan, Italy from February 21, 2020 to May 1, 2020. All had positive RT–PCR testing for SARS-CoV-2 on nasal or pharyngeal swab with bilateral lung infiltrates on chest radiography. The institutional review board approved the study (# 9890/2017). All measurements were performed under rigorously controlled conditions. Patients were deeply sedated and paralysed, and then ventilated by volume-control ventilation with tidal volumes of 6–8 ml kg<sup>-1</sup> of ideal body weight. An oesophageal balloon catheter (SmartCath, Bicare, Irvine, CA, USA) was inserted in the lower part of the oesophagus; its correct position was assessed by external compression during an end-expiratory hold.

A PEEP trial was performed within 48 h of ICU admission. With stable haemodynamic conditions, a recruitment manoeuvre was performed with pressure control ventilation to a plateau pressure of 45 cm H<sub>2</sub>O with PEEP of 5 cm H<sub>2</sub>O, I:E ratio of 1:1, and ventilatory frequency of 10 bpm for 2 min. Subsequently, PEEP was set to 5 cm H<sub>2</sub>O, and airway and oesophageal pressures were recorded during both end-inspiratory and end-expiratory holds. Values of respiratory system, lung and chest-wall elastance and of end-inspiratory transpulmonary pressure were derived accordingly. Arterial and central venous blood gas analyses were also collected.

**Table 1** Respiratory mechanics and gas exchange variables at different levels of PEEP. Continuous variables are expressed as median [inter-quartile range]; Mann–Whitney test was used for continuous variable comparison. Two-tailed P-values <0.05 were considered statistically significant.

Characteristic	5 cm H <sub>2</sub> O	15 cm H <sub>2</sub> O	P-value
Driving pressure (cm H <sub>2</sub> O)	10 [8–12]	11 [10–13]	0.004
Respiratory system elastance (cm H <sub>2</sub> O L <sup>-1</sup> )	18 [15–25]	20 [19–28]	0.007
Lung elastance (cm H <sub>2</sub> O L <sup>-1</sup> )	13 [11–18]	17 [14–24]	0.001
Chest-wall elastance (cm H <sub>2</sub> O L <sup>-1</sup> )	6 [4–7]	4 [3–6]	0.276
End-inspiratory airway pressure (cm H <sub>2</sub> O)	17 [14–18]	27 [25–29]	<0.001
End-inspiratory transpulmonary pressure (cm H <sub>2</sub> O)	11 [10–14]	22 [19–24]	<0.001
Pa <sub>co2</sub> (kPa)	6.4 [5.9–6.8]	6.3 [5.7–7.3]	0.357
Pa <sub>o2</sub> /FiO <sub>2</sub> (mm Hg)	84 [67–102]	119 [107–144]	<0.001

After 15 min, PEEP was set to 15 cm H<sub>2</sub>O, and these measurements were repeated.

Baseline characteristics were median age of 65 (56–69) yr, BMI 27 (25–30) kg m<sup>-2</sup>, and Pa<sub>o2</sub>/FiO<sub>2</sub> 108 (82–141) mm Hg with a clinical PEEP of 10 (10–12) cm H<sub>2</sub>O. Respiratory mechanics and gas exchange data at 5 and 15 cm H<sub>2</sub>O of PEEP are reported in Table 1. At 15 cm H<sub>2</sub>O of PEEP, the driving pressure, lung elastance, and end-inspiratory transpulmonary pressure significantly increased compared with 5 cm H<sub>2</sub>O of PEEP. Arterial oxygenation also significantly increased without change in arterial carbon dioxide.

Our results show that increasing PEEP from a low (5 cm H<sub>2</sub>O) to higher (15 cm H<sub>2</sub>O) level led to significant deterioration in lung mechanics in critically ill COVID-19 patients. In addition, at 15 cm H<sub>2</sub>O of PEEP, end-inspiratory transpulmonary pressure was dangerously close to the physiological limit of 20–25 cm H<sub>2</sub>O. These data suggest that the lung in COVID-19 is particularly prone to over-distension and to ventilator-induced lung injury. In this regard, oesophageal manometry represents an invaluable tool in the ventilatory management of C-ARDS, allowing computation of the partitioned respiratory mechanics between lung and chest wall. Although standard ventilatory management would imply use of conventional PEEP/FiO<sub>2</sub> tables with resulting high PEEP levels caused by the severity of hypoxaemia in COVID-19 patients, our data support use of a PEEP trial and with oesophageal manometry to provide an individualised ventilatory strategy.

### Declarations of interest

The authors declare that they have no conflicts of interest.

### References

1. Wiersinga WJ, Rhodes A, Cheng AC, Peacock SJ, Prescott HC. Pathophysiology, transmission, diagnosis, and treatment of coronavirus disease 2019 (COVID-19): a review. *JAMA* 2020. <https://doi.org/10.1001/jama.2020.12839>. Advance Access published on July 10
2. Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 2020; **395**: 1054–62 (published correction appears in *Lancet* 2020; **395**: 1038)
3. Grasselli G, Zangrillo A, Zanella A, et al. Baseline characteristics and outcomes of 1591 patients infected with SARS-CoV-2 admitted to ICUs of the Lombardy region, Italy. *JAMA* 2020; **323**: 1574–81
4. Alhazzani W, Møller MH, Arabi YM, et al. Surviving Sepsis Campaign: guidelines on the management of critically ill adults with coronavirus disease 2019 (COVID-19). *Intensive Care Med* 2020; **46**: 854–87
5. Mauri T, Yoshida T, Bellani G, et al. Esophageal and transpulmonary pressure in the clinical setting: meaning, usefulness and perspectives. *Intensive Care Med* 2016; **42**: 1360–73
6. Gattinoni L, Coppola S, Cressoni M, Busana M, Rossi S, Chiumello D. COVID-19 does not lead to a “typical” acute respiratory distress syndrome. *Am J Respir Crit Care Med* 2020; **201**: 1299–300
7. Pan C, Chen L, Lu C, et al. Lung recruitability in COVID-19-associated acute respiratory distress syndrome: a single-center observational study. *Am J Respir Crit Care Med* 2020; **201**: 1294–7
8. Gattinoni L, Meissner K, Marini JJ. The baby lung and the COVID-19 era. *Intensive Care Med* 2020; **46**: 1438–40
9. Gattinoni L, Chiumello D, Caironi P, et al. COVID-19 pneumonia: different respiratory treatments for different phenotypes? *Intensive Care Med* 2020; **46**: 1099–102
10. Bos LD, Paulus F, Vlaar APJ, Beenen LFM, Schultz MJ. Subphenotyping ARDS in COVID-19 patients: consequences for ventilator management. *Ann Am Thorac Soc* 2020. <https://doi.org/10.1513/AnnalsATS.202004-376RL>. Advance Access published on May 12
11. Roesthuis L, van den Berg M, van der Hoeven H. Advanced respiratory monitoring in COVID-19 patients: use less PEEP! *Crit Care* 2020; **24**: 230

doi: 10.1016/j.bja.2020.07.045

Advance Access Publication Date: 7 August 2020

© 2020 British Journal of Anaesthesia. Published by Elsevier Ltd. All rights reserved.

## A low-cost off-the-shelf pressure-controlled mechanical ventilator for a mass respiratory failure scenario

Alcendino C. Jardim-Neto\* and Carrie E. Perlman

Stevens Institute of Technology, Department of Biomedical Engineering, Hoboken, NJ, USA

\*Corresponding author. E-mail: [alcendinoneto@gmail.com](mailto:alcendinoneto@gmail.com)

**Keywords:** acute respiratory failure; COVID-19; mechanical ventilator; pandemic; ventilator design