

EurJOC

European Journal of Organic Chemistry

 **Chemistry
Europe**
European Chemical
Societies Publishing

Accepted Article

Title: Formal Aza-Diels-Alder Reactions of Spiroindolenines with Electronrich Dienes

Authors: Elisabetta Rossi, Elisa Brambilla, Sara Leoni, Giorgio Abbiati, and Valentina Pirovano

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: *Eur. J. Org. Chem.* 10.1002/ejoc.202100251

Link to VoR: <https://doi.org/10.1002/ejoc.202100251>

WILEY-VCH

FULL PAPER

Formal Aza-Diels-Alder Reactions of Spiroindolenines with Electronrich Dienes

Elisa Brambilla,^{*,[a]} Sara Leoni,^[a] Giorgio Abbiati,^[a] Valentina Pirovano^[a] and Elisabetta Rossi^{*,[a]}

Dedicated to Professor Franco Cozzi on the occasion of his 70th birthday.

[a] Dr. E. Brambilla, Dr. S. Leoni, Prof. G. Abbiati, Dr. V. Pirovano, Prof. E. Rossi
Dipartimento di Scienze Farmaceutiche - Sezione di Chimica Generale e Organica "A. Marchesini"
Università degli Studi di Milano
Via Venezian, 21 - 20133 Milano -Italy
E-mail: elisabetta.rossi@unimi.it
<https://www.unimi.it/it/ugov/person/elisabetta-rossi>
E-mail: elisa.brambilla@unimi.it
<https://www.unimi.it/it/ugov/person/elisa-brambilla>

Supporting information for this article is given via a link at the end of the document.

Abstract: Spiroindolenines were employed as cyclic imine substrates in formal aza-Diels-Alder reactions with Danishefsky's diene or silyloxy-substituted electron-rich dienes for the synthesis of the corresponding tetrahydropyrido[1,2-a]spiroindolinones. The reactions occur under mild conditions in the presence of ytterbium triflate as Lewis acidic catalyst delivering the desired compounds in good yield. The reaction results in the preparation of a small library of a new class of conformational constrained heterocyclic derivatives that easily undergo selective and efficient manipulations.

Introduction

The constant search for new lead compounds in medicinal chemistry recently resulted in the identification of new privileged scaffolds that encompass the presence of innovative and original nitrogen heterocycles with rigid three-dimensional skeletons. [1] Spirocyclic heterocycles fall in this frame and among these, spirocyclic indolines/indolenines deserve particular attention because they are the core motif of many bioactive natural compounds and occur in some synthetic derivatives selected for the development of new drugs, figure 1. [2]




Figure 1. Biologically Active Natural and Synthetic Spiro(poly)cyclic Indolines/Indolenines.

Looking at the structures of the synthetic compounds we noticed that a good strategy to increase their complexity and in particular the rigid 3D structure could be the construction of a new ring at N1-C2 bond. In this way, spirotricyclic derivatives could be efficiently obtained, figure 2. Retrosynthetic and literature analysis permits to select spirocyclic indolenines as useful starting compounds for the target compounds. [3]

FULL PAPER





Figure 2. Planned Construction of New Spirotricyclic Indolines.


In particular, the imine type nature of N1-C2 bond suggests the possibility to involve this bond in cycloaddition/cyclization reactions with suitable partners. [4] Literature reports involving indolenines as reactants in these reactions comprise the one-step synthesis of pyrroline and piperidine derivatives as reported in scheme 1. [5]



Scheme 1. Reported Cycloaddition/Cyclization reactions at N1-C2 Bond of Spiroindolines.

In particular, spiroindolenines were involved in [3+2] cyclization reactions with donor-acceptor cyclopropanes, in the Castagnoli-Cushman reaction with homophthalic anhydride and in organocatalyzed formal aza-Diels-Alder reactions with α,β -unsaturated ketones.

Taking into account these premises and our previous works on the cycloaddition/cyclization reactions of indole derivatives [6], we decided to test the reactivity of spiroindolenines **1** with 1,4-dienes **2** in aza-Diels-Alder reactions for the synthesis of tetrahydropyrido[1,2-*a*]spiroindolinones **3**, scheme 2.



Scheme 2. Planned [4+2] Cycloaddition Reactions.

Results and Discussion

At the outset, spiroindolenine **1a** was chosen as model compound for the reactions with Danishefsky's diene **2a** in the presence of different Lewis acid (LA) catalysts, table 1. An electronrich dienes was selected based on previously reported results on the cycloaddition reactions of both acyclic and cyclic imines. [4] Moreover, our former outcomes on LAs catalyzed [4+2] cycloadditions directed the selection of appropriate catalysts and reaction parameters. [6]

Table 1. Optimization of the reaction conditions.^[a]


Entry	Catalysts	Solvent	T, °C	Time, h	Yield, % ^[b]
1	Sc(OTf) ₃	DCM	rt	2	51 ^[c]
2	Zn(OTf) ₂	DCM	rt	2	46 ^[c]
3	Ni(ClO ₄) ₂ ·6H ₂ O	DCM	rt	2	38 ^[c]
4	Mg(ClO ₄) ₂	DCM	rt	2	45 ^[c]
5	BF ₃ ·Et ₂ O	DCM	-20	2	- ^[d]
6	Yb(OTf) ₃	DCM	rt	1	83
7	Yb(OTf) ₃	Toluene	rt	1	80
8	Yb(OTf) ₃	MeCN	rt	1	61 ^[e]
9	Yb(OTf) ₃	THF	rt	1	63 ^[e]
10	Yb(OTf) ₃ ^[f]	DCM	rt	2	49 ^[e]
11	Yb(OTf) ₃ ^[g]	DCM	rt	1	29 ^[e]
12	none	DCM	rt	8	- ^[d]

[a] All reactions were carried out using **1a** (0.2 mmol) and **2a** (0.4 mmol) in the stated solvent (0.125 M) and catalyst (20 mol%) and quenched by treatment with 1N HCl solution. [b] Isolated yield. [c] Beside starting materials (traces) and unidentified compounds. [d] Complex reaction mixture. [e] Beside unreacted starting materials. [f] Catalyst 10 mol%. [g] **2a**, 0.2 mmol.

At the beginning, several classical LAs including Sc(OTf)₃, Zn(OTf)₂, Ni(ClO₄)₂·6H₂O and Mg(ClO₄)₂ were examined in dichloromethane (DCM) at room temperature (table 1, entries 1-4). Under these conditions, the expected cycloadduct **3a** was isolated in comparable moderate yields (38-51%) beside small quantities of unreacted starting materials and several degradation compounds. Switching to a more electrophilic LA such as BF₃·Et₂O, we observed complete degradation of the starting materials and the formation of complex reaction mixture (table 1, entry 5). Recalling that both imines and Danishefsky's diene are acid sensitive substrates, we moved our attention to Yb(OTf)₃, an efficient and mild LA catalyst for [4+2] cycloadditions involving acid labile substrates. [7] As reported in entry 6, in the presence of Yb(OTf)₃, **3a** was obtained in 83% yield. Comparable yields

FULL PAPER


were obtained moving to toluene as solvent, whereas the use of MeCN or THF resulted in poorer yields, entries 7-9. Using a small amount of catalyst (10% instead of 20%) or lowering the reactants ratio, **1a:2a**, from 1:2 to 1:1 gave worse results, entries 10-11. Finally, the reaction did not occur in the absence of the catalyst and complete degradation of the starting materials was observed after 8 hours, entry 12. Therefore, the best reaction conditions are those reported in entries 6 and 7 giving rise to **3a** in 83 and 80% yield, respectively. The two methods differ for the solvent employed and during the evaluation of the reaction scope, we choose to use toluene as solvent because it dissolves the starting materials better than DCM. The structure of **3a** was confirmed by 1D and 2D NMR analysis. Thus, we started our evaluation by reacting different indolenines **1b-k** with diene **2a**, scheme 3.



Scheme 3. Scope of the Reaction between **1a-k** and **2a**.

Starting from parent model compound **3a**, substitution at position 5' of the indolenines core with a soft electron-donating methyl group or with an electron-withdrawing fluorine atom did not affect the reaction course and the corresponding compounds **3b-c** were isolated in 81 and 76 % yield, respectively. Variations at the spirocyclohexyl moiety by introduction of an oxygen or of a benzyloxycarbonylamino group (NCbz) in position 4 were tolerated as well. In particular, a moderate decrease in the reaction yields was observed using NCbz derivatives **1f-h**. In this case, the average yield was 65% compared to 80% observed in previous cases. Finally, several limitations were encountered using spiroindolenines **1i-k**. The spiroindolenine **1i** bearing a cyclopentyl moiety was unstable and, under usual reaction conditions, gave only a small 15% amount of the carbazole **4** arising from a ring expansion reaction. Moreover, the indolenine **1j** bearing linear ethyl substituents at C3 decomposed under our standard reaction conditions whereas the C2 methyl substituted spiroindolenine **1k** was recovered unreacted. We then evaluated

the influence of dienes other than Danishefsky's diene **2a** on the reaction course. In particular, we selected two electron-rich dienes, namely 2-trimethylsiloxy-1,3-butadiene **2b** and 2-trimethylsiloxy-1,3-cyclohexadiene **2c** and the obtained results are reported in scheme 4.



Scheme 4. Scope of the Reaction between **1** and **2b-d**.

The employment of the acyclic diene **2b** was well tolerated and the corresponding indolines **3i-m** were isolated in good yields, comparable to those obtained with Danishefsky's diene **2a**. However, the reactions performed with the corresponding cyclic diene **2c** gave rise to spiroindolines **3n-r** in slightly lower yields. Moreover, compounds **3n-r** were isolated as a couple of diastereoisomers in almost equimolecular amounts. The isomers could be easily separated by flash column chromatography and the structures of both diastereoisomers were assigned based on 2D NMR analysis conducted on **3n**, figure 3.





Figure 3. 2D NMR Analysis of compound **3n**.

Finally, spiroindolenine **1a** was tested with 2-benzyloxy-1,3-butadiene **2d** bearing an alkoxy instead of a siloxy substituent at

FULL PAPER

C2. The reaction resulted in the isolation of the unreacted starting materials after 3h.


Synthesized spiroindolines **3a-r** were finally investigated as useful substrates for selective transformations. Thus, compound **3j** was synthesized on a gram scale and subjected to selected transformations, scheme 5.



^a Reaction performed on a 1.0 mmol scale

Scheme 5. Selective Functional Group Transformations on **3j**.


Initially, chemical reduction of the carbonyl group was attempted with sodium borohydride in ethanol at room temperature affording the corresponding alcohol **5** in 84% yield. Then, the formation of new carbon-carbon bonds was realized following two different strategies. As first reaction, we realized the Wittig olefination reaction by reacting **3j** with triphenylmethyl phosphonium bromide under basic conditions in dry diethyl ether at 0°C. The corresponding terminal alkene **6** was isolated in 76% yield. Finally, the formation of a new single carbon-carbon bond was achieved by Grignard addition of ethyl magnesium bromide on the ketone moiety of **3j** in dry tetrahydrofuran at 0°C. The corresponding compound **7** was isolated in 65% yield. Compounds **5** and **7** were isolated as single diastereoisomers, however the reciprocal positions of the substituents could be assigned only for compound **5** on the basis of 1D and 2D NMR analysis. In compound **7**, the overlapping of ethyl group signals with the cyclohexyl and piperidinyll moieties prevents to observe the diagnostic nOe interactions with the angular hydrogen. Finally, to complete our investigations, we realized the conjugate addition of the same Grignard reagent at the α,β -unsaturated carbonyl function of **3a**, scheme 6.



Scheme 6. Grignard Addition on **3a**.

The reaction was realized as reported for the synthesis of **7** resulting in the formation of 1,4-conjugate addition compound **8** in 67% yield as single diastereoisomer (see SI for 2D NMR analysis).

Selective 1,4-addition of Grignard reagent to **3a** could be related to the assistance of the nitrogen atom in the stabilization of the enolate form of the starting compound. Several related reactions involving push-pull ethylene derivatives are reported in literature.[9] Several useful considerations on the reaction mechanism can be done according to literature reports, scheme 7. As stated by several authors,[4] the reaction occurs upon coordination of the LA to imine nitrogen followed by polarity-driven addition of the diene in the final cyclization step. The addition/cyclization could happen via Mannich-type reaction involving enolate addition to LA activated imine function, followed by a cyclization step that occurs during the hydrolytic stage. Beside, an alternative mechanism involving aza-Diels-Alder cycloaddition reaction directly delivers the cyclic compound and, after hydrolysis, the final indolines **3**. [8]



Scheme 7. Mechanistic Hypothesis.

During our study, we evaluated the possibility to detect some of the proposed intermediates in the crude reaction mixture prior to the hydrolytic workup. However, although some spots other than that of the final product were evident in the thin layer chromatographic analysis of the crude reaction mixture, any attempts to isolate or identify them through NMR analysis were unsuccessful. Despite that, considering the high polar character of both reactants, a reaction mechanism involving a Mannich addition and a stepwise non-concerted cycloaddition mechanism is more plausible. Further investigations on this topic could be useful in the perspective to develop an asymmetric version for these transformations.

Conclusion

In summary, we designed an alternative approach for the efficient synthesis of tetrahydropyrido[1,2-*a*]spiroindolinones. The main advantages of the proposed methodology are the use of simple, cheap and easy affordable starting materials and catalyst, the mild reaction conditions employed and the wide range of tolerated substituents. Moreover, tetrahydropyrido[1,2-*a*]spiroindolinones possess useful features such as the structural rigidity and the relationship with known bioactive compounds that makes them interesting for drug discovery processes. With respect to already known syntheses, the only reported method involving [4+2] cycloadditions results in the isolation of enantiomerically enriched but structurally less powerful derivatives, see scheme 1 and ref.

FULL PAPER

5c. The reaction works well with silyoxy-substituted dienes whereas simple less electronrich alkyloxy diene **2d** resulted unreactive. Despite that, as reported in scheme 5 and 6, tetrahydropyrido[1,2-*a*]spiroindolinones **3** can deliver more complex or differently decorated derivatives by means of simple high yielding procedures.

Experimental Section

For experimental details, copies of ^1H and ^{13}C NMR spectra and other supplementary material see the SI section. [10]

General procedure for the synthesis of tetrahydropyridine[1,2-*a*]-indoles **3a-r**

To a nitrogen fluxed solution of indolenines **1a-k** (0.2 mmol) and $\text{Yb}(\text{OTf})_3$ (20 mol%) in anhydrous toluene (1.6 ml), the diene **2a-d** (0.4 mmol) was added. The reaction was stirred at rt for 1h and monitored by TLC. Then 1 ml of HCl 1 N was added and the reaction was stirred at rt for 30 min. The organic layer was extracted with EtOAc (3 x 5 ml). The combined organic extracts were dried over Na_2SO_4 and the solvent was removed under vacuum. The crude was purified by flash chromatography to yield the corresponding indoline derivatives **3a-r**.

9',9a'-dihydro-8'-H-spiro[cyclohexane-1,10'-pyrido[1,2-*a*]indol]-8'-one (**3a**)

General procedure was followed using spiro[cyclohexane-1,3'-indole] **1a** (37 mg, 0.2 mmol), $\text{Yb}(\text{OTf})_3$ (25 mg, 0.04 mmol) in toluene (1.6 ml) and Dainshfsky's diene **2a** (78 μl , 0.4 mmol). Purification by flash chromatography (SiO_2 , hexane/EtOAc 2:1 to 1:1) yielded **3a** (40.5 mg, 80%) as orange solid (m.p. 105.8-108.5 $^\circ\text{C}$). $^1\text{H-NMR}$ (300 MHz, CDCl_3): 7.70 (d, $J = 7.4$ Hz, 1H), 7.45 (d, $J = 7.5$ Hz, 1H), 7.24 (t, $J = 7.9$ Hz, 1H), 7.00 (t, $J = 7.5$ Hz, 1H), 6.92 (d, $J = 8.0$ Hz, 1H), 5.29 (d, $J = 5.7$ Hz, 1H), 3.95 (dd, $J = 16.5$, 4.8 Hz, 1H), 2.86 – 2.62 (m, 2H), 2.01 – 1.86 (m, 2H), 1.79 – 1.43 (m, 8H). $^{13}\text{C-NMR}$ (75 MHz, CDCl_3): 193.0 (C), 141.4 (CH), 140.9 (C), 139.7 (C), 128.0 (CH), 125.5 (CH), 122.6 (CH), 107.8 (CH), 100.6 (CH), 69.1 (CH), 46.0 (C), 38.9 (CH₂), 36.2 (CH₂), 32.8 (CH₂), 25.3 (CH₂), 22.9 (CH₂), 21.9 (CH₂). **ESI(+)-MS**: m/z (%) = 254 (100) [$\text{M}+1$]⁺. Elemental analysis calcd for $\text{C}_{17}\text{H}_{19}\text{NO}$: C 80.60, H 7.56, N 5.53, found: C 80.48, H 7.54, N 5.55

2'-methyl-9',9a'-dihydro-8'-H-spiro[cyclohexane-1,10'-pyrido[1,2-*a*]indol]-8'-one (**3b**)

General procedure was followed using: 5'-methylspiro[cyclohexane-1,3'-indole] **1b** (40 mg, 0.2 mmol), $\text{Yb}(\text{OTf})_3$ (25 mg, 0.04 mmol) in toluene (1.6 ml) and Dainshfsky's diene **2a** (78 μl , 0.4 mmol). Purification by flash chromatography (SiO_2 , cyclohexane/EtOAc 2:1) yielded **3b** (43 mg, 81%) as yellow solid (m.p. dec $T > 120$ $^\circ\text{C}$) $^1\text{H-NMR}$ (300 MHz, CD_2Cl_2): 7.67 (dd, $J = 7.4$, 0.9 Hz, 1H), 7.26 (s, 1H), 7.04 (ddd, $J = 8.1$, 1.7, 0.7 Hz, 1H), 6.83 (d, $J = 8.1$ Hz, 1H), 5.16 (d, $J = 7.3$ Hz, 1H), 3.92 (dd, $J = 16.7$, 4.0 Hz, 1H), 2.75 (m, 1H), 2.61 (ddd, $J = 15.6$, 4.7, 1.0 Hz, 1H), 2.34 (s, 3H), 2.02 – 1.86 (m, 2H), 1.78 – 1.40 (m, 8H). $^{13}\text{C-NMR}$ (75 MHz, CD_2Cl_2): 192.0 (C), 141.1 (CH), 139.9 (C), 138.7 (C), 132.0 (C), 128.2 (CH), 126.0 (CH), 107.3 (CH), 99.6 (CH), 69.4 (CH), 45.9 (C), 39.1 (CH₂), 36.2 (CH₂), 32.7 (CH₂), 25.3 (CH₂), 22.9 (CH₂), 22.0 (CH₂), 20.8 (CH₃). **ESI(+)-MS**: m/z (%) = 268 (100) [$\text{M}+1$]⁺. Elemental analysis calcd for $\text{C}_{18}\text{H}_{21}\text{NO}$: C 80.86, H 7.92, N 5.24, found: C 81.03, H 7.94, N 5.26.

2'-fluoro-9',9a'-dihydro-8'-H-spiro[cyclohexane-1,10'-pyrido[1,2-*a*]indol]-8'-one (**3c**)

General procedure was followed using: 5'-fluorospiro[cyclohexane-1,3'-indole] **1c** (41 mg, 0.2 mmol), $\text{Yb}(\text{OTf})_3$ (25 mg, 0.04 mmol) in toluene (1.6

ml) and Dainshfsky's diene **2a** (78 μl , 0.4 mmol). Purification by flash chromatography (SiO_2 , cyclohexane/EtOAc 2:1) yielded **3c** (41 mg, 76%) as yellow solid (m.p. 160.3-162.4). $^1\text{H-NMR}$ (300 MHz, CD_2Cl_2) 7.67 (dd, $J = 7.5$, 0.7 Hz, 1H), 7.21 (dd, $J = 8.9$, 2.5 Hz, 1H), 6.97 (td, $J = 8.7$, 2.5 Hz, 1H), 6.88 (dd, $J = 8.7$, 4.4 Hz, 1H), 5.21 (d, $J = 7.4$ Hz, 1H), 3.98 (dd, $J = 16.7$, 4.4 Hz, 1H), 2.78 (m, 1H), 2.62 (ddd, $J = 15.6$, 4.6, 0.8 Hz, 1H), 1.99 – 1.84 (m, 2H), 1.79 – 1.45 (m, 8H). $^{13}\text{C-NMR}$ (75 MHz, CD_2Cl_2) 191.9 (C), 158.6 (d, $J = 239.7$ Hz, C), 141.5 (d, $J = 7.4$ Hz, C), 141.2 (CH), 137.5 (C), 114.2 (d, $J = 24.2$ Hz, CH), 113.0 (d, $J = 24.6$ Hz, CH), 108.1 (d, $J = 8.7$ Hz CH), 100.2 (CH), 69.6 (CH), 46.0 (C), 38.9 (CH₂), 35.8 (CH₂), 32.5 (CH₂), 25.2 (CH₂), 22.8 (CH₂), 21.8 (CH₂). **ESI(+)-MS**: m/z (%) = 272 (100) [$\text{M}+1$]⁺. Elemental analysis calcd for $\text{C}_{17}\text{H}_{18}\text{FNO}$: C 75.25, H 6.69, N 5.16, found: C 75.46, H 6.67, N 5.15.

2,3,5,6,9',9a'-hexahydro-8'-H-spiro[pyran-4,10'-pyrido[1,2-*a*]indol]-8'-one (**3d**)

General procedure was followed using: 2',3',5',6'-tetrahydrospiro[indole-3,4'-pyran] **1d** (37 mg, 0.2 mmol), $\text{Yb}(\text{OTf})_3$ (25 mg, 0.04 mmol) in toluene (1.6 ml) and Dainshfsky's diene **2a** (78 μl , 0.4 mmol). Purification by flash chromatography (SiO_2 , cyclohexane/EtOAc 2:1) yielded **3d** (41 mg, 80%) as yellow solid (m.p. 134.7-136.8 $^\circ\text{C}$). $^1\text{H-NMR}$ (300 MHz, CD_2Cl_2) 7.73 (dd, $J = 7.5$, 0.9 Hz, 1H), 7.56 (d, $J = 7.6$ Hz, 1H), 7.27 (td, $J = 7.7$, 1.1 Hz, 1H), 7.03 (td, $J = 7.6$, 1.0 Hz, 1H), 6.97 (d, $J = 8.0$ Hz, 1H), 5.23 (d, $J = 7.4$ Hz, 1H), 4.10 (m, 1H), 3.98 (dd, $J = 15.1$, 6.1 Hz, 1H), 3.88 – 3.71 (m, 3H), 2.83 – 2.63 (m, 2H), 2.06 (m, 1H), 1.98 – 1.81 (m, 2H), 1.57 (m, 1H). $^{13}\text{C-NMR}$ (75 MHz, CD_2Cl_2) 191.6 (C), 141.2 (C), 141.1 (CH), 138.2 (C), 128.3 (CH), 125.2 (CH), 122.4 (CH), 107.9 (CH), 100.8 (CH), 68.8 (CH), 64.5 (CH₂), 63.7 (CH₂), 43.6 (C), 38.7 (CH₂), 35.6 (CH₂), 32.6 (CH₂). **ESI(+)-MS**: m/z (%) = 256 (100) [$\text{M}+1$]⁺. Elemental analysis calcd for $\text{C}_{16}\text{H}_{17}\text{NO}_2$: C 75.27, H 6.71, N 5.49, found: C 75.11, H 6.70, N 5.51.

2'-methyl-2,3,5,6,9',9a'-hexahydro-8'-H-spiro[pyran-4,10'-pyrido[1,2-*a*]indol]-8'-one (**3e**)

General procedure was followed using: 5-methyl-2',3',5',6'-tetrahydrospiro[indole-3,4'-pyran] **1e** (40 mg, 0.2 mmol), $\text{Yb}(\text{OTf})_3$ (25 mg, 0.04 mmol) in toluene (1.6 ml) and Dainshfsky's diene **2a** (78 μl , 0.4 mmol). Purification by flash chromatography (SiO_2 , cyclohexane/EtOAc 1:1 to 3:1) yielded **3e** (45 mg, 84%) as yellow solid (m.p. $T > 130$ Dec.). $^1\text{H-NMR}$ (300 MHz, CD_2Cl_2) 7.72 (d, $J = 7.5$ Hz, 1H), 7.39 (s, 1H), 7.11 (d, $J = 8.1$ Hz, 1H), 6.88 (d, $J = 8.1$ Hz, 1H), 5.21 (d, $J = 7.5$ Hz, 1H), 4.14 (m, 1H), 3.99 (dd, $J = 15.3$, 5.9 Hz, 1H), 3.92 – 3.73 (m, 3H), 2.84 – 2.64 (m, 2H), 2.37 (s, 3H), 2.08 (m, 1H), 1.98 – 1.82 (m, 2H), 1.59 (ddd, $J = 13.7$, 6.8, 3.9 Hz, 1H). $^{13}\text{C-NMR}$ (75 MHz, CD_2Cl_2) 191.7 (C), 141.3 (CH), 138.9 (C), 138.4 (C), 132.3 (C), 128.7 (CH), 126.0 (CH), 107.7 (CH), 100.0 (CH), 68.9 (CH), 64.5 (CH₂), 63.7 (CH₂), 43.6 (C), 38.7 (CH₂), 35.5 (CH₂), 32.6 (CH₂), 20.9 (CH₃). **ESI(+)-MS**: m/z (%) = 270 (100) [$\text{M}+1$]⁺. Elemental analysis calcd for $\text{C}_{17}\text{H}_{19}\text{NO}_2$: C 75.81, H 7.11, N 5.20, found: C 75.39, H 7.13, N 5.21.

Benzyl 8'-oxo-9',9a'-dihydro-8'-H-spiro[piperidine-4,10'-pyrido[1,2-*a*]indole]-1-carboxylate (**3f**)

General procedure was followed using: benzyl spiro[indole-3,4'-piperidine]-1-carboxylate **1f** (78 mg, 0.2 mmol), $\text{Yb}(\text{OTf})_3$ (25 mg, 0.04 mmol) in toluene (1.6 ml) and Dainshfsky's diene **2a** (78 μl , 0.4 mmol). Purification by flash chromatography (SiO_2 , cyclohexane/EtOAc 1:1 to 1:2) yielded **3f** (51 mg, 66%) as yellow solid (m.p. 69.3-72.8 $^\circ\text{C}$). $^1\text{H-NMR}$ (300 MHz, CD_2Cl_2) 7.73 (dd, $J = 7.5$, 0.9 Hz, 1H), 7.46 (d, $J = 7.6$ Hz, 1H), 7.42 – 7.33 (m, 5H), 7.28 (td, $J = 7.7$, 1.2 Hz, 1H), 7.02 (dd, $J = 7.6$, 1.0 Hz, 1H), 6.98 (m, 1H), 5.23 (d, $J = 7.5$ Hz, 1H), 5.16 (s, 2H), 3.99 (dd, $J = 16.3$, 4.5 Hz, 1H), 3.93 – 3.66 (m, 3H), 3.55 (m, 1H), 2.73 (m, 1H), 2.59 (m, 1H), 2.02 (m, 1H, overlapped with EtOAc), 1.91 – 1.79 (m, 2H), 1.60 (m, 1H). $^{13}\text{C-NMR}$ (75 MHz, CD_2Cl_2) 191.5 (C), 155.2 (C), 141.3 (C), 141.0 (CH), 137.5 (C), 137.1 (C), 128.42 (CH), 128.39 (2xCH), 127.9 (CH), 127.8 (2xCH), 125.3 (CH), 122.4 (CH), 108.0 (CH), 100.8 (CH), 68.5 (CH), 67.0 (CH₂), 44.3 (C), 40.9 (CH₂), 40.0 (CH₂), 38.4 (CH₂), 34.7 (CH₂), 31.8 (CH₂).

FULL PAPER

ESI(+)-MS: m/z (%) = 411 (100) [M+Na]⁺. Elemental analysis calcd for C₂₄H₂₄N₂O₃: C 74.21, H 6.23, N 7.21, found: C 74.06, H 6.21, N 7.23.

Benzyl 2'-methyl-8'-oxo-9',9a'-dihydro-8'H-spiro[piperidine-4,10'-pyrido[1,2-a]indole]-1-carboxylate (3g)

General procedure was followed using: 5-methylspiro[indole-3,4'-piperidine]-1'-carboxylate **1g** (67 mg, 0.2 mmol), Yb(OTf)₃ (25 mg, 0.04 mmol) in toluene (1.6 ml) and Dainshfsky's diene **2a** (78 μl, 0.4 mmol). Purification by flash chromatography (SiO₂, cyclohexane/EtOAc 1:1) yielded **3g** (50 mg, 62%) as yellow solid (m.p. 68.9-71.3° C). **¹H-NMR** (300 MHz, CDCl₃) 7.65 (d, J = 7.3 Hz, 1H), 7.40 – 7.31 (m, 5H), 7.20 (s, 1H), 7.07 (d, J = 8.1 Hz, 1H), 6.83 (d, J = 8.1 Hz, 1H), 5.25 (d, J = 7.4 Hz, 1H), 5.17 (s, 2H), 3.99 – 3.81 (m, 2H), 3.78 – 3.63 (m, 2H), 3.55 (m, 1H), 2.79 – 2.53 (m, 2H), 2.33 (s, 3H), 2.01 (m, 1H, overlapped with EtOAc), 1.86-1.74 (m, 2H), 1.58 (m, 1H). **¹³C-NMR** (75 MHz, CDCl₃) 191.8 (C), 155.4 (C), 141.2 (CH), 138.9 (C), 137.5 (C), 136.6 (C), 132.5 (C), 129.1 (CH), 128.5 (2xCH), 128.1 (CH), 128.0 (2xCH), 125.9 (CH), 107.8 (CH), 100.5 (CH), 68.6 (CH), 67.3 (CH₂), 44.3 (C), 41.0 (CH₂), 40.1 (CH₂), 38.4 (CH₂), 34.7 (CH₂), 31.8 (CH₂), 21.2 (CH₃). **ESI(+)-MS:** m/z (%) = 425 (100) [M+Na]⁺. Elemental analysis calcd for C₂₅H₂₆N₂O₃: C 74.60, H 6.51, N 6.69, found: C 74.77, H 6.49, N 6.70.

Benzyl 2'-fluoro-8'-oxo-9',9a'-dihydro-8'H-spiro[piperidine-4,10'-pyrido[1,2-a]indole]-1-carboxylate (3h)

General procedure was followed using benzyl 5-fluorospiro[indole-3,4'-piperidine]-1'-carboxylate **1h** (68 mg, 0.2 mmol), Yb(OTf)₃ (25 mg, 0.04 mmol) in toluene (1.6 ml) and Dainshfsky's diene **2a** (78 μl, 0.4 mmol). Purification by flash chromatography (SiO₂, heptane/EtOAc 1:1) yielded **3h** (55 mg, 68%) as yellow solid (m.p. 57.6-59.9° C). **¹H-NMR** (300 MHz, CDCl₃) 7.64 (d, J = 7.5 Hz, 1H), 7.43 – 7.32 (m, 5H), 7.14 (dd, J = 8.5, 2.2 Hz, 1H), 6.99 (td, J = 8.6, 2.3 Hz, 1H), 6.87 (dd, J = 8.7, 4.2 Hz, 1H), 5.29 (d, J = 7.5 Hz, 1H), 5.18 (s, 2H), 4.00 (dd, J = 16.3, 4.5 Hz, 1H), 3.90 – 3.67 (m, 3H), 3.55 (m, 1H), 2.81 – 2.57 (m, 2H), 1.98 (m, 1H, overlapped with EtOAc), 1.92 – 1.77 (m, 2H), 1.60 (m, 1H). **¹³C-NMR** (75 MHz, CDCl₃) 191.59 (C), 158.68 (d, J = 242.1 Hz, C), 155.36 (C), 141.24 (CH), 138.99 (d, J = 7.3 Hz, C), 137.36 (C), 136.53 (C), 128.58 (2xCH), 128.20 (CH), 128.05 (2xCH), 115.22 (d, J = 24.0 Hz, CH), 112.98 (d, J = 24.7 Hz, CH), 108.62 (d, J = 8.6 Hz, CH), 101.11 (CH), 68.85 (CH), 67.42 (CH₂), 44.38 (C), 40.77 (CH₂), 39.92 (CH₂), 38.33 (CH₂), 34.46 (CH₂), 31.66 (CH₂). **ESI(+)-MS:** m/z (%) = 429 (100) [M+Na]⁺. Elemental analysis calcd for C₂₄H₂₃FN₂O₃: C 70.92, H 5.70, N 6.89, found: C 70.72, H 5.72, N 6.88.

6',7',9',9a'-tetrahydro-8'H-spiro[cyclohexane-1,10'-pyrido[1,2-a]indol]-8'-one (3i)

General procedure was followed using: spiro[cyclohexane-1,3'-indole] **1a** (37 mg, 0.2 mmol), Yb(OTf)₃ (25 mg, 0.04 mmol) in toluene (1.6 ml) and (buta-1,3-dien-2-yloxy)trimethylsilane **2b** (69 μl, 0.4 mmol). Purification by flash chromatography (SiO₂, cyclohexane/EtOAc 95:5 to 9:1) yielded **3i** (38 mg, 75%) as white solid (m.p. 114.5-116.9° C). **¹H-NMR** (300 MHz, CDCl₃) 7.21-7.14 (m, 2H), 6.81 (t, J = 7.1 Hz, 1H), 6.62 (d, J = 7.7 Hz, 1H), 4.05 (ddd, J = 13.5, 6.7, 1.3 Hz, 1H), 3.69 (m, 1H), 3.27 (m, 1H), 2.54 (m, 1H), 2.39 – 2.31 (m, 3H), 1.89 – 1.61 (m, 6H), 1.58 – 1.27 (m, 4H). **¹³C-NMR** (75 MHz, CDCl₃) 209.8 (C), 147.9 (C), 137.8 (C), 127.8 (CH), 123.7 (CH), 118.8 (CH), 107.0 (CH), 70.4 (CH), 47.7 (C), 44.1 (CH₂), 41.9 (CH₂), 39.9 (CH₂), 35.8 (CH₂), 29.8 (CH₂), 25.9 (CH₂), 23.3 (CH₂), 23.1 (CH₂). **ESI(+)-MS:** m/z (%) = 411 (60) [M+1]⁺. Elemental analysis calcd for C₁₇H₂₁NO: C 79.96, H 8.29, N 5.49, found: C 80.16, H 8.32, N 5.48.

2'-fluoro-6',7',9',9a'-tetrahydro-8'H-spiro[cyclohexane-1,10'-pyrido[1,2-a]indol]-8'-one (3j)

General procedure was followed using: 5'-fluorospiro[cyclohexane-1,3'-indole] **1c** (41 mg, 0.2 mmol), Yb(OTf)₃ (25 mg, 0.04 mmol) in toluene (1.6 ml) and (buta-1,3-dien-2-yloxy)trimethylsilane **2b** (69 μl, 0.4 mmol).

Purification by flash chromatography (SiO₂, cyclohexane/EtOAc 9:1 to 8:2) yielded **3j** (44 mg, 80%) as white solid (m.p. 119.8-123.9° C). **¹H-NMR** (300 MHz, CDCl₃) 6.91 (dd, J = 8.6, 2.6 Hz, 1H), 6.83 (m, 1H), 6.48 (dd, J = 8.4, 4.2 Hz, 1H), 3.94 (ddd, J = 13.5, 6.6, 1.9 Hz, 1H), 3.64 (dd, J = 9.5, 5.2 Hz, 1H), 3.22 (m, 1H), 2.53 (m, 1H), 2.42 – 2.29 (m, 3H), 1.87 – 1.23 (m, 10H overlapped with cyclohexane). **¹³C-NMR** (75 MHz, CDCl₃) 209.0 (C), 157.1 (d, J = 235.8 Hz, C), 144.1 (C), 139.6 (d, J = 7.2 Hz, C), 113.4 (d, J = 23.3 Hz, CH), 111.6 (d, J = 24.3 Hz, CH), 107.1 (d, J = 8.2 Hz, CH), 71.1 (CH), 47.6 (C), 44.5 (CH₂), 41.7 (CH₂), 39.6 (CH₂), 35.5 (CH₂), 29.7 (CH₂), 25.7 (CH₂), 22.99 (CH₂), 22.95 (CH₂). **ESI(+)-MS:** m/z (%) = 274 (100) [M+1]⁺. Elemental analysis calcd for C₁₇H₂₀FNO: C 74.70, H 7.38, N 5.12, found: C 74.89, H 7.36, N 5.11.

2,3,5,6,6',7',9',9a'-octahydro-8'H-spiro[pyran-4,10'-pyrido[1,2-a]indol]-8'-one (3k)

General procedure was followed using: 2',3',5',6'-tetrahydrospiro[indole-3,4'-pyran] **1d** (38 mg, 0.2 mmol), Yb(OTf)₃ (25 mg, 0.04 mmol) in toluene (1.6 ml) and (buta-1,3-dien-2-yloxy)trimethylsilane **2b** (69 μl, 0.4 mmol). Purification by flash chromatography (SiO₂, cyclohexane/EtOAc 2:1) yielded **3k** (37 mg, 72%) as white solid (m.p. 160.7-162.5° C). **¹H-NMR** (300 MHz, CDCl₃) 7.27 (d, J = 7.1 Hz, 1H), 7.21 (t, J = 7.7 Hz, 1H), 6.84 (d, J = 7.5 Hz, 1H), 6.65 (d, J = 7.8 Hz, 1H), 4.07 – 2.92 (m, 3H), 3.81 – 3.69 (m, 2H), 3.58 (m, 1H), 3.27 (td, J = 13.4, 3.2 Hz, 1H), 2.56 (m, 1H), 2.46 – 2.28 (m, 3H), 2.06 (m, 1H), 1.87 (m, 1H), 1.78 – 1.62 (m, 2H). **¹³C-NMR** (75 MHz, CDCl₃) 208.9 (C), 148.0 (C), 136.0 (C), 128.3 (CH), 123.9 (CH), 119.1 (CH), 107.3 (CH), 70.3 (CH), 65.0 (CH₂), 45.3 (C), 44.1 (CH₂), 41.9 (CH₂), 39.8 (CH₂), 35.5 (CH₂), 29.8 (CH₂). One CH₂ is missing, probably overlapped. **ESI(+)-MS:** m/z (%) = 258 (100) [M+1]⁺. Elemental analysis calcd for C₁₆H₁₉NO₂: C 74.68, H 7.44, N 5.44, found: C 74.54, H 7.41, N 5.42.

2'-methyl-2,3,5,6,6',7',9',9a'-octahydro-8'H-spiro[pyran-4,10'-pyrido[1,2-a]indol]-8'-one (3l)

General procedure was followed using: 5-methyl-2',3',5',6'-tetrahydrospiro[indole-3,4'-pyran] **1e** (40 mg, 0.2 mmol), Yb(OTf)₃ (25 mg, 0.04 mmol) in toluene (1.6 ml) and (buta-1,3-dien-2-yloxy)trimethylsilane **2b** (69 μl, 0.4 mmol). Purification by flash chromatography (SiO₂, cyclohexane/EtOAc 3:1) yielded **3l** (40 mg, 73%) as white solid (m.p. 110.6-112.8° C). **¹H-NMR** (300 MHz, CDCl₃) 7.06 (s, 1H), 6.99 (d, J = 7.9 Hz, 1H), 6.53 (d, J = 7.9 Hz, 1H), 4.05 – 3.89 (m, 3H), 3.79 – 3.65 (m, 2H), 3.55 (ddd, J = 12.1, 9.5, 2.8 Hz, 1H), 3.20 (m, 1H), 2.54 (m, 1H), 2.42 – 2.24 (m, 6H), 2.00 (ddd, J = 13.6, 9.5, 3.9 Hz, 1H), 1.84 (m, 1H), 1.75 – 1.63 (m, 2H). **¹³C-NMR** (75 MHz, CDCl₃) 208.6 (C), 145.8 (C), 136.4 (C), 128.5 (CH), 128.4 (C), 124.7 (CH), 107.2 (CH), 70.7 (CH), 65.0 (CH₂), 64.9 (CH₂), 45.3 (C), 44.4 (CH₂), 41.8 (CH₂), 39.7 (CH₂), 35.5 (CH₂), 29.9 (CH₂), 20.9 (CH₃). **ESI(+)-MS:** m/z (%) = 272 (100) [M+1]⁺. Elemental analysis calcd for C₁₇H₂₁NO₂: C 75.25, H 7.80, N 5.16, found: C 75.08, H 7.82, N 5.17.

Benzyl 2'-fluoro-8'-oxo-7',8',9',9a'-tetrahydro-6'H-spiro[piperidine-4,10'-pyrido[1,2-a]indole]-1-carboxylate (3m)

General procedure was followed using: benzyl 5-fluorospiro[indole-3,4'-piperidine]-1'-carboxylate **1h** (68 mg, 0.2 mmol), Yb(OTf)₃ (25 mg, 0.04 mmol) in toluene (1.6 ml) and (buta-1,3-dien-2-yloxy)trimethylsilane **2b** (69 μl, 0.4 mmol). Purification by flash chromatography (SiO₂, heptane/EtOAc 2:1) yielded **3m** (65 mg, 79%) as white solid (m.p. 54.2-56.7° C). **¹H-NMR** (300 MHz, CDCl₃) 7.40 – 7.30 (m, 5H), 6.93 – 6.81 (m, 2H), 6.50 (dd, J = 8.4, 4.1 Hz, 1H), 5.15 (s, 2H), 3.99 – 3.81 (m, 3H), 3.62 (dd, J = 11.2, 3.5 Hz, 1H), 3.42 (m, 1H), 3.29 – 3.11 (m, 2H), 2.53 (td, J = 12.9, 6.7 Hz, 1H), 2.41 – 2.19 (m, 3H), 1.90 – 1.77 (m, 2H), 1.75 – 1.53 (m, 2H). **¹³C-NMR** (75 MHz, CDCl₃) 207.76 (C), 157.05 (d, J = 236.8 Hz, C), 155.26 (C), 144.26 (C), 137.38 (d, J = 7.1 Hz, C), 136.70 (C), 128.49 (2xCH), 128.04 (CH), 127.91 (2xCH), 114.18 (d, J = 23.2 Hz, CH), 111.73 (d, J = 24.6 Hz, CH), 107.66 (d, J = 8.2 Hz, CH), 70.72 (CH), 67.22 (CH₂), 45.89 (C), 44.46 (CH₂), 41.57 (CH₂), 40.95 (2xCH₂), 39.49 (CH₂), 34.25 (CH₂), 29.02 (CH₂).

FULL PAPER

ESI(+)-MS: *m/z* (%) = 409 (100) [M+1]⁺. Elemental analysis calcd for C₂₄H₂₅FN₂O₃: C 70.57, H 6.17, N 6.86, found: C 70.69, H 6.17, N 6.89.

6',7',9',9a'-tetrahydro-8'-H-spiro[cyclohexane-1,10'-[6,9]ethanopyrido[1,2-*a*]indol]-8'-one (3n/3'n)

General procedure was followed using: spiro[cyclohexane-1,3'-indole] **1a** (37 mg, 0.2 mmol), Yb(OTf)₃ (25 mg, 0.04 mmol) in toluene (1.6 ml) and (cyclohexa-1,5-dien-1-yloxy)trimethylsilane **2c** (74 μl, 0.4 mmol). Purification by flash chromatography (SiO₂ cyclohexane/EtOAc 9:1 to 8:2) yielded the two diastereoisomers **3n** (24 mg, 43%) and **3'n** (14 mg, 25%) as white solids (m.p. 132.3-134.7°C; 61.8-63.3°C respectively). **3n:** **¹H-NMR** (300 MHz, CD₂Cl₂) 7.18 – 7.07 (m, 2H), 6.82 (t, J = 7.3 Hz, 1H), 6.69 (d, J = 7.8 Hz, 1H), 4.09 (d, J = 3.3 Hz, 1H), 4.04 (s, 1H), 2.99 (d, J = 18.3 Hz, 1H), 2.53 (s, 1H), 2.43 (dd, J = 18.3, 3.3 Hz, 1H), 2.18 (m, 1H), 1.98 – 1.81 (m, 3H), 1.77 – 1.55 (m, 5H), 1.50 – 1.22 (m, 5H). **¹³C-NMR** (75 MHz, CD₂Cl₂) 212.9 (C), 150.7 (C), 140.7 (C), 127.2 (CH), 122.1 (CH), 119.3 (CH), 112.1 (CH), 67.1 (CH), 50.8 (CH), 46.2 (C), 44.3 (CH₂), 44.2 (CH), 41.7 (CH₂), 32.0 (CH₂), 25.5 (CH₂), 24.3 (CH₂), 22.4 (CH₂), 21.8 (CH₂), 17.5 (CH₂). **ESI(+)-MS:** *m/z* (%) = 282 (100) [M+1]⁺. Elemental analysis calcd for C₁₉H₂₃NO: C 81.10, H 8.24, N 4.98, found: C 80.93, H 8.23, N 4.98. **3'n:** **¹H-NMR** (300 MHz, CDCl₃) 7.08 (td, J = 7.8, 1.2 Hz, 1H), 7.01 (d, J = 6.8 Hz, 1H), 6.76 (td, J = 7.4, 0.9 Hz, 1H), 6.64 (d, J = 7.9 Hz, 1H), 4.08 (m, 1H), 3.96 (s, 1H), 2.70 – 2.75 (m, 2H), 2.31 (m, 1H), 2.17 – 1.80 (m, 6), 1.71 – 1.58 (m, 4H), 1.46 – 1.23 (m, 4H). **¹³C-NMR** (75 MHz, CDCl₃) 213.54 (C), 148.98 (C), 139.86 (C), 127.78 (CH), 123.19 (CH), 120.15 (CH), 112.41 (CH), 73.03 (CH), 49.90 (CH), 47.04 (C), 44.31 (CH), 41.63 (CH₂), 41.30 (CH₂), 31.42 (CH₂), 25.42 (CH₂), 24.38 (CH₂), 23.90 (CH₂), 23.31 (CH₂), 22.61 (CH₂). **ESI(+)-MS:** *m/z* (%) = 282 (70) [M+1]⁺. Elemental analysis calcd for C₁₉H₂₃NO: C 81.10, H 8.24, N 4.98, found: 80.85, H 8.26, N 4.96.

2'-fluoro-6',7',9',9a'-tetrahydro-8'-H-spiro[cyclohexane-1,10'-[6,9]ethanopyrido[1,2-*a*]indol]-8'-one (3o/3'o)

General procedure was followed using: 5'-fluorospiro[cyclohexane-1,3'-indole] **1c** (41 mg, 0.2 mmol), Yb(OTf)₃ (25 mg, 0.04 mmol) in toluene (1.6 ml) and (cyclohexa-1,5-dien-1-yloxy)trimethylsilane **2c** (74 μl, 0.4 mmol). Purification by flash chromatography (SiO₂ cyclohexane/EtOAc 8:2) yielded the two diastereoisomers **3o** (13 mg, 22%) and **3'o** (21 mg, 36%) as white solids (m.p. T>68° C (Dec), 78.3-81.6° C, respectively). **3o:** **¹H-NMR** (300 MHz, CDCl₃) 6.86 – 6.75 (m, 2H), 6.55 (dd, J = 8.3, 4.3 Hz, 1H), 4.06 – 3.93 (m, 2H), 2.97 (dt, J = 18.3, 2.7 Hz, 1H), 2.51 (t, J = 3.0 Hz, 1H), 2.42 (dd, J = 18.4, 3.4 Hz, 1H), 2.14 (m, 1H), 1.85 – 1.76 (m, 3H), 1.73 – 1.55 (m, 5H), 1.41 – 1.21 (m, 5H). **¹³C-NMR** (75 MHz, CDCl₃) 213.0 (C), 157.5 (d, J = 236.5 Hz, C), 146.3 (C), 142.2 (d, J = 6.9 Hz, C), 113.6 (d, J = 23.4 Hz, CH), 112.4 (d, J = 8.4 Hz, CH), 109.6 (d, J = 23.9 Hz, CH), 67.4 (CH), 51.1 (CH), 46.4 (C), 44.3 (CH₂), 43.9 (CH), 41.5 (CH₂), 31.9 (CH₂), 25.3 (CH₂), 24.1 (CH₂), 22.2 (CH₂), 21.6 (CH₂), 17.3 (CH₂). **ESI(+)-MS:** *m/z* (%) = 300 (100) [M+1]⁺. Elemental analysis calcd for C₁₉H₂₂FNO: C 76.22, H 7.41, N 4.68, found: C 76.44, H 7.43, N 4.67. **3'o:** **¹H-NMR** (300 MHz, CDCl₃) 6.80 (dd, J = 8.7, 2.6 Hz, 1H), 6.73 (m, 1H), 6.54 (dd, J = 8.5, 4.4 Hz, 1H), 4.05 – 3.96 (m, 2H), 2.67 – 2.53 (m, 2H), 2.29 (m, 1H), 2.16 – 1.96 (m, 3H), 1.72 – 1.57 (m, 4H), 1.45 – 1.26 (m, 4H). **¹³C-NMR** (75 MHz, CDCl₃) 213.3 (C), 157.8 (d, J = 237.2 Hz, C), 144.9 (C), 141.5 (d, J = 7.2 Hz, C), 114.2 (d, J = 23.6 Hz, CH), 112.7 (d, J = 8.3 Hz, CH), 110.5 (d, J = 23.7 Hz, CH), 73.3 (CH), 50.2 (CH), 47.2 (C), 44.1 (CH), 41.4 (CH₂), 41.1 (CH₂), 31.2 (CH₂), 25.3 (CH₂), 24.3 (CH₂), 23.7 (CH₂), 23.0 (CH₂), 22.5 (CH₂). **ESI(+)-MS:** *m/z* (%) = 300 (100) [M+1]⁺. Elemental analysis calcd for C₁₉H₂₂FNO: C 76.22, H 7.41, N 4.68, found: C 76.13, H 7.40, N 4.66.

2,3,5,6,6',7',9',9a'-octahydro-8'-H-spiro[pyran-4,10'-[6,9]ethanopyrido[1,2-*a*]indol]-8'-one (3p/3'p)

General procedure was followed using: 2',3',5',6'-tetrahydrospiro[indole-3,4'-pyran] **1d** (38 mg, 0.2 mmol), Yb(OTf)₃ (25 mg, 0.04 mmol) in toluene (1.6 ml) and (cyclohexa-1,5-dien-1-yloxy)trimethylsilane **2c** (74 μl, 0.4 mmol). Purification by flash chromatography (SiO₂ cyclohexane/EtOAc 3:1

to 2:1) yielded the two diastereoisomers **3p** (18 mg, 32%) and **3'p** 19 mg, 34%) as white solids (m.p. T>150° C (Dec), 118.6-120.9° C, respectively). **3p:** **¹H-NMR** (300 MHz, CDCl₃) 7.19 – 7.10 (m, 2H), 6.85 (td, J = 7.4, 0.8 Hz, 1H), 6.69 (d, J = 7.8 Hz, 1H), 4.13 – 3.99 (m, 3H), 3.89 (m, 1H), 3.60 (td, J = 11.9, 2.5 Hz, 1H), 3.51 (td, J = 12.0, 2.4 Hz, 1H), 3.00 (dt, J = 18.3, 2.6 Hz, 1H), 2.41-2.51 (m, 2H) 2.34 – 2.12 (m, 2H), 1.82 – 1.69 (m, 2H), 1.66 – 1.54 (m, 3H), 1.31 (m, 1H). **¹³C-NMR** (75 MHz, CDCl₃) 212.6 (C), 150.5 (C), 138.8 (C), 127.9 (CH), 122.3 (CH), 119.9 (CH), 112.4 (CH), 67.1 (CH), 65.9 (CH₂), 64.3 (CH₂), 50.9 (CH), 44.3 (CH₂), 44.2 (CH), 44.0 (C), 41.3 (CH₂), 32.3 (CH₂), 21.9 (CH₂), 17.2 (CH₂). **ESI(+)-MS:** *m/z* (%) = 284 (100) [M+1]⁺. Elemental analysis calcd for C₁₈H₂₁NO₂: C 76.30, H 7.47, N 4.94, found: C 76.48, H 7.45, N 4.92. **3'p:** **¹H-NMR** (300 MHz, CDCl₃) 7.14 (t, J = 7.6 Hz, 1H), 7.08 (d, J = 7.4 Hz, 1H), 6.82 (t, J = 7.4 Hz, 1H), 6.69 (d, J = 7.9 Hz, 1H), 4.17 – 4.02 (m, 3H), 3.96 (dt, J = 12.0, 3.6 Hz, 1H), 3.64 (td, J = 11.8, 2.4 Hz, 1H), 3.54 (td, J = 11.7, 2.8 Hz, 1H), 2.69 – 2.57 (m, 2H), 2.34 (m, 1H), 2.26 – 1.89 (m, 6H), 1.79 (m, 1H), 1.63 (m, 1H). **¹³C-NMR** (75 MHz, CDCl₃) 213.3 (C), 149.0 (C), 138.1 (C), 128.4 (CH), 123.2 (CH), 120.6 (CH), 112.6 (CH), 72.7 (CH), 66.0 (CH₂), 64.5 (CH₂), 49.8 (CH), 44.7 (C), 44.4 (CH), 41.6 (CH₂), 40.9 (CH₂), 31.5 (CH₂), 24.3 (CH₂), 23.0 (CH₂). **ESI(+)-MS:** *m/z* (%) = 284 (100) [M+1]⁺. Elemental analysis calcd for C₁₈H₂₁NO₂: C 76.30, H 7.47, N 4.94, found: C 76.07, H 7.46, N 4.95.

2'-methyl-2,3,5,6,6',7',9',9a'-octahydro-8'-H-spiro[pyran-4,10'-[6,9]ethanopyrido[1,2-*a*]indol]-8'-one (3q/3'q)

General procedure was followed using: 5-methyl-2',3',5',6'-tetrahydrospiro[indole-3,4'-pyran] **1e** (40 mg, 0.2 mmol), Yb(OTf)₃ (25 mg, 0.04 mmol) in toluene (1.6 ml) and (cyclohexa-1,5-dien-1-yloxy)trimethylsilane **2c** (74 μl, 0.4 mmol). Purification by flash chromatography (SiO₂ cyclohexane/EtOAc 3:1 to 2:1) yielded the two diastereoisomers **3q** (14 mg, 24%) and **3'q** (10 mg, 16%) as white solid (m.p. T>157.9° C (Dec); T>107° C (Dec), respectively). **¹H-NMR** (300 MHz, CDCl₃) 6.99 – 6.91 (m, 2H), 6.59 (d, J = 7.9 Hz, 1H), 4.10 (s, 1H), 4.08 – 4.00 (m, 2H), 3.89 (m, 1H), 3.65 – 3.45 (m, 2H), 2.98 (m, 1H), 2.44 (dd, J = 18.6, 3.2 Hz, 2H), 2.31 (s, 3H), 2.30 – 2.11 (m, 2H), 1.81 – 1.68 (m, 2H), 1.64 – 1.53 (m, 3H), 1.30 (m, 1H). **¹³C-NMR** (75 MHz, CDCl₃) 212.7 (C), 148.1 (C), 138.9 (C), 129.3 (C), 128.4 (CH), 122.9 (CH), 112.3 (CH), 67.1 (CH), 65.9 (CH₂), 64.4 (CH₂), 51.0 (CH), 44.3 (CH₂), 44.2 (CH), 44.0 (C), 41.4 (CH₂), 32.3 (CH₂), 21.7 (CH₂), 20.9 (CH₃), 17.2 (CH₂). **ESI(+)-MS:** *m/z* (%) = 298 (100) [M+1]⁺. Elemental analysis calcd for C₁₉H₂₃NO₂: C 76.74, H 7.80, N 4.71, found: C 76.51, H 7.83, N 4.73. **3'q:** **¹H-NMR** (300 MHz, CDCl₃) 6.92 (dd, J = 8.0, 1.0 Hz, 1H), 6.85 (s, 1H), 6.56 (d, J = 8.0 Hz, 1H), 4.08 – 3.99 (m, 3H), 3.92 (dt, J = 11.9, 3.6 Hz, 1H), 3.61 (td, J = 11.8, 2.5 Hz, 1H), 3.51 (td, J = 11.6, 2.9 Hz, 1H), 2.67 – 2.55 (m, 2H), 2.31 (m, 1H), 2.25 (s, 3H), 2.16 – 1.84 (m, 6H), 1.76 (m, 1H), 1.60 (m, 1H). **¹³C-NMR** (75 MHz, CDCl₃) 213.1 (C), 146.6 (C), 138.2 (C), 129.9 (C), 128.9 (CH), 123.8 (CH), 112.5 (CH), 72.9 (CH), 65.9 (CH₂), 64.5 (CH₂), 50.0 (CH), 44.7 (C), 44.4 (CH), 41.5 (CH₂), 41.0 (CH₂), 31.4 (CH₂), 24.3 (CH₂), 23.0 (CH₂), 20.9 (CH₃). **ESI(+)-MS:** *m/z* (%) = 298 (100) [M+1]⁺. Elemental analysis calcd for C₁₉H₂₃NO₂: C 76.74, H 7.80, N 4.71, found: C 76.86, H 7.79, N 4.70.

Benzyl 2'-fluoro-8'-oxo-7',8',9',9a'-tetrahydro-6'-H-spiro[piperidine-4,10'-[6,9]ethanopyrido[1,2-*a*]indole]-1-carboxylate (3r/3'r)

General procedure was followed using: benzyl 5-fluorospiro[indole-3,4'-piperidine]-1-carboxylate **1h** (68 mg, 0.2 mmol), Yb(OTf)₃ (25 mg, 0.04 mmol) in toluene and (1.6 ml) (cyclohexa-1,5-dien-1-yloxy)trimethylsilane **2c** (74 μl, 0.4 mmol). Purification by flash chromatography (SiO₂ cyclohexane/EtOAc 3:1 to 2:1) yielded the two diastereoisomers **3r** (16 mg, 18%) and **3'r** (25 mg, 29%) as white solid (m.p. T>157.9° C (Dec); T>107° C (Dec), respectively). **¹H-NMR** (300 MHz, CDCl₃) 7.39 – 7.30 (m, 5H), 6.86 (td, J = 8.7, 2.6 Hz, 1H), 6.77 (dd, J = 8.6, 2.6 Hz, 1H), 6.59 (dd, J = 8.6, 4.3 Hz, 1H), 5.16 (s, 1H), 4.21 – 3.99 (m, 4H), 3.16 – 2.92 (m, 3H), 2.50 – 2.39 (m, 2H), 2.22 – 1.96 (m, 2H, overlapped with EtOAc), 1.87 (m, 1H), 1.74 – 1.49 (m, 4H), 1.44 – 1.21 (m, 2H). **¹³C-NMR** (75 MHz, CDCl₃) 212.0 (C), 157.5 (d, J = 237.5 Hz, C), 155.2 (C), 146.4 (C), 139.8 (d, J = 7.1 Hz, C), 136.7 (C), 128.5 (2xCH), 128.1 (CH), 127.9 (2xCH), 114.4 (d, J = 23.4 Hz, CH), 112.7 (d, J = 8.3 Hz, CH), 109.7 (d, J = 24.1 Hz, CH),

FULL PAPER

67.3 (CH₂), 67.1 (CH), 51.1 (CH), 44.8 (C), 44.2 (CH₂), 44.1 (CH), 41.9 (CH₂), 40.3 (CH₂), 40.2 (CH₂), 31.4 (CH₂), 21.6 (CH₂), 17.2 (CH₂). **ESI(+)-MS**: m/z (%) = 457 (100) [M+Na]⁺. Elemental analysis calcd for C₂₆H₂₇FN₂O₃: C 71.87, H 6.26, N 6.45, found: C 72.04, H 6.28, N 6.42. **3^r**: **¹H-NMR** (300 MHz, CD₂Cl₂) 7.46 – 7.33 (m, 5H), 6.86 (td, J = 8.9, 2.5 Hz, 1H), 6.75 (dd, J = 8.8, 2.2 Hz, 1H), 6.62 (dd, J = 8.5, 4.4 Hz, 1H), 5.17 (s, 2H), 4.19 – 3.97 (m, 4H), 3.25 – 2.98 (m, 2H), 2.67 – 2.50 (m, 2H), 2.22 – 1.80 (m, 6H), 1.74 – 1.53 (m, 3H). **¹³C-NMR** (75 MHz, CD₂Cl₂) 212.6 (C), 157.5 (d, J = 236.7 Hz, C), 155.1 (C), 145.4 (C), 139.4 (d, J = 7.1 Hz, C), 137.1 (C), 128.4 (2xCH), 127.9 (2xCH), 127.8 (CH), 114.7 (d, J = 23.6 Hz, CH), 113.0 (d, J = 8.3 Hz, CH), 110.3 (d, J = 24.0 Hz, CH), 72.6 (CH), 66.9 (CH₂), 50.0 (CH), 45.5 (C), 44.2 (CH), 41.8 (CH₂), 41.4 (CH₂), 40.4 (CH₂), 39.7 (CH₂), 30.5 (CH₂), 24.2 (CH₂), 22.5 (CH₂). **ESI(+)-MS**: m/z (%) = 457 (100) [M+Na]⁺. Elemental analysis calcd for C₂₆H₂₇FN₂O₃: C 71.87, H 6.26, N 6.45, found: C 72.95, H 6.29, N 6.46.

Acknowledgements


The authors acknowledge MIUR-Italy (Ph.D. fellowship to E.B.) for financial support. Donatella Nava and Stefano Pandini (University of Milan) are thanked for some NMR and mass analyses.

Keywords: • imine, • aza-Diels-Alder, • nitrogen heterocycles, • spiroindolenine, • synthetic methods.

- [1] a) *Privileged Scaffolds in Medicinal Chemistry: Design, Synthesis, Evaluation* (Ed.: S. Bräse) RSC, Cambridge, **2016**; b) H. Zhao, J. Dietrich, *Expert Opin. Drug Discov.* **2015**, *10*, 781-790; c) E. K. Davison, M. A., *Curr. Opin. Chem. Biol.*, **2019**, *52*, 1-8.
- [2] a) *Comprehensive Natural Products II - Chemistry and Biology* (Eds: H.-W. Liu, L. Mander) Elsevier, Amsterdam, **2010**; b) N. A. Powell, J. T. Kohrt, K. J. Filipowski, M. Kaufman, D. Sheehan, J. E. Edmunds, A. Delaney, *Bioorg. Med. Chem. Lett.* **2012**, *22* 190-193; c) F. Shirai, T. Tsumura, Y. Yashiroda, H. Yuki, H. Niwa, S. Sato, T. Chikada, Y. Koda, K. Washizuka, N. Yoshimoto, M. Abe, T. Onuki, Y. Mazaki, C. Hiramata, T. Fukami, H. Watanabe, T. Honma, T. Umehara, M. Shirouzu, M. Okue, Y. Kano, T. Watanabe, K. Kitamura, E. Shitara, Y. Muramatsu, H. Yoshida, A. Mizutani, H. Seimiya, M. Yoshida, H. Koyama, *J. Med. Chem.* **2019**, *62*, 3407-3427.
- [3] a) M. J. James, P. O'Brien, R. J. K. Taylor, W. P. Unsworth, *Chem. Eur. J.* **2016**, *22*, 2856-2881; b) C. Zheng, S.-L. You, *Acc. Chem. Res.* **2020**, *53*, 974-987; c) J. Bariwal, L. G. Voskressensky, E. V. Van der Eycken, *Chem. Soc. Rev.* **2018**, *47*, 3831-3848.
- [4] Recent examples of [4+2] cycloaddition reactions with acyclic imines: a) K. Cheng, L. Lin, S. Chen, X. Feng, *Tetrahedron* **2005**, *61*, 9594-9599; b) D. Shang, J. Xin, Y. Liu, X. Zhou, X. Liu, X. Feng, *J. Org. Chem.* **2008**, *73*, 630-637; c) Y. Park, E. Park, H. Jung, Y.-J. Lee, S.-s. Jew, H.-g. Park, *Tetrahedron* **2011**, *67*, 1166-1170; d) T. Kagawa, K. Fujita, K. Kawada, *J. Fluor. Chem.* **2013**, *152*, 77-80; e) J. Kollmann, Y. Zhang, W. Schilling, T. Zhang, D. Riemer, S. Das, *Green Chem.* **2019**, *21*, 1916-1920; Review on cyclic imines: f) J. Iwanejko, E. Wojaczyńska, *Org. Biomol. Chem.* **2018**, *16*, 7296-7314; Recent examples of [4+2] cycloaddition reactions with cyclic imines: g) J. Shao, J.-S. Yang, *J. Org. Chem.* **2012**, *77*, 7891-7900; h) P. Szczesniak, S. Stecko, E. Maziarz, Olga S.-K., B. Furman, *J. Org. Chem.* **2014**, *79*, 10487-10503.
- [5] a) J. Li, J.-A. Xiao, S.-J. Zhao, H.-Y. Xiang, H. Yang, *Synthesis* **2017**, *49*, 4292-4298; b) O. Bakulina, A. Ivanov, V. Suslonov, D. Dar'in, M. Krasavin, *Beilstein J. Org. Chem.* **2017**, *13*, 1413-1424; c) H. Hu, C. Meng, Y. Dong, X. Li, J. Ye, *ACS Catal.* **2015**, *5*, 3700-3703.
- [6] a) G. Abbiati, V. Canevari, D. Facoetti, E. Rossi, *Eur. J. Org. Chem.* **2007**, 517-525; b) G. Abbiati, M. Dell'Acqua, D. Facoetti, V. Pirovano, M. Giordano, E. Rossi, *Synlett* **2012**, *23*, 2913-2918; c) V. Pirovano, M. Dell'Acqua, D. Facoetti, S. Rizzato, G. Abbiati, E. Rossi, *Eur. J. Org. Chem.* **2013**, 6267-6279; d) E. Rossi, V. Pirovano, M. Negrato, G. Abbiati, M. Dell'Acqua, *Beilstein J. Org. Chem.* **2015**, *11*, 1997-2006; e) E. Rossi, G. Abbiati, V. Pirovano, *Eur. J. Org. Chem.* **2017**, 4512-4529.
- [7] R. Sakhujia, K. Pericherla, K. Bajaj, B. Khungar, A. Kumar, *Synthesis* **2016**, *48*, 4305-4346.
- [8] a) J. F. Kerwin, Jr., S. Danishefsky, *Tetrahedron Lett.* **1982**, *23*, 3739-3742; b) X. Garcia-Mera, M. J. Alves, A. Goth, M. L. do Vale, J. E. Rodríguez-Borges, *Tetrahedron* **2013**, *69*, 2909-2919; c) K. Cheng, L. Lin, S. Chen, X. Feng, *Tetrahedron*, **2005**, *61*, 9594-9599; d) V. Jurčik, R. Wilhelm, *Org. Biomol. Chem.* **2005**, *3*, 239-244; e) Y. Yuan, X. Li, K. Ding, *Org. Lett.* **2002**, *4*, 3309-3311; f) L. Di Bari, S. Guillaume, J. Hanan, A. P. Henderson, J. A. K. Howard, G. Pescitelli, M. R. Probert, P. Salvadori, A. Whiting, *Eur. J. Org. Chem.* **2007**, 5771-5779.
- [9] a) B. G. Kelly, D. G. Gilheany, *Tetrahedron Lett.* **2002**, *43*, 887-890; b) H. Kitagawa, K. Kumura, S. Takahata, M. Iida, K. Atsumi, *Bioorg. Med. Chem.* **2007**, *15*, 1106-1116; c) F. Guo, R. C. Dhakal, R. K. Dieter, *J. Org. Chem.* **2013**, *78*, 8451-8464.
- [10] Supporting information file contains detailed procedure for the synthesis of starting compounds and compounds **5**, **6**, **7**, **8**; 2D NMR analysis for compounds **3a**, **3n**, **5** and **8** and copies of 1D NMR spectra for all unknown compounds.

FULL PAPER

Entry for the Table of Contents



Spiroindolenine serve as imine moiety for formal aza-Diels-Alder reactions with electronrich dienes. The reaction delivers tetrahydropyrido[1,2-*a*]spiroindolinones as in interesting class of rigid tridimensional polyheterocycles for drug discovery processes. Manipulation of the synthesized compounds encompasses transformation of the functional groups and formation of new carbon-carbon bonds.