

An overdetermined problem associated to the Finsler Laplacian

Giulio Ciraolo and Antonio Greco

Abstract

We prove a rigidity result for the anisotropic Laplacian. More precisely, the domain of the problem is bounded by an unknown surface supporting a Dirichlet condition together with a Neumann-type condition which is not translation-invariant. Using a comparison argument, we show that the domain is in fact a Wulff shape. We also consider the more general case when the unknown surface is required to have its boundary on a given conical surface: in such a case, the domain of the problem is bounded by the unknown surface and by a portion of the given conical surface, which supports a homogeneous Neumann condition. We prove that the unknown surface lies on the boundary of a Wulff shape.

1 Introduction

In this manuscript we study an overdetermined boundary value problem for elliptic equations. In these kinds of problems, a well posed elliptic PDEs problem is overdetermined by adding a further condition on the solution at the boundary and, for this reason, a solution may exist only if the domain and the solution itself satisfy some suitable symmetry. For instance, the well-known Serrin's overdetermined problem deals with the torsion problem in a bounded domain $\Omega \subseteq \mathbb{R}^N$

$$\begin{cases} \Delta u = -1 & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (1.1)$$

with the overdetermining condition

$$|Du| = c \quad \text{on } \partial\Omega, \quad (1.2)$$

for some positive constant c . Hence, Problem (1.1)-(1.2) is not well-posed and a solution may exist only if the domain (and the solution itself) satisfies some special symmetry (radial symmetry in this case, see [37]). There are many other results concerning overdetermined problems and, in particular, many generalizations of problem (1.1)-(1.2) have been considered in recent years, such as for quasilinear operators, for domains in convex cones, and in a Finsler (or anisotropic) setting (see for instance [1, 9, 16, 22, 24, 40] and references therein).

The anisotropic setting that we are considering can be described in terms of a norm H_0 in \mathbb{R}^N . Let H be the dual norm of H_0 (see Section 2), and consider the Finsler Laplacian Δ_H , whose definition is recalled in Section 3. Under convenient assumptions, Cianchi and Salani [9, Theorem 2.2] generalized Serrin's result to this setting and proved that the translation-invariant overdetermined problem

$$\begin{cases} -\Delta_H u = 1 & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega \\ H(Du(x)) = \text{const.} & \text{on } \partial\Omega \end{cases} \quad (1.3)$$

is solvable if and only if $\Omega = B_R(x_0, H_0)$ for some $x_0 \in \mathbb{R}^N$ and $R > 0$ (see also [1, Theorem 1.1] for the generalization to anisotropic p -Laplace equations).

If the overdetermining condition is not prescribed on the whole boundary, then the problem is called partially overdetermined. In this case, one can say less on the solution and a large variety of situations may occur. For instance, if we relax problem (1.1)-(1.2) by prescribing the Dirichlet condition $u = 0$ on a proper subset $\Gamma_0 \subseteq \partial\Omega$ instead that on the whole boundary, then the existence of a solution does not imply that Ω is a ball: the simplest counterexample is given by the annulus, and more refined counterexamples are found in [23]. Nevertheless, under convenient additional assumptions, a partially overdetermined problem turns out to be globally overdetermined and the conclusion can be recovered (see [17, 18, 21]).

In this paper we consider an anisotropic overdetermined problem in cones. Let $\Omega \subseteq \mathbb{R}^N$, $N \geq 2$, be a bounded domain (i.e. an open, connected, nonempty subset) containing the origin O , and let $\Sigma \subseteq \mathbb{R}^N$ be a cone

$$\Sigma = \{tx : x \in \omega, t \in (0, +\infty)\} \quad (1.4)$$

for some domain $\omega \subseteq S^{N-1}$. We mention that the equality $\omega = S^{N-1}$ (which implies $\Sigma = \mathbb{R}^N$) is allowed throughout the paper. In the case when $\omega \subsetneq S^{N-1}$ we require that $\partial\Sigma \setminus \{O\}$ is a hypersurface of class C^1 and therefore possesses an outward normal ν . Define

$$\Gamma_0 = \Sigma \cap \partial\Omega \quad \text{and} \quad \Gamma_1 = \partial\Omega \setminus \bar{\Gamma}_0. \quad (1.5)$$

Several problems in convex cones have been considered recently, like the isoperimetric and Sobolev inequalities in convex cones (see [5, 10, 30, 31]) and overdetermined and Liouville type problems in [10, 12, 32].

Here we extend the approach in [26] to the more general anisotropic setting and by considering a (possibly) mixed boundary-value problem. The starting point lies in the observation (done in [12, p. 28]) that the solution of (1.1) in the Euclidean ball $\Omega = B_R(O, |\cdot|)$ obviously satisfies (being a radial function) $u_\nu = 0$ along $\Gamma_1 \setminus \{O\}$ for every smooth cone $\Sigma \subsetneq \mathbb{R}^N$. Our main result is the following.

Theorem 1.1. *Let Ω and $\Sigma \subseteq \mathbb{R}^N$ be as above, and let H be a norm of class $C^1(\mathbb{R}^N \setminus \{O\})$ such that the function $V(\xi) = \frac{1}{2} H^2(\xi)$ is strictly convex. Let $q(r)$ be a positive, real-valued function such that the ratio $q(r)/r$ is increasing in $r > 0$. If there exists a weak solution $u \in C^1((\Sigma \cap \Omega) \cup (\Gamma_1 \setminus \{O\})) \cap C^0(\bar{\Sigma \cap \Omega} \setminus \{O\})$ of the problem*

$$\begin{cases} -\Delta_H u = 1 & \text{in } \Sigma \cap \Omega \\ u = 0 & \text{on } \Gamma_0 \\ DV(Du(x)) \cdot \nu = 0 & \text{on } \Gamma_1 \setminus \{O\}, \end{cases} \quad (1.6)$$

satisfying the condition

$$\lim_{x \rightarrow z} H(Du(x)) = q(H_0(z)) \quad \text{for all } z \in \bar{\Gamma}_0, \quad (1.7)$$

then $\Sigma \cap \Omega = \Sigma \cap B_R(O, H_0)$ for some $R > 0$.

In the case when $\Sigma = \mathbb{R}^N$ we have $\Gamma_1 = \emptyset$ and the third condition in (1.6) is trivially satisfied. If, in addition to $\Sigma = \mathbb{R}^N$, we also have $H(\cdot) = |\cdot|$, then Theorem 1.1 was proved by the second author in [26] under the weaker assumption that $q(r)/r$ is non-decreasing (see also [25, 27]). We mention that the rate of growth of q is crucial to obtain the rigidity result. A counterexample for the Euclidean norm can be found in [25, p. 488]. We also notice that, in the Euclidean case, the boundary condition on Γ_1 is simply $u_\nu = 0$.

We stress that problem (1.6)-(1.7) can be seen as a partially overdetermined problem, since the overdetermining condition is given only on the part $\bar{\Gamma}_0$ of the boundary. Accordingly, we are able to determine the shape of Γ_0 , while Γ_1 depends on the choice of the cone Σ .

We emphasize that no regularity assumption is imposed on Γ_0 . For this reason, we have to consider condition (1.7) instead of the simpler

$$H(Du(z)) = q(H_0(z)) \quad \text{on } \Gamma_0$$

(as, for instance, in [24]).

We also mention that Theorem 1.1 could be extended to the case in which the ratio $q(r)/r$ is non-decreasing in $r > 0$ by using Hopf's boundary point lemma (see [25]), as well as to more general anisotropic quasilinear operators (see for instance [34]). More precisely, one has to prove a Hopf's boundary point comparison principle between the solution and the solution in the Wulff shape. In this direction, the results in [7] can be a starting point for this investigation, and one can expect to prove a symmetry result in cones in the spirit of Theorem 1.1 for a class of anisotropic equations of the form $\operatorname{div}(DV(Du)) + f(u) = 0$.

The paper is organized as follows. In Section 2 we recall some well-known facts about norms in \mathbb{R}^N . In Section 3 we recall the definition of Finsler Laplacian and prove some basic properties of (1.6). Sections 2 and 3 will be the occasion to give full details of some basic facts, and for this reason we give a detailed description which is readable also at a beginner level. In Section 4 we give the proof of Theorem 1.1. In Appendix A we provide an example of a *smooth* norm having *non-smooth* dual norm (see also [6, Example A. 1.19]).

2 Norms, dual norms and Wulff shapes

In this section we collect the definitions and properties needed in the sequel. Further details are found in many recent papers: see, for instance, [1, Section 2.2], [3, Section 2.1], and [15, Section 2.3]. Standard references on convex analysis are [33] and [36] (see also [38, Section 5.3]).

2.1 Norms, convexity, and the Wulff shape

Let $H_0: \mathbb{R}^N \rightarrow \mathbb{R}$ be a norm on \mathbb{R}^N , $N \geq 1$, i.e. let H_0 be a nonnegative function such that

$$H_0(x) = 0 \text{ if and only if } x = 0; \tag{2.1}$$

$$H_0(tx) = |t| H_0(x) \text{ for all } t \in \mathbb{R} \text{ and } x \in \mathbb{R}^N; \tag{2.2}$$

$$H_0(x_1 + x_2) \leq H_0(x_1) + H_0(x_2) \text{ for all } x_1, x_2 \in \mathbb{R}^N. \tag{2.3}$$

The last inequality, known as *the triangle inequality*, may be equivalently replaced by the requirement that H_0 is a *convex function*, as in [1, 2, 9]. Indeed, from (2.2)-(2.3) it follows that every norm satisfies $H_0(\lambda x_1 + (1 - \lambda)x_2) \leq \lambda H_0(x_1) + (1 - \lambda)H_0(x_2)$ for all $\lambda \in (0, 1)$ and $x_1, x_2 \in \mathbb{R}^N$, i.e. H_0 is a convex function. Conversely, every nonnegative, convex function $H_0: \mathbb{R}^N \rightarrow \mathbb{R}$ satisfying (2.1) and (2.2) also satisfies (2.3): indeed, we may write $H_0(x_1 + x_2) = H_0((2x_1 + 2x_2)/2)$, and by convexity $H_0(x_1 + x_2) \leq \frac{1}{2}H_0(2x_1) + \frac{1}{2}H_0(2x_2)$. Now using (2.2) we arrive at (2.3) and hence H_0 is a norm (cf. [38, Theorem 5.3.8]).

We denote by $B_R(x_0, H_0) = \{x \in \mathbb{R}^N : H_0(x - x_0) < R\}$ the ball centered at x_0 and with radius $R > 0$ with respect to the norm H_0 (also called the *Wulff shape*).

2.2 Dual norms

As usual, the dual norm $H(\xi)$ of the norm $H_0(x)$ is defined for $\xi \in \mathbb{R}^N$ by

$$H(\xi) = \sup_{x \neq 0} \frac{x \cdot \xi}{H_0(x)}. \quad (2.4)$$

It is well known that the supremum above is indeed a maximum, i.e., it is attained with a particular $x \neq 0$. Furthermore, any given norm H_0 turns out to be the dual norm of its dual norm H : see, for instance, [4, Corollary 1.4].

2.3 Properties of the gradient of a norm

Let us recall some essential properties of the gradient DH_0 of a (differentiable) norm H_0 .

Lemma 2.1. *If H_0 is differentiable at some $x \in \mathbb{R}^N \setminus \{O\}$, then*

1. *The scalar product $x \cdot DH_0(x)$ satisfies*

$$x \cdot DH_0(x) = H_0(x). \quad (2.5)$$

2. *H_0 is differentiable at tx for every $t \in \mathbb{R} \setminus \{0\}$, and satisfies*

$$DH_0(tx) = (\text{sgn } t) DH_0(x). \quad (2.6)$$

3. *The gradient $DH_0(x)$ is a unit vector with respect to the dual norm H in the sense that*

$$H(DH_0(x)) = 1. \quad (2.7)$$

Proof. Property (2.5) is already found in the seminal dissertation by Finsler (cf. [20, (28)]) as well as in several recent papers: see, for instance, [3, (2.10)], [19, (1.8)] and [40, Proposition 1, (i)]. Equality (2.6), instead, can be derived from [3, (2.9)]. Let us give a proof, for completeness. Both (2.5) and (2.6) are obtained by differentiating the equality (2.2): more precisely, (2.5) follows by differentiating (2.2) in t at $t = 1$, while (2.6) is obtained by differentiation in x_i for $i = 1, \dots, N$. Equality 2.7 corresponds to [9, (3.12)], [19, (1.7)] and [40, Proposition 1, (iii)]. Let us prove the assertion and give a geometrical interpretation.




Figure 1: Maximizing the scalar product $y \cdot DH_0(x)$ under $H_0(y) = R$

Define $R = H_0(x)$ and consider the ball $B_R(O, H_0)$. By (2.4), in order to compute the dual norm $H(DH_0(x))$ it suffices to find a point $y \in \partial B_R(O, H_0)$ that maximizes the ratio

$$\frac{y \cdot DH_0(x)}{H_0(y)} = \frac{y \cdot DH_0(x)}{R}.$$

The hyperplane passing through x and orthogonal to $DH_0(x)$ is a supporting hyperplane for the convex set $B_R(O, H_0)$, and hence we may take $y = x$ to maximize the numerator (see Figure 1). Consequently we have

$$H(DH_0(x)) = \frac{x \cdot DH_0(x)}{R}.$$

Now using (2.5) the conclusion follows. \square

2.4 Differentiability of a norm

Let H_0 be a norm, and denote by H its dual norm. Because of (2.2), $H_0(x)$ is never differentiable at $x = 0$. By [36, Corollary 1.7.3], instead, differentiability of H_0 at $x \neq 0$ is related to the strict convexity of the unit ball $B_1(0, H)$ of the dual norm H : the next lemma collects several equivalent conditions.

Lemma 2.2. *The following conditions are equivalent.*

- (1) $H_0(x)$ is differentiable at every $x \neq 0$.
- (2) $H_0 \in C^1(\mathbb{R}^N \setminus \{O\})$.
- (3) $B_1(0, H_0)$ is a domain of class C^1 .
- (4) $B_1(0, H)$ is strictly convex.
- (5) The function $V(\xi) = \frac{1}{2} H^2(\xi)$ is strictly convex.

Proof. (2) \Rightarrow (1) is obvious. The converse implication (2) \Leftarrow (1) follows by [33, Corollary 25.5.1] because H_0 is a convex function. The implication (2) \Rightarrow (3) holds because $DH_0(x) \neq 0$ when $x \neq 0$ by homogeneity (2.2), and hence the level surface $\Gamma_1 = \partial B_1(0, H_0) = \{x \in \mathbb{R}^N : H_0(x) = 1\}$ is of class C^1 by the implicit function theorem. To prove the converse implication (2) \Leftarrow (3), observe that every $x \neq 0$ can be represented in polar coordinates ρ, η given by $\rho = |x|$ and $\eta = |x|^{-1}x$. In such a coordinate system, by (3) and by the convexity of $B_1(0, H_0)$ the surface $\Gamma_1 = \partial B_1(0, H_0)$ is the graph of a C^1 -function $\rho = \rho(\eta)$ whose domain is the Euclidean unit sphere $\mathbb{S}^{N-1} = \{\eta \in \mathbb{R}^N : |\eta| = 1\}$. Since $x/H_0(x) \in \Gamma_1$, we may write

$$\frac{x}{H_0(x)} = \rho(\eta) \eta = \rho(|x|^{-1}x) \frac{x}{|x|}$$

and hence $H_0(x) = |x|/\rho(|x|^{-1}x)$. Consequently, $H_0 \in C^1(\mathbb{R}^N \setminus \{O\})$ and (2) holds, hence (2) \Leftarrow (3). Let us check that (1) is equivalent to (4). As a consequence of [36, Corollary 1.7.3], the unit ball $B_1(0, H)$ of the dual norm H is strictly convex (its boundary does not contain any segment) if and only if H_0 is differentiable at every $x \in \mathbb{R}^N \setminus \{O\}$, namely (1) \Leftrightarrow (4), as claimed. The preceding arguments imply that the first four conditions in the statement are equivalent to each other. To complete the proof we now verify that (4) is equivalent to (5). Before proceeding further, observe that $2V(\xi)$ (hence $V(\xi)$ as well) is convex because it is the square of the convex, nonnegative function $H(\xi)$. Hence $V(\xi)$ is not strictly convex if and only if there exists a line

segment $\ell \subseteq \mathbb{R}^N$ such that the restriction $V|_\ell$ is a linear function. But then $H(\xi) = \sqrt{2V(x)}$ is concave along ℓ . Since $H(\xi)$ is also convex, it follows that $H(\xi)$ is constant along ℓ . Hence $V(\xi)$ fails to be strictly convex if and only if $H(\xi)$ takes a constant value (say c) along some segment ℓ . Now recall that $H(\xi)$ is homogeneous of degree 1: this has two relevant consequences. The first consequence is that $H(\xi)$ is not constant in the radial direction, hence the segment ℓ is not aligned with the origin. The second consequence is that we may find a new segment, say ℓ' , parallel to ℓ and such that $H(\xi) = 1$ on ℓ' : indeed, we may take

$$\ell' = \{ \xi \in \mathbb{R}^N : c\xi \in \ell \}.$$

In short, there is no loss of generality if we assume $c = 1$. But then we may assert that $V(\xi)$ is not strictly convex if and only if $\partial B_1(0, H)$ contains a line segment, i.e., if and only if $B_1(0, H)$ is not strictly convex. This proves the equivalence between (4) and (5), and the lemma follows. \square

It may well happen that $H_0 \in C^1(\mathbb{R}^N \setminus \{O\})$ and $H \notin C^1(\mathbb{R}^N \setminus \{O\})$: see Section A or [6, Example A. 1.19]. If both H_0 and H are smooth, then it is relevant for our purposes to notice that the gradient $DH(\xi)$ evaluated at $\xi = DH_0(x)$ is radial in the following sense (see also [3, Lemma 2.2] and [19, c), p. 249]):

Lemma 2.3. *If $H_0, H \in C^1(\mathbb{R}^N \setminus \{O\})$ then $x = H_0(x) DH(DH_0(x))$ for all $x \in \mathbb{R}^N \setminus \{O\}$. Furthermore, for every $\xi \neq O$, a point $x \neq O$ realizes the supremum in (2.4) if and only if*

$$x = H_0(x) DH(\xi). \quad (2.8)$$

Proof. The unit ball $B_1(0, H_0)$ is a strictly convex domain of class C^1 , hence for every $\xi \in \partial B_1(0, H)$ there exists a unique $x \in \partial B_1(0, H_0)$ that maximizes the linear function $L(x) = x \cdot \xi$ under the constraint $H_0(x) = 1$. Taking the definition (2.4) of $H(\xi)$ into account, we may say that for every $\xi \in \partial B_1(0, H)$ there exists a unique $x \in \partial B_1(0, H_0)$ satisfying the equality

$$1 = x \cdot \xi.$$

Furthermore there exists $\lambda \in \mathbb{R}$ (the Lagrange multiplier) such that $\xi = \lambda DH_0(x)$. More precisely, since $x \cdot DH_0(x) > 0$ by the convexity of $B_1(0, H_0)$, we have $\lambda > 0$. This and (2.7) imply $H(\xi) = \lambda$, and therefore $\xi = H(\xi) DH_0(x)$. By reverting the roles of x and ξ we get (2.8) and the lemma follows using (2.6). \square

We conclude this section with the following proposition:

Proposition 2.4 (Regularity of the Lagrangian). *Let H be a norm of class $C^1(\mathbb{R}^N \setminus \{O\})$. Then the Lagrangian $V(\xi) = \frac{1}{2} H^2(\xi)$ belongs to the class $C^1(\mathbb{R}^N)$*

Proof. Since all norms on \mathbb{R}^N are equivalent, there exist two positive constants σ, γ such that

$$\sigma |\xi| \leq H(\xi) \leq \gamma |\xi| \text{ for all } \xi \in \mathbb{R}^N \quad (2.9)$$

(the notation is taken from [9, (3.4)]). In particular, $H(\xi)$ is continuous at $\xi = 0$. Concerning the differentiability, for $\xi \neq 0$ we may apply the standard rules of calculus and get

$$DV(\xi) = H(\xi) DH(\xi). \quad (2.10)$$

Furthermore, the right-hand side admits a continuous extension to $\xi = 0$ because $DH(\xi)$ is bounded on the compact surface $\mathbb{S}^{N-1} = \{ \xi : |\xi| = 1 \}$, and by (2.6) $DH(\xi)$ is also bounded in the whole punctured space $\mathbb{R}^N \setminus \{O\}$. Hence $DV(\xi) \rightarrow 0$ as $\xi \rightarrow 0$. This and the continuity of $V(\xi)$ at $\xi = 0$ imply that V is also differentiable at $\xi = 0$, and $DV(0) = 0$. The proposition follows. \square

3 The Finsler Laplacian

Given a norm H of class $C^1(\mathbb{R}^N \setminus \{O\})$, the function $V(\xi) = \frac{1}{2} H^2(\xi)$ belongs to the class $C^1(\mathbb{R}^N)$ by Proposition 2.4. The *Finsler Laplacian* associated to H is the differential operator Δ_H which is formally defined by

$$\Delta_H u(x) = \operatorname{div}\left(DV(Du(x))\right).$$

Notation. In the present paper it is understood that the gradient operator D takes precedence over the composition of functions: thus, the notation $DV(Du(x))$ represents the vector field $DV(\xi)$ evaluated at the point $\xi = Du(x)$. Such a vector field differs, in general, from the field whose components are the derivatives of $V(Du(x))$ with respect to x_i , $i = 1, \dots, N$. Clearly, if $H(\xi)$ is the Euclidean norm $|\xi|$ then $V(\xi) = \frac{1}{2} |\xi|^2$ and therefore $DV(\xi) = \xi$. Thus, the operator Δ_H reduces to the standard Laplacian Δ .

Let Ω be a bounded domain in \mathbb{R}^N , $N \geq 2$, containing the origin O . Let Σ and Γ_0, Γ_1 be as in (1.4) and (1.5), respectively. We define the function space

$$W_{\Gamma_0}^{1,2}(\Omega \cap \Sigma) = \{v: \Omega \cap \Sigma \rightarrow \mathbb{R} \text{ s.t. } v = w \chi_{\Omega \cap \Sigma} \text{ for some } w \in W_0^{1,2}(\Omega)\},$$

where $\chi_{\Omega \cap \Sigma}$ denotes the characteristic function of $\Omega \cap \Sigma$. Notice that a function v in $W_{\Gamma_0}^{1,2}(\Omega \cap \Sigma)$ has zero trace on Γ_0 .

Definition 3.1 (Weak solution). Let Ω be as above and let f be a function in $L^2(\Omega \cap \Sigma)$. A *weak solution* of

$$\begin{cases} -\Delta_H u = f & \text{in } \Omega \cap \Sigma; \\ u = 0 & \text{on } \Gamma_0 \\ DV(Du) \cdot \nu = 0 & \text{on } \Gamma_1 \setminus \{O\} \end{cases} \quad (3.1)$$

is a function $u \in W_{\Gamma_0}^{1,2}(\Omega \cap \Sigma)$ such that

$$\int_{\Omega \cap \Sigma} Dv(x) \cdot DV(Du(x)) dx = \int_{\Omega \cap \Sigma} f(x) v(x) dx \quad (3.2)$$

for every $v \in W_{\Gamma_0}^{1,2}(\Omega \cap \Sigma)$.

Theorem 3.2 (Existence). *Let Ω , Σ and f be as above. If H is a norm of class $C^1(\mathbb{R}^N \setminus \{O\})$, then Problem (3.1) has a weak solution.*

Proof. Define $V(\xi) = \frac{1}{2} H^2(\xi)$. By (2.9), and by the Poincaré inequality in $W_{\Gamma_0}^{1,2}(\Omega \cap \Sigma)$ (see [35, Theorem 7.91]), the functional

$$F[u] = \int_{\Omega \cap \Sigma} \left(V(Du(x)) - f(x) u(x) \right) dx \quad (3.3)$$

is well defined and coercive over the Sobolev space $W_{\Gamma_0}^{1,2}(\Omega \cap \Sigma)$, hence there exists a minimizer. Since the functional F is differentiable (as a consequence of Proposition 2.4), each minimizer is a weak solution of the Euler equation $-\Delta_H u = f(x)$. \square

Remark 3.3. If, in addition to the assumption of Theorem 3.2, the function $V(\xi)$ is strictly convex, then the functional F in (3.3) is also strictly convex, and the minimizer is unique. Uniqueness of the weak solution to Problem (3.1) also follows by letting $\Omega^1 = \Omega^2 = \Omega$ in Lemma 3.5. Several conditions equivalent to the strict convexity of $V(\xi)$ are given in Lemma 2.2.

In view of our subsequent application we now prepare the following comparison principle, which asserts that if $f \geq 0$ then the solution of (3.1) is not only unique but also nonnegative and monotonically increasing with respect to set inclusion.

Lemma 3.4 (Nonnegativity). *Let H be a norm of class $C^1(\mathbb{R}^N \setminus \{O\})$, and let Ω and Σ be as above. If $f \in L^2(\Omega \cap \Sigma)$ is nonnegative, then any weak solution u to (3.1) is also nonnegative.*

Proof. By (2.10) and (2.5) we find $\xi \cdot DV(\xi) = 2V(\xi)$ for $\xi \neq 0$. The equality continues to hold at $\xi = 0$ by Proposition 2.4. Hence, using $v(x) = -u^-(x) = \min\{u(x), 0\}$ as a test-function in (3.2) we get

$$0 \leq 2 \int_{\Omega \cap \Sigma} V(Dv(x)) dx = 2 \int_{\{u < 0\}} V(Du(x)) dx = \int_{\{u < 0\}} f(x) u(x) dx \leq 0,$$

which implies $Dv(x) = 0$ almost everywhere in $\Omega \cap \Sigma$. Since $v \in W_{\Gamma_0}^{1,2}(\Omega \cap \Sigma)$, by the Poincaré inequality [35, Theorem 7.91] it follows that $v = 0$, hence $u \geq 0$ a.e. in $\Omega \cap \Sigma$. \square

Lemma 3.5 (Monotonicity). *Let H be a norm of class $C^1(\mathbb{R}^N \setminus \{O\})$ such that the function $V(\xi) = \frac{1}{2} H^2(\xi)$ is strictly convex. Let Σ be as in (1.4), and let Ω^i , $i = 1, 2$, be two bounded domains in \mathbb{R}^N , $N \geq 2$, containing the origin and satisfying $\Omega^1 \cap \Sigma \subseteq \Omega^2 \cap \Sigma$. Choose a nonnegative $f \in L^2(\Omega^2 \cap \Sigma)$, and denote by u_i any weak solution of Problem (3.1) with $\Omega = \Omega^i$. Then $u_1 \leq u_2$ almost everywhere in Ω^1 .*

Proof. Let $\Gamma_0^i = \Sigma \cap \partial\Omega^i$, $i = 1, 2$. Since $f \geq 0$, from Lemma 3.4 we have $u_2 \geq 0$ a.e. in Ω_2 . Hence the function $v = (u_1 - u_2)^+$ belongs to $W_{\Gamma_0^1}^{1,2}(\Omega^1 \cap \Sigma)$ and has an extension, still denoted by v , to $W_{\Gamma_0^2}^{1,2}(\Omega^2 \cap \Sigma)$ vanishing identically outside $\Omega^1 \cap \Sigma$. Therefore v is an admissible test-function in Definition 3.1 for $\Omega = \Omega^i$, $i = 1, 2$, and we may write

$$\begin{aligned} \int_{\Omega^1 \cap \Sigma} Dv(x) \cdot DV(Du_1(x)) dx &= \int_{\Omega^1 \cap \Sigma} f(x) v(x) dx, \\ \int_{\Omega^1 \cap \Sigma} Dv(x) \cdot DV(Du_2(x)) dx &= \int_{\Omega^1 \cap \Sigma} f(x) v(x) dx. \end{aligned}$$

By subtracting the second equality from the first one we obtain

$$\int_{\{v > 0\}} \left(Du_1(x) - Du_2(x) \right) \cdot \left(DV(Du_1(x)) - DV(Du_2(x)) \right) dx = 0$$

Since V is strictly convex by assumption, the Lebesgue measure of the set $\{v > 0\}$ must be zero, and the lemma follows. \square

In the case when $\Omega = B_R(O, H_0)$ for some $R > 0$ and $f \equiv 1$, Problem (3.1) is explicitly solvable:

Proposition 3.6 (Solution in the Wulff shape). *Let H be a norm of class $C^1(\mathbb{R}^N \setminus \{O\})$, and suppose that its dual norm H_0 also belongs to $C^1(\mathbb{R}^N \setminus \{O\})$. Let Σ be as in (1.4). The function $u_R \in C^1(\mathbb{R}^N)$ given by $u_R(x) = \frac{1}{2N} (R^2 - H_0^2(x))$ is a weak solution of the problem*

$$\begin{cases} -\Delta_H u = 1 & \text{in } B_R(O, H_0) \cap \Sigma, \\ u = 0 & \text{on } \Sigma \cap \partial B_R(O, H_0), \\ DV(Du) \cdot \nu = 0 & \text{on } B_R(O, H_0) \cap \partial\Sigma \setminus \{O\}. \end{cases} \quad (3.4)$$

Furthermore, the gradient Du_R is given by $Du_R(x) = -\frac{1}{N} H_0(x) DH_0(x)$ for $x \neq 0$ and satisfies $H(Du_R(x)) = \frac{1}{N} H_0(x)$ for all $x \in \mathbb{R}^N$.

Proof. By differentiation we find $Du_R(x) = -\frac{1}{N} H_0(x) DH_0(x)$ for $x \neq 0$, and therefore $H(Du_R(x)) = \frac{1}{N} H_0(x)$ by (2.7). The last equality continues to hold at the origin by Proposition 2.4. Let us check that u_R satisfies (3.4) in the weak sense. Of course, u_R vanishes by definition when $H_0(x) = R$. Since $u_R \in C^1(\mathbb{R}^N)$ and the boundary of $\Omega = B_R(0, H_0)$ also belongs to the class C^1 , it follows that $u_R \in W_{\Gamma_0}^{1,2}(\Omega \cap \Sigma)$, where Γ_0 is as in (1.5). Furthermore, by (2.10), (2.6) and (2.7) we have $DV(Du_R(x)) = H(Du_R(x)) DH(Du_R(x)) = -\frac{1}{N} H_0(x) DH(DH_0(x))$. But then by Lemma 2.3 it follows that $DV(Du_R(x)) = -\frac{1}{N} x$. We note in passing that $DV(Du_R(x)) \cdot \nu = 0$ pointwise on $\Omega \cap \partial\Sigma \setminus \{O\}$. To complete the proof we have to show that (3.2) holds. This is peculiar because, although u_R may fail to have second derivatives, the compound function $DV(Du_R(x)) = -\frac{1}{N} x$ belongs to $C^\infty(\mathbb{R}^N, \mathbb{R}^N)$, and therefore by the divergence theorem we have

$$\int_{\Omega \cap \Sigma} Dv(x) \cdot DV(Du(x)) dx = -\frac{1}{N} \int_{\Omega \cap \Sigma} Dv(x) \cdot x dx = \int_{\Omega \cap \Sigma} v(x) dx$$

for every $v \in W_{\Gamma_0}^{1,2}(\Omega \cap \Sigma)$, as claimed. \square

Remark 3.7. In the case when $\Sigma = \mathbb{R}^N$, the solution in the Wulff shape is considered, for instance, in [1, (1.8)] and [19, Theorem 2.1].

4 Proof of Theorem 1.1

Roughly speaking, Theorem 1.1 asserts that if $q(r)$ grows faster than r then the solvability of Problem (1.6)-(1.7) implies that Ω is a Wulff shape centered at the origin. The minimal rate of increase of $q(r)$ in order to get the result is discovered by letting R vary in Problem (3.4): more precisely, using Proposition 3.6 we find that $H(Du_R(x)) = R/N$ for every $x \in \partial B_R(O, H_0)$, hence the value of $H(Du_R(x))$ at $x \in \partial B_R(O, H_0)$ is proportional to R . This information is transferred to Problem (1.6) by means of the following comparison argument.

Proof of Theorem 1.1. Preliminaries. Define

$$R_1 = \min_{z \in \bar{\Gamma}_0} H_0(z), \quad R_2 = \max_{z \in \bar{\Gamma}_0} H_0(z)$$

and let u_i , $i = 1, 2$, be the solution of the Dirichlet problem (3.4) in the Wulff shape $\Omega_i = B_{R_i}(0, H_0)$. Thus, $\Sigma \cap \Omega_1 \subseteq \Sigma \cap \Omega \subseteq \Sigma \cap \Omega_2$. We aim to prove that $\Omega_1 = \Omega_2$, which implies the claim of the theorem. To this purpose, pick $z_i \in \bar{\Gamma}_0 \cap \partial\Omega_i$ and observe that $R_i = H_0(z_i)$, $i = 1, 2$. Using Lemma 3.4 and Lemma 3.5 twice, we get

$$u_1 \leq u \text{ a.e. in } \Sigma \cap \Omega_1, \quad u \leq u_2 \text{ a.e. in } \Sigma \cap \Omega. \quad (4.1)$$

Part 1. Taking into account that $u_1(z_1) = u(z_1) = 0$ and u_1 is continuously differentiable up to z_1 , let us check that the first inequality in (4.1) implies

$$\frac{R_1}{N} = H(Du_1(z_1)) \leq q(R_1). \quad (4.2)$$

Letting $x(t) = z_1 - t|z_1|^{-1} z_1 \in \bar{\Sigma} \cap \Omega_1$ for $t \in (0, |z_1|)$, we compute the limit

$$\ell = \lim_{t \rightarrow 0^+} \frac{u_1(x(t))}{t}$$

following two different arguments. On the one side, the limit ℓ is the radial derivative $\ell = -|z_1|^{-1} z_1 \cdot Du_1(z_1)$, and using Proposition 3.6 and equality (2.5) we may write

$$\ell = \frac{1}{N} |z_1|^{-1} R_1^2.$$

On the other side, by the mean-value theorem we have $u(x(t)) = -t |z_1|^{-1} z_1 \cdot Du(\tilde{x})$ for a convenient point \tilde{x} on the segment from z_1 to $x(t)$. Letting $\xi = Du(\tilde{x})$ and $x = t |z_1|^{-1} z_1$ in (2.4), and since $H_0(z_1) = R_1$, we may estimate $u(x(t)) \leq t R_1 |z_1|^{-1} H(Du(\tilde{x}))$. Recalling that $u_1 \leq u$ by (4.1), and using assumption (1.7) we arrive at $\ell \leq R_1 |z_1|^{-1} q(R_1)$ and (4.2) follows.

Part 2. By using the second inequality in (4.1), and since u_2 is continuously differentiable, taking assumption (1.7) into account we now prove the inequality

$$q(R_2) \leq H(Du_2(z_2)) = \frac{R_2}{N}. \quad (4.3)$$

The argument is by contradiction: suppose there exists $\epsilon_0 \in (0, |z_2|)$ such that $H(Du(x)) > H(Du_2(z_2)) + \epsilon_0$ for all $x \in U_0 = \{x \in \Sigma \cap \Omega : |x - z_2| < \epsilon_0\}$, and choose $x_0 \in U_0$. Observe that $w_0 = u_2(x_0) - u(x_0) > 0$ because the equality $w_0 = 0$ together with $u \leq u_2$ implies $Du_2(x_0) = Du(x_0)$, which is not the case. While ϵ_0 is kept fixed, the point x_0 will tend to z_2 in the end of the argument. Since the vector field $DH(\xi)$ is continuous by assumption in $\mathbb{R}^N \setminus \{O\}$, for every choice of $x_0 \in U_0$ there exists a local solution $x(t)$, $t > 0$, of the initial-value problem

$$\begin{cases} x'(t) = DH(Du(x(t))), \\ x(0) = x_0. \end{cases}$$

Since the Euclidean norm $|x'(t)|$ is bounded from above by some constant M_0 , the length of the arc γ described by $x(\tau)$ when τ ranges in the interval $(0, t)$ satisfies $|\gamma| \leq M_0 t$, and therefore

$$|x(t) - z_2| \leq M_0 t. \quad (4.4)$$

We claim that the curve γ can be extended until $|x(t_0) - z_2| = \epsilon_0$ for some finite $t_0 > 0$. Indeed, by differentiation of $u(x(t))$ we find $du/dt = Du(x(t)) \cdot x'(t)$. Letting $\xi = Du(x(t))$ in Lemma 2.3 we see that the vector $x'(t)$ realizes the supremum in (2.4), i.e., we may write the equality

$$H(Du(x(t))) = \frac{Du(x(t)) \cdot x'(t)}{H_0(x'(t))}.$$

This and (2.7) imply $du/dt = H(Du(x(t)))$. Since $H(Du(x)) > \epsilon_0$ in U_0 , it follows that u increases along γ and therefore the curve, which starts at $x_0 \in U_0$, cannot end on $\bar{\Gamma}_0$ where $u = 0$. Similarly, we find $du_2/dt = Du_2(x(t)) \cdot x'(t) \leq H(Du_2(x(t)))$ and therefore

$$\frac{du_2}{dt} + \epsilon_0 < \frac{du}{dt}, \quad (4.5)$$

hence the difference $w(t) = u_2(x(t)) - u(x(t))$ satisfies $dw/dt < -\epsilon_0$. Since $w(t)$ must be positive as long as $x(t) \in U_0$, it follows that

$$t < w_0/\epsilon_0 \quad (4.6)$$

and the length of γ is bounded from above by

$$|\gamma| \leq M_0 w_0/\epsilon_0. \quad (4.7)$$

In the case when γ reaches prematurely $\Gamma_1 \subseteq \partial\Sigma$, i.e., if $x(t) \in U_0$ for $t \in [0, t_1)$ and $x(t_1) = x_1 \in \Gamma_1$, with $|x_1 - z_2| < \epsilon_0$, the assumption that $\partial\Sigma$ is of class C^1 ensures the existence of a local solution $x(t) \in \partial\Sigma$, $t > t_1$, to the following initial-value problem:

$$\begin{cases} x'(t) = DH(Du(x(t))), \\ x(t_1) = x_1. \end{cases} \quad (4.8)$$

Indeed, the third condition in (1.6) implies that $DH(Du(x))$ is a tangent vector to $\partial\Sigma$ as long as $x \in \partial\Sigma$, and therefore problem (4.8) admits a local solution lying on the hypersurface $\partial\Sigma$ and extending the curve γ . Since the curve γ , possibly extended as above, has a finite length by (4.7), and cannot end either on $\bar{\Gamma}_0$ nor on Γ_1 as long as $|x(t) - z_2| < \epsilon_0$, nor can it have a limiting point in U_0 because $x'(t)$ keeps far from zero and the parameter t is bounded by (4.6), there must be some finite $t_0 > 0$ such that $|x(t_0) - z_2| = \epsilon_0$, and therefore $|\gamma| \geq \epsilon_0$. This and (4.7) yield the estimate $\epsilon_0^2 \leq M_0 w_0$, which is uniform in the sense that M_0 and ϵ_0 do not depend on the choice of $x_0 \in U_0$. To conclude the argument, we now let $x_0 \rightarrow z_2$: thus, $w_0 \rightarrow 0$ while ϵ_0 and M_0 do not change, and a contradiction is reached.

Conclusion. By (4.2) and (4.3) we deduce

$$\frac{q(R_2)}{R_2} \leq \frac{q(R_1)}{R_1}.$$

Since the ratio $q(r)/r$ strictly increases, we must have $R_1 = R_2$ and $\Omega_1 = \Omega_2 = \Omega$, as claimed. \square

A A smooth norm with a non-smooth dual norm

A simple example of norm in \mathbb{R}^N is the p -norm $|x|_p$ given by

$$|x|_p = \left(\sum_{k=1}^N |x_k|^p \right)^{\frac{1}{p}} \text{ for } p \in [1, +\infty).$$

In the special case when $p \in (1, +\infty)$, the dual norm of $|x|_p$ is the q -norm $|\xi|_q$, where q is related to p by the equality $\frac{1}{p} + \frac{1}{q} = 1$. Both norms belong to the class $C^1(\mathbb{R}^N \setminus \{O\})$. The dual norm of $|x|_1$, instead, is $|\xi|_\infty = \max\{|\xi_1|, \dots, |\xi_N|\}$. Neither of the last two norms belongs to the class $C^1(\mathbb{R}^N \setminus \{O\})$. In this section we construct an explicit example of a norm $H_0 \in C^1(\mathbb{R}^2 \setminus \{O\})$ whose dual norm H does not belong to the same class. The example below should be compared with [6, Example A. 1.19].

Definition A.1. The norm H_0 is defined as the *gauge function*, also called *the Minkowski functional* (see [28, Definition 6.3.11] or [38, Remark 1, p. 380]), of a convenient, convex, plane domain which plays the role of the unit ball $B_1(0, H_0)$. Such a ball is defined as the convex envelope of the four Euclidean discs of radius $\frac{1}{2}$ centered at $(\pm\frac{1}{2}, 0)$, $(0, \pm\frac{1}{2})$ (see Figure 2). Notice that the origin belongs to the boundary of each of the given discs.

Since the boundary of the ball $B_1(0, H_0)$ defined above is a C^1 -curve, by Lemma 2.2 the norm H_0 belongs to the class $C^1(\mathbb{R}^2 \setminus \{O\})$. However, we have:

Lemma A.2. *Denote by $H(\xi)$ the dual norm of the norm $H_0(x)$ in Definition A.1. The unit ball $B_1(0, H)$ is the intersection of four convex open sets, each bounded by a parabola with focus at the origin and vertex at one of the four points $(\pm 1, 0)$, $(0, \pm 1)$.*

Before proving the lemma, we notice that the boundary $\partial B_1(0, H)$ of the unit ball described in the statement has a corner at each of the four points $(\pm\bar{\xi}, \pm\bar{\xi})$, $\bar{\xi} = 2(\sqrt{2} - 1)$. To see this, let us consider the parabola γ with focus at O and vertex at $(1, 0)$. This line is the graph of the function $\xi_1(\xi_2) = 1 - \frac{1}{4}\xi_2^2$ whose derivative satisfies $\xi_1'(\bar{\xi}) = -\frac{1}{2}\bar{\xi} = 1 - \sqrt{2} > -1$. Hence γ is not orthogonal to the straight line $\xi_1 = \xi_2$ at the point of intersection $(\bar{\xi}, \bar{\xi})$, and consequently the boundary $\partial B_1(0, H)$ must have a corner there. Thus, the dual norm H of the given norm H_0 does not belong to the class $C^1(\mathbb{R}^2 \setminus \{O\})$.




Figure 2: The ball $B_1(0, H_0)$ (left) is smooth, its dual (right) is not.

Proof of Lemma A.2. Let us describe the boundary of the ball $B_1(0, H)$ in parametric form. Passing to polar coordinates ρ, ϑ related to ξ_1, ξ_2 by $\xi_1 = \rho \cos \vartheta$, $\xi_2 = \rho \sin \vartheta$, for every $\vartheta \in (-\frac{\pi}{4}, \frac{\pi}{4})$ we compute the Euclidean norm $\rho(\vartheta) = |v_\vartheta|$ of the unique vector $v_\vartheta = \rho(\vartheta) (\cos \vartheta, \sin \vartheta)$ satisfying $H(v_\vartheta) = 1$. To this purpose it is enough to locate the point $P_\vartheta \in \partial B_1(0, H_0)$ where the outer normal ν equals $(\cos \vartheta, \sin \vartheta)$: indeed, due to (2.7), we have $H(DH_0(P_\vartheta)) = 1$ and therefore we may take $v_\vartheta = DH_0(P_\vartheta)$. Recalling that $H_0(P_\vartheta) = 1$, the radial derivative $\partial H_0 / \partial r$, $r = \sqrt{x_1^2 + x_2^2}$, is easily computed at P_ϑ by (2.5) and (2.6):

$$\frac{\partial H_0}{\partial r}(P_\vartheta) = |P_\vartheta|^{-1}.$$

Now the construction of $B_1(0, H_0)$ comes into play: since the origin O and the point P_ϑ belong to the circumference of radius $\frac{1}{2}$ centered at $(\frac{1}{2}, 0)$, by a classical theorem in Euclidean geometry we get that the line segment OP_ϑ makes an angle $\alpha = \vartheta/2$ with the x_1 -axis (see Figure 3), and therefore $|P_\vartheta| = \cos \alpha = \cos(2\vartheta)$. Finally, since the radial derivative is the projection of the gradient in the radial direction, we have


$$\frac{\partial H_0}{\partial r}(P_\vartheta) = |DH_0(P_\vartheta)| \cos \alpha,$$

and hence

$$|DH_0(P_\vartheta)| = \frac{1}{\cos \alpha} \frac{\partial H_0}{\partial r}(P_\vartheta) = \frac{2}{1 + \cos \vartheta}.$$

Thus, the components $\xi_1(\vartheta), \xi_2(\vartheta)$ of v_ϑ are given by

$$\begin{cases} \xi_1(\vartheta) = \frac{2 \cos \vartheta}{1 + \cos \vartheta}; \\ \xi_2(\vartheta) = \frac{2 \sin \vartheta}{1 + \cos \vartheta}. \end{cases}$$

Figure 3: Finding the Euclidean norm of $DH_0(P_\vartheta)$

The parametric equations given above describe the parabola whose Cartesian equation is $\xi_1 = 1 - \frac{1}{4} \xi_2^2$, which passes through the points $(0, \pm 2(\sqrt{2} - 1))$ and has focus at the origin and vertex at $(1, 0)$. The remaining parts of $\partial B_1(0, H)$ are managed similarly, and the lemma follows. \square

Acknowledgments

The authors are members of the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM). G. Ciraolo has been partially supported by the PRIN 2017 project “Qualitative and quantitative aspects of nonlinear PDEs”. A. Greco is partially supported by the research project *Evolutionary and stationary Partial Differential Equations with a focus on bio-mathematics*, funded by [Fondazione di Sardegna](#) (2019).

References

- [1] C. Bianchini, G. Ciraolo. *Wulff shape characterizations in overdetermined anisotropic elliptic problems*. Comm. Partial Differential Equations **43** (2018), 790–820.
- [2] C. Bianchini, G. Ciraolo, P. Salani. *An overdetermined problem for the anisotropic capacity*. Calc. Var. **55** (2016), Paper No. 84, 24 pp.
- [3] G. Bellettini, M. Paolini. *Anisotropic motion by mean curvature in the context of Finsler geometry*. Hokkaido Math. J. **25** (1996), 537–566.
- [4] H. Brézis. *Functional Analysis, Sobolev Spaces and Partial Differential Equations*. Springer, 2011.
- [5] X. Cabré, X. Ros-Oton, J. Serra. *Sharp isoperimetric inequalities via the ABP method*. J. Eur. Math. Soc. (JEMS) **18** (2016), 2971–2998.
- [6] P. Cannarsa, C. Sinestrari. *Semiconcave functions, Hamilton-Jacobi equations, and optimal control*. Progress in Nonlinear Differential Equations and their Applications **58**. Birkhäuser, 2004.

- [7] D. Castorina, G. Riey, B. Sciunzi. *Hopf Lemma and regularity results for quasilinear anisotropic elliptic equations*. Calc. Var. Partial Differ. Equ. **58** (2019), Paper No. 95, 18 pp.
- [8] A. Cianchi, V. Maz'ya. *Global Lipschitz Regularity for a Class of Quasilinear Elliptic Equations*. Commun. Partial Differ. Equations **36** (2011), 100–133.
- [9] A. Cianchi, P. Salani. *Overdetermined anisotropic elliptic problems*. Math. Ann. **345** (2009), 859–881.
- [10] G. Ciraolo, A. Figalli, A. Roncoroni. *Symmetry results for critical anisotropic p -Laplacian equations in convex cones*. Geom. Funct. Anal., **30** (2020), 770–803.
- [11] G. Ciraolo, R. Magnanini, S. Sakaguchi. *Symmetry of minimizers with a level surface parallel to the boundary*. J. Eur. Math. Soc. (JEMS) **17** (2015), 2789–2804.
- [12] G. Ciraolo, A. Roncoroni. *Serrin's type overdetermined problems in convex cones*. Calc. Var. Partial Differential Equations **59** (2020), Paper No. 28, 21 pp.
- [13] M. Cozzi, A. Farina, E. Valdinoci. *Monotonicity formulae and classification results for singular, degenerate, anisotropic PDEs*. Adv. Math. **293** (2016), 343–381.
- [14] G. Crasta. *A symmetry problem in the calculus of variations*. J. Eur. Math. Soc. (JEMS) **8** (2006), 139–154.
- [15] G. Crasta, A. Malusa. *The distance function from the boundary in a Minkowski space*. Trans. Amer. Math. Soc. **359** (2007), 5725–5759.
- [16] A. Farina, B. Kawohl. *Remarks on an overdetermined boundary value problem*. Calc. Var. Partial Differential Equations **31** (2008), 351–357.
- [17] A. Farina, E. Valdinoci. *Partially and globally overdetermined problems of elliptic type*. Adv. Nonlinear Anal. **1** (2012), 27–45.
- [18] A. Farina, E. Valdinoci. *On partially and globally overdetermined problems of elliptic type*. Amer. J. Math. **135** (2013), 1699–1726.
- [19] E. Ferone, B. Kawohl. *Remarks on a Finsler-Laplacian*. Proc. Amer. Math. Soc. **137** (2009), 247–253.
- [20] P. Finsler. *Ueber Kurven und Flächen in allgemeinen Räumen*. Dissertation. Georg-August-Universität zu Göttingen, 1918.
- [21] I. Fragalà, F. Gazzola. *Partially overdetermined elliptic boundary value problems*. J. Differ. Equations **245** (2008), 1299–1322.
- [22] I. Fragalà, F. Gazzola, B. Kawohl. *Overdetermined problems with possibly degenerate ellipticity, a geometric approach*. Math. Z. **254** (2006), 117–132.
- [23] I. Fragalà, F. Gazzola, J. Lamboley, M. Pierre. *Counterexamples to symmetry for partially overdetermined elliptic problems*. Analysis **29** (2009), 85–93.
- [24] N. Garofalo, J.L. Lewis. *A symmetry result related to some overdetermined boundary value problems*. Amer. J. Math. **111** (1989), 9–33.

- [25] A. Greco. *Comparison principle and constrained radial symmetry for the subdiffusive p -Laplacian*. Publ. Mat. **58** (2014), 485–498.
- [26] A. Greco. *Symmetry around the origin for some overdetermined problems*. Adv. Math. Sci. Appl. **13** (2003), 387–399.
- [27] A. Greco and V. Mascia. *Non-local sublinear problems: existence, comparison, and radial symmetry*. Discrete Contin. Dyn. Syst. **39** (2019), 503–519.
- [28] S. G. Krantz, H. R. Parks. *The geometry of domains in space*. Birkäuser, 1999.
- [29] G. Lieberman. *Boundary regularity for solutions of degenerate elliptic equations*. Nonlinear Anal. **12** (1988), 1203–1219.
- [30] P.-L. Lions, F. Pacella. *Isoperimetric inequalities for convex cones*. Proc. Amer. Math. Soc. **109** (1990), 477–485.
- [31] P.-L. Lions, F. Pacella, M. Tricarico. *Best constants in Sobolev inequalities for functions vanishing on some part of the boundary and related questions*. Indiana Univ. Math. J. **37** (1998), 301–324.
- [32] F. Pacella, G. Tralli. *Overdetermined problems and constant mean curvature surfaces in cones*. Rev. Mat. Iberoam. **36** (2020), 841–867.
- [33] R. T. Rockafellar. *Convex Analysis*. Princeton University Press, 1970.
- [34] A. Roncoroni, *A symmetry result for the φ -Laplacian in model manifolds*. In preparation.
- [35] S. Salsa. *Partial Differential Equations in Action. From Modelling to Theory*, 3rd ed. Springer International Publishing, 2016.
- [36] R. Schneider. *Convex Bodies: The Brunn-Minkowski Theory. Encyclopedia of Mathematics and its Applications* **44**. Cambridge University Press, 1993.
- [37] J. Serrin. *A symmetry problem in potential theory*. Arch. Rational Mech. Anal. **43** (1971), 304–318.
- [38] B. Simon. *A Comprehensive Course in Analysis, Part 1*. AMS, 2015.
- [39] P. Tolksdorf. *Regularity for a more general class of quasilinear elliptic equations*. J. Differential Equations **51** (1984), 126–150.
- [40] G. Wang, C. Xia. *A characterization of the Wulff shape by an overdetermined anisotropic PDE*. Arch. Rational Mech. Anal. **199** (2011) 99–115.
- [41] G. Wulff. *Zur Frage der Geschwindigkeit des Wachstums und der Auflösung der Kristallflächen*. Zeitschrift für Kristallographie - Crystalline Materials **34** (1901), 449–530.

Giulio Ciraolo
 Department of Mathematics “Federigo Enriques”
 Università degli Studi di Milano
 Italy
 e-mail: giulio.ciraolo@unimi.it

Antonio Greco
 Department of Mathematics and Computer Science
 Università degli Studi di Cagliari
 Italy
 e-mail: greco@unica.it