



ELSEVIER



Earth observation for drought risk financing in pastoral systems of sub-Saharan Africa

Francesco Fava¹ and Anton Vrieling²

As climate-related crises increase globally, climate risk financing is becoming an integral part of financial protection and resilience building strategies of African countries. Drought-induced crises result in devastating human impacts and high costs for vulnerable countries, threatening longer-term investments and development efforts. While earth observation (EO) has been widely used for drought early warning, new opportunities emerge from integrating EO data and methods into index-based drought risk financing (IBDRF) instruments. Such instruments aim at supporting an effective and timely response during drought shocks and improving the resilience of small-holder farmers and livestock keepers. This review documents the current status, and discusses future prospects and potential challenges for EO utilization in IBDRF applications in sub-Saharan Africa. We focus on pastoral systems, which are hotspots in terms of vulnerability to climate and environmental change, food insecurity, poverty, and conflicts. In these systems, EO-based IBDRF interventions are rapidly scaling up as part of national and international risk management strategies.

Addresses

¹ International Livestock Research Institute (ILRI), P.O. Box 30709, Nairobi 00100, Kenya

² University of Twente, Faculty of Geo-information Science and Earth Observation, P.O. Box 217, 7500 AE Enschede, The Netherlands

Corresponding author: Fava, Francesco (f.fava@cgiar.org)

Current Opinion in Environmental Sustainability 2021, **48**:44–52

This review comes from a themed issue on **The dryland social-ecological systems in changing environments**

Edited by **Bojie Fu**, **Mark Stafford-Smith** and **Chao Fu**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 18th October 2020

Received: 01 June 2020; Accepted: 14 September 2020

<https://doi.org/10.1016/j.cosust.2020.09.006>

1877-3435/ 2020 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Introduction

Crisis risk financing refers to mechanisms that aim at reducing adverse socio-economic or ecological impacts of potential crises [1^{*}]. This can include paying to prevent

and reduce the risk, or to prepare for and respond to a shock. Climate risk financing (CRF) targets climate-related shocks (e.g. drought, floods and heat waves) and is becoming an integral part of climate risk management frameworks as key components of financial protection strategic planning for low and middle income countries [2]. Multiple CRF approaches exist, including market-based instruments (e.g. insurance schemes, catastrophic bonds and swaps), contingent financing (e.g. credit), or budgetary tools (i.e. dedicated reserve funds or contingency budgets). These approaches are all designed to increase financial resilience to climate-related crises, linking the response actions to pre-defined mechanisms for timely release of financial resources. In this way, they aim at ensuring rapid and cost-effective preparation, assistance, recovery or reconstruction efforts.

Droughts, defined as periods with water deficit relative to normal conditions, are one of the most disrupting natural disasters, each year affecting millions of people worldwide with devastating impacts [3,4]. Severe droughts cause massive disruptions to national economies and dramatic impacts on the livelihood and food security of small-holder farmers and livestock keepers. Standard responses to drought in African countries, such as humanitarian support in the form of cash or food transfers, are important instruments to support drought-affected vulnerable populations. However, these responses have proven to be often too slow, cost-ineffective, and to foster dependency rather than resilience, especially when not integrated into holistic risk management strategies [2].

Among the different CRF instruments, index-based approaches have gained considerable traction over the last two decades, particularly for targeting drought shock impacts on African small-holder farming systems. Index-based drought risk financing (IBDRF) uses trigger mechanisms that rely on a transparent and objectively measured indicator of drought (i.e. the index). The underlying index must be highly correlated with drought-related economic losses to be useful in tracking, and, therefore, transferring, the risk. Payouts are made when the index values fall below a pre-defined threshold, normally derived from historic index realizations. When compared to *ex-post* loss verification (e.g. traditional insurance), IBDRF limits information asymmetry issues, such as adverse selection and moral hazards,³ reduces transaction and verification costs, and

³ This terminology originates from insurance literature: adverse selection occurs when potential policyholders make decisions based on information about their risk exposure that is not available to the insurance provider. Moral hazard occurs when the policyholders engage in activities that increase their exposure to risk, leaving the insurer exposed to higher risk than had been assessed for premium rate determination.

enables more effective and timely distribution of payouts. However, these advantages come at the cost of the inherently imperfect correlation between index and actual loss, also known as basis risk [5].

Pastoral production systems dominate the African drylands, cover about 43% of Africa's land mass, and are the main livelihood for about 268 million people in these areas [6]. Drought is a distinctive feature and over millennia the pastoral livelihood developed to deal with this [7]. However, particularly in SSA, the combination of increasing rainfall variability [8] and other pressures (e.g. changes in land use and tenure, population growth, rangeland degradation [9]) is weakening the resilience of pastoral communities and the effectiveness of traditional drought coping mechanisms (e.g. mobility, restocking). Severe droughts can lead to catastrophic herd losses in pastoral regions, and as such cause food insecurity and poverty trap dynamics, and dramatically reduce national GDPs of SSA countries [5]. Consequently, the development and implementation of drought risk management strategies for pastoral regions, including IBDRF initiatives, is a key component of the policy agendas of development institutions and national governments.

The scarcity and poor quality of ground data and national statistics in SSA pastoral regions [10] has made in most cases Earth observation (EO) by satellites the only viable option for designing IBDRF instruments for large-scale implementation. The growing availability and increased quality of long-term EO datasets of rainfall products and vegetation indices [11*] have been instrumental in designing indices that are thought to reliably represent drought risks. Furthermore, the close interlinkage between rangeland condition and impacts on productive assets (i.e. livestock) and livelihoods has facilitated the design of financial triggering mechanisms that link impact to payouts. Therefore, datasets obtained through multi-temporal satellite imagery are currently a key component of IBDRF initiatives in pastoral regions.

However, whilst the scientific literature on drought monitoring is vast, only a few studies reviewed the contribution of EO to insurance [12,13*], and currently no overview exists of the IBDRF initiatives, challenges, and prospects in sub-Saharan Africa.

Evolution of IBDRF in pastoral Africa

During the last decade IBDRF initiatives in African pastoral drylands have gained significant momentum, also thanks to large international initiatives such as the InsuResilience Investment Fund [14] and the Global Index-Insurance Facility (GIIF). While for several years IBDRF schemes have largely remained at the pilot level, constrained by the limited demand for retail micro-insurance products [15,16], these schemes are now gradually covering larger areas, given their growing integration in

country-wide social protection and drought risk management programs. This caused a significant increase in volume of financial transactions and a growing participation of international re-insurance companies, which facilitates risk transfer toward international financial markets, and promotes larger investments by governments and international organizations.

Table 1 summarizes operational IBDRF initiatives in SSA. These initiatives, initially launched as retail micro-level index-insurance schemes (IBLI), more recently have expanded their scope and modality of implementation, including fully subsidized insurance programs targeting vulnerable pastoralists (KLIP, SIIPE), sovereign-level insurance schemes (ARC), and scalability mechanisms of shock responsive safety nets programs (HSNP, NUSAF). At the same time, IBDRF product design evolved from indices designed to assess an observed loss (i.e. the IBLI livestock mortality scheme), thus triggering when drought is already impacting pastoralist assets, to indices designed to identify deteriorating forage condition, thus triggering at the earlier drought stages, with the aim of supporting pastoralists to protect their assets (i.e. livestock or livelihood) or countries implementing early response actions.

While fully operational initiatives in pastoral areas are concentrated largely in East Africa and, to a lesser extent, in the Sahel, several insurance providers and organizations are launching similar IBDRF solutions on a retail basis across SSA, including in Niger, Zambia and South Africa. In addition, feasibility studies are being conducted in Somalia, Senegal, Burkina Faso, and Mali. Thus lessons gathered from early implementation efforts are of key relevance for sustainable scaling of IBDRF in SSA.

EO-based index design in operational IBDRF schemes

Similar to most drought early warning systems [e.g. Refs. 17*,18,19], existing IBDRF schemes for pastoral systems predominantly use vegetation indices or rainfall estimates (RFE) as input (**Table 1**). These EO products should meet the fundamental operational requirements for IBDRF, which include 1) full transparency and accessibility of the source data, 2) availability of historical records for financial risk modelling and pricing of ideally 20 years or more, 3) near real-time availability, and 4) an expected remaining lifetime of at least a few years. Satellite-derived RFE can provide a direct indication of meteorological drought, which largely determines water availability in rangelands. While availability of drinking water is important for livestock health, monitoring this for large areas is cumbersome, given that much water for livestock comes from wells or small water bodies that cannot easily be monitored with EO data. RFEs are instead typically used in IBDRF to estimate the available water for vegetation, for example using simple water-balance models.

Table 1

Summary of operational IBDRF programs targeting pastoral drylands, ordered chronologically with insurance programs listed first and non-insurance programs at the bottom

Program ^a	Years	Scope	Satellite ^b sensor/indicator	Index ^c	Area	Households (nr)
IBLI	2010–2015	Retail micro-insurance for drought related livestock mortality	AVHRR NDVI	Livestock mortality	Marsabit (Kenya), Borana (Ethiopia)	5983
IBLI	2015 – present	Retail micro-insurance for asset protection	MODIS NDVI	z-score seasonal NDVI	Northern Kenya and Borana (Ethiopia)	7000
KLIP	2015 – present	Subsidized insurance for asset protection	MODIS NDVI	z-score seasonal NDVI	Northern and Eastern Kenya	18 000
SIPE	2017- present	Subsidized insurance for asset protection	MODIS NDVI	z-score seasonal NDVI	Somali region (Ethiopia)	5000
ARC	2017- present	Sovereign level insurance	RFE – multiple datasets	WRSI (RFE based)	East Africa and Sahel	N/a ^d
ARC	2019- present	Sovereign level insurance	MODIS NDVI	z-score NDVI or VCI	East Africa and Sahel	N/a ^d
HSNP	2015- present	Social protection scalability mechanism	MODIS NDVI	VCI running average	Northern Kenya	>100 000
NUSAF	2017- present	Social protection scalability mechanism	MODIS NDVI	NDVI percent anomaly	Karamoja (Uganda)	25 000

^a IBLI = Index-based Livestock Insurance, KLIP = Kenyan Livestock Insurance Program, SIPE = Satellite Index-Insurance for Pastoralists in Ethiopia, ARC = African Risk Capacity, HSNP = Hunger Safety Net Program, NUSAF = Northern Uganda Social Action Fund.

^b AVHRR = Advanced Very High Resolution Radiometer. MODIS = Moderate Resolution Imaging Spectroradiometer, NDVI = Normalized Difference Vegetation Index, RFE = Rainfall Estimates (using satellite data).

^c Z-score is sometimes referred to as standard score, WRSI = Water Requirement Satisfaction Index, VCI = Vegetation Condition Index.

^d As ARC is a sovereign-level insurance scheme, the recipient of the payout is the country and the number of households covered will depend on the country's contingency plans. ARC is offering insurance cover for rangelands in the entire Sahel and Horn of Africa and plans to offer it also in Central and Southern Africa. The NDVI product has been launched in Chad, Niger, Mali, Mauritania, and Kenya.

The water requirement satisfaction index (WRSI) used by ARC is such a model; it estimates evapotranspiration demands of vegetation and compares this with a simple 'bucket' model of the soil that gets filled with rainfall [20,21]. Two main drawbacks exist when using rainfall-based indices for large-scale pastoral IBDRF: 1) despite the growing availability of gridded rainfall products, the quality, accessibility, and density of rainfall station data in pastoral SSA is generally low [22,23], resulting in unknown or poor accuracies in these areas [24]; 2) the link between rainfall and the vegetation's water availability is complex, and depends on vegetation characteristics, soil, and rainfall distribution [25,26], which cannot be characterized sufficiently with 10-daily rainfall sums [27].

For extensive pastoral systems, forage availability is a key determinant of livestock health, as alternative feed resources are largely unavailable or inaccessible. To overcome drawbacks of RFEs, optical sensors onboard satellites can be used to measure the vegetation's reflectance. Dense healthy vegetation reflects much in near infrared (NIR), and little in red wavelengths, and spectral vegetation indices, like the normalized difference

vegetation index (NDVI), use this to monitor vegetation condition. For drought monitoring, NDVI images from coarse-resolution satellites (250 m and up) are generally used, because 1) long time series exist to describe long-term variability in forage conditions, 2) their daily data collections allow for more cloud-free observations to describe vegetation changes throughout the season, and 3) documented evidence exists for a strong relationship between rangeland biomass and NDVI [28,29]. Required pre-processing steps to effectively use NDVI for anomaly analysis include temporal compositing [transforming daily to, e.g. 10-day data, keeping the best-quality observation for each pixel; 30], and smoothing to further reduce atmospheric effects [31].

To transform NDVI or alternative drought indices into a useful index for pastoral IBDRF schemes, three steps are required:

- 1) Spatial aggregation; geographic units are normally larger than grid cells, both for operational reasons and to reflect that herds move. Aggregation within units generally incorporates a mask of where rangelands occur.

- 2) Temporal aggregation; most schemes aim to assess seasonal forage scarcity, requiring expert or EO-derived [32] knowledge on rainfall/vegetation seasonality.
- 3) Normalization to compare the current index value against historic index realizations in past years. Multiple options exist, such as for example z-scoring (subtract mean and divide by standard deviation), linear scaling between minimum and maximum historic values [i.e. the vegetation condition index, or VCI; 33], or percentile calculation.

The term ‘index design’ in IBDRF refers to: a) the choice of input data, b) the precise methods for performing the above-mentioned three steps, c) the sequencing of these steps, and d) the testing of the resulting index against drought-related loss estimates. Eventually, the choice for a design will at least partly depend on the IBDRF instrument’s purpose, satisfaction of stakeholders on historic and current index readings, and (scientific) background of the ‘designers’. Given that droughts normally affect large regions and are simultaneously characterized by reduced precipitation and vegetation growth, it is well possible that alternative designs provide similar index outcomes during main drought events.

Opportunities and challenges for EO contribution to IBDRF

Recent targeted research efforts for EO support to innovative design of IBDRF solutions, made in the framework of the programs listed in Table 1, helped to remove critical barriers for scaling IBDRF initiatives in pastoral drylands. Here, we discuss four emerging trends in IBDRF where the demand for innovative solutions from EO research is strong. While we argue that the quality of IBDRF product design differs and that EO-aided solutions can play an important role to improve operational programs, we also emphasize that each solution is context-specific and needs to consider the trade-off between a) timing and accuracy of the assessment, b) cost-effectiveness of the response action, and c) the complexity of the interaction between drought shocks and the socio-ecological pastoral systems (Figure 1) with the goal of mitigating impacts and speeding-up recovery in the short term, while building long-term social and environmental resilience.

Emerging EO data products for IBDRF

EO data products used for IBDRF index design in pastoral areas have evolved over time in response to stakeholder feedbacks and with emerging technologies. For example, the ARC product for extensive rangelands has recently transitioned from WRSI to NDVI (Table 1), meeting the demand of key stakeholders. However, besides vegetation index (e.g. NDVI) products that provide a direct measure of forage status, drought characteristics can be obtained, for example, from EO-derived

precipitation [35], soil moisture [36,37], or evapotranspiration [38] products [for reviews on EO drought monitoring options see also Refs. 11*,39]. A variety of these products have already been proposed and in some cases integrated into drought index-insurance pilot projects for crops in Africa, for example, through data service provision by commercial EO companies [e.g. Refs. 40–43]. These efforts provide promising alternative drought indices for innovative IBDRF design by addressing multiple phases of drought progression and thus merit further analysis.


IBDRF design could also take advantage of the continuous stream of free 10–30 m resolution EO data that are provided by satellites such as Sentinel-1, Sentinel-2, and Landsat-8. Given their high observation frequency, timely fine-scale estimates of seasonality and forage conditions can be provided [44], even if cloud cover remains a concern for short vegetation seasons [45]. Arguably pastoral IBDRF may not require fine-scale data because droughts generally affect large areas, but where drought impacts differ due to greater landscape variability (e.g. in agro-pastoral systems) they could prove beneficial. As historical archives of 10–30 m resolution data are building, and tools to analyze resulting large data amounts become commonplace [46], finer-scale drought analysis will likely find its way into IBDRF.

Remote sensing advances for improved product design

The spatial component of basis risk (i.e. impacts are not equally distributed within a geographic insurance unit) is a recurrent issue for IBDRF products in pastoral areas, as administrative units cannot fully reflect the agro-ecological variability and herd mobility patterns. This type of basis risk has been reported for some insurance units in Kenya and Ethiopia, especially at the fringe between agro-pastoral and extensive pastoral systems. To deal with this issue, administrative units could be replaced by more meaningful ecological units, for example, defined based on similar temporal behavior of NDVI [47]. In addition, high resolution EO data can help to improve rangeland mapping and characterization [48], allowing to better spatially focus coarser-resolution drought indices on the areas within insurance units that matter most for forage production.

Payout timing and temporal aggregation is another critical component of product design. Triggering early action for expected adverse events can make disaster response more cost-effective [49*,61]. Because drought is a slow-onset shock, indicators can be designed that detect early stages of the drought progression [11*]. EO has supported the anticipation of the response by effectively characterizing between-year variability of forage conditions earlier in the season through shortening temporal aggregation windows, with the aim of designing asset protection insurance products [49*,62,63]. This has been a fundamental

Figure 1



Ground and satellite EO applications can contribute to monitor drought progression and its ecological and socio-economic impacts to support context-specific design of IBDRF instruments. The upper part of the Figure illustrates the progression of drought impacts following Mishra and Singh [34]. While drought evolves from meteorological to socio-economic, IBDRF design needs to consider trade-offs to maximize the social-ecological benefits, while accounting for economic costs to ascertain long-term sustainability of the instruments. For example, an EO-based index designed with an early trigger is expected to be more accurate in detecting meteorological or agricultural drought, but might be less effective in assessing socio-economic impacts. This source of basis risk may (or may not) be counterbalanced by the savings in economic costs associated with asset protection and early response. These assumptions are context-specific and product quality assessment is thus fundamental to evaluate trade-offs.

step for the IBLI and KLIP programs to increase the uptake and gain political support toward geographic scaling. However, indices that allow for early triggering must also be backed by efficient payout delivery mechanisms to avoid negative impacts on program sustainability.

More recently, forecasting approaches based on time series analysis and machine learning techniques have been developed to predict time-integrated (seasonal) NDVI anomalies from lagged NDVI, rainfall, and climate indices [64–68]. While research on drought forecasting models is gaining momentum, particularly for early warning systems (e.g. HSNP, Table 1), the implementation of forecasts into IBDRF instruments is challenging given the forecasts’ large uncertainty at local scales and longer time lags [69]. Using forecasts as a triggering index could thus increase basis risk and lead to operational implementation challenges (e.g. in case of false alarm and unneeded response) [61,70]. Moreover, deciding on thresholds for response triggering could be complex because this relates not only to the level of the impact, but also to its probability. A different challenge emerges when forecasts are not directly used but could nonetheless affect IBDRF instruments such as insurance.

For example, since insurance premiums are based on historical realizations of the index, reasonably accurate forecasts made before insurance sales [possibly through indigenous indicators: 71] could lead to adverse selection if prospective purchasers have information about expected payouts that the product’s pricing does not account for [e.g. Ref. 15]. Overall, while anticipatory risk financing based on forecasts is a promising innovation in IBDRF, its operational implementation would require a careful assessment of associated risks and effective harmonization with other IBDRF instruments.

Quality assurance of IBDRF

Basis risk remains a critical concern for the quality and sustainability of IBDRF schemes. For operational initiatives a few recent studies compared results from multiple index designs [41,47,49*,50]. Notwithstanding, these studies highlighted that evaluation of the resulting indices remains complex given the scarcity of and/or difficulty to collect good-quality *in-situ* data on drought outcomes. While EO product development and improved index design have potential to reduce basis risk, ultimately it remains an empirical question whether this potential is

realized. This question goes beyond the traditional accuracy assessment approaches in the EO domain (e.g. an evaluation if soil moisture is accurately represented in a soil moisture product), as it should encompass also the socio-economic value of the proposed solution [49], and be formalized through quality metrics, minimum standards and robust assessment frameworks [1,51]. The overarching goal should be to design rigorous quantitative metrics and approaches to assess and compare the utility of IBDRF interventions.

A main challenge is collecting and analyzing relevant reference data on drought outcomes, given the limited formal data sets [52]. Potential data sources include multi-year forage biomass measurements [53], drought recall exercises [52], and repetitive household surveys on drought outcomes such as livestock mortality [49] or child nutrition [54]. Given the cost-intensive and labor-intensive nature of collecting such data, ground-based EO approaches can provide a useful addition. A good example is the repetitive observation of the same vegetation, either by permanent cameras [55,56], or through crowdsourcing platforms [57]. For crops, this has already led to the idea and implementation of ‘picture-based-insurance’ whereby farmers repeatedly collect pictures of their fields for verification of insurance claims [58]. This idea could be extended to rangeland and livestock conditions, possibly taking advantage of computer-vision based automation of body scoring techniques [59,60]. Increased efforts in reference data collection are urgently needed to answer the empirical question and provide quality insurance products.

Socio-economic and environmental impact evaluation

With the geographic expansion and growing number of households covered, the need for impact evaluation of IBDRF initiatives has increased. Traditionally impact evaluation of drought crises on small-holder farmers and livestock keepers largely focused on socio-economic factors through mixed qualitative-quantitative household-level surveys [72]. However, the high costs and complexity of such surveys have prevented a systematic integration of impact assessment studies in IBDRF initiatives, so that only few robust assessments are available in pastoral regions [e.g. Ref. 73]. In addition, environmental impacts have been poorly considered so far, while scaling IBDRF interventions to a large number of households might have relevant ecological impacts. For example, reduced herd losses may result in increased grazing pressure on rangelands [74,75], although empirical studies suggest on the contrary that insured pastoralists keep smaller herds because they reduce their use of livestock as precautionary savings [76].

EO approaches for rangeland health monitoring are well established [e.g. Ref. 77]. However, only few studies, focused on land restoration, have integrated these approaches into impact assessment of large programs in drylands [78,79]. Recent literature also showed potential

for assessing food security and household wealth [80,81,82], but pointed out that satellite EO capacity to directly monitor socio-economic indicators of household wellbeing and rapid land use dynamics during drought events (e.g. land accessibility, land tenure changes, migration) is limited. Impact evaluation could be improved by combining satellite EO information with *in-situ* collected environmental and social-economic data [83], which are increasingly available via mobile and crowdsourcing platforms also in rural Africa [e.g. Refs. 84,85,86]. This can be supported by advances in machine learning algorithms, allowing to extract patterns and understand spatial connections from diverse data sources [87]. Data driven approaches should be, however, used with caution and framed within a robust set of causal hypotheses, taking into account the cross-scale interactions between physical and socio-economic factors [81].

Conclusions

IBDRF initiatives are scaling up as part of the policy agendas of SSA countries and development organizations on resilience building, poverty reduction, and sustainable development. EO plays a key role in sustaining this trend in pastoral drylands, with the potential of significant societal and policy impact in the region. Innovations from ground and satellite EO technologies could contribute to design more accurate, cost-effective, and harmonized drought risk financing programs, as well as to assess their effectiveness during shocks and their longer-term impacts. However, this can only be achieved if the EO community does not limit its role to providing technological solutions and services, but rather becomes an integral part of the interdisciplinary framework to address drought risk management in complex socio-ecological systems, understanding synergies and trade-offs between research, operational implementation, and policy formulation.

Conflict of interest statement

Nothing declared.

Acknowledgements

The CGIAR Research program on Livestock and contributors to the CGIAR Trust Fund are acknowledged for supporting this research. Anton Vrieling was funded by the Dutch Research Council (NWO), Space for Global Development (WOTRO) programme, as part of the CGIAR-Netherlands partnership. We thank Nathaniel Jensen (ILRI) and James Hassell (Smithsonian GHP) for their valuable comments and inputs on the manuscript.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest

1. Poole L, Clarke D, Swithern S: *The Future of Crisis Financing: A Call to Action*. London, UK: Centre for Disaster Protection; 2020 <https://www.disasterprotection.org/crisisfinance>.

The report provides an overview of crisis financing and a reference for terminology, setting out a new vision for effective crisis financing packages and instruments.

2. The World Bank: *Disaster Risk Finance as a Tool for Development: A Summary of Findings from the Disaster Risk Finance Impact Analytics Project*. Washington DC: The World Bank; 2016 <https://openknowledge.worldbank.org/handle/10986/24374>.
3. Carrão H, Naumann G, Barbosa P: **Mapping global patterns of drought risk: an empirical framework based on sub-national estimates of hazard, exposure and vulnerability**. *Global Environ Change* 2016, **39**:108-124.
4. Meza I, Siebert S, Döll P, Kusche J, Herbert C, Rezaei EE, Nouri H, Gerdener H, Popat E, Frischen J *et al.*: **Global-scale drought risk assessment for agricultural systems**. *Nat Hazards Earth Syst Sci* 2020, **20**:695-712.
5. Barrett CB, Barnett BJ, Carter MR, Chantarat S, Hansen JW, Mude AG, Osgood D, Skees JR, Turvey CG, Ward MN: *Poverty Traps and Climate Risk: Limitations and Opportunities of Index-Based Risk Financing IRI Technical Report No. 07-02*. 2007.
6. FAO: *Pastoralism in Africa's Drylands; Reducing Risks, Addressing Vulnerability and Enhancing Resilience*. Rome, Italy: Food and Agriculture Organization of the United Nations; 2018.
7. Butt B, Shortridge A, WinklerPrins AMGA: **Pastoral herd management, drought coping strategies, and cattle mobility in Southern Kenya**. *Ann Assoc Am Geogr* 2009, **99**:309-334.
8. Sloat LL, Gerber JS, Samberg LH, Smith WK, Herrero M, Ferreira LG, Godde CM, West PC: **Increasing importance of precipitation variability on global livestock grazing lands**. *Nat Climate Change* 2018, **8**:214-218.
9. Haller T, Van Dijk H, Bollig M, Greiner C, Schareika N, Gabbert C: **Conflicts, security and marginalisation: institutional change of the pastoral commons in a 'glocal' world**. *OIE Revue Scientifique et Technique* 2016, **35**:405-416.
10. Zezza A, Federighi G, Kalilou AA, Hiernaux P: **Milking the data: measuring milk off-take in extensive livestock systems. Experimental evidence from Niger**. *Food Policy* 2016, **59**:174-186.
11. West H, Quinn N, Horswell M: **Remote sensing for drought monitoring & impact assessment: progress, past challenges and future opportunities**. *Remote Sens Environ* 2019, **232** 111291.
- A comprehensive review summarizing the latest developments of remote sensing technologies for drought monitoring and assessment, with a clear vision toward the future role of new missions, sensor technologies, algorithms and analytics.
12. de Leeuw J, Vrieling A, Shee A, Atzberger C, Hadgu K, Biradar C, Keah H, Turvey C: **The potential and uptake of remote sensing in insurance: a review**. *Remote Sens* 2014, **6**:10888-10912.
13. Vroege W, Dalhaus T, Finger R: **Index insurances for grasslands – a review for Europe and North-America**. *Agric Syst* 2019, **168**:101-111.
- This reviews provides an overview of emerging trends in index-insurance for grasslands in Europe and North America, setting the stage for future opportunities.
14. InsuResilience: Joint Statement on the InsuResilience Global Partnership. 14 November 2017. Bonn <https://www.insuresilience.org/wp-content/uploads/2017/11/20171205-Joint-Statement-InsuResilience-Global-Partnership.pdf>. Accessed: 29 May 2020.
15. Jensen ND, Mude AG, Barrett CB: **How basis risk and spatiotemporal adverse selection influence demand for index insurance: evidence from northern Kenya**. *Food Policy* 2018, **74**:172-198.
16. Takahashi K, Ikegami M, Sheahan M, Barrett CB: **Experimental evidence on the drivers of index-based livestock insurance demand in southern Ethiopia**. *World Dev* 2016, **78**:324-340.
17. Rembold F, Meroni M, Urbano F, Csak G, Kerdeles H, Perez-Hoyos A, Lemoine G, Leo O, Negre T: **ASAP: a new global early warning system to detect anomaly hot spots of agricultural production for food security analysis**. *Agric Syst* 2019, **168**:247-257.
- This study describes a new global early warning system, which provides timely warnings through a user-friendly public online webGIS. State-of-the-art EO methods are used to derive agricultural production deficit warning categories, which are integrated by comprehensive expert analyses.
18. Fritz S, See L, Bayas JCL, Waldner F, Jacques D, Becker-Reshef I, Whitcraft A, Baruth B, Bonifacio R, Crutchfield J *et al.*: **A comparison of global agricultural monitoring systems and current gaps**. *Agric Syst* 2019, **168**:258-272.
19. Becker-Reshef I, Justice C, Barker B, Humber M, Rembold F, Bonifacio R, Zappacosta M, Budde M, Magadzire T, Shitote C *et al.*: **Strengthening agricultural decisions in countries at risk of food insecurity: the GEOGLAM crop monitor for early warning**. *Remote Sens Environ* 2020, **237**.
20. Verdin J, Klaver R: **Grid-cell-based crop water accounting for the famine early warning system**. *Hydrol Process* 2002, **16**:1617-1630.
21. Tarnavsky E, Chavez E, Boogaard H: **Agro-meteorological risks to maize production in Tanzania: sensitivity of an adapted Water Requirements Satisfaction Index (WRSI) model to rainfall**. *Int J Appl Earth Observ Geoinf* 2018, **73**:77-87.
22. Nicholson SE, Funk C, Fink AH: **Rainfall over the African continent from the 19th through the 21st century**. *Global Planetary Change* 2018, **165**:114-127.
23. Maidment RI, Grimes D, Black E, Tarnavsky E, Young M, Greatrex H, Allan RP, Stein T, Nkonde E, Senkunda S *et al.*: **A new, long-term daily satellite-based rainfall dataset for operational monitoring in Africa**. *Sci Data* 2017, **4** 170063.
24. Nicholson SE, Klotter D, Zhou L, Hua W: **Validation of satellite precipitation estimates over the Congo Basin**. *J Hydrometeorol* 2019, **20**:631-656.
25. Wilcox KR, Shi Z, Gherardi LA, Lemoine NP, Koerner SE, Hoover DL, Bork E, Byrne KM, Cahill J Jr, Collins SL *et al.*: **Asymmetric responses of primary productivity to precipitation extremes: a synthesis of grassland precipitation manipulation experiments**. *Global Change Biol* 2017, **23**:4376-4385.
26. Zeppel MJB, Wilks JV, Lewis JD: **Impacts of extreme precipitation and seasonal changes in precipitation on plants**. *Biogeosciences* 2014, **11**:3083-3093.
27. Kimani MW, Hoedjes JCB, Su Z: **An assessment of satellite-derived rainfall products relative to ground observations over East Africa**. *Remote Sens* 2017, **9**:430.
28. Schucknecht A, Meroni M, Kayitakire F, Boureima A: **Phenology-based biomass estimation to support rangeland management in semi-arid environments**. *Remote Sens* 2017, **9**.
29. Dinga MNV, Tsubo M: **Improved assessment of pasture availability in semi-arid grassland of South Africa**. *Environ Monit Assess* 2019, **191**.
30. Fensholt R, Anyamba A, Stisen S, Sandholt I, Pak E, Small J: **Comparisons of compositing period length for vegetation index data from polar-orbiting and geostationary satellites for the cloud-prone region of West Africa**. *Photogramm Eng Remote Sens* 2007, **73**:297-309.
31. Cai Z, Jönsson P, Jin H, Eklundh L: **Performance of smoothing methods for reconstructing NDVI time-series and estimating vegetation phenology from MODIS data**. *Remote Sens* 2017, **9**.
32. Zeng L, Wardlow BD, Xiang D, Hu S, Li D: **A review of vegetation phenological metrics extraction using time-series, multispectral satellite data**. *Remote Sens Environ* 2020, **237** 111511.
33. Klisch A, Atzberger C: **Operational drought monitoring in Kenya using MODIS NDVI time series**. *Remote Sens* 2016, **8**:267.
34. Mishra AK, Singh VP: **A review of drought concepts**. *J Hydrol* 2010, **391**:204-216.
35. Sun Q, Miao C, Duan Q, Ashouri H, Sorooshian S, Hsu KL: **A review of global precipitation data sets: data sources, estimation, and intercomparisons**. *Rev Geophys* 2018, **56**:79-107.
36. Babaeian E, Sadeghi M, Jones SB, Montzka C, Vereecken H, Tuller M: **Ground, proximal, and satellite remote sensing of soil moisture**. *Rev Geophys* 2019, **57**:530-616.

37. Dorigo W, Wagner W, Albergel C, Albrecht F, Balsamo G, Brocca L, Chung D, Ertl M, Forkel M, Gruber A *et al.*: **ESA CCI soil moisture for improved Earth system understanding: state-of-the-art and future directions**. *Remote Sens Environ* 2017, **203**:185-215.
38. Liou YA, Kar SK: **Evapotranspiration estimation with remote sensing and various surface energy balance algorithms-a review**. *Energies* 2014, **7**:2821-2849.
39. AghaKouchak A, Farahmand A, Melton FS, Teixeira J, Anderson MC, Wardlow BD, Hain CR: **Remote sensing of drought: progress, challenges and opportunities**. *Rev Geophys* 2015, **53**:452-480.
40. Rosema A, van Huystee J, Foppes S, van der Woerd J, Klaassen E, Barendse J, Asseldonk MV, Dubreuil M, Régent S, Weber S *et al.*: **FESA Micro-insurance: Crop Insurance Reaching Every Farmer in Africa**. Delft, The Netherlands: EARS Earth Environment Monitoring; 2014.
41. IFAD: **Remote Sensing for Index Insurance; Findings and Lessons Learned for Smallholder Agriculture**. Rome, Italy: International Fund of Agricultural Development; 2017.
42. Enenkel M, Farah C, Hain C, White A, Anderson M, You L, Wagner W, Osgood D: **What rainfall does not tell us; enhancing financial instruments with satellite-derived soil moisture and evaporative stress**. *Remote Sens* 2018, **10**:1819.
43. VanderSat How VanderSat is helping smallholder farmers with satellites. <https://www.vandersat.com/blog/how-vandersat-is-helping-smallholder-farmers-with-satellites>: Accessed: 28 May 2020.
44. Bolton DK, Gray JM, Melaas EK, Moon M, Eklundh L, Friedl MA: **Continental-scale land surface phenology from harmonized Landsat 8 and Sentinel-2 imagery**. *Remote Sens Environ* 2020, **240** 111685.
45. Cheng Y, Vrieling A, Fava F, Meroni M, Marshall M, Gachoki S: **Phenology of short vegetation cycles in a Kenyan rangeland from PlanetScope and Sentinel-2**. *Remote Sens Environ* 2020, **248** 112004.
46. Gorelick N, Hancher M, Dixon M, Ilyushchenko S, Thau D, Moore R: **Google Earth Engine: planetary-scale geospatial analysis for everyone**. *Remote Sens Environ* 2017, **202**:18-27.
47. De Oto L, Vrieling A, Fava F, de Bie K: **Exploring improvements to the design of an operational seasonal forage scarcity index from NDVI time series for livestock insurance in East Africa**. *Int J Appl Earth Observ Geoinf* 2019, **82** 101885.
48. Zhang W, Brandt M, Wang Q, Prishchepov AV, Tucker CJ, Li Y, Lyu H, Fensholt R: **From woody cover to woody canopies: how Sentinel-1 and Sentinel-2 data advance the mapping of woody plants in savannas**. *Remote Sens Environ* 2019, **234** 111465.
49. Jensen N, Stoeffler Q, Fava F, Vrieling A, Atzberger C, Meroni M, Mude A, Carter M: **Does the design matter? Comparing satellite-based indices for insuring pastoralists against drought**. *Ecol Econ* 2019, **162**:59-73.
- A quality assessment of multiple remote sensing-based product design options for index-based livestock insurance is conducted using a utility framework. This study bridges the remote sensing and development economics literature.
50. Roumigué A, Sigel G, Poilvé H, Bouchard B, Vrieling A, Jacquin A: **Insuring forage through satellites: testing alternative indices against grassland production estimates for France**. *Int J Remote Sens* 2017, **38**:1912-1939.
51. Carter M, Chiu T: **A Minimum Standard to Ensure Index Insurance Contracts Do No Harm**. Davis, California: Feed the Future Innovation Lab for Assets and Market Access; 2018.
52. Osgood D, Powell B, Diro R, Farah C, Enenkel M, Brown M, Husak G, Blakeley S, Hoffman L, McCarty J: **Farmer perception, recollection, and remote sensing in weather index insurance: an Ethiopia case study**. *Remote Sens* 2018, **10**:1887.
53. Roumigué A, Jacquin A, Sigel G, Poilvé H, Lepoivre B, Hagolle O: **Development of an index-based insurance product: validation of a forage production index derived from medium spatial resolution fCover time series**. *GIScience Remote Sens* 2015, **52**:94-113.
54. Mude AG, Barrett CB, McPeak JG, Kaitho R, Kristjanson P: **Empirical forecasting of slow-onset disasters for improved emergency response: an application to Kenya's arid north**. *Food Policy* 2009, **34**:329-339.
55. Inoue T, Nagai S, Kobayashi H, Koizumi H: **Utilization of ground-based digital photography for the evaluation of seasonal changes in the aboveground green biomass and foliage phenology in a grassland ecosystem**. *Ecol Inf* 2015, **25**:1-9.
56. Browning DM, Snyder KA, Herrick JE: **Plant phenology: taking the pulse of rangelands**. *Rangelands* 2019, **41**:129-134.
57. Jensen N, Toth R, Xue Y, Bernstein R, Chebelyon EK, Mude AG, Barrett C, Gomes C: **Don't follow the crowd: incentives for directed spatial sampling**. *2017 Annual Meeting; Chicago, Illinois: Agricultural and Applied Economics Association* 2017.
58. Ceballos F, Kramer B, Robles M: **The feasibility of picture-based insurance (PBI): Smartphone pictures for affordable crop insurance**. *Dev Eng* 2019, **4**.
59. Liu D, He D, Norton T: **Automatic estimation of dairy cattle body condition score from depth image using ensemble model**. *Biosyst Eng* 2020, **194**:16-27.
60. Rodríguez Alvarez J, Arroqui M, Mangudo P, Toloza J, Jatip D, Rodríguez JM, Teyseyre A, Sanz C, Zunino A, Machado C *et al.*: **Body condition estimation on cows from depth images using Convolutional Neural Networks**. *Comput Electron Agric* 2018, **155**:12-22.
61. Guimarães Nobre G, Davenport F, Bischiniotis K, Veldkamp T, Jongman B, Funk CC, Husak G, Ward PJ, Aerts JCJH: **Financing agricultural drought risk through ex-ante cash transfers**. *Sci Total Environ* 2019, **653**:523-535.
62. Vrieling A, Meroni M, Mude AG, Chantarat S, Ummenhofer CC, de Bie CAJM: **Early assessment of seasonal forage availability for mitigating the impact of drought on East African pastoralists**. *Remote Sens Environ* 2016, **174**:44-55.
63. Meroni M, Fasbender D, Kayitakire F, Pini G, Rembold F, Urbano F, Verstraete M: **Early detection of production deficit hot-spots in semi-arid environment using FAPAR time series and a probabilistic approach**. *Remote Sens Environ* 2014, **142**:57-68.
64. Adede C, Oboko R, Wagacha PW, Atzberger C: **A mixed model approach to vegetation condition prediction using artificial neural networks (ANN): case of Kenya's operational drought monitoring**. *Remote Sens* 2019, **11**:1099.
65. Adede C, Oboko R, Wagacha PW, Atzberger C: **Model ensembles of artificial neural networks and support vector regression for improved accuracy in the prediction of vegetation conditions and droughts in four northern Kenya counties**. *ISPRS Int J Geo-Inf* 2019, **8**.
66. Matere J, Simpkin P, Angerer J, Olesambu E, Ramasamy S, Fasina F: **Predictive Livestock Early Warning System (PLEWS): monitoring forage condition and implications for animal production in Kenya**. *Weather Climate Extremes* 2020, **27** 100209.
67. Barrett AB, Duivenvoorden S, Salakpi E, Muthoka JM, Mwangi J, Oliver S, Rowhani P: **Forecasting vegetation condition for drought early warning systems in pastoral communities in Kenya**. *Remote Sens Environ* 2020. accepted.
68. Meroni M, Rembold F, Fasbender D, Vrieling A: **Evaluation of the Standardized Precipitation Index as an early predictor of seasonal vegetation production anomalies in the Sahel**. *Remote Sens Lett* 2017, **8**:301-310.
69. Hao Z, Singh VP, Xia Y: **Seasonal drought prediction: advances, challenges, and future prospects**. *Rev Geophys* 2018, **56**:108-141.
70. Lopez A, Coughlan de Perez E, Bazo J, Suarez P, van den Hurk B, van Aalst M: **Bridging forecast verification and humanitarian decisions: a valuation approach for setting up action-oriented early warnings**. *Weather Climate Extremes* 2020, **27**.

71. Andersson L, Wilk J, Graham LP, Wikner J, Mokwatlo S, Petja B: **Local early warning systems for drought – could they add value to nationally disseminated seasonal climate forecasts?** *Weather Climate Extremes* 2020, **28**.
72. Levine S, Gray B: *How to assess the impact of a Drought Risk Financing facility: a guide: Start Network*. 2017.
73. Banerjee R, Hall A, Mude A, Wandera B, Kelly J: **Emerging research practice for impact in the CGIAR: the case of Index-Based Livestock Insurance (IBLI)**. *Outlook Agric* 2019, **48**:255-267.
- The Index-Based Livestock Insurance program is discussed and good practices in linking research and implementation for the achievement of development impacts are highlighted.
74. John F, Toth R, Frank K, Groeneveld J, Müller B: **Ecological vulnerability through insurance? Potential unintended consequences of livestock drought insurance**. *Ecol Econ* 2019, **157**:357-368.
- This study analyzes potential long-term ecological impacts of livestock drought insurance using model simulations, and suggests how land degradation dynamics could potentially be induced by scaling-up insurance programs.
75. Müller B, Johnson L, Kreuer D: **Maladaptive outcomes of climate insurance in agriculture**. *Global Environ Change* 2017, **46**:23-33.
76. Matsuda A, Takahashi K, Ikegami M: **Direct and indirect impact of index-based livestock insurance in Southern Ethiopia**. *Geneva Papers Risk Insur: Issues Pract* 2019, **44**:481-502.
77. Herrick JE, Shaver P, Pyke DA, Pellant M, Toledo D, Lepak N: **A strategy for defining the reference for land health and degradation assessments**. *Ecol Indic* 2019, **97**:225-230.
78. del Río-Mena T, Willemsen L, Tesfamariam GT, Beukes O, Nelson A: **Remote sensing for mapping ecosystem services to support evaluation of ecological restoration interventions in an arid landscape**. *Ecol Indic* 2020, **113**.
79. Meroni M, Schucknecht A, Fasbender D, Rembold F, Fava F, Mauclaire M, Goffner D, Di Lucchio LM, Leonardi U: **Remote sensing monitoring of land restoration interventions in semi-arid environments with a before–after control-impact statistical design**. *Int J Appl Earth Observ Geoinf* 2017, **59**:42-52.
80. Watmough GR, Marcinko CLJ, Sullivan C, Tschirhart K, Mutuo PK, Palm CA, Svenning JC: **Socioecologically informed use of remote sensing data to predict rural household poverty**. *Proc Natl Acad Sci U S A* 2019, **116**:1213-1218.
81. He X, Estes L, Konar M, Tian D, Anghileri D, Baylis K, Evans TP, Sheffield J: **Integrated approaches to understanding and reducing drought impact on food security across scales**. *Curr Opin Environ Sustain* 2019, **40**:43-54.
- This review discusses how drought impacts food security, with a focus on resource availability, access, and stability, and links data and models with the decision making process.
82. Cooper MW, Brown ME, Hochrainer-Stigler S, Pflug G, McCallum I, Fritz S, Silva J, Zvoleff A: **Mapping the effects of drought on child stunting**. *Proc Natl Acad Sci U S A* 2019, **116**:17219-17224.
83. Dong J, Metternicht G, Hostert P, Fensholt R, Chowdhury RR: **Remote sensing and geospatial technologies in support of a normative land system science: status and prospects**. *Curr Opin Environ Sustain* 2019, **38**:44-52.
84. Knippenberg E, Jensen N, Constanas M: **Quantifying household resilience with high frequency data: temporal dynamics and methodological options**. *World Dev* 2019, **121**:1-15.
- This study presents multiple approaches to measure household resilience to shocks, outlining a methodological approach for collecting monthly 'high-frequency' household data.
85. Giroux SA, Kouper I, Estes LD, Schumacher J, Waldman K, Greenshields JT, Dickinson SL, Caylor KK, Evans TP: **A high-frequency mobile phone data collection approach for research in social-environmental systems: applications in climate variability and food security in sub-Saharan Africa**. *Environ Model Softw* 2019, **119**:57-69.
86. Kimiti DW, Ganguli AC, Herrick JE, Karl JW, Bailey DW: **A decision support system for incorporating land potential information in the evaluation of restoration outcomes**. *Ecol Restor* 2020, **38**:94-104.
87. Reichstein M, Camps-Valls G, Stevens B, Jung M, Denzler J, Carvalhais N: **Prabhat: Deep learning and process understanding for data-driven Earth system science**. *Nature* 2019, **566**:195-204.