

# Higgs Rapidity Distribution in $b\bar{b}$ Annihilation at Threshold in N<sup>3</sup>LO QCD

---

Taushif Ahmed,<sup>a</sup> M. K. Mandal,<sup>a</sup> Narayan Rana<sup>a</sup> and V. Ravindran<sup>b</sup>

<sup>a</sup>*Regional Centre for Accelerator-based Particle Physics,  
Harish-Chandra Research Institute, Allahabad, India*

<sup>b</sup>*The Institute of Mathematical Sciences, Chennai, India*

*E-mail:* [taushif@hri.res.in](mailto:taushif@hri.res.in), [mandal@hri.res.in](mailto:mandal@hri.res.in), [narayan@hri.res.in](mailto:narayan@hri.res.in),  
[ravindra@imsc.res.in](mailto:ravindra@imsc.res.in)

**ABSTRACT:** We present the rapidity distribution of the Higgs boson produced through bottom quark annihilation at third order in QCD using the threshold approximation. We provide a framework, based on the factorization properties of the QCD amplitudes along with Sudakov resummation and the renormalization group invariance, that allows one to perform the computation of the threshold corrections in a consistent, systematic and accurate way. The recent results on threshold N<sup>3</sup>LO correction in QCD for the Drell-Yan production and on three loop QCD correction to Higgs form factor with bottom anti-bottom quark are used to achieve this task. We also demonstrate the numerical impact of these corrections at the LHC.

**KEYWORDS:** QCD, Higgs, Threshold corrections, Rapidity

---

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Differential Distribution with Respect to Rapidity</b>	<b>3</b>
2.1	Threshold Correction at NLO	4
2.2	Threshold Corrections Beyond NLO	10
2.3	Numerical Results	25
<b>3</b>	<b>Conclusions</b>	<b>30</b>

---

## 1 Introduction

With the spectacular discovery of the Higgs boson at CERN LHC [1, 2], the full spectrum of matter particles and force carriers of the Standard Model (SM) has been established very successfully. Though the mass of the newly discovered boson is already pinned down with an impressive experimental uncertainty of just a few hundred MeV in the range of 125 - 126 GeV, to fully validate the mechanism of electroweak symmetry breaking and to shed light on the possible potential deviations from its SM apprehension, it is indispensable to study the inclusive as well as exclusive observables associated with the production and decay channels of the Higgs boson to a very high accuracy.

Within the framework of SM, the production mechanism of the Higgs boson is dominated by gluon fusion, whereas one of the alternative channels, namely, bottom quark annihilation is severely suppressed by the small Yukawa coupling of bottom quark to the Higgs boson. However, in extensions of the SM with an enlarged spectrum of Higgs sector, as in the case of two-Higgs doublet model, the Yukawa coupling of bottom quark to some of the Higgs bosons can be enhanced significantly, such that the production channel of bottom quark annihilation could be the dominant one. Moreover, the contribution from gluon fusion channel decreases due to enhanced negative top-bottom interference diagrams. Furthermore, the bottom quark initiated processes at hadron colliders are of much theoretical interest on account of the freedom in treating the initial state bottom-quarks. In the four flavor scheme (4FS), alternatively known as the fixed flavor number scheme (FFS), the mass of the bottom quarks is considered to be non-zero throughout and they are excluded from the proton constituents, whereas, in the framework of five flavor scheme (5FS), also known as the variable flavor number scheme (VFS), the bottom quarks are considered as massless partons, except in the Yukawa coupling, with their own parton distribution functions (PDF).

The inclusive productions of the Higgs boson in gluon-gluon fusion [3–11], vector boson fusion [12] and associated production with vector bosons [13] are known to next-to-next-to

leading order (NNLO) accuracy in QCD. The Higgs boson production through bottom-antibottom ( $b\bar{b}$ ) annihilation is also known to NNLO accuracy in the VFS [14–19], whereas it is known to NLO in the fixed FFS [20–25].

While the theoretical predictions at NNLO and next-to-next-to leading log (NNLL) [26] QCD corrections and of two loop electroweak effects [27–32] played an important role in the discovery of the Higgs boson, the theoretical uncertainties resulting from the unphysical factorization and renormalization scales are not fully under control. In addition, the interpretation of the experimental data with higher accuracy from the upcoming run at the LHC demands the inclusion of higher order terms in QCD in the theoretical computation. Hence, the efforts to go beyond NNLO are going on intensively in past few decades. The computation of N<sup>3</sup>LO corrections is underway and some of the crucial ingredients, like the quark and gluon form factors [33–37], the mass factorization kernels [38] and the renormalization constant [39] for the effective operator describing the coupling between the Higgs boson and the SM fields in the infinite top quark mass limit are available up to three loop level in dimensional regularization. In addition, NNLO soft contributions are also known [40] in  $n$  dimensions. These results were already used to compute the partial threshold contributions at N<sup>3</sup>LO to the production cross-section of di-leptons in Drell-Yan (DY) and of the Higgs boson in gluon fusion as well as in  $b\bar{b}$  annihilation, see [41–45]. Since then, there have been several advances [46–49] towards obtaining the complete N<sup>3</sup>LO result for the inclusive Higgs production. The milestone in this direction was achieved by Anastasiou et al. in [50] to obtain the complete threshold N<sup>3</sup>LO corrections. This result provided a crucial input in [51] to obtain the corresponding N<sup>3</sup>LO threshold corrections to DY production. Independently, in [52], using light-like Wilson lines threshold corrections to the Higgs boson as well as Drell-Yan productions up to N<sup>3</sup>LO were obtained. Catani et al. in [53] used the universality of soft gluon contributions near threshold and the results of [50] to obtain general expression of the hard-virtual coefficient relevant for N<sup>3</sup>LO threshold as well as threshold resummation at next- to-next-to-next-to-leading-logarithmic (N<sup>3</sup>LL) accuracy for the production cross section of a colourless heavy particle at hadron colliders. There have been several attempts to go beyond threshold corrections [54, 55] for the inclusive Higgs production at N<sup>3</sup>LO. Recently, [56], the full next to soft as well as the exact results for the coefficients of the first three leading logarithms at this order have been obtained for the first time. For the Higgs boson production through  $b\bar{b}$  annihilation, the recent results of the Higgs form factor with bottom-antibottom by Gehrmann and Kara [57] and the universal soft distribution obtained for the Drell-Yan production [51] enabled us to obtain the missing  $\delta(1-z)$  contribution (see [44, 45, 58] for the partial results to this order) to the production cross-section at threshold at N<sup>3</sup>LO [59].

Like the inclusive one, the differential rapidity distributions are computed for the dilepton pair in DY [60] and the Higgs boson produced through gluon fusion in [61, 62], the Higgs boson through  $b\bar{b}$  annihilation in [63] and associated production of the Higgs with vector boson in [64, 65] to NNLO in QCD. Using the formalism developed in [44, 45], the partial N<sup>3</sup>LO threshold correction to the rapidity distributions of the dileptons in DY and the Higgs boson in gluon fusion as well as bottom quark annihilation were computed in [66]. Following the same technique, we obtained the complete N<sup>3</sup>LO threshold correction

to the rapidity distributions of both dilepton pair in DY and the Higgs boson in gluon fusion [67]. We had seen the dominance of the threshold contribution to the rapidity distribution in these processes. A significant amount of reduction in the dependence on the unphysical renormalization and factorization scale of the rapidity distribution takes place upon inclusion of the N<sup>3</sup>LO threshold corrections. In addition, these computations provide first results beyond NNLO level and will serve as a non-trivial check for a complete N<sup>3</sup>LO results. Keeping these motivations in mind, we intend to extend the existing result of the rapidity distribution of the Higgs boson produced through  $b\bar{b}$  annihilation to higher accuracy, namely the inclusion of complete N<sup>3</sup>LO threshold correction.

In Sec. 2.1, we perform an explicit calculation of threshold correction to the rapidity distribution of the Higgs boson in  $b\bar{b}$  annihilation at NLO, using the factorization properties of QCD amplitude, Sudakov resummation of soft gluons and renormalization group invariance. This helps us to build an elegant framework to calculate the rapidity distribution at threshold, of a colorless state produced at hadron colliders, to all orders in QCD perturbation theory. In Sec. 2.2, we use that general framework to achieve the goal of computing the complete analytic expression for the threshold corrections beyond NLO and provide the result up to N<sup>3</sup>LO. Sec. 2.3 contains the discussion on the numerical impacts of our results. Finally, we conclude with our findings in Sec. 3.

## 2 Differential Distribution with Respect to Rapidity

The interaction of bottom quarks and the Higgs boson is encapsulated in the following action

$$S_I^b = -\frac{\lambda}{\sqrt{2}} \int d^4x \phi(x) \bar{\psi}_b(x) \psi_b(x) \quad (2.1)$$

where,  $\psi_b(x)$  and  $\phi(x)$  denote the bottom quark and scalar field, respectively. The Yukawa coupling  $\lambda$  is given by  $\sqrt{2}m_b/v$ , with the bottom quark mass  $m_b$  and the vacuum expectation value  $v \approx 246$  GeV. Throughout our calculation, we consider five active flavours (VFS scheme), hence except in the Yukawa coupling,  $m_b$  is taken to be zero like other light quarks in the theory.

We study infrared safe differential distribution, namely rapidity distribution of the Higgs boson at hadron colliders, in particular those produced through bottom anti-bottom annihilation. Our findings are very well suited for similar observables where the rapidity distribution is for any colorless state produced at hadron colliders. We will set up a framework that can provide threshold corrections to rapidity distribution of the Higgs boson to all orders in perturbation theory. It is then straightforward to obtain fixed order perturbative results in the threshold limit.

The general frame work that we set up for the computation of threshold corrections beyond leading order in the perturbation theory for such observables is based on the factorization property of the QCD amplitudes. Sudakov resummation of soft gluons, renormalization group equations and most importantly the infrared safety of the observable play important role in achieving this task. QCD amplitudes that contribute to hard scattering

cross sections exhibit rich infra-red structure through cusp and collinear anomalous dimensions due to the factorization property of soft and collinear configurations. Massless gluons and light quarks are responsible for soft and collinear singularities in these amplitudes and also in partonic subprocesses. Singularities resulting from soft gluons cancel between virtual and real emission diagrams in infrared safe observables. While the final state collinear singularities cancel among themselves if the summation over degenerate states are appropriately carried out in such observables, the initial state collinear singular configurations remain until they are absorbed into bare parton distribution functions. In the upcoming section, we present one loop computation for the rapidity distribution in order to demonstrate how the various soft singularities cancel and also to give a pedagogical derivation of how the most general resummed threshold correction to the rapidity distribution can be obtained.

## 2.1 Threshold Correction at NLO

The process under consideration is the production of the Higgs boson through bottom quark annihilation in hadron colliders. The leading order process is

$$b(k_1) + \bar{b}(k_2) \rightarrow H(q) \quad (2.2)$$

where,  $k_i$ 's are the momenta of the incoming bottom and anti-bottom quarks involved in partonic reaction and  $q$  is the momentum of the Higgs boson. The hadronic center of mass energy squared is defined by  $S \equiv (p_1 + p_2)^2$ , where  $p_i$ 's are the hadronic momenta and the corresponding one for the incoming partons is given as  $\hat{s} = (k_1 + k_2)^2$ . The fraction of the initial state hadron momentum carried by the parton is denoted by  $x_i$  *i.e.*  $k_i = x_i p_i$ . The rapidity of the Higgs boson is defined through

$$y = \frac{1}{2} \ln \left( \frac{p_2 \cdot q}{p_1 \cdot q} \right). \quad (2.3)$$

The differential distribution with respect to rapidity of the Higgs boson can be expressed as

$$\frac{d}{dy} \sigma^b(\tau, q^2, y) = \sigma^{b,(0)}(\tau, q^2, \mu_R^2) W^b(\tau, y, q^2, \mu_R^2), \quad \sigma^{b,(0)} = \frac{\pi}{4SN} \lambda^2(\mu_R^2) \quad (2.4)$$

with  $\tau \equiv q^2/S$ ,  $q^2 = m_H^2$ ,  $m_H$ -the mass of the Higgs boson.  $\lambda(\mu_R^2)$  is the Yukawa coupling defined at the renormalization scale  $\mu_R$ ,  $N = 3$  is the number of QCD colors and  $\sigma^{b,(0)}$  is the leading order cross-section. Defining  $z \equiv q^2/\hat{s}$ , we find

$$W^b(\tau, y, q^2, \mu_R^2) = \frac{(Z^b(\mu_R^2))^2}{\sigma^{b,(0)}} \sum_{a,c=b,\bar{b},g} \int_0^1 dx_1 \int_0^1 dx_2 \hat{\mathcal{H}}_{ac}(x_1, x_2) \int_0^1 dz \delta(\tau - zx_1 x_2) \\ \times \int dPS_{1+X} |\mathcal{M}_{ac \rightarrow H+X}|^2 \delta \left( y - \frac{1}{2} \ln \left( \frac{p_2 \cdot q}{p_1 \cdot q} \right) \right). \quad (2.5)$$

In this expression,  $X$  is the remnants other than the Higgs boson,  $Z^b(\mu_R^2)$  is the ultraviolet (UV) renormalization constant for the Yukawa coupling  $\lambda$  and  $dPS_{1+X}$  is the phase

space element for the  $H + X$  system.  $\mathcal{M}_{ac \rightarrow H+X}$  denotes the scattering amplitude at partonic level. The function  $\hat{\mathcal{H}}_{ac}(x_1, x_2)$  is the product of unrenormalized parton distribution functions (PDF)  $\hat{f}_a(x_1)$  and  $\hat{f}_c(x_2)$ ,

$$\hat{\mathcal{H}}_{ac}(x_1, x_2) \equiv \hat{f}_a(x_1)\hat{f}_c(x_2). \quad (2.6)$$

The PDF  $f_a(x_1, \mu_F^2)$ , renormalized at the factorization scale  $\mu_F$ , is related to the unrenormalized ones through Altarelli-Parisi (AP) kernel  $\Gamma_{ad}$  as follows:

$$f_a(x_i, \mu_F^2) = \sum_{d=b, \bar{b}, g} \int_{x_i}^1 \frac{dz}{z} \Gamma_{ad}(\hat{a}_s, \mu^2, \mu_F^2, z, \epsilon) \hat{f}_d\left(\frac{x_i}{z}\right), \quad a = b, \bar{b}, g \quad (2.7)$$

where, the scale  $\mu$  is introduced to keep the strong coupling constant  $\hat{g}_s$  dimensionless in space-time dimensions  $n = 4 + \epsilon$ , regulating the theory and  $\hat{a}_s \equiv \hat{g}_s^2/16\pi^2$ . Expanding the AP kernel in powers of  $\hat{a}_s$ , we get

$$\Gamma_{ad}(\hat{a}_s, \mu^2, \mu_F^2, z, \epsilon) = \delta_{ad}\delta(1-z) + \hat{a}_s S_\epsilon \left(\frac{\mu_F^2}{\mu^2}\right)^{\frac{\epsilon}{2}} \frac{1}{\epsilon} P_{ad}^{(0)}(z) + \mathcal{O}(\hat{a}_s^2) \quad (2.8)$$

where,  $P_{ad}^{(0)}(z)$  is the leading order AP splitting function.  $S_\epsilon = \exp\left((\gamma_E - \ln 4\pi)\frac{\epsilon}{2}\right)$  where  $\gamma_E$  is the Euler-Mascheroni constant. Using  $\Gamma_{ad}$ ,  $W^b$  can be written in terms of renormalized  $\mathcal{H}$ , given by

$$\begin{aligned} \mathcal{H}_{ac}(x_1, x_2, \mu_F^2) &\equiv f_a(x_1, \mu_F^2) f_c(x_2, \mu_F^2) \\ &= \int_{x_1}^1 \frac{dy_1}{y_1} \int_{x_2}^1 \frac{dy_2}{y_2} \Gamma_{aa'}(\hat{a}_s, \mu^2, \mu_F^2, y_1, \epsilon) \hat{\mathcal{H}}_{a'c'}\left(\frac{x_1}{y_1}, \frac{x_2}{y_2}\right) \Gamma_{cc'}(\hat{a}_s, \mu^2, \mu_F^2, y_2, \epsilon). \end{aligned} \quad (2.9)$$

The LO contribution arises from the Born process  $b + \bar{b} \rightarrow H$  and the NLO ones are from one loop virtual contributions to born process and from the real emission processes, namely  $b + \bar{b} \rightarrow H + g$ ,  $b(\bar{b}) + g \rightarrow H + b(\bar{b})$ . For LO and virtual contributions,  $dPS_{1+X} = dPS_1$  and for real emission processes we have two body phase space element  $dPS_{1+X} = dPS_2$ . In order to define the threshold limit at the partonic level and to express the hadronic cross-section in terms of the partonic one through convolution integrals, we choose to work with the symmetric scaling variables  $x_1^0$  and  $x_2^0$  instead of  $y$  and  $\tau$  which are related through

$$y = \frac{1}{2} \ln\left(\frac{x_1^0}{x_2^0}\right), \quad \tau = x_1^0 x_2^0. \quad (2.10)$$

In terms of these new variables, the partonic subprocess contributions can be shown to depend on the ratios  $z_j = \frac{x_j^0}{x_j}$  which take the role of scaling variables at the partonic level. The dimensionless partonic differential cross-section denoted by  $\hat{\Delta}_{d,ac}^b$  through

$$\frac{1}{x_1 x_2} \hat{\Delta}_{d,ac}^b\left(\frac{x_1^0}{x_1}, \frac{x_2^0}{x_2}, \hat{a}_s, \mu^2, q^2, \mu_R^2\right) = \frac{(Z^b(\mu_R^2))^2}{\sigma^{b,(0)}} \int dPS_{1+X} \int dz |\mathcal{M}_{ac \rightarrow H+X}|^2$$

$$\times \delta(\tau - zx_1x_2) \delta\left(y - \frac{1}{2} \ln\left(\frac{p_2 \cdot q}{p_1 \cdot q}\right)\right) \quad (2.11)$$

is UV finite. Here subscript  $d$  stands for differential distribution. The collinear singularities that arise due to the initial state light partons are removed through the AP kernels resulting in the following finite  $\Delta_{d,ac}^b$

$$\begin{aligned} \Delta_{d,ac}^b(z_1, z_2, a_s(\mu_R^2), q^2, \mu_F^2, \mu_R^2) &= \int_{z_1}^1 \frac{dy_1}{y_1} \int_{z_2}^1 \frac{dy_2}{y_2} \Gamma_{aa'}^{-1}(\hat{a}_s, \mu^2, \mu_F^2, y_1, \epsilon) \\ &\times \hat{\Delta}_{d,a'c'}^b\left(\frac{z_1}{y_1}, \frac{z_2}{y_2}, \hat{a}_s, \mu^2, q^2, \mu_R^2, \epsilon\right) \Gamma_{cc'}^{-1}(\hat{a}_s, \mu^2, \mu_F^2, y_2, \epsilon). \end{aligned} \quad (2.12)$$

Therefore, expressing  $W^b$  in terms of renormalized  $\mathcal{H}_{ac}$  and finite  $\Delta_{d,ac}^b$ , we get

$$\begin{aligned} W^b(x_1^0, x_2^0, q^2, \mu_R^2) &= \sum_{ac=b, \bar{b}, g} \int_{x_1^0}^1 \frac{dz_1}{z_1} \int_{x_2^0}^1 \frac{dz_2}{z_2} \mathcal{H}_{ac}\left(\frac{x_1^0}{z_1}, \frac{x_2^0}{z_2}, \mu_F^2\right) \\ &\times \Delta_{d,ac}^b(z_1, z_2, a_s(\mu_R^2), q^2, \mu_F^2, \mu_R^2). \end{aligned} \quad (2.13)$$

Since,  $W^b$  involves convolutions of various functions, it becomes normal multiplication in the Mellin space of the Mellin moments of renormalized PDFs, AP kernels and bare differential partonic cross-section. The double Mellin moment of  $W^b(x_1^0, x_2^0)$  is defined by

$$\begin{aligned} \widetilde{W}^b(N_1, N_2) &\equiv \int dx_1^0 (x_1^0)^{N_1-1} \int dx_2^0 (x_2^0)^{N_2-1} W^b(x_1^0, x_2^0) \\ &= \widetilde{\mathcal{H}}_{ac}(N_1, N_2) \widetilde{\Delta}_{d,ac}^b(N_1, N_2) \end{aligned} \quad (2.14)$$

where

$$\widetilde{\Delta}_{d,ac}^b(N_1, N_2) = \widetilde{\Gamma}_{ae}^{-1}(N_1) \widetilde{\Gamma}_{cf}^{-1}(N_2) \widetilde{\Delta}_{d,ef}^b(N_1, N_2). \quad (2.15)$$

The threshold limit is defined by  $N_i \rightarrow \infty$ , which in  $z_j$  variables corresponds to  $z_j \rightarrow 1$ . In this limit, only diagonal terms in the AP kernel  $\widetilde{\Gamma}^{-1}$  and  $\widetilde{\Delta}_d^b$  contribute to the differential cross-section. Hence,  $\ln \widetilde{\Delta}_d^b$  is simply a sum of the contributions from 1) diagonal terms of the AP kernels and 2) bare differential partonic cross-section. Due to the born kinematics, the form factor contribution can be further factored out from the differential partonic cross sections to all orders in perturbation theory. Hence, the remaining part of the differential partonic cross-sections contains contributions from only real emission processes, namely those involving only soft gluons. Taking into account the renormalization constant of the Yukawa coupling  $\hat{\lambda}$ , we find

$$\begin{aligned} \ln \widetilde{\Delta}_d^b(N_1, N_2, q^2, \mu_R^2, \mu_F^2) &= \ln\left(Z^b(\mu_R^2)\right)^2 - \ln \widetilde{\Gamma}_{bb}(N_1, \mu_F^2) - \ln \widetilde{\Gamma}_{bb}(N_2, \mu_F^2) \\ &+ \ln |\hat{F}^b(q^2)|^2 + \ln \widetilde{S}^b(N_1, N_2, q^2) \end{aligned} \quad (2.16)$$

where  $\hat{F}^b$  and  $\tilde{S}^b(N_1, N_2)$  are bare form factor and real emission contributions of partonic subprocesses, respectively. The inverse Mellin transform will bring back the expressions in terms of the variables  $z_j$  and they will contain besides regular functions, the distributions namely  $\delta(1 - z_j)$ ,  $\mathcal{D}_i$  and  $\bar{\mathcal{D}}_i$ , defined as

$$\mathcal{D}_i = \left[ \frac{\ln^i(1 - z_1)}{(1 - z_1)} \right]_+, \quad \bar{\mathcal{D}}_i = \left[ \frac{\ln^i(1 - z_2)}{(1 - z_2)} \right]_+ \quad i = 0, 1, \dots \quad (2.17)$$

The subscript ‘+’ denotes the customary ‘plus-distribution’  $f_+(z)$  which acts on functions regular in  $z \rightarrow 1$  limit as

$$\int_0^1 dz f_+(z) g(z) = \int_0^1 dz f(z) (g(z) - g(1)) \quad (2.18)$$

where,  $g(z)$  is any well behaved function in the region  $0 \leq z \leq 1$ . In the threshold limit, we drop all the regular terms and keep only these distributions.

In the following, we perform NLO computation in the threshold limit. The overall renormalization constant  $(Z^b)^2$  is found to be

$$(Z^b(\mu_R^2))^2 = 1 + \hat{a}_s S_\epsilon \left( \frac{\mu_R^2}{\mu^2} \right)^\epsilon C_F \left( \frac{12}{\epsilon} \right) + \mathcal{O}(\hat{a}_s^2). \quad (2.19)$$

The form factor contribution  $|\hat{F}^b|^2$  at one loop level gives

$$|\hat{F}^b(q^2)|^2 = 1 + \hat{a}_s S_\epsilon \left( \frac{q^2}{\mu^2} \right)^\epsilon C_F \left( -\frac{16}{\epsilon} - 4 + 14\zeta_2 + \mathcal{O}(\epsilon) \right) + \mathcal{O}(\hat{a}_s^2) \quad (2.20)$$

The contribution from  $\Gamma_{bb}$  in the threshold limit is found to be

$$\begin{aligned} \int_{z_1}^1 \frac{dy_1}{y_1} \int_{z_2}^1 \frac{dy_2}{y_2} \Gamma_{bb}^{-1}(y_1, \mu_F^2) \delta\left(1 - \frac{z_1}{y_1}\right) \delta\left(1 - \frac{z_2}{y_2}\right) \Gamma_{bb}^{-1}(y_2, \mu_F^2) = \delta(1 - z_1) \delta(1 - z_2) \\ - \hat{a}_s S_\epsilon \left( \frac{\mu_F^2}{\mu^2} \right)^\epsilon C_F \frac{1}{\epsilon} \left[ \left( 8 \frac{1}{(1 - z_1)_+} + 6\delta(1 - z_1) \right) \delta(1 - z_2) \right. \\ \left. + \left( 8 \frac{1}{(1 - z_2)_+} + 6\delta(1 - z_2) \right) \delta(1 - z_1) \right] + \mathcal{O}(\hat{a}_s^2). \end{aligned} \quad (2.21)$$

Note that the regular terms in the limit  $z_j \rightarrow 1$  in  $\Gamma_{bb}$  do not contribute in the threshold limit and hence dropped.

The inverse Mellin transform of  $\tilde{S}^b(N_1, N_2)$ , namely  $S^b(z_1, z_2)$  can be obtained directly from the real gluon emission processes in bottom anti-bottom annihilation processes:  $b + \bar{b} \rightarrow H + g$ . The two body phase space is given by

$$dPS_{H+g} = \frac{1}{8\pi x_1 x_2} \frac{1}{\Gamma(1 + \frac{\epsilon}{2})} \left( \frac{m_H^2}{4\pi} \right)^{\frac{\epsilon}{2}} \frac{2z_1 z_2 (1 + z_1 z_2)}{(z_1 + z_2)^{2-\epsilon}} ((1 - z_1^2)(1 - z_2^2))^{\frac{\epsilon}{2}}. \quad (2.22)$$



The phase space in the limit  $z_j \rightarrow 1$  becomes

$$dPS_{H+g}|_{z_i \rightarrow 1} = \frac{1}{8\pi x_1 x_2} \frac{1}{\Gamma(1 + \frac{\epsilon}{2})} \left( \frac{m_H^2}{4\pi} \right)^{\frac{\epsilon}{2}} ((1 - z_1^2)(1 - z_2^2))^{\frac{\epsilon}{2}}. \quad (2.23)$$

The spin and color averaged matrix element square in threshold limit is found to be

$$|\overline{M}_{b\bar{b} \rightarrow H+g}|_{z_j \rightarrow 1}^2 = \sigma_0^b \frac{\hat{a}_s}{\mu^\epsilon} C_F \left[ \frac{32}{(1 - z_1)(1 - z_2)} + \mathcal{O}(\epsilon^3) \right] \quad (2.24)$$

where terms that are regular in  $z_j$  as  $z_j \rightarrow 1$  have been dropped. It is then straightforward to obtain the threshold contribution resulting from the real gluon emission process:

$$\begin{aligned} S^b(z_1, z_2) &= \delta(1 - z_1)\delta(1 - z_2) + \hat{a}_s \left( \frac{q^2}{4\pi\mu^2} \right)^{\frac{\epsilon}{2}} \frac{1}{\Gamma(1 + \frac{\epsilon}{2})} \\ &\quad \times 4C_F \left[ \frac{(1 - z_1)^{\frac{\epsilon}{2}}(1 - z_2)^{\frac{\epsilon}{2}}}{(1 - z_1)(1 - z_2)} + \mathcal{O}(\epsilon^3) \right]. \end{aligned} \quad (2.25)$$

Using the identity

$$\frac{(1 - z_j)^{a\frac{\epsilon}{2}}}{(1 - z_j)} = \frac{2}{a\epsilon} \delta(1 - z_j) + \left( \frac{(1 - z_j)^{a\frac{\epsilon}{2}}}{(1 - z_j)} \right)_+, \quad (2.26)$$

it can be shown that  $\Delta_d^b(z_1, z_2)$  in the threshold limit contains only the distributions such as  $\delta(1 - z_j)$ ,  $\mathcal{D}_i$  and  $\overline{\mathcal{D}}_i$ . Decomposing  $\Delta_{d,ac}^b$  into hard and soft parts,

$$\Delta_{d,ac}^b(z_1, z_2, q^2, \mu_F^2, \mu_R^2) = \Delta_{d,ac}^{b,\text{hard}}(z_1, z_2, q^2, \mu_F^2, \mu_R^2) + \delta_{ac} \Delta_{d,b}^{\text{SV}}(z_1, z_2, q^2, \mu_F^2, \mu_R^2), \quad (2.27)$$

and setting  $\mu_F = \mu_R = m_H$ , we find

$$\begin{aligned} \Delta_{d,b}^{\text{SV},(0)} &= \delta(1 - z_1)\delta(1 - z_2), \\ \Delta_{d,b}^{\text{SV},(1)} &= \delta(1 - z_1)\delta(1 - z_2)C_F \left( -2 + 6\zeta_2 \right) + \mathcal{D}_0 \overline{\mathcal{D}}_0 \left( 2C_F \right) \\ &\quad + \mathcal{D}_1 \delta(1 - z_2) \left( 4C_F \right) + \left\{ z_1 \leftrightarrow z_2 \right\} \end{aligned} \quad (2.28)$$

At the hadronic level, decomposing  $W^b$  as

$$W^b(x_1^0, x_2^0, q^2, \mu_R^2, \mu_F^2) = W_b^{\text{hard}}(x_1^0, x_2^0, q^2, \mu_R^2, \mu_F^2) + W_b^{\text{SV}}(x_1^0, x_2^0, q^2, \mu_R^2, \mu_F^2), \quad (2.29)$$

similar to  $\Delta_{d,b}$  and putting  $\Delta_{d,b}^{\text{SV}}$  we get, to order  $a_s = a_s(m_H^2)$

$$\begin{aligned} W_b^{\text{SV}}(x_1^0, x_2^0, q^2, m_H^2) &= \mathcal{H}_{b\bar{b}}(x_1^0, x_2^0) + a_s 4C_F \left[ \mathcal{H}_{b\bar{b}}(x_1^0, x_2^0) \left( -1 + \zeta_2 + li_2(x_1^0) + li_2(x_2^0) \right) \right. \\ &\quad \left. + \frac{1}{2} \ln^2((1 - x_1^0)(1 - x_2^0)) + \ln \left( \frac{(1 - x_1^0)}{x_2^0} \right) \ln \left( \frac{(1 - x_2^0)}{x_1^0} \right) \right] \end{aligned}$$

$$\begin{aligned}
& + \int dx_1 \mathcal{H}_{b\bar{b},1}(x_1, x_2^0) \frac{1}{x_1 - x_1^0} \ln \left( \frac{(1 - x_2^0)(x_1 - x_1^0)}{x_1 x_2^0} \right) \\
& + \int dx_2 \mathcal{H}_{b\bar{b},1}(x_1^0, x_2) \frac{1}{x_2 - x_2^0} \ln \left( \frac{(1 - x_1^0)(x_2 - x_2^0)}{x_1^0 x_2} \right) \\
& + \int dx_1 \int dx_2 \mathcal{H}_{b\bar{b},12}(x_1, x_2) \frac{1}{(x_1 - x_1^0)(x_2 - x_2^0)} \Big]. \tag{2.30}
\end{aligned}$$

where all the parton densities are defined at  $\mu_F = m_H$ . In general,

$$\begin{aligned}
\mathcal{H}_{b\bar{b},12}(x_1, x_2, \mu_F^2) &\equiv \mathcal{H}_{b\bar{b}}(x_1, x_2, \mu_F^2) - \mathcal{H}_{b\bar{b}}(x_1^0, x_2, \mu_F^2) - \mathcal{H}_{b\bar{b}}(x_1, x_2^0, \mu_F^2) + \mathcal{H}_{b\bar{b}}(x_1^0, x_2^0, \mu_F^2), \\
\mathcal{H}_{b\bar{b},1}(x_1, x_2, \mu_F^2) &\equiv \mathcal{H}_{b\bar{b}}(x_1, x_2, \mu_F^2) - \mathcal{H}_{b\bar{b}}(x_1^0, x_2, \mu_F^2), \\
\mathcal{H}_{b\bar{b},2}(x_1, x_2, \mu_F^2) &\equiv \mathcal{H}_{b\bar{b}}(x_1, x_2, \mu_F^2) - \mathcal{H}_{b\bar{b}}(x_1, x_2^0, \mu_F^2) \tag{2.31}
\end{aligned}$$

The Spence function ( $li_2(x)$ ) is defined as

$$li_2(x) \equiv - \int_0^x \frac{dz}{z} \ln(1 - z). \tag{2.32}$$

The exact result computed at NLO level confirms our expectations, see for example [68] where the rapidity distribution of di-leptons in the Drell-Yan production for a physics beyond the SM (BSM) involving a generic Yukawa type interaction was obtained to NLO level. After the suitable replacement of the BSM coupling in [68], we obtain

$$\frac{d\sigma^b}{dy}(\tau, y, Q^2) = \sigma^{b,(0)}(\mu_F) \left[ W_{b\bar{b}}(x_1^0, x_2^0, \mu_F^2) + W_{bg}(x_1^0, x_2^0, \mu_F^2) + W_{gb}(x_1^0, x_2^0, \mu_F^2) \right] \tag{2.33}$$

where  $W$ 's can be expanded in the strong coupling constant  $a_s(\mu_F^2)$  as

$$W_{ac}(x_1^0, x_2^0, \mu_F^2) = W_{ac}^{(0)}(x_1^0, x_2^0, \mu_F^2) + a_s(\mu_F^2) W_{ac}^{(1)}(x_1^0, x_2^0, \mu_F^2) + \mathcal{O}(a_s^2) \tag{2.34}$$

and the corresponding coefficients are given by

$$\begin{aligned}
W_{b\bar{b}}^{(0)}(x_1^0, x_2^0, \mu_F^2) &= \mathcal{H}_{b\bar{b}}(x_1^0, x_2^0, \mu_F^2) \\
W_{b\bar{b}}^{(1)}(x_1^0, x_2^0, \mu_F^2) &= 2 C_F \left\{ \varphi_0^{b\bar{b}} + \int dx_1 \varphi_1^{b\bar{b}} + \int dx_1 dx_2 \varphi_2^{b\bar{b}} \right\} + (1 \leftrightarrow 2) \\
W_{gb}^{(1)}(x_1^0, x_2^0, \mu_F^2) &= 2 T_f \int \frac{dx_1}{x_1^3} \left[ \varphi_1^{g\bar{b}} + \int dx_2 \left\{ \varphi_2^{g\bar{b}} - \frac{\varphi_3^{g\bar{b}} \mathcal{H}_{gb}(x_1, x_2, \mu_F^2)}{x_2^2 (x_2 + x_2^0) (x_1 x_2^0 + x_2 x_1^0)^3} \right\} \right] \\
W_{bg}^{(1)}(x_1^0, x_2^0, \mu_F^2) &= W_{gb}^{(1)}(x_1^0, x_2^0, \mu_F^2)|_{(1 \leftrightarrow 2)} \tag{2.35}
\end{aligned}$$

with

$$\varphi_0^{b\bar{b}} = \frac{1}{2} \mathcal{H}_{b\bar{b}}(x_1^0, x_2^0, \mu_F^2) \left( -2 + \kappa_{12}^2 + 6 \zeta_2 + 2 \kappa_{12} \ln \frac{q^2}{\mu_F^2} \right)$$

$$\begin{aligned}
\varphi_1^{\bar{b}\bar{b}} &= \frac{2\kappa_{b_1}}{x_1 - x_1^0} \mathcal{H}_{\bar{b}\bar{b},1}(x_1, x_2^0, \mu_F^2) + \mathcal{H}_{\bar{b}\bar{b}}(x_1, x_2^0, \mu_F^2) \left( \frac{1 - \kappa_{a_1}}{x_1} + \frac{2\kappa_{c_1}}{x_1 - x_1^0} - \frac{1 + \kappa_{a_1}}{x_1^2} x_1^0 \right) \\
\varphi_2^{\bar{b}\bar{b}} &= \frac{\mathcal{H}_{\bar{b}\bar{b},12}(x_1, x_2, \mu_F^2)}{(x_1 - x_1^0)(x_2 - x_2^0)} - \frac{x_2 + x_2^0}{(x_1 - x_1^0)x_2^2} \mathcal{H}_{\bar{b}\bar{b},1}(x_1, x_2, \mu_F^2) \\
&\quad + \frac{\mathcal{H}_{\bar{b}\bar{b}}(x_1, x_2, \mu_F^2)}{2x_1^2 x_2^2} \left( (x_1 + x_1^0)(x_2 + x_2^0) + \frac{x_1^2 x_2^2 + x_1^0 x_2^0}{(x_1 + x_1^0)(x_2 + x_2^0)} \right) \\
\varphi_1^{g\bar{b}} &= \mathcal{H}_{g\bar{b}}(x_1, x_2^0, \mu_F^2) \left( 2x_1^0(x_1 - x_1^0) + \kappa_{a_1} (x_1^0 + (x_1 - x_1^0)^2) \right) \\
\varphi_2^{g\bar{b}} &= \frac{\mathcal{H}_{g\bar{b},2}(x_1, x_2, \mu_F^2)}{x_2 - x_2^0} (x_1^0 + (x_1 - x_1^0)^2) \\
\varphi_3^{g\bar{b}} &= -x_1^5 x_2^2 x_2^0 + x_1^4 x_1^0 x_2^2 x_2^0 (3x_2 + 4x_2^0) + x_1^3 x_1^0 x_2 x_2^0 (3x_2^3 + 2x_2^0) + 2x_1^0 x_2^2 (x_2^3 + 2x_2^2 x_2^0 \\
&\quad + 2x_2 x_2^0 + 2x_2^0) + 2x_1 x_1^0 x_2 (-x_2^4 + x_2^3 x_2^0 + 4x_2^2 x_2^0 + 2x_2 x_2^0 + 2x_2^0) \\
&\quad + x_1^2 x_1^0 (x_2^5 - 4x_2^4 x_2^0 - 4x_2^3 x_2^0 + 2x_2^2 x_2^0 + 2x_2 x_2^0 + 2x_2^0) \tag{2.36}
\end{aligned}$$

and

$$\begin{aligned}
\kappa_{a_1} &= \ln \frac{2q^2(1-x_2^0)(x_1-x_1^0)}{\mu_F^2(x_1+x_1^0)x_2^0}, & \kappa_{b_1} &= \ln \frac{q^2(1-x_2^0)(x_1-x_1^0)}{\mu_F^2 x_1^0 x_2^0} \\
\kappa_{c_1} &= \ln \frac{2x_1^0}{x_1+x_1^0}, & \kappa_{12} &= \ln \frac{(1-x_1^0)(1-x_2^0)}{x_1^0 x_2^0}. \tag{2.37}
\end{aligned}$$

In the threshold limit, after setting  $\mu_F = m_H$ , we find that the above result reduces to given in Eq. 2.30.

## 2.2 Threshold Corrections Beyond NLO

Following the factorization approach that we used in the previous section to obtain the threshold correction to NLO rapidity distribution, we now set up a framework to compute threshold corrections to rapidity distribution to all orders in strong coupling constant. Our approach is based on the fact that the rapidity distribution in the threshold limit can be systematically factorized into 1) the exact form factor, 2) overall UV renormalization constant, 3) soft gluon contributions from real emission partonic subprocesses and 4) the diagonal collinear subtraction terms involving only  $\delta(1-z)$  and  $\mathcal{D}_0(z)$  terms of AP splitting functions. We call such a combination soft-virtual (SV) part of the rapidity distribution and the remaining part as hard. Hence, we propose that

$$\Delta_{d,b}^{\text{SV}}(z_1, z_2, q^2, \mu_R^2, \mu_F^2) = \mathcal{C} \exp \left( \Psi_d^b(q^2, \mu_R^2, \mu_F^2, z_1, z_2, \epsilon) \right) \Big|_{\epsilon=0}. \tag{2.38}$$

The symbol ‘ $\mathcal{C}$ ’ means convolution with the following definition

$$\begin{aligned}
\mathcal{C}e f(z_1, z_2) &= \delta(1-z_1)\delta(1-z_2) + \frac{1}{1!} f(z_1, z_2) + \frac{1}{2!} f(z_1, z_2) \otimes f(z_1, z_2) \\
&\quad + \frac{1}{3!} f(z_1, z_2) \otimes f(z_1, z_2) \otimes f(z_1, z_2) + \dots, \tag{2.39}
\end{aligned}$$

where,  $\otimes$  indicates double Mellin convolution with respect to the variables  $z_1$  and  $z_2$  and the function  $f(z_1, z_2)$  is a distribution of the kind  $\delta(1-z_j)$  and/or  $\mathcal{D}_i(z_j)$ . The finite distribution  $\Psi_d^b$  in dimensional regularization contains  $Hb\bar{b}$  unrenormalized form factor  $\hat{F}^b(\hat{a}_s, q^2 = -Q^2, \mu^2, \epsilon)$ , UV overall operator renormalization constant  $Z^b(\hat{a}_s, \mu_R^2, \mu^2, \epsilon)$ , soft distribution functions  $\Phi_d^b(\hat{a}_s, q^2, \mu^2, z_1, z_2, \epsilon)$  and the mass factorization kernels  $\Gamma_{bb}(\hat{a}_s, \mu^2, \mu_F^2, z_j, \epsilon)$ :

$$\begin{aligned} \Psi_d^b &= \left( \ln \left( Z^b(\hat{a}_s, \mu_R^2, \mu^2, \epsilon) \right)^2 + \ln |\hat{F}^b(\hat{a}_s, Q^2, \mu^2, \epsilon)|^2 \right) \delta(1-z_1)\delta(1-z_2) \\ &+ 2 \Phi_d^b(\hat{a}_s, q^2, \mu^2, z_1, z_2, \epsilon) - \mathcal{C} \ln \Gamma_{bb}(\hat{a}_s, \mu^2, \mu_F^2, z_1, \epsilon) \delta(1-z_2) \\ &- \mathcal{C} \ln \Gamma_{bb}(\hat{a}_s, \mu^2, \mu_F^2, z_2, \epsilon) \delta(1-z_1). \end{aligned} \quad (2.40)$$

We have expressed all the quantities in the above equation in terms of unrenormalized strong coupling constant  $\hat{a}_s$  related to the standard  $\hat{\alpha}_s$  through  $\hat{a}_s = \hat{\alpha}_s/4\pi$  and the dimensional regularization scale  $\mu$ . The UV renormalization of  $\hat{a}_s$  is done at the renormalization scale  $\mu_R$  through  $Z(\mu_R^2)$  giving the renormalized  $a_s(\mu_R^2)$ , that is

$$\hat{a}_s = \left( \frac{\mu}{\mu_R} \right)^\epsilon Z(\mu_R^2) S_\epsilon^{-1} a_s(\mu_R^2). \quad (2.41)$$

The renormalization group equation (RGE) for  $a_s(\mu_R^2)$

$$\mu_R^2 \frac{da_s(\mu_R^2)}{d\mu_R^2} = \frac{\epsilon a_s(\mu_R^2)}{2} + \beta(a_s(\mu_R^2)) \quad (2.42)$$

with

$$\beta(a_s(\mu_R^2)) = a_s(\mu_R^2) \mu_R^2 \frac{d \ln Z(\mu_R^2)}{d\mu_R^2} = - \sum_{i=0}^{\infty} a_s^{i+2}(\mu_R^2) \beta_i, \quad (2.43)$$

determines the structure of the  $Z(\mu_R^2)$ , up to  $\mathcal{O}(a_s^3)$ , we find

$$Z(\mu_R^2) = 1 + a_s(\mu_R^2) \frac{2}{\epsilon} \beta_0 + a_s^2(\mu_R^2) \left( \frac{4}{\epsilon^2} \beta_0^2 + \frac{1}{\epsilon} \beta_1 \right) + a_s^3(\mu_R^2) \left( \frac{8}{\epsilon^3} \beta_0^3 + \frac{14}{3\epsilon^2} \beta_0 \beta_1 + \frac{2}{3\epsilon} \beta_2 \right). \quad (2.44)$$

The first three coefficients of the QCD  $\beta$  function,  $\beta_0$ ,  $\beta_1$  and  $\beta_2$  are given by [69]

$$\begin{aligned} \beta_0 &= \frac{11}{3} C_A - \frac{4}{3} T_F n_f, \\ \beta_1 &= \frac{34}{3} C_A^2 - 4 T_F n_f C_F - \frac{20}{3} T_F n_f C_A, \\ \beta_2 &= \frac{2857}{54} C_A^3 - \frac{1415}{27} C_A^2 T_F n_f + \frac{158}{27} C_A T_F^2 n_f^2 \\ &+ \frac{44}{9} C_F T_F^2 n_f^2 - \frac{205}{9} C_F C_A T_F n_f + 2 C_F^2 T_F n_f \end{aligned} \quad (2.45)$$

with the  $SU(N)$  color factors

$$C_A = N, \quad C_F = \frac{N^2 - 1}{2N}, \quad T_F = \frac{1}{2} \quad (2.46)$$

and  $n_f$  is the number of active flavours.

The overall operator renormalization constant  $Z^b$  renormalizes the bare Yukawa coupling  $\hat{\lambda}$  resulting  $\lambda(\mu_R^2)$  through the relation

$$\hat{\lambda} = \left(\frac{\mu}{\mu_R}\right)^{\frac{\epsilon}{2}} Z^b(\mu_R^2) S_\epsilon^{-1} \lambda(\mu_R^2). \quad (2.47)$$

In  $\overline{\text{MS}}$  scheme,  $Z^b(\mu_R^2)$  is identical to quark mass renormalization constant. The RGE for  $\lambda(\mu_R^2)$  takes the form

$$\mu_R^2 \frac{d}{d\mu_R^2} \ln Z^b(\hat{a}_s, \mu_R^2, \mu^2, \epsilon) = \sum_{i=1}^{\infty} a_s^i(\mu_R^2) \gamma_{i-1}^b, \quad (2.48)$$

with the anomalous dimensions  $\gamma_i^b$  given by [70–72]

$$\begin{aligned} \gamma_0^b &= 3C_F, \\ \gamma_1^b &= \frac{3}{2}C_F^2 + \frac{97}{6}C_F C_A - \frac{10}{3}C_F T_F n_f, \\ \gamma_2^b &= \frac{129}{2}C_F^3 - \frac{129}{4}C_F^2 C_A + \frac{11413}{108}C_F C_A^2 + \left(-46 + 48\zeta_3\right)C_F^2 T_F n_f \\ &\quad + \left(-\frac{556}{27} - 48\zeta_3\right)C_F C_A T_F n_f - \frac{140}{27}C_F T_F^2 n_f^2. \end{aligned} \quad (2.49)$$

Upon solving the above RGE in  $4 + \epsilon$  space-time dimensions, we obtain

$$\begin{aligned} \ln Z^b(\mu_R^2) &= a_s(\mu_R^2) \frac{1}{\epsilon} \left(2\gamma_0^b\right) + a_s^2(\mu_R^2) \left[\frac{1}{\epsilon^2} \left(2\beta_0\gamma_0^b\right) + \frac{1}{\epsilon} \left(\gamma_1^b\right)\right] \\ &\quad + a_s^3(\mu_R^2) \left[\frac{1}{\epsilon^3} \left(\frac{8}{3}\beta_0^2\gamma_0^b\right) + \frac{1}{\epsilon^2} \left(\frac{4}{3}\beta_1\gamma_0^b + \frac{4}{3}\beta_0\gamma_1^b\right) + \frac{1}{\epsilon} \left(\frac{2}{3}\gamma_2^b\right)\right] \end{aligned} \quad (2.50)$$

up to  $\mathcal{O}(a_s^3)$ .

The bare form factor  $\hat{F}^b(\hat{a}_s, Q^2, \mu^2, \epsilon)$  satisfies the following differential equation which follows from the gauge as well as renormalization group invariances [73–76]

$$Q^2 \frac{d}{dQ^2} \ln \hat{F}^b = \frac{1}{2} \left[ K^b(\hat{a}_s, \frac{\mu_R^2}{\mu^2}, \epsilon) + G^b(\hat{a}_s, \frac{Q^2}{\mu_R^2}, \frac{\mu_R^2}{\mu^2}, \epsilon) \right] \quad (2.51)$$

where, all the poles in  $\epsilon$  are encapsulated within  $K^b$  and  $G^b$  contains the terms finite in  $\epsilon$ . Renormalization group invariance of  $\hat{F}^b(\hat{a}_s, Q^2, \mu^2, \epsilon)$  leads

$$\mu_R^2 \frac{d}{d\mu_R^2} K^b = -\mu_R^2 \frac{d}{d\mu_R^2} G^b = -\sum_{i=1}^{\infty} a_s^i(\mu_R^2) A_i^q, \quad (2.52)$$

where,  $A_i^q$ 's are the cusp anomalous dimensions, found to be [38, 77–80]

$$A_1^q = 4C_F,$$

$$\begin{aligned}
A_2^q &= 8C_F C_A \left\{ \frac{67}{18} - \zeta_2 \right\} + 8C_F n_f \left\{ -\frac{5}{9} \right\}, \\
A_3^q &= 16C_F C_A^2 \left\{ \frac{245}{24} - \frac{67}{9}\zeta_2 + \frac{11}{6}\zeta_3 + \frac{11}{5}\zeta_2^2 \right\} + 16C_F^2 n_f \left\{ -\frac{55}{24} + 2\zeta_3 \right\} \\
&\quad + 16C_F C_A n_f \left\{ -\frac{209}{108} + \frac{10}{9}\zeta_2 - \frac{7}{3}\zeta_3 \right\} + 16C_F n_f^2 \left\{ -\frac{1}{27} \right\}. \tag{2.53}
\end{aligned}$$

Being flavor independent,  $A_i^b$ 's are same as  $A_i^q$ . Solving the RGE 2.52 satisfied by  $K^b$  we get

$$K^b(\hat{a}_s, \mu^2, \mu_R^2, \epsilon) = \sum_{i=1}^{\infty} \hat{a}_s^i \left( \frac{\mu_R^2}{\mu^2} \right)^{i\frac{\epsilon}{2}} S_\epsilon^i K^{b,(i)}(\epsilon) \tag{2.54}$$

with

$$\begin{aligned}
K^{b,(1)}(\epsilon) &= \frac{1}{\epsilon} \left\{ -2A_1^b \right\}, \quad K^{b,(2)}(\epsilon) = \frac{1}{\epsilon^2} \left\{ 2\beta_0 A_1^b \right\} + \frac{1}{\epsilon} \left\{ -A_2^b \right\}, \\
K^{b,(3)}(\epsilon) &= \frac{1}{\epsilon^3} \left\{ -\frac{8}{3}\beta_0^2 A_1^b \right\} + \frac{1}{\epsilon^2} \left\{ \frac{2}{3}\beta_1 A_1^b + \frac{8}{3}\beta_0 A_2^b \right\} + \frac{1}{\epsilon} \left\{ -\frac{2}{3}A_3^b \right\}. \tag{2.55}
\end{aligned}$$

Similarly upon solving the RGE 2.52 for  $G^b$ , we obtain

$$\begin{aligned}
G^b(\hat{a}_s, \frac{Q^2}{\mu_R^2}, \frac{\mu_R^2}{\mu^2}, \epsilon) &= G^b(a_s(\mu_R^2), \frac{Q^2}{\mu_R^2}, \epsilon) \\
&= G^b(a_s(Q^2), 1, \epsilon) + \int_{Q^2/\mu_R^2}^1 \frac{d\lambda^2}{\lambda^2} A^b(a_s(\lambda^2 \mu_R^2)) \\
&= G^b(a_s(Q^2), 1, \epsilon) + \sum_{i=1}^{\infty} S_\epsilon^i \hat{a}_s^i \left( \frac{\mu_R^2}{\mu^2} \right)^{i\frac{\epsilon}{2}} \left[ \left( \frac{Q^2}{\mu_R^2} \right)^{i\frac{\epsilon}{2}} - 1 \right] K^{b,(i)}(\epsilon). \tag{2.56}
\end{aligned}$$

Expanding the finite function  $G^b(a_s(Q^2), 1, \epsilon)$  in powers of  $a_s(Q^2)$  as

$$G^b(a_s(Q^2), 1, \epsilon) = \sum_{i=1}^{\infty} a_s^i(Q^2) G_i^b(\epsilon), \tag{2.57}$$

one finds that  $G_i^b$  can be expressed in terms of collinear  $B_i^q$  and soft  $f_i^q$  anomalous dimensions through the relation [81–83]

$$G_i^b(\epsilon) = 2(B_i^q - \gamma_i^b) + f_i^q + C_i^b + \sum_{k=1}^{\infty} \epsilon^k g_i^{b,k}. \tag{2.58}$$

Note that the single pole term of the form factor depends on three different anomalous dimensions, namely the collinear anomalous dimension  $B_i^q$ , anomalous dimension of the coupling constant  $\gamma_i^b$  and the soft anomalous dimension  $f_i^q$ .  $B_i^q$  can be obtained from the

$\delta(1-z)$  part of the diagonal splitting function known up to three loop level [38, 77] which are

$$\begin{aligned}
B_1^q &= 3C_F, \\
B_2^q &= \frac{1}{2} \left( C_F^2 \left\{ 3 - 24\zeta_2 + 48\zeta_3 \right\} + C_A C_F \left\{ \frac{17}{3} + \frac{88}{3}\zeta_2 - 24\zeta_3 \right\} + n_f T_F C_F \left\{ -\frac{4}{3} - \frac{32}{3}\zeta_2 \right\} \right), \\
B_3^q &= -16C_A^2 C_F \left\{ \frac{1}{8}\zeta_2^2 - \frac{281}{27}\zeta_2 + \frac{97}{9}\zeta_3 - \frac{5}{2}\zeta_5 + \frac{1657}{576} \right\} + 16C_A C_F^2 \left\{ -\frac{247}{60}\zeta_2^2 + \zeta_2\zeta_3 \right. \\
&\quad \left. - \frac{205}{24}\zeta_2 + \frac{211}{12}\zeta_3 + \frac{15}{2}\zeta_5 + \frac{151}{64} \right\} + 16C_A C_F n_f \left\{ \frac{1}{20}\zeta_2^2 - \frac{167}{54}\zeta_2 + \frac{25}{18}\zeta_3 + \frac{5}{4} \right\} \\
&\quad + 16C_F^3 \left\{ \frac{18}{5}\zeta_2^2 - 2\zeta_2\zeta_3 + \frac{9}{8}\zeta_2 + \frac{17}{4}\zeta_3 - 15\zeta_5 + \frac{29}{32} \right\} \\
&\quad - 16C_F^2 n_f \left\{ -\frac{29}{30}\zeta_2^2 - \frac{5}{12}\zeta_2 + \frac{17}{6}\zeta_3 + \frac{23}{16} \right\} - 16C_F n_f^2 \left\{ -\frac{5}{27}\zeta_2 + \frac{1}{9}\zeta_3 + \frac{17}{144} \right\}.
\end{aligned} \tag{2.59}$$

The  $f_i^q$  for  $i = 1, 2$  can be found in [81] and in [38] for  $i = 3$ . We list them below:

$$\begin{aligned}
f_1^q &= 0, \\
f_2^q &= C_A C_F \left\{ -\frac{22}{3}\zeta_2 - 28\zeta_3 + \frac{808}{27} \right\} + C_F n_f T_F \left\{ \frac{8}{3}\zeta_2 - \frac{224}{27} \right\}, \\
f_3^q &= C_A^2 C_F \left\{ \frac{352}{5}\zeta_2^2 + \frac{176}{3}\zeta_2\zeta_3 - \frac{12650}{81}\zeta_2 - \frac{1316}{3}\zeta_3 + 192\zeta_5 + \frac{136781}{729} \right\} \\
&\quad + C_A C_F n_f \left\{ -\frac{96}{5}\zeta_2^2 + \frac{2828}{81}\zeta_2 + \frac{728}{27}\zeta_3 - \frac{11842}{729} \right\} \\
&\quad + C_F^2 n_f \left\{ \frac{32}{5}\zeta_2^2 + 4\zeta_2 + \frac{304}{9}\zeta_3 - \frac{1711}{27} \right\} + C_F n_f^2 \left\{ -\frac{40}{27}\zeta_2 + \frac{112}{27}\zeta_3 - \frac{2080}{729} \right\}.
\end{aligned} \tag{2.60}$$

Since  $B_i^q$  and  $f_i^q$  are flavour independent, we have used  $B_i^b \equiv B_i^q$  and  $f_i^b \equiv f_i^q$  in  $G_i^b$ . The constants  $C_i^b$  are controlled by the beta function of the strong coupling constant through renormalization group invariance of the bare form factor:

$$C_1^b = 0, \quad C_2^b = -2\beta_0 g_1^{b,1}, \quad C_3^b = -2\beta_1 g_1^{b,1} - 2\beta_0 (g_2^{b,1} + 2\beta_0 g_1^{b,2}). \tag{2.61}$$

The coefficients  $g_i^{b,k}$  can be extracted from the finite part of the form factor. Up to two loop level, we use [19, 44, 45] and at three loop level the recent computation by Gehrmann and Kara [57] enable us to compute the relevant  $g_3^{b,1}$  in [59] where  $g_3^{b,1}$  was already used to obtain threshold correction to inclusive Higgs production in bottom anti-bottom annihilation process:

$$g_1^{b,1} = C_F \left\{ -2 + \zeta_2 \right\}, \quad g_1^{b,2} = C_F \left\{ 2 - \frac{7}{3}\zeta_3 \right\}, \quad g_1^{b,3} = C_F \left\{ -2 + \frac{1}{4}\zeta_2 + \frac{47}{80}\zeta_2^2 \right\},$$

$$\begin{aligned}
g_2^{b,1} &= C_{Fnf} \left\{ \frac{616}{81} + \frac{10}{9} \zeta_2 - \frac{8}{3} \zeta_3 \right\} + C_F C_A \left\{ -\frac{2122}{81} - \frac{103}{9} \zeta_2 + \frac{88}{5} \zeta_2^2 + \frac{152}{3} \zeta_3 \right\} \\
&\quad + C_F^2 \left\{ 8 + 32 \zeta_2 - \frac{88}{5} \zeta_2^2 - 60 \zeta_3 \right\}, \\
g_2^{b,2} &= C_{Fnf} \left\{ \frac{7}{12} \zeta_2^2 - \frac{55}{27} \zeta_2 + \frac{130}{27} \zeta_3 - \frac{3100}{243} \right\} + C_A C_F \left\{ -\frac{365}{24} \zeta_2^2 + \frac{89}{3} \zeta_2 \zeta_3 + \frac{1079}{54} \zeta_2 \right. \\
&\quad \left. - \frac{2923}{27} \zeta_3 - 51 \zeta_5 + \frac{9142}{243} \right\} + C_F^2 \left\{ \frac{96}{5} \zeta_2^2 - 28 \zeta_2 \zeta_3 - 44 \zeta_2 + 116 \zeta_3 + 12 \zeta_5 - 24 \right\}, \\
g_3^{b,1} &= C_A^2 C_F \left\{ -\frac{6152}{63} \zeta_2^3 + \frac{2738}{9} \zeta_2^2 + \frac{976}{9} \zeta_2 \zeta_3 - \frac{342263}{486} \zeta_2 - \frac{1136}{3} \zeta_3^2 + \frac{19582}{9} \zeta_3 \right. \\
&\quad \left. + \frac{1228}{3} \zeta_5 + \frac{4095263}{8748} \right\} + C_A C_F^2 \left\{ -\frac{15448}{105} \zeta_2^3 - \frac{3634}{45} \zeta_2^2 - \frac{2584}{3} \zeta_2 \zeta_3 + \frac{13357}{9} \zeta_2 \right. \\
&\quad \left. + 296 \zeta_3^2 - \frac{11570}{9} \zeta_3 - \frac{1940}{3} \zeta_5 - \frac{613}{3} \right\} + C_A C_{Fnf} \left\{ -\frac{1064}{45} \zeta_2^2 + \frac{392}{9} \zeta_2 \zeta_3 + \frac{44551}{243} \zeta_2 \right. \\
&\quad \left. - \frac{41552}{81} \zeta_3 - 72 \zeta_5 - \frac{6119}{4374} \right\} + C_F^2 n_f \left\{ \frac{772}{45} \zeta_2^2 - \frac{152}{3} \zeta_2 \zeta_3 - \frac{3173}{18} \zeta_2 + \frac{15956}{27} \zeta_3 - \frac{368}{3} \zeta_5 \right. \\
&\quad \left. + \frac{32899}{324} \right\} + C_F n_f^2 \left\{ -\frac{40}{9} \zeta_2^2 - \frac{892}{81} \zeta_2 + \frac{320}{81} \zeta_3 - \frac{27352}{2187} \right\} + C_F^3 \left\{ \frac{21584}{105} \zeta_2^3 - \frac{1644}{5} \zeta_2^2 \right. \\
&\quad \left. + 624 \zeta_2 \zeta_3 - 275 \zeta_2 + 48 \zeta_3^2 - 2142 \zeta_3 + 1272 \zeta_5 + 603 \right\}. \tag{2.62}
\end{aligned}$$

Using the expressions for  $K^b$  and  $G^b$  given in Eq. 2.54 and Eq. 2.58, respectively, we obtain the renormalized form factor up to order  $\mathcal{O}(a_s^3)$  as

$$\begin{aligned}
\ln |\hat{F}^b|^2(Q^2, \epsilon) &= a_s(q^2) \left[ \frac{1}{\epsilon^2} \left( -4A_1^q \right) + \frac{1}{\epsilon} \left( 2f_1^q + 4B_1^q - 4\gamma_0^b \right) + \left( 2g_1^{b,1} + 3\zeta_2 A_1^q \right) \right] \\
&\quad + a_s^2(q^2) \left[ \frac{1}{\epsilon^3} \left( -6\beta_0 A_1^q \right) + \frac{1}{\epsilon^2} \left( -A_2^q + 2\beta_0 \left( f_1^q + 2B_1^q - 2\gamma_0^b \right) \right) + \frac{1}{\epsilon} \left( f_2^q \right. \right. \\
&\quad \left. \left. + 2B_2^q - 2\gamma_1^b \right) + \left( g_2^{b,1} + 2\beta_0 g_1^{b,2} + 3\zeta_2 A_2^q + 3\zeta_2 \beta_0 \left( f_1^q + 2B_1^q - 2\gamma_0^b \right) \right) \right] \\
&\quad + a_s^3(q^2) \left[ \frac{1}{\epsilon^4} \left( -\frac{88}{9} \beta_0^2 A_1^q \right) + \frac{1}{\epsilon^3} \left( -\frac{32}{9} \beta_1 A_1^q - \frac{20}{9} \beta_0 A_2^q + \frac{8}{3} \beta_0^2 \left( f_1^q + 2B_1^q \right. \right. \right. \\
&\quad \left. \left. - 2\gamma_0^b \right) \right) + \frac{1}{\epsilon^2} \left( -\frac{4}{9} A_3^q + \frac{4}{3} \beta_1 \left( f_1^q + 2B_1^q - 2\gamma_0^b \right) + \frac{4}{3} \beta_0 \left( f_2^q + 2B_2^q - 2\gamma_1^b \right) \right) \\
&\quad + \frac{1}{\epsilon} \left( \frac{2}{3} f_3^q + \frac{4}{3} B_3^q - \frac{4}{3} \gamma_2^b \right) + \left( \frac{2}{3} g_3^{b,1} + \frac{4}{3} \beta_1 g_1^{b,2} + \frac{4}{3} \beta_0 g_2^{b,2} + \frac{8}{3} \beta_0^2 g_1^{b,3} + 3\zeta_2 A_3^q \right. \\
&\quad \left. + 3\zeta_2 \beta_1 \left( f_1^q + 2B_1^q - 2\gamma_0^b \right) + 6\zeta_2 \beta_0 \left( f_2^q + 2B_2^q - 2\gamma_1^b \right) \right)
\end{aligned}$$



$$\left. - 12\zeta_2\beta_0^2g_1^{b,1} - 3\zeta_2^2\beta_0^2A_1^q \right]. \quad (2.63)$$

Note that the poles of  $\ln|\hat{F}^b|^2$  are fully controlled by the universal anomalous dimensions  $A^q, \gamma^b, B^q$  and  $f^q$  while the constant terms require vertex dependent constants  $g_i^{b,k}$ .

In  $\overline{\text{MS}}$  scheme, the mass factorization kernels  $\Gamma_{bb}(\hat{a}_s, \mu^2, \mu_F^2, z_j, \epsilon)$  remove the collinear singularities which arise due to massless partons. These kernels satisfy the following RG equation :

$$\mu_F^2 \frac{d}{d\mu_F^2} \Gamma_{bb}(z_j, \mu_F^2, \epsilon) = \frac{1}{2} \sum_c P_{bc}(z_j, \mu_F^2) \otimes \Gamma_{cb}(z_j, \mu_F^2, \epsilon), \quad (2.64)$$

where  $P_{bc}(z_j, \mu_F^2)$  are AP splitting functions. We can expand the  $P_{bc}(z_j, \mu_F^2)$  in powers of  $a_s$  as

$$P_{bc}(z_j, \mu_F^2) = \sum_{i=1}^{\infty} a_s^i(\mu_F^2) P_{bc}^{(i-1)}(z_j). \quad (2.65)$$

The off diagonal splitting functions are regular as  $z_j \rightarrow 1$ . The diagonal ones contain in addition distributions such as  $\delta(1 - z_j)$  and  $\mathcal{D}_0$  multiplied by the universal anomalous dimensions  $B_i^q$  and  $A_i^q$ , respectively:

$$P_{bb}^{(i)}(z_j) = 2 \left( B_{i+1}^b \delta(1 - z_j) + A_{i+1}^b \mathcal{D}_0 \right) + P_{reg,bb}^{(i)}(z_j). \quad (2.66)$$

As we are interested in results from the threshold region, we can ignore all the non-diagonal splitting functions and also the regular part  $P_{reg,bb}^{(i)}$  arising from the diagonal terms. Hence, the solution to Eq. 2.64 takes the following form:

$$\begin{aligned} \ln \Gamma_{bb}(z_j, \mu_F^2) = & a_s(\mu_F^2) \left[ \delta(1 - z_j) \left( \frac{1}{\epsilon} (2B_1^q) \right) + \mathcal{D}_0 \left( \frac{1}{\epsilon} (2A_1^q) \right) \right] \\ & + a_s^2(\mu_F^2) \left[ \delta(1 - z_j) \left( \frac{1}{\epsilon^2} (2\beta_0 B_1^q) + \frac{1}{\epsilon} (B_2^q) \right) + \mathcal{D}_0 \left( \frac{1}{\epsilon^2} (2\beta_0 A_1^q) + \frac{1}{\epsilon} (A_2^q) \right) \right] \\ & + a_s^3(\mu_F^2) \left[ \delta(1 - z_j) \left( \frac{1}{\epsilon^3} \left( \frac{8}{3} \beta_0^2 B_1^q \right) + \frac{1}{\epsilon^2} \left( \frac{4}{3} \beta_1 B_1^q + \frac{4}{3} \beta_0 B_2^q \right) + \frac{1}{\epsilon} \left( \frac{2}{3} B_3^q \right) \right) \right. \\ & \left. + \mathcal{D}_0 \left( \frac{1}{\epsilon^3} \left( \frac{8}{3} \beta_0^2 A_1^q \right) + \frac{1}{\epsilon^2} \left( \frac{4}{3} \beta_1 A_1^q + \frac{4}{3} \beta_0 A_2^q \right) + \frac{1}{\epsilon} \left( \frac{2}{3} A_3^q \right) \right) \right]. \quad (2.67) \end{aligned}$$

Finally, we need to determine the soft distribution function  $\Phi_d^b(\hat{a}_s, q^2, \mu^2, z_1, z_2, \epsilon)$  in  $\Delta_{d,b}^{\text{SV}}$ . Its most general form can be systematically constructed if  $\Phi_d^b$  also satisfies a differential equation similar to the form factor. It is indeed the case because the  $q^2$  dependence and pole structure of  $\Phi_d^b$  have to be similar to those of  $\ln|\hat{F}^b|^2$  in order to obtain finite distribution  $\Psi$  in the limit  $\epsilon \rightarrow 0$  [44, 45]. Hence, we propose that  $\Phi_d^b$  satisfies

$$q^2 \frac{d}{dq^2} \Phi_d^b = \frac{1}{2} \left[ \overline{K}_d^b(\hat{a}_s, \frac{\mu_R^2}{\mu^2}, z_1, z_2, \epsilon) + \overline{G}_d^b(\hat{a}_s, \frac{q^2}{\mu_R^2}, \frac{\mu_R^2}{\mu^2}, z_1, z_2, \epsilon) \right]. \quad (2.68)$$

It is natural to move all the singular terms in  $\epsilon$  of  $\Phi_d^b$  to  $\overline{K}_d^b$  and keep  $\overline{G}_d^b$  finite as  $\epsilon \rightarrow 0$  similar to  $K_d^b$  and  $G_d^b$  of the logarithm of the form factor,  $\ln \hat{F}^b$ . The RG invariance of  $\Phi_d^b(\hat{a}_s, q^2, \mu^2, z_1, z_2, \epsilon)$  leads to

$$\mu_R^2 \frac{d}{d\mu_R^2} \Phi_d^b(\hat{a}_s, q^2, \mu^2, z_1, z_2, \epsilon) = 0 \quad (2.69)$$

and consequently

$$\mu_R^2 \frac{d}{d\mu_R^2} \overline{K}_d^b = -\mu_R^2 \frac{d}{d\mu_R^2} \overline{G}_d^b = -\delta(1-z_1)\delta(1-z_2)a_s(\mu_R^2)\overline{A}^q. \quad (2.70)$$

The right hand side of the above equation is proportion to  $\delta(1-z_1)\delta(1-z_2)$  as the most singular terms resulting from  $\overline{K}_d^b$  should cancel with those from the form factor contribution which is proportional to only pure delta functions. To make the  $\Delta_{d,b}^{\text{SV}}$  finite, the poles from  $\Phi_d^b(\hat{a}_s, q^2, \mu^2, z_1, z_2, \epsilon)$  have to cancel those coming from  $\hat{F}^b$  and  $\Gamma_{bb}$ . Hence the constants  $\overline{A}^q$  should satisfy

$$\overline{A}^q = -A^q. \quad (2.71)$$

The RGE 2.70 for  $\overline{G}_d^b$  can be solved using the above relation to get

$$\begin{aligned} \overline{G}_d^b \left( \hat{a}_s, \frac{q^2}{\mu_R^2}, \frac{\mu_R^2}{\mu^2}, z_1, z_2, \epsilon \right) &= \overline{G}_d^b \left( a_s(\mu_R^2), \frac{q^2}{\mu_R^2}, z_1, z_2, \epsilon \right) \\ &= \overline{G}_d^b(a_s(q^2), 1, z_1, z_2, \epsilon) - \delta(1-z_1)\delta(1-z_2) \int_{\frac{q^2}{\mu_R^2}}^1 \frac{d\lambda^2}{\lambda^2} A^q(a_s(\lambda^2 \mu_R^2)). \end{aligned} \quad (2.72)$$

With these solutions, it is now straightforward to solve the above differential equations 2.68 for  $\Phi_d^b$  to get

$$\begin{aligned} \Phi_d^b &= \Phi_d^b(\hat{a}_s, q^2(1-z_1)(1-z_2), \mu^2, \epsilon) \\ &= \sum_{i=1}^{\infty} \hat{a}_s^i \left( \frac{q^2(1-z_1)(1-z_2)}{\mu^2} \right)^{i\frac{\epsilon}{2}} S_\epsilon^i \left( \frac{(i\epsilon)^2}{4(1-z_1)(1-z_2)} \right) \hat{\phi}_d^{b,(i)}(\epsilon), \end{aligned} \quad (2.73)$$

where,

$$\hat{\phi}_d^{b,(i)}(\epsilon) = \frac{1}{i\epsilon} \left[ \overline{K}_d^{b,(i)}(\epsilon) + \overline{G}_d^{b,(i)}(\epsilon) \right]. \quad (2.74)$$

The form of  $z_j$  dependence part of the solution in the above solution is inspired by our one loop computation in the previous section and it can be justified from the factorization property of the QCD amplitudes and the corresponding partonic cross sections. The constants  $\overline{K}_d^{b,(i)}(\epsilon)$  are determined by expanding  $\overline{K}_d^b$  in powers of  $\hat{a}_s$  as follows

$$\overline{K}_d^b \left( \hat{a}_s, \frac{\mu_R^2}{\mu^2}, z_1, z_2, \epsilon \right) = \delta(1-z_1)\delta(1-z_2) \sum_{i=1}^{\infty} \hat{a}_s^i \left( \frac{\mu_R^2}{\mu^2} \right)^{i\frac{\epsilon}{2}} S_\epsilon^i \overline{K}_d^{b,(i)}(\epsilon) \quad (2.75)$$

and solving the RGE 2.70 for  $\overline{K}_d^b$ . The constants  $\overline{K}_d^{b,(i)}(\epsilon)$  are identical to  $\overline{K}^{b,(i)}(\epsilon)$  given in [44, 45].  $\overline{G}_d^{b,(i)}(\epsilon)$  are related to the finite functions  $\overline{G}_d^b(a_s(q^2), 1, z_1, z_2, \epsilon)$ . In terms of renormalized coupling constant, we find

$$\sum_{i=1}^{\infty} \hat{a}_s^i \left( \frac{q^2(1-z_1)(1-z_2)}{\mu^2} \right)^{i\frac{\epsilon}{2}} S_\epsilon^i \overline{G}_d^{b,(i)}(\epsilon) = \sum_{i=1}^{\infty} a_s^i (q^2(1-z_1)(1-z_2)) \overline{\mathcal{G}}_{d,i}^b(\epsilon) \quad (2.76)$$

where the constants  $\overline{\mathcal{G}}_{d,i}^b(\epsilon)$  are flavour independent and they satisfy the following structure similar to  $G_i^b(\epsilon)$  of the form factor, i.e.,

$$\overline{\mathcal{G}}_{d,i}^b(\epsilon) = -f_i^q + \mathcal{C}_{d,i}^b + \sum_{k=1}^{\infty} \epsilon^k \overline{\mathcal{G}}_{d,i}^{b,k}, \quad (2.77)$$

where

$$\mathcal{C}_{d,1}^b = 0, \quad \mathcal{C}_{d,2}^b = -2\beta_0 \overline{\mathcal{G}}_{d,1}^{b,1}, \quad \mathcal{C}_{d,3}^b = -2\beta_1 \overline{\mathcal{G}}_{d,1}^{b,1} - 2\beta_0 (\overline{\mathcal{G}}_{d,2}^{b,1} + 2\beta_0 \overline{\mathcal{G}}_{d,1}^{b,2}). \quad (2.78)$$

Using  $\overline{K}_d^{b,(i)}$  from Eq. 2.75 and  $\overline{G}_d^{b,(i)}$  from Eq. 2.76 and using Eq. 2.26, we find that the soft distribution function up to third order in  $a_s(q^2)$  takes the form

$$\begin{aligned} \Phi_d^b = a_s(q^2) & \left[ \delta(1-z_1)\delta(1-z_2) \left( \frac{1}{\epsilon^2} (2A_1^q) + \frac{1}{\epsilon} (-f_1^q + \overline{\mathcal{G}}_{d,1}^{q,1}) \right) + \mathcal{D}_0 \delta(1-z_2) \left( \frac{1}{\epsilon} (A_1^q) \right. \right. \\ & + \left. \left. (-\frac{1}{2}f_1^q) \right) + \mathcal{D}_0 \overline{\mathcal{D}}_0 \left( \frac{1}{2}A_1^q \right) + \mathcal{D}_1 \delta(1-z_2) \left( \frac{1}{2}A_1^q \right) + \overline{\mathcal{D}}_0 \delta(1-z_1) \left( \frac{1}{\epsilon} (A_1^q) + (-\frac{1}{2}f_1^q) \right) \right. \\ & + \left. \overline{\mathcal{D}}_1 \delta(1-z_1) \left( \frac{1}{2}A_1^q \right) \right] + a_s^2(q^2) \left[ \delta(1-z_1)\delta(1-z_2) \left( \frac{1}{\epsilon^3} (3\beta_0 A_1^q) + \frac{1}{\epsilon^2} \left( \frac{1}{2}A_2^q - \beta_0 f_1^q \right) \right. \right. \\ & + \left. \frac{1}{\epsilon} \left( -\frac{1}{2}f_2^q + \left( \frac{1}{2}\overline{\mathcal{G}}_{d,2}^{q,1} + \beta_0 \overline{\mathcal{G}}_{d,1}^{q,2} \right) \right) + \mathcal{D}_0 \delta(1-z_2) \left( \frac{1}{\epsilon^2} (\beta_0 A_1^q) + \frac{1}{\epsilon} \left( \frac{1}{2}A_2^q \right) \right. \right. \\ & + \left. \left. \left( -\frac{1}{2}f_2^q - \beta_0 \overline{\mathcal{G}}_{d,1}^{q,1} \right) \right) + \mathcal{D}_0 \overline{\mathcal{D}}_0 \left( \frac{1}{2}A_2^q + \frac{1}{2}\beta_0 f_1^q \right) + \mathcal{D}_0 \overline{\mathcal{D}}_1 \left( -\frac{1}{2}\beta_0 A_1^q \right) \right. \\ & + \left. \mathcal{D}_1 \delta(1-z_2) \left( \frac{1}{2}A_2^q + \frac{1}{2}\beta_0 f_1^q \right) + \mathcal{D}_1 \overline{\mathcal{D}}_0 \left( -\frac{1}{2}\beta_0 A_1^q \right) + \mathcal{D}_2 \delta(1-z_2) \left( -\frac{1}{4}\beta_0 A_1^q \right) \right. \\ & + \left. \overline{\mathcal{D}}_0 \delta(1-z_1) \left( \frac{1}{\epsilon^2} (\beta_0 A_1^q) + \frac{1}{\epsilon} \left( \frac{1}{2}A_2^q \right) + \left( -\frac{1}{2}f_2^q - \beta_0 \overline{\mathcal{G}}_{d,1}^{q,1} \right) \right) \right. \\ & + \left. \overline{\mathcal{D}}_1 \delta(1-z_1) \left( \frac{1}{2}A_2^q + \frac{1}{2}\beta_0 f_1^q \right) + \overline{\mathcal{D}}_2 \delta(1-z_1) \left( -\frac{1}{4}\beta_0 A_1^q \right) \right] \\ & + a_s^3(q^2) \left[ \delta(1-z_1)\delta(1-z_2) \left( \frac{1}{\epsilon^4} \left( \frac{44}{9}\beta_0^2 A_1^q \right) + \frac{1}{\epsilon^3} \left( \frac{16}{9}\beta_1 A_1^q + \frac{10}{9}\beta_0 A_2^q - \frac{4}{3}\beta_0^2 f_1^q \right) \right. \right. \\ & + \left. \frac{1}{\epsilon^2} \left( \frac{2}{9}A_3^q - \frac{2}{3}\beta_1 f_1^q - \frac{2}{3}\beta_0 f_2^q \right) - \frac{1}{\epsilon} \left( \frac{1}{3}f_3^q \right) + \left( \frac{1}{3}\overline{\mathcal{G}}_{d,3}^{q,1} + \frac{2}{3}\beta_1 \overline{\mathcal{G}}_{d,1}^{q,2} + \frac{2}{3}\beta_0 \overline{\mathcal{G}}_{d,2}^{q,2} + \frac{4}{3}\beta_0^2 \overline{\mathcal{G}}_{d,1}^{q,3} \right) \right] \end{aligned}$$

$$\begin{aligned}
& + \mathcal{D}_0 \delta(1 - z_2) \left( \frac{1}{\epsilon^3} \left( \frac{4}{3} \beta_0^2 A_1^q \right) + \frac{1}{\epsilon^2} \left( \frac{2}{3} \beta_1 A_1^q + \frac{2}{3} \beta_0 A_2^q \right) + \frac{1}{\epsilon} \left( \frac{1}{3} A_3^q \right) + \left( -\frac{1}{2} f_3^q - \beta_1 \bar{\mathcal{G}}_{d,1}^{q,1} \right. \right. \\
& \left. \left. - \beta_0 \bar{\mathcal{G}}_{d,2}^{q,1} - 2\beta_0^2 \bar{\mathcal{G}}_{d,1}^{q,2} \right) \right) + \mathcal{D}_0 \bar{\mathcal{D}}_0 \left( \frac{1}{2} A_3^q + \frac{1}{2} \beta_1 f_1^q + \beta_0 f_2^q + 2\beta_0^2 \bar{\mathcal{G}}_{d,1}^{q,1} \right) + \mathcal{D}_0 \bar{\mathcal{D}}_1 \left( -\frac{1}{2} \beta_1 A_1^q \right. \\
& \left. - \beta_0 A_2^q - \beta_0^2 f_1^q \right) + \mathcal{D}_0 \bar{\mathcal{D}}_2 \left( \frac{1}{2} \beta_0^2 A_1^q \right) + \mathcal{D}_1 \delta(1 - z_2) \left( \frac{1}{2} A_3^q + \frac{1}{2} \beta_1 f_1^q + \beta_0 f_2^q + 2\beta_0^2 \bar{\mathcal{G}}_{d,1}^{q,1} \right) \\
& + \mathcal{D}_1 \bar{\mathcal{D}}_0 \left( -\frac{1}{2} \beta_1 A_1^q - \beta_0 A_2^q - \beta_0^2 f_1^q \right) + \mathcal{D}_1 \bar{\mathcal{D}}_1 \left( \beta_0^2 A_1^q \right) \\
& + \mathcal{D}_2 \delta(1 - z_2) \left( -\frac{1}{4} \beta_1 A_1^q - \frac{1}{2} \beta_0 A_2^q - \frac{1}{2} \beta_0^2 f_1^q \right) + \mathcal{D}_2 \bar{\mathcal{D}}_0 \left( \frac{1}{2} \beta_0^2 A_1^q \right) \\
& + \mathcal{D}_3 \delta(1 - z_2) \left( \frac{1}{6} \beta_0^2 A_1^q \right) + \bar{\mathcal{D}}_0 \delta(1 - z_1) \left( \frac{1}{\epsilon^3} \left( \frac{4}{3} \beta_0^2 A_1^q \right) + \frac{1}{\epsilon^2} \left( \frac{2}{3} \beta_1 A_1^q + \frac{2}{3} \beta_0 A_2^q \right) \right. \\
& \left. + \frac{1}{\epsilon} \left( \frac{1}{3} A_3^q \right) + \left( -\frac{1}{2} f_3^q - \beta_1 \bar{\mathcal{G}}_{d,1}^{q,1} - \beta_0 \bar{\mathcal{G}}_{d,2}^{q,1} - 2\beta_0^2 \bar{\mathcal{G}}_{d,1}^{q,2} \right) \right) \\
& + \bar{\mathcal{D}}_1 \delta(1 - z_1) \left( \frac{1}{2} A_3^q + \frac{1}{2} \beta_1 f_1^q + \beta_0 f_2^q + 2\beta_0^2 \bar{\mathcal{G}}_{d,1}^{q,1} \right) \\
& + \bar{\mathcal{D}}_2 \delta(1 - z_1) \left( -\frac{1}{4} \beta_1 A_1^q - \frac{1}{2} \beta_0 A_2^q - \frac{1}{2} \beta_0^2 f_1^q \right) \\
& \left. + \bar{\mathcal{D}}_3 \delta(1 - z_1) \left( \frac{1}{6} \beta_0^2 A_1^q \right) \right]. \tag{2.79}
\end{aligned}$$

In the above expression, we have used  $\bar{\mathcal{G}}_{d,i}^{b,k} = \bar{\mathcal{G}}_{d,i}^{q,k}$ , being flavour independent. The soft distribution function depends in addition to the universal anomalous dimensions  $A_i^q, B_i^q, \gamma_i^q$  and  $f_i^q$ , the constants  $\bar{\mathcal{G}}_{d,i}^{q,k}$  which need to be determined. At  $\mathcal{O}(a_s)$  level  $\bar{\mathcal{G}}_{d,1}^{q,1}, \bar{\mathcal{G}}_{d,2}^{q,1}, \bar{\mathcal{G}}_{d,3}^{q,1}$ , at  $\mathcal{O}(a_s^2)$   $\bar{\mathcal{G}}_{d,1}^{q,2}, \bar{\mathcal{G}}_{d,2}^{q,2}$  and at  $\mathcal{O}(a_s^3)$   $\bar{\mathcal{G}}_{d,1}^{q,3}$  are needed to obtain  $\Phi_d^b$ . We achieve this using the following identity:

$$\int_0^1 dx_1^0 \int_0^1 dx_2^0 (x_1^0 x_2^0)^{N-1} \frac{d\sigma^b}{dy} = \int_0^1 d\tau \tau^{N-1} \sigma^b, \tag{2.80}$$

where  $\sigma^b$  is known to NNLO level [19] exactly and to N<sup>3</sup>LO level in the threshold limit [59]. In large N limit *i.e.*  $N \rightarrow \infty$  the above Eq. 2.80 relates  $\hat{\phi}_d^{q,(i)}(\epsilon)$  to  $\hat{\phi}_d^{b,(i)}(\epsilon)$  that appears in inclusive threshold corrections to Drell-Yan process (see [44, 45, 51, 59]) as follows

$$\hat{\phi}_d^{b,(i)}(\epsilon) = \frac{\Gamma(1+i\epsilon)}{\Gamma^2(1+i\frac{\epsilon}{2})} \hat{\phi}_d^{q,(i)}(\epsilon) \tag{2.81}$$

and

$$\hat{\phi}_d^{b,(i)}(\epsilon) = \hat{\phi}_d^{q,(i)}(\epsilon) \tag{2.82}$$

since  $\hat{\phi}^{q,(i)}(\epsilon)$  is flavour independent. Hence

$$\hat{\phi}_d^{b,(i)}(\epsilon) = \hat{\phi}_d^{q,(i)}(\epsilon) \quad (2.83)$$

and all the relevant constants  $\bar{\mathcal{G}}_{d,i}^{q,k}$  required for threshold prediction up to  $\mathcal{O}(a_s^3)$  can be obtained from  $\bar{\mathcal{G}}_i^{q,k}$  which are analogous to these factors appeared in the computation of inclusive threshold cross-section to Drell-Yan process. The relevant  $\bar{\mathcal{G}}_i^{q,k}$ 's at  $\mathcal{O}(a_s)$  and  $\mathcal{O}(a_s^2)$  [44, 45] are

$$\begin{aligned} \bar{\mathcal{G}}_1^{q,1} &= C_F(-3\zeta_2), \\ \bar{\mathcal{G}}_1^{q,2} &= C_F\left(\frac{7}{3}\zeta_3\right), \\ \bar{\mathcal{G}}_1^{q,3} &= C_F\left(-\frac{3}{16}\zeta_2^2\right), \\ \bar{\mathcal{G}}_2^{q,1} &= C_F n_f \left(-\frac{328}{81} + \frac{70}{9}\zeta_2 + \frac{32}{3}\zeta_3\right) + C_A C_F \left(\frac{2428}{81} - \frac{469}{9}\zeta_2 + 4\zeta_2^2 - \frac{176}{3}\zeta_3\right), \\ \bar{\mathcal{G}}_2^{q,2} &= C_A C_F \left(\frac{11}{40}\zeta_2^2 - \frac{203}{3}\zeta_2\zeta_3 + \frac{1414}{27}\zeta_2 + \frac{2077}{27}\zeta_3 + 43\zeta_5 - \frac{7288}{243}\right) \\ &\quad + C_F n_f \left(-\frac{1}{20}\zeta_2^2 - \frac{196}{27}\zeta_2 - \frac{310}{27}\zeta_3 + \frac{976}{243}\right) \end{aligned}$$

and at  $\mathcal{O}(a_s^3)$  [51]

$$\begin{aligned} \bar{\mathcal{G}}_3^{q,1} &= C_F \left\{ C_A^2 \left( \frac{152}{63}\zeta_2^3 + \frac{1964}{9}\zeta_2^2 + \frac{11000}{9}\zeta_2\zeta_3 - \frac{765127}{486}\zeta_2 + \frac{536}{3}\zeta_3^2 - \frac{59648}{27}\zeta_3 \right. \right. \\ &\quad \left. \left. - \frac{1430}{3}\zeta_5 + \frac{7135981}{8748} \right) + C_A n_f \left( -\frac{532}{9}\zeta_2^2 - \frac{1208}{9}\zeta_2\zeta_3 + \frac{105059}{243}\zeta_2 + \frac{45956}{81}\zeta_3 \right. \right. \\ &\quad \left. \left. + \frac{148}{3}\zeta_5 - \frac{716509}{4374} \right) + C_F n_f \left( \frac{152}{15}\zeta_2^2 - 88\zeta_2\zeta_3 + \frac{605}{6}\zeta_2 + \frac{2536}{27}\zeta_3 + \frac{112}{3}\zeta_5 \right. \right. \\ &\quad \left. \left. - \frac{42727}{324} \right) + n_f^2 \left( \frac{32}{9}\zeta_2^2 - \frac{1996}{81}\zeta_2 - \frac{2720}{81}\zeta_3 + \frac{11584}{2187} \right) \right\}. \quad (2.84) \end{aligned}$$

These lead to the following expressions of  $\bar{\mathcal{G}}_{d,i}^{q,k}$ 's at  $\mathcal{O}(a_s)$ ,  $\mathcal{O}(a_s^2)$  [66] and  $\mathcal{O}(a_s^3)$  [67] :

$$\begin{aligned} \bar{\mathcal{G}}_{d,1}^{q,1} &= -C_F\zeta_2, \quad \bar{\mathcal{G}}_{d,2}^{q,1} = C_F \left\{ \frac{1}{3}\zeta_3 \right\}, \quad \bar{\mathcal{G}}_{d,3}^{q,1} = C_F \left\{ \frac{1}{80}\zeta_2^2 \right\}, \\ \bar{\mathcal{G}}_{d,1}^{q,2} &= C_A C_F \left\{ -4\zeta_2^2 - \frac{67}{3}\zeta_2 - \frac{44}{3}\zeta_3 + \frac{2428}{81} \right\} + C_F n_f \left\{ \frac{8}{3}\zeta_3 + \frac{10}{3}\zeta_2 - \frac{328}{81} \right\}, \\ \bar{\mathcal{G}}_{d,2}^{q,2} &= C_A C_F \left\{ -\frac{319}{120}\zeta_2^2 - \frac{71}{3}\zeta_2\zeta_3 + \frac{202}{9}\zeta_2 + \frac{469}{27}\zeta_3 + 43\zeta_5 - \frac{7288}{243} \right\} \\ &\quad + C_F n_f \left\{ \frac{29}{60}\zeta_2^2 - \frac{28}{9}\zeta_2 - \frac{70}{27}\zeta_3 + \frac{976}{243} \right\}, \\ \bar{\mathcal{G}}_{d,1}^{q,3} &= C_A^2 C_F \left\{ \frac{17392}{315}\zeta_2^3 + \frac{1538}{45}\zeta_2^2 + \frac{4136}{9}\zeta_2\zeta_3 - \frac{379417}{486}\zeta_2 + \frac{536}{3}\zeta_3^2 - 936\zeta_3 \right\} \end{aligned}$$

$$\begin{aligned}
& -\frac{1430}{3}\zeta_5 + \frac{7135981}{8748} \Big\} + C_A C_F n_f \left\{ -\frac{1372}{45}\zeta_2^2 - \frac{392}{9}\zeta_2\zeta_3 + \frac{51053}{243}\zeta_2 \right. \\
& + \frac{12356}{81}\zeta_3 + \frac{148}{3}\zeta_5 - \frac{716509}{4374} \Big\} + C_F n_f^2 \left\{ \frac{152}{45}\zeta_2^2 - \frac{316}{27}\zeta_2 - \frac{320}{81}\zeta_3 + \frac{11584}{2187} \right\} \\
& + C_F^2 n_f \left\{ \frac{152}{15}\zeta_2^2 - 40\zeta_2\zeta_3 + \frac{275}{6}\zeta_2 + \frac{1672}{27}\zeta_3 + \frac{112}{3}\zeta_5 - \frac{42727}{324} \right\}. \tag{2.85}
\end{aligned}$$

With all these information available at hand, it is now straightforward to obtain threshold corrections to rapidity distribution of Higgs boson in the bottom quark annihilation processes. We substitute Eq. 2.50, 2.63, 2.67, 2.79 in Eq. 2.40 to obtain  $\Psi_d^b(\epsilon)$ . Since all the UV and IR singularities cancel among various terms, we can set  $\epsilon = 0$  in the the distribution  $\Psi_d^b(\epsilon)$  to obtain  $\Delta_{d,b}^{\text{SV}}$ . Expanding the finite distribution  $\Psi_d^b(\epsilon = 0)$  in Eq. 2.38 in terms of convolutions Eq. 2.39 and performing all those convolutions using the formula given in Eq. 52 of [45], we obtain  $\Delta_b^{\text{SV},(i)}$  defined by

$$\Delta_{d,b}^{\text{SV}}(z_1, z_2, q^2, \mu_R^2, \mu_F^2) = \sum_{i=0}^{\infty} a_s^i(q^2) \Delta_{d,b}^{\text{SV},(i)}(z_1, z_2, q^2, \mu_R^2, \mu_F^2) \tag{2.86}$$

We present below our results for  $\Delta_{d,b}^{\text{SV},(i)}$  up to N<sup>3</sup>LO level in terms of of the constants  $A_j^q$ ,  $B_j^q$ ,  $f_j^q$ ,  $\gamma_j^b$ ,  $\beta_j$ ,  $g_j^{b,k}$  and  $\bar{\mathcal{G}}_{d,j}^{q,k}$ :

$$\begin{aligned}
\Delta_{d,b}^{\text{SV},(1)} &= \delta(1-z_1)\delta(1-z_2) \left[ \bar{\mathcal{G}}_{d,1}^{q,1} + g_1^{b,1} + \frac{3}{2}\zeta_2 A_1^q \right] + \mathcal{D}_0 \delta(1-z_2) \left[ -f_1^q \right] \\
&+ \mathcal{D}_0 \bar{\mathcal{D}}_0 \left[ \frac{1}{2} A_1^q \right] + \mathcal{D}_1 \delta(1-z_2) \left[ A_1^q \right] + \left\{ z_1 \leftrightarrow z_2 \right\} \\
\Delta_{d,b}^{\text{SV},(2)} &= \delta(1-z_1)\delta(1-z_2) \left[ \frac{1}{2} \bar{\mathcal{G}}_{d,2}^{q,1} + \bar{\mathcal{G}}_{d,1}^{q,1^2} + \frac{1}{2} g_2^{b,1} + 2g_1^{b,1} \bar{\mathcal{G}}_{d,1}^{q,1} + g_1^{b,1^2} + \beta_0 \left( \bar{\mathcal{G}}_{d,1}^{q,2} + g_1^{b,2} \right) \right. \\
&- \zeta_3 A_1^q f_1^q + \zeta_2 \left( -\frac{1}{2} (f_1^q)^2 + \frac{3}{2} A_2^q + 3\bar{\mathcal{G}}_{d,1}^{q,1} A_1^q + 3g_1^{b,1} A_1^q \right) + \zeta_2 \beta_0 \left( \frac{3}{2} f_1^q + 3B_1^q - 3\gamma_0^b \right) \\
&+ \frac{49}{20} \zeta_2^2 (A_1^q)^2 \Big] + \mathcal{D}_0 \delta(1-z_2) \left[ -f_2^q - 2\bar{\mathcal{G}}_{d,1}^{q,1} f_1^q - 2g_1^{b,1} f_1^q - 2\beta_0 \bar{\mathcal{G}}_{d,1}^{q,1} + 2\zeta_3 (A_1^q)^2 \right. \\
&- \zeta_2 A_1^q f_1^q \Big] + \mathcal{D}_0 \bar{\mathcal{D}}_0 \left[ \frac{1}{2} (f_1^q)^2 + \frac{1}{2} A_2^q + \bar{\mathcal{G}}_{d,1}^{q,1} A_1^q + g_1^{b,1} A_1^q + \frac{1}{2} \beta_0 f_1^q + \frac{1}{2} \zeta_2 (A_1^q)^2 \right] \\
&+ \mathcal{D}_1 \delta(1-z_2) \left[ (f_1^q)^2 + A_2^q + 2\bar{\mathcal{G}}_{d,1}^{q,1} A_1^q + 2g_1^{b,1} A_1^q + \beta_0 f_1^q + \zeta_2 (A_1^q)^2 \right] \\
&+ \mathcal{D}_1 \bar{\mathcal{D}}_0 \left[ -3A_1^q f_1^q - \beta_0 A_1^q \right] + \mathcal{D}_1 \bar{\mathcal{D}}_1 \left[ \frac{3}{2} (A_1^q)^2 \right] + \mathcal{D}_2 \delta(1-z_2) \left[ -\frac{3}{2} A_1^q f_1^q - \frac{1}{2} \beta_0 A_1^q \right] \\
&+ \mathcal{D}_2 \bar{\mathcal{D}}_0 \left[ \frac{3}{2} (A_1^q)^2 \right] + \mathcal{D}_3 \delta(1-z_2) \left[ \frac{1}{2} (A_1^q)^2 \right] + \left\{ z_1 \leftrightarrow z_2 \right\} \tag{2.87}
\end{aligned}$$

$$\begin{aligned}
\Delta_{d,b}^{\text{SV},(3)} = & \delta(1-z_1)\delta(1-z_2) \left[ \frac{1}{3}\overline{\mathcal{G}}_{d,3}^{q,1} + \overline{\mathcal{G}}_{d,1}^{q,1}\overline{\mathcal{G}}_{d,2}^{q,1} + \frac{2}{3}\overline{\mathcal{G}}_{d,1}^{q,1^3} + \frac{1}{3}g_3^{b,1} + g_2^{b,1}\overline{\mathcal{G}}_{d,1}^{q,1} + g_1^{b,1}\overline{\mathcal{G}}_{d,2}^{q,1} \right. \\
& + 2g_1^{b,1}\overline{\mathcal{G}}_{d,1}^{q,1^2} + g_1^{b,1}g_2^{b,1} + 2g_1^{b,1^2}\overline{\mathcal{G}}_{d,1}^{q,1} + \frac{2}{3}g_1^{b,1^3} + \frac{2}{3}\beta_1\left(\overline{\mathcal{G}}_{d,1}^{q,2} + g_1^{b,2}\right) + 2\beta_0\left(\frac{1}{3}\overline{\mathcal{G}}_{d,2}^{q,2} + \overline{\mathcal{G}}_{d,1}^{q,1}\overline{\mathcal{G}}_{d,1}^{q,2} \right. \\
& + \frac{1}{3}g_2^{b,2} + g_1^{b,2}\overline{\mathcal{G}}_{d,1}^{q,1} + g_1^{b,1}\overline{\mathcal{G}}_{d,1}^{q,2} + g_1^{b,1}g_1^{b,2}\left. \right) + \frac{4}{3}\beta_0^2\left(\overline{\mathcal{G}}_{d,1}^{q,3} + g_1^{b,3}\right) - 3\zeta_5(A_1^q)^2 f_1^q - 2\zeta_5\beta_0(A_1^q)^2 \\
& - \zeta_3\left(\frac{1}{3}(f_1^q)^3 + A_2^q f_1^q + A_1^q f_2^q + 2\overline{\mathcal{G}}_{d,1}^{q,1}A_1^q f_1^q + 2g_1^{b,1}A_1^q f_1^q + \beta_0(f_1^q)^2 + 2\beta_0\overline{\mathcal{G}}_{d,1}^{q,1}A_1^q\right) \\
& + \frac{5}{3}\zeta_3^2(A_1^q)^3 + \zeta_2\left(-f_1^q f_2^q + \frac{3}{2}A_3^q + \frac{3}{2}\overline{\mathcal{G}}_{d,2}^{q,1}A_1^q - \overline{\mathcal{G}}_{d,1}^{q,1}(f_1^q)^2 + 3\overline{\mathcal{G}}_{d,1}^{q,1}A_2^q + 3\overline{\mathcal{G}}_{d,1}^{q,1^2}A_1^q \right. \\
& + \frac{3}{2}g_2^{b,1}A_1^q - g_1^{b,1}(f_1^q)^2 + 3g_1^{b,1}A_2^q + 6g_1^{b,1}\overline{\mathcal{G}}_{d,1}^{q,1}A_1^q + 3g_1^{b,1^2}A_1^q\left. \right) + \frac{3}{2}\zeta_2\beta_1\left(f_1^q + 2B_1^q - 2\gamma_0^b\right) \\
& + 3\zeta_2\beta_0\left(f_2^q + 2B_2^q + \overline{\mathcal{G}}_{d,1}^{q,2}A_1^q + \frac{1}{3}\overline{\mathcal{G}}_{d,1}^{q,1}f_1^q + 2\overline{\mathcal{G}}_{d,1}^{q,1}B_1^q + g_1^{b,2}A_1^q + g_1^{b,1}f_1^q + 2g_1^{b,1}B_1^q - 2\gamma_1^b \right. \\
& - 2\gamma_0^b\overline{\mathcal{G}}_{d,1}^{q,1} - 2\gamma_0^b g_1^{b,1}\left. \right) - 6\zeta_2\beta_0^2 g_1^{b,1} + \zeta_2\zeta_3(A_1^q)^2 f_1^q + 2\zeta_2\zeta_3\beta_0(A_1^q)^2 + \frac{49}{10}\zeta_2^2\left(-\frac{11}{49}A_1^q(f_1^q)^2 \right. \\
& + A_1^q A_2^q + \overline{\mathcal{G}}_{d,1}^{q,1}(A_1^q)^2 + g_1^{b,1}(A_1^q)^2\left. \right) + \frac{9}{2}\zeta_2^2\beta_0\left(A_1^q f_1^q + 2A_1^q B_1^q - 2\gamma_0^b A_1^q - \frac{1}{3}A_1^q\right) \\
& + \frac{1181}{420}\zeta_2^3(A_1^q)^3\left. \right] + \mathcal{D}_0\delta(1-z_2) \left[ -f_3^q - \overline{\mathcal{G}}_{d,2}^{q,1}f_1^q - 2\overline{\mathcal{G}}_{d,1}^{q,1}f_2^q - 2\overline{\mathcal{G}}_{d,1}^{q,1^2}f_1^q - g_2^{b,1}f_1^q - 2g_1^{b,1}f_2^q \right. \\
& - 4g_1^{b,1}\overline{\mathcal{G}}_{d,1}^{q,1}f_1^q - 2g_1^{b,1^2}f_1^q - 2\beta_1\overline{\mathcal{G}}_{d,1}^{q,1} - 2\beta_0\left(\overline{\mathcal{G}}_{d,2}^{q,1} + \overline{\mathcal{G}}_{d,1}^{q,2}f_1^q + 2\overline{\mathcal{G}}_{d,1}^{q,1^2} + g_1^{b,2}f_1^q + 2g_1^{b,1}\overline{\mathcal{G}}_{d,1}^{q,1}\right) \\
& - 4\beta_0^2\overline{\mathcal{G}}_{d,1}^{q,2} + 6\zeta_5(A_1^q)^3 + 4\zeta_3\left(A_1^q(f_1^q)^2 + A_1^q A_2^q + \overline{\mathcal{G}}_{d,1}^{q,1}(A_1^q)^2 + g_1^{b,1}(A_1^q)^2 + \frac{3}{2}\beta_0 A_1^q f_1^q\right) \\
& + \zeta_2\left((f_1^q)^3 - A_2^q f_1^q - A_1^q f_2^q - 2\overline{\mathcal{G}}_{d,1}^{q,1}A_1^q f_1^q - 2g_1^{b,1}A_1^q f_1^q\right) - \zeta_2\beta_0\left((f_1^q)^2 + 6B_1^q f_1^q + 2\overline{\mathcal{G}}_{d,1}^{q,1}A_1^q \right. \\
& - 6\gamma_0^b f_1^q\left. \right) - 2\zeta_2\zeta_3(A_1^q)^3 - \frac{1}{2}\zeta_2^2(A_1^q)^2 f_1^q\left. \right] + \mathcal{D}_0\overline{\mathcal{D}}_0 \left[ f_1^q f_2^q + \frac{1}{2}A_3^q + \frac{1}{2}\overline{\mathcal{G}}_{d,2}^{q,1}A_1^q + \overline{\mathcal{G}}_{d,1}^{q,1}(f_1^q)^2 \right. \\
& + \overline{\mathcal{G}}_{d,1}^{q,1}A_2^q + \overline{\mathcal{G}}_{d,1}^{q,1^2}A_1^q + \frac{1}{2}g_2^{b,1}A_1^q + g_1^{b,1}(f_1^q)^2 + g_1^{b,1}A_2^q + 2g_1^{b,1}\overline{\mathcal{G}}_{d,1}^{q,1}A_1^q + g_1^{b,1^2}A_1^q + \frac{1}{2}\beta_1 f_1^q \\
& + \beta_0\left(f_2^q + \overline{\mathcal{G}}_{d,1}^{q,2}A_1^q + 3\overline{\mathcal{G}}_{d,1}^{q,1}f_1^q + g_1^{b,2}A_1^q + g_1^{b,1}f_1^q\right) + 2\beta_0^2\overline{\mathcal{G}}_{d,1}^{q,1} - 5\zeta_3(A_1^q)^2 f_1^q - 3\zeta_3\beta_0(A_1^q)^2 \\
& + \zeta_2\left(A_1^q A_2^q - A_1^q(f_1^q)^2 + \overline{\mathcal{G}}_{d,1}^{q,1}(A_1^q)^2 + g_1^{b,1}(A_1^q)^2\right) + 3\zeta_2\beta_0\left(A_1^q B_1^q - \gamma_0^b A_1^q\right) + \frac{1}{4}\zeta_2^2(A_1^q)^3\left. \right] \\
& + \mathcal{D}_1\delta(1-z_2) \left[ 2f_1^q f_2^q + A_3^q + \overline{\mathcal{G}}_{d,2}^{q,1}A_1^q + 2\overline{\mathcal{G}}_{d,1}^{q,1}(f_1^q)^2 + 2\overline{\mathcal{G}}_{d,1}^{q,1}A_2^q + 2\overline{\mathcal{G}}_{d,1}^{q,1^2}A_1^q + g_2^{b,1}A_1^q \right. \\
& + 2g_1^{b,1}(f_1^q)^2 + 2g_1^{b,1}A_2^q + 4g_1^{b,1}\overline{\mathcal{G}}_{d,1}^{q,1}A_1^q + 2g_1^{b,1^2}A_1^q + \beta_1 f_1^q + 2\beta_0\left(f_2^q + \overline{\mathcal{G}}_{d,1}^{q,2}A_1^q + 3\overline{\mathcal{G}}_{d,1}^{q,1}f_1^q \right. \\
& + g_1^{b,2}A_1^q + g_1^{b,1}f_1^q\left. \right) + 4\beta_0^2\overline{\mathcal{G}}_{d,1}^{q,1} - 10\zeta_3(A_1^q)^2 f_1^q - 6\zeta_3\beta_0(A_1^q)^2 + 2\zeta_2\left(-A_1^q(f_1^q)^2 + A_1^q A_2^q \right. \\
& + \overline{\mathcal{G}}_{d,1}^{q,1}(A_1^q)^2 + g_1^{b,1}(A_1^q)^2\left. \right) + 6\zeta_2\beta_0\left(A_1^q B_1^q - \gamma_0^b A_1^q\right) + \frac{1}{2}\zeta_2^2(A_1^q)^3\left. \right] + \mathcal{D}_1\overline{\mathcal{D}}_0 \left[ -(f_1^q)^3 \right. \\
& - 3A_2^q f_1^q - 3A_1^q f_2^q - 6\overline{\mathcal{G}}_{d,1}^{q,1}A_1^q f_1^q - 6g_1^{b,1}A_1^q f_1^q - \beta_1 A_1^q - \beta_0\left(3(f_1^q)^2 + 2A_2^q + 8\overline{\mathcal{G}}_{d,1}^{q,1}A_1^q \right.
\end{aligned}$$

$$\begin{aligned}
& + 2g_1^{b,1}A_1^q) - 2\beta_0^2 f_1^q + 10\zeta_3(A_1^q)^3 + 3\zeta_2(A_1^q)^2 f_1^q + 3\zeta_2\beta_0(A_1^q)^2 \Big] + \mathcal{D}_1\overline{\mathcal{D}}_1 \Big[ 3A_1^q(f_1^q)^2 \\
& + 3A_1^q A_2^q + 3\overline{\mathcal{G}}_{d,1}^{q,1}(A_1^q)^2 + 3g_1^{b,1}(A_1^q)^2 + 5\beta_0 A_1^q f_1^q + \beta_0^2 A_1^q - \frac{3}{2}\zeta_2(A_1^q)^3 \Big] \\
& + \mathcal{D}_2\delta(1-z_2) \Big[ -\frac{1}{2}(f_1^q)^3 - \frac{3}{2}A_2^q f_1^q - \frac{3}{2}A_1^q f_2^q - 3\overline{\mathcal{G}}_{d,1}^{q,1}A_1^q f_1^q - 3g_1^{b,1}A_1^q f_1^q - \frac{1}{2}\beta_1 A_1^q \\
& - \beta_0 \left( \frac{3}{2}(f_1^q)^2 + A_2^q + 4\overline{\mathcal{G}}_{d,1}^{q,1}A_1^q + g_1^{b,1}A_1^q \right) - \beta_0^2 f_1^q + 5\zeta_3(A_1^q)^3 + \frac{3}{2}\zeta_2(A_1^q)^2 f_1^q \\
& + \frac{3}{2}\zeta_2\beta_0(A_1^q)^2 \Big] + \mathcal{D}_2\overline{\mathcal{D}}_0 \Big[ 3A_1^q(f_1^q)^2 + 3A_1^q A_2^q + 3\overline{\mathcal{G}}_{d,1}^{q,1}(A_1^q)^2 + 3g_1^{b,1}(A_1^q)^2 + 5\beta_0 A_1^q f_1^q \\
& + \beta_0^2 A_1^q - \frac{3}{2}\zeta_2(A_1^q)^3 \Big] + \mathcal{D}_2\overline{\mathcal{D}}_1 \Big[ -\frac{15}{2}(A_1^q)^2 f_1^q - 5\beta_0(A_1^q)^2 \Big] + \mathcal{D}_2\overline{\mathcal{D}}_2 \Big[ \frac{15}{8}(A_1^q)^3 \Big] \\
& + \mathcal{D}_3\delta(1-z_2) \Big[ A_1^q(f_1^q)^2 + A_1^q A_2^q + \overline{\mathcal{G}}_{d,1}^{q,1}(A_1^q)^2 + g_1^{b,1}(A_1^q)^2 + \frac{5}{3}\beta_0 A_1^q f_1^q + \frac{1}{3}\beta_0^2 A_1^q \\
& - \frac{1}{2}\zeta_2(A_1^q)^3 \Big] + \mathcal{D}_3\overline{\mathcal{D}}_0 \Big[ -\frac{5}{2}(A_1^q)^2 f_1^q - \frac{5}{3}\beta_0(A_1^q)^2 \Big] + \mathcal{D}_3\overline{\mathcal{D}}_1 \Big[ \frac{5}{2}(A_1^q)^3 \Big] \\
& + \mathcal{D}_4\delta(1-z_2) \Big[ -\frac{5}{8}(A_1^q)^2 f_1^q - \frac{5}{12}\beta_0(A_1^q)^2 \Big] + \mathcal{D}_4\overline{\mathcal{D}}_0 \Big[ \frac{5}{8}(A_1^q)^3 \Big] \\
& + \mathcal{D}_5\delta(1-z_2) \Big[ \frac{1}{8}(A_1^q)^3 \Big] + \left\{ z_1 \leftrightarrow z_2 \right\}. \tag{2.88}
\end{aligned}$$

At the stage, we can demonstrate that integration over the rapidity correctly reproduces inclusive threshold contribution to the Higgs production in bottom anti-bottom annihilation reported in [59] :

$$\int dy \frac{d}{dy} \sigma_b(\tau, y, q^2) = \sigma_b(\tau, q^2). \tag{2.89}$$

The integration over the rapidity  $y$  leads to the following relation between  $\Delta_{d,b}^{\text{SV}}(z_1, z_2)$  obtained in this paper and  $\Delta_b^{\text{SV}}(z)$  in [59]:

$$\Delta_b^{\text{SV}}(z) = \int dz_1 \int dz_2 \delta(z - z_1 z_2) \Delta_{d,b}^{\text{SV}}(z_1, z_2). \tag{2.90}$$

We have explicitly checked that the results presented here for  $\Delta_{d,b}^{\text{SV}}$  and those for  $\Delta_b^{\text{SV}}$  in the [59] up to N<sup>3</sup>LO level satisfy the above relation confirming the consistency of the formalism used. For completeness, we present the results for  $\Delta_{d,b}^{\text{SV},(i)}$  up to N<sup>3</sup>LO after substituting all the constants that are required to this order:

$$\begin{aligned}
\Delta_{d,b}^{\text{SV},(1)} &= \delta(1-z_1)\delta(1-z_2)C_F(-2+6\zeta_2) + \mathcal{D}_0\overline{\mathcal{D}}_0(2C_F) + \mathcal{D}_1\delta(1-z_2)(4C_F) + \left\{ z_1 \leftrightarrow z_2 \right\}, \\
\Delta_{d,b}^{\text{SV},(2)} &= \delta(1-z_1)\delta(1-z_2) \left[ C_F C_A \left( \frac{83}{9} + \frac{32}{3}\zeta_3 + \frac{250}{9}\zeta_2 - \frac{26}{5}\zeta_2^2 \right) + C_F^2 \left( 8 - 30\zeta_3 - 8\zeta_2 \right) \right.
\end{aligned}$$



$$\begin{aligned}
& + \frac{152}{5}\zeta_2^2) + n_f C_F \left( \frac{4}{9} + \frac{4}{3}\zeta_3 - \frac{40}{9}\zeta_2 \right) \Big] + \mathcal{D}_0 \delta(1-z_2) \left[ C_F C_A \left( -\frac{808}{27} + 28\zeta_3 + \frac{44}{3}\zeta_2 \right) \right. \\
& + C_F^2 (32\zeta_3) + n_f C_F \left( \frac{112}{27} - \frac{8}{3}\zeta_2 \right) \Big] + \mathcal{D}_0 \bar{\mathcal{D}}_0 \left[ C_F C_A \left( \frac{134}{9} - 4\zeta_2 \right) + C_F^2 (-8 + 8\zeta_2) \right. \\
& + n_f C_F \left( -\frac{20}{9} \right) \Big] + \mathcal{D}_1 \delta(1-z_2) \left[ C_F C_A \left( \frac{268}{9} - 8\zeta_2 \right) + C_F^2 (-16 + 16\zeta_2) \right. \\
& + n_f C_F \left( -\frac{40}{9} \right) \Big] + \mathcal{D}_1 \bar{\mathcal{D}}_0 \left[ C_F C_A \left( -\frac{44}{3} \right) + n_f C_F \left( \frac{8}{3} \right) \right] + \mathcal{D}_1 \bar{\mathcal{D}}_1 \left[ C_F^2 (24) \right] \\
& + \mathcal{D}_2 \delta(1-z_2) \left[ C_F C_A \left( -\frac{22}{3} \right) + n_f C_F \left( \frac{4}{3} \right) \right] + \mathcal{D}_2 \bar{\mathcal{D}}_0 \left[ C_F^2 (24) \right] \\
& + \mathcal{D}_3 \delta(1-z_2) \left[ C_F^2 (8) \right] + \left\{ z_1 \leftrightarrow z_2 \right\}
\end{aligned}$$

and

$$\begin{aligned}
\Delta_{d,b}^{\text{SV},(3)} = & \delta(1-z_1)\delta(1-z_2) \left[ C_F C_A^2 \left( \frac{34495}{81} - 42\zeta_5 + \frac{14254}{81}\zeta_3 - \frac{200}{3}\zeta_3^2 + \frac{4487}{81}\zeta_2 - 324\zeta_2\zeta_3 \right. \right. \\
& - \frac{2446}{135}\zeta_2^2 + \frac{12176}{315}\zeta_2^3) + C_F^2 C_A \left( -\frac{491}{3} - \frac{2732}{9}\zeta_5 - \frac{922}{3}\zeta_3 + \frac{632}{3}\zeta_3^2 + \frac{10441}{27}\zeta_2 \right. \\
& + \frac{4288}{9}\zeta_2\zeta_3 + \frac{21302}{135}\zeta_2^2 - \frac{39136}{315}\zeta_2^3) + C_F^3 \left( \frac{539}{3} + 424\zeta_5 - 594\zeta_3 + \frac{368}{3}\zeta_3^2 - \frac{179}{3}\zeta_2 \right. \\
& - 152\zeta_2\zeta_3 - \frac{196}{5}\zeta_2^2 + \frac{45008}{315}\zeta_2^3) + n_f C_F C_A \left( -\frac{5770}{81} - 4\zeta_5 - \frac{5660}{81}\zeta_3 - \frac{806}{27}\zeta_2 \right. \\
& + \frac{136}{3}\zeta_2\zeta_3 - \frac{380}{27}\zeta_2^2) + n_f C_F^2 \left( -\frac{35}{9} - \frac{112}{9}\zeta_5 + 180\zeta_3 - \frac{1507}{27}\zeta_2 - \frac{736}{9}\zeta_2\zeta_3 - \frac{1604}{135}\zeta_2^2) \\
& + n_f^2 C_F \left( \frac{8}{27} - \frac{80}{81}\zeta_3 + \frac{184}{81}\zeta_2 + \frac{296}{135}\zeta_2^2) \Big] + \mathcal{D}_0 \delta(1-z_2) \left[ C_F C_A^2 \left( -\frac{297029}{729} - 192\zeta_5 \right. \right. \\
& + \frac{14264}{27}\zeta_3 + \frac{27752}{81}\zeta_2 - \frac{176}{3}\zeta_2\zeta_3 - \frac{616}{15}\zeta_2^2) + C_F^2 C_A \left( \frac{3232}{27} + \frac{3280}{9}\zeta_3 - \frac{4816}{27}\zeta_2 - 16\zeta_2\zeta_3 \right. \\
& + \frac{176}{3}\zeta_2^2) + C_F^3 \left( 384\zeta_5 - 128\zeta_3 - 128\zeta_2\zeta_3) + n_f C_F C_A \left( \frac{62626}{729} - \frac{536}{9}\zeta_3 - \frac{7760}{81}\zeta_2 \right. \\
& + \frac{208}{15}\zeta_2^2) + n_f C_F^2 \left( \frac{421}{9} - \frac{944}{9}\zeta_3 + \frac{520}{27}\zeta_2 - \frac{256}{15}\zeta_2^2) + n_f^2 C_F \left( -\frac{1856}{729} - \frac{32}{27}\zeta_3 \right. \\
& + \frac{160}{27}\zeta_2) \Big] + \mathcal{D}_0 \bar{\mathcal{D}}_0 \left[ C_F C_A^2 \left( \frac{15503}{81} - 88\zeta_3 - \frac{340}{3}\zeta_2 + \frac{88}{5}\zeta_2^2) + C_F^2 C_A \left( -\frac{68}{3} - \frac{400}{3}\zeta_3 \right. \right. \\
& + \frac{608}{9}\zeta_2 - \frac{24}{5}\zeta_2^2) + C_F^3 \left( 32 - 120\zeta_3 + 32\zeta_2 - \frac{96}{5}\zeta_2^2) + n_f C_F C_A \left( -\frac{4102}{81} + \frac{256}{9}\zeta_2 \right) \right. \\
& + n_f C_F^2 \left( -\frac{23}{3} + \frac{160}{3}\zeta_3 - \frac{80}{9}\zeta_2) + n_f^2 C_F \left( \frac{200}{81} - \frac{16}{9}\zeta_2) \Big] + \mathcal{D}_1 \delta(1-z_2) \left[ C_F C_A^2 \left( \frac{31006}{81} \right. \right. \right. \\
& - 176\zeta_3 - \frac{680}{3}\zeta_2 + \frac{176}{5}\zeta_2^2) + C_F^2 C_A \left( -\frac{136}{3} - \frac{800}{3}\zeta_3 + \frac{1216}{9}\zeta_2 - \frac{48}{5}\zeta_2^2) + C_F^3 (64
\end{aligned}$$

$$\begin{aligned}
& -240\zeta_3 + 64\zeta_2 - \frac{192}{5}\zeta_2^2) + n_f C_F C_A \left( -\frac{8204}{81} + \frac{512}{9}\zeta_2 \right) + n_f C_F^2 \left( -\frac{46}{3} + \frac{320}{3}\zeta_3 \right. \\
& \left. - \frac{160}{9}\zeta_2 \right) + n_f^2 C_F \left( \frac{400}{81} - \frac{32}{9}\zeta_2 \right) \Big] + \mathcal{D}_1 \bar{\mathcal{D}}_0 \left[ C_F C_A^2 \left( -\frac{7120}{27} + \frac{176}{3}\zeta_2 \right) \right. \\
& + C_F^2 C_A \left( -\frac{2704}{9} + 336\zeta_3 + 352\zeta_2 \right) + C_F^3 (640\zeta_3) + n_f C_F C_A \left( \frac{2312}{27} - \frac{32}{3}\zeta_2 \right) \\
& + n_f C_F^2 \left( \frac{424}{9} - 64\zeta_2 \right) + n_f^2 C_F \left( -\frac{160}{27} \right) \Big] + \mathcal{D}_1 \bar{\mathcal{D}}_1 \left[ C_F C_A^2 \left( \frac{484}{9} \right) + C_F^2 C_A \left( \frac{1072}{3} \right. \right. \\
& \left. \left. - 96\zeta_2 \right) + C_F^3 \left( -96 - 96\zeta_2 \right) + n_f C_F C_A \left( -\frac{176}{9} \right) + n_f C_F^2 \left( -\frac{160}{3} \right) + n_f^2 C_F \left( \frac{16}{9} \right) \right] \\
& + \mathcal{D}_2 \delta(1 - z_2) \left[ C_F C_A^2 \left( -\frac{3560}{27} + \frac{88}{3}\zeta_2 \right) + C_F^2 C_A \left( -\frac{1352}{9} + 168\zeta_3 + 176\zeta_2 \right) \right. \\
& + C_F^3 (320\zeta_3) + n_f C_F C_A \left( \frac{1156}{27} - \frac{16}{3}\zeta_2 \right) + n_f C_F^2 \left( \frac{212}{9} - 32\zeta_2 \right) + n_f^2 C_F \left( -\frac{80}{27} \right) \Big] \\
& + \mathcal{D}_2 \bar{\mathcal{D}}_0 \left[ C_F C_A^2 \left( \frac{484}{9} \right) + C_F^2 C_A \left( \frac{1072}{3} - 96\zeta_2 \right) + C_F^3 \left( -96 - 96\zeta_2 \right) \right. \\
& + n_f C_F C_A \left( -\frac{176}{9} \right) + n_f C_F^2 \left( -\frac{160}{3} \right) + n_f^2 C_F \left( \frac{16}{9} \right) \Big] + \mathcal{D}_2 \bar{\mathcal{D}}_1 \left[ C_F^2 C_A \left( -\frac{880}{3} \right) \right. \\
& \left. + n_f C_F^2 \left( \frac{160}{3} \right) \right] + \mathcal{D}_2 \bar{\mathcal{D}}_2 \left[ C_F^3 (120) \right] + \mathcal{D}_3 \delta(1 - z_2) \left[ C_F C_A^2 \left( \frac{484}{27} \right) + C_F^2 C_A \left( \frac{1072}{9} \right. \right. \\
& \left. \left. - 32\zeta_2 \right) + C_F^3 \left( -32 - 32\zeta_2 \right) + n_f C_F C_A \left( -\frac{176}{27} \right) + n_f C_F^2 \left( -\frac{160}{9} \right) + n_f^2 C_F \left( \frac{16}{27} \right) \right] \\
& + \mathcal{D}_3 \bar{\mathcal{D}}_0 \left[ C_F^2 C_A \left( -\frac{880}{9} \right) + n_f C_F^2 \left( \frac{160}{9} \right) \right] + \mathcal{D}_3 \bar{\mathcal{D}}_1 \left[ C_F^3 (160) \right] \\
& + \mathcal{D}_4 \delta(1 - z_2) \left[ C_F^2 C_A \left( -\frac{220}{9} \right) + n_f C_F^2 \left( \frac{40}{9} \right) \right] + \mathcal{D}_4 \bar{\mathcal{D}}_0 \left[ C_F^3 (40) \right] \\
& + \mathcal{D}_5 \delta(1 - z_2) \left[ C_F^3 (8) \right] + \left\{ z_1 \leftrightarrow z_2 \right\}. \tag{2.91}
\end{aligned}$$


Substituting  $\Delta_{d,b}^{\text{SV},(1)}$ ,  $\Delta_{d,b}^{\text{SV},(2)}$  and  $\Delta_{d,b}^{\text{SV},(3)}$  in the Eq. 2.13, we obtain  $W_b^{\text{SV},(i)}$  or equivalently  $\frac{d}{dy}\sigma^{b,\text{SV},(i)}$  (Eq. 2.4) at the hadronic level order by order up to  $\mathcal{O}(a_s^3)$ .

### 2.3 Numerical Results

In this section, we present the numerical impact of the rapidity distribution of the Higgs boson, produced via bottom anti-bottom annihilation subprocess at the LHC. The rapidity distribution can be expanded in powers of the strong coupling constant  $a_s$  as

$$\frac{d\sigma^b}{dY} = \frac{d\sigma^{b,(0)}}{dY} + \sum_{i=1}^{\infty} a_s^i \frac{d\sigma^{b,(i)}}{dY}. \tag{2.92}$$

Beyond LO, the distribution is split into hard and SV parts as



**Figure 1.** The comparison between NLO and NLO<sub>SV</sub> with the renormalization scale  $\mu_R = m_H$  and factorization scale  $\mu_F = m_H/4$  at 8 TeV(left panel) and 13 TeV (right panel) LHC.

$$\frac{d\sigma^{b,(i)}}{dY} = \frac{d\sigma^{\text{hard},b,(i)}}{dY} + \frac{d\sigma^{\text{SV},b,(i)}}{dY}. \quad (2.93)$$

In the following, for our numerical study we will use the exact results up to NLO level but at NNLO, we use exact NLO and only threshold contribution at  $\mathcal{O}(a_s^2)$  as we do not have access to the hard part at  $\mathcal{O}(a_s^2)$  computed in [63]<sup>1</sup>. We call it NNLO(SV). Similarly at N<sup>3</sup>LO level, we will use NNLO(SV) and threshold contribution at  $\mathcal{O}(a_s^3)$ , denoted by N<sup>3</sup>LO(SV) hereafter. We present results for the center of mass energies 8 and 13 TeV at the LHC. The standard model parameters which enter into our computation are the Z boson mass  $m_Z = 91.1876$  GeV, top quark mass  $m_t = 173.4$  GeV and mass of the Higgs boson  $m_H = 125$  GeV. The strong coupling constant is evolved using the 4-loop RG equations with  $\alpha_s^{\text{N}^3\text{LO}}(m_Z) = 0.117$ . Following the Ref. [84], the solution to RGE 2.47 for  $\lambda(\mu_R^2)$  is given by,

$$\lambda(\mu_R^2) = \lambda(\mu_0^2) \frac{M(a_s(\mu_R^2))}{M(a_s(\mu_0^2))} \quad (2.94)$$

with

$$M(a_s) = a_s^{A_0} \sum_{i=0}^{\infty} a_s^i M_i. \quad (2.95)$$

The  $K_i$  are given by

$$M_0 = 1, \quad M_1 = A_1,$$

<sup>1</sup>The authors informed us that the code is not yet ready for public distribution

Y	0.0	0.4	0.8	1.2	1.6	2.0	2.4	2.8	3.2
10 <sup>2</sup> LO	4.137	4.027	3.705	3.196	2.549	1.828	1.126	5.427	1.686
10 <sup>2</sup> NLO	6.485	6.225	5.495	4.429	3.217	2.054	1.097	4.419	1.065
10 <sup>2</sup> NNLO(SV)	6.921	6.650	5.879	4.731	3.407	2.135	1.113	4.417	1.118
10 <sup>2</sup> N <sup>3</sup> LO(SV)	6.984	6.707	5.922	4.757	3.415	2.130	1.105	4.340	1.084

**Table 1.** Contributions at LO, NLO, NNLO(SV) and N<sup>3</sup>LO(SV) with the renormalization scale  $\mu_R = m_H$  and factorization scale  $\mu_F = m_H/4$  at 8 TeV LHC.

Y	0.0	0.4	0.8	1.2	1.6	2.0	2.4	2.8	3.2
10 <sup>2</sup> LO	8.465	8.293	7.787	6.981	5.925	4.686	3.371	2.115	1.068
10 <sup>2</sup> NLO	13.466	13.063	11.903	10.133	7.985	5.737	3.671	2.001	0.849
10 <sup>2</sup> NNLO(SV)	14.284	13.875	12.689	10.844	8.549	6.099	3.833	2.035	0.848
10 <sup>2</sup> N <sup>3</sup> LO(SV)	14.475	14.057	12.843	10.959	8.620	6.131	3.837	2.025	0.838

**Table 2.** Contributions at LO, NLO, NNLO(SV) and N<sup>3</sup>LO(SV) with the renormalization scale  $\mu_R = m_H$  and factorization scale  $\mu_F = m_H/4$  at 13 TeV LHC.

$$M_2 = \frac{1}{2}(A_1^2 + A_2), \quad M_3 = \frac{1}{6}(A_1^3 + 3A_1A_2 + 2A_3), \quad (2.96)$$

with

$$A_0 = c_0, \quad A_1 = c_1 - b_1c_0, \quad A_2 = c_2 - b_1c_1 + c_0(b_1^2 - b_2),$$


$$A_3 = c_3 - b_1c_2 + c_1(b_1^2 - b_2) + c_0(b_1b_2 - b_1(b_1^2 - b_2) - b_3), \quad (2.97)$$

and


$$c_i = \frac{\gamma_i^b}{\beta_0}, \quad b_i = \frac{\gamma_i^b}{\beta_0}, \quad (2.98)$$

where  $\mu_0$  is some reference scale at which  $\lambda$  is known. We have numerically evaluated  $\lambda(\mu_R^2)$  to relevant order namely LO, NLO, NNLO and N<sup>3</sup>LO by truncating the terms in the RHS of Eq. 2.47. We have used  $\lambda(\mu_0^2) = \sqrt{2}m_b(\mu_0)/v$  and  $m_b(\mu_0) = 3.63$  GeV with the choice  $\mu_0 = 10$  GeV. We use the MSTW2008 [85] parton density sets with errors estimated at 68% confidence level with five active flavours. Parton densities and  $\alpha_s$  are evaluated at each corresponding perturbative order. Specifically, we use  $(n+1)$ -loop  $\alpha_s$  at N<sup>n</sup>LO, with  $n = 0, 1, 2, 3$ . However, we use MSTW2008NNLO PDFs at N<sup>3</sup>LO, the N<sup>3</sup>LO kernels not being available at the moment. We set the renormalization scale  $\mu_R = m_H$  and factorization scale  $\mu_F = m_H/4$  [16] as their central values.

Several checks have been performed on our numerical code. We have found complete agreement with the literature on the inclusive Higgs production rate [19, 59] after performing an additional numerical integration over the rapidity Y of our distribution. The check was also performed at the analytical level. However, we were not able to reproduce the




**Figure 2.** The rapidity distribution of the Higgs boson at NLO, NNLO(SV) and  $N^3\text{LO(SV)}$  at 8 TeV(left panel) and 13 TeV (right panel) LHC. The band indicates the uncertainty due to renormalization scale.



**Figure 3.** The rapidity distribution of the Higgs boson at NLO, NNLO(SV) and  $N^3\text{LO(SV)}$  at 8 TeV(left panel) and 13 TeV (right panel) LHC. The band indicates the uncertainty due to factorization scale.

plot given in [63], after using the same set of values of the input parameters. We begin our discussion with the results at NLO level. In Sec. 2.1, we presented the contributions coming from the exact results, containing the regular as well as pure threshold ones to the rapidity distribution at  $\mathcal{O}(a_s)$ . In Fig. 1, we plot both the NLO(SV) and exact NLO rapidity distri-




**Figure 4.** The distribution of  $K_1^{(SV)}$ ,  $K_1$ ,  $K_2^{(SV)}$  and  $K_3^{(SV)}$  at different perturbative order at 8 TeV(left panel) and 13 TeV (right panel) LHC.

butions to exhibit the dominance of threshold over the entire rapidity range after setting the values of the renormalization and factorization scales to their central values. From now onward, we adopt a consistent representation to display the figures corresponding to our results. In every figure, the left panel shows the result for 8 TeV whereas the right panel corresponds to 13 TeV at the LHC. We observe that the exact NLO contribution is well approximated by the NLO(SV), thanks to the intrinsic property of the matrix element, where the phase-space points corresponding to the born kinematics contribute towards the largest radiative corrections for the low  $\tau$  ( $m_H^2/s \approx 10^{-4}$ ) values. So, we expect that the trend of approximating the exact results by threshold corrections at that order to remain same after the inclusion of higher order terms also.

With this in mind, we present the results at LO, NLO, NNLO(SV), N<sup>3</sup>LO(SV) for different values of the rapidity  $Y$  after setting the central values for renormalization and factorization scales for 8 TeV in Table 1 and for 13 TeV in Table 2 at LHC. The hadronic cross-section, obtained by the convolution of the partonic cross section with the parton densities, suffers from the theoretical uncertainties, arising from the missing higher order corrections, through the renormalization ( $\mu_R$ ) and factorization ( $\mu_F$ ) scales. These can be estimated through the variation of the differential hadronic cross section with  $\mu_R$  and  $\mu_F$ , thereby exhibiting the size of the higher order effects.

In Fig. 2, we plot two curves for each order for the predictions at NLO, NNLO(SV), N<sup>3</sup>LO(SV) corresponding to two different choices of the renormalization scale,  $\mu_R = 0.1m_H$  and  $\mu_R = 10m_H$ , keeping the factorization scale fixed at  $\mu_F = m_H/4$ , whereas in Fig. 3, we plot the predictions at each order corresponding to two different choices of the factorization scale,  $\mu_F = m_H/8$  and  $\mu_F = m_H/2$ , keeping the renormalization scale fixed at  $\mu_R = m_H$ . We observe a consistent improvement in the accuracy of the predictions with the inclusion



**Figure 5.** The distribution of  $K_{\text{NLO}}$ ,  $K_{\text{NNLO}}^{(\text{SV})}$  and  $K_{\text{N}^3\text{LO}}^{(\text{SV})}$  at different perturbative order at 8 TeV(left panel) and 13 TeV (right panel) LHC.

of the higher order terms, the width of the bands being an clear indicator of the theoretical uncertainties. Moreover, we can see that the dependence on the renormalization scale for this process is very mild. Another way to assess the reliability of the prediction is to study the rate of convergence of the perturbation series, represented by the K-factor.

In the Fig. 4, we plot the K-factors defined as  $K_1 = d\sigma^{NLO}/d\sigma^{LO}$  and  $K_i^{(\text{SV})} = d\sigma^{N^iLO(\text{SV})}/d\sigma^{LO}$ ,  $i = 2, 3$  as a function of  $Y$ . For 8 TeV LHC, we see that the  $K_1$  varies from 1.57 to 0.63 over the entire rapidity range, while the value of  $K_1$  for the inclusive rate is 1.37. Similarly, for  $K_2^{(\text{SV})}$ , the variation is from 1.67 to 0.66, while for the inclusive rate it is 1.35. It shows, particularly, that the shape at higher orders can not be rescaled from lower orders as the differential K-factor varies significantly over the full rapidity range. In the Fig. 5 we plot K factors defined by  $K_{\text{NLO}}^{(\text{SV})} = d\sigma^{NLO(\text{SV})}/d\sigma^{LO}$ ,  $K_{\text{NNLO}}^{(\text{SV})} = d\sigma^{\text{NNLO}(\text{SV})}/d\sigma^{\text{NLO}}$  and  $K_{\text{N}^3\text{LO}}^{(\text{SV})} = d\sigma^{\text{N}^3\text{LO}(\text{SV})}/d\sigma^{\text{NNLO}(\text{SV})}$ . The values of the K-factors with the inclusion of higher order terms decrease, thereby implying a considerable amount of improvement in the rate of convergence.

### 3 Conclusions

To summarize, we present threshold enhanced  $\text{N}^3\text{LO}$  QCD correction to rapidity distribution of the Higgs boson produced through bottom quark annihilation at the LHC. We show in detail the infra-red structure of the QCD amplitudes at NLO level as well as the cancellation of the various soft and collinear singularities through the summation of all possible degenerate states and the renormalization of the PDFs in order to demonstrate a general framework to obtain threshold corrections to rapidity distributions to all orders in perturbation theory. We have used factorization properties, along with Sudakov resuma-

tion of soft gluons and renormalization group invariance to achieve this. The recent result on three loop form factor by Gehrmann and Kara [57] and the universal soft distribution obtained in [51] provide the last missing information to obtain threshold correction to N<sup>3</sup>LO for the rapidity distribution of Higgs boson in bottom quark annihilation. We find the dominance of the threshold contribution over the entire rapidity range at NLO. We extend this approximation beyond NLO to make predictions for center of mass energies 8 and 13 TeV. We observe that the inclusion of N<sup>3</sup>LO contributions reduces the scale dependency further, as expected, through the variation of the renormalization and factorization scales around their central values and that K-factors show stability at higher orders.

## Acknowledgement

The work of T.A., M.K.M. and N.R. has been partially supported by funding from Regional Center for Accelerator-based Particle Physics (RECAPP), Department of Atomic Energy, Govt. of India. V.R. would like to thank T. Gehrmann for useful discussion.

## References

- [1] **ATLAS Collaboration** Collaboration, G. Aad *et al.*, *Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC*, *Phys.Lett.* **B716** (2012) 1–29, [arXiv:1207.7214 \[hep-ex\]](#).
- [2] **CMS Collaboration** Collaboration, S. Chatrchyan *et al.*, *Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC*, *Phys.Lett.* **B716** (2012) 30–61, [arXiv:1207.7235 \[hep-ex\]](#).
- [3] H. Georgi, S. Glashow, M. Machacek, and D. V. Nanopoulos, *Higgs Bosons from Two Gluon Annihilation in Proton Proton Collisions*, *Phys.Rev.Lett.* **40** (1978) 692.
- [4] A. Djouadi, M. Spira, and P. Zerwas, *Production of Higgs bosons in proton colliders: QCD corrections*, *Phys.Lett.* **B264** (1991) 440–446.
- [5] S. Dawson, *Radiative corrections to Higgs boson production*, *Nucl.Phys.* **B359** (1991) 283–300.
- [6] M. Spira, A. Djouadi, D. Graudenz, and P. Zerwas, *Higgs boson production at the LHC*, *Nucl.Phys.* **B453** (1995) 17–82, [arXiv:hep-ph/9504378 \[hep-ph\]](#).
- [7] S. Catani, D. de Florian, and M. Grazzini, *Higgs production in hadron collisions: Soft and virtual QCD corrections at NNLO*, *JHEP* **0105** (2001) 025, [arXiv:hep-ph/0102227 \[hep-ph\]](#).
- [8] R. V. Harlander and W. B. Kilgore, *Soft and virtual corrections to proton proton  $\rightarrow gt$ ;  $H + x$  at NNLO*, *Phys.Rev.* **D64** (2001) 013015, [arXiv:hep-ph/0102241 \[hep-ph\]](#).
- [9] R. V. Harlander and W. B. Kilgore, *Next-to-next-to-leading order Higgs production at hadron colliders*, *Phys.Rev.Lett.* **88** (2002) 201801, [arXiv:hep-ph/0201206 \[hep-ph\]](#).
- [10] C. Anastasiou and K. Melnikov, *Higgs boson production at hadron colliders in NNLO QCD*, *Nucl.Phys.* **B646** (2002) 220–256, [arXiv:hep-ph/0207004 \[hep-ph\]](#).



- [11] V. Ravindran, J. Smith, and W. L. van Neerven, *NNLO corrections to the total cross-section for Higgs boson production in hadron hadron collisions*, *Nucl.Phys.* **B665** (2003) 325–366, [arXiv:hep-ph/0302135](#) [hep-ph].
- [12] P. Bolzoni, F. Maltoni, S.-O. Moch, and M. Zaro, *Higgs production via vector-boson fusion at NNLO in QCD*, *Phys.Rev.Lett.* **105** (2010) 011801, [arXiv:1003.4451](#) [hep-ph].
- [13] T. Han and S. Willenbrock, *QCD correction to the  $p p \rightarrow gt$ ;  $W H$  and  $Z H$  total cross-sections*, *Phys.Lett.* **B273** (1991) 167–172.
- [14] D. A. Dicus and S. Willenbrock, *Higgs Boson Production from Heavy Quark Fusion*, *Phys.Rev.* **D39** (1989) 751.
- [15] D. Dicus, T. Stelzer, Z. Sullivan, and S. Willenbrock, *Higgs boson production in association with bottom quarks at next-to-leading order*, *Phys.Rev.* **D59** (1999) 094016, [arXiv:hep-ph/9811492](#) [hep-ph].
- [16] F. Maltoni, Z. Sullivan, and S. Willenbrock, *Higgs-boson production via bottom-quark fusion*, *Phys.Rev.* **D67** (2003) 093005, [arXiv:hep-ph/0301033](#) [hep-ph].
- [17] F. I. Olness and W.-K. Tung, *When Is a Heavy Quark Not a Parton? Charged Higgs Production and Heavy Quark Mass Effects in the QCD Based Parton Model*, *Nucl.Phys.* **B308** (1988) 813.
- [18] J. Gunion, H. Haber, F. Paige, W.-K. Tung, and S. Willenbrock, *Neutral and Charged Higgs Detection: Heavy Quark Fusion, Top Quark Mass Dependence and Rare Decays*, *Nucl.Phys.* **B294** (1987) 621.
- [19] R. V. Harlander and W. B. Kilgore, *Higgs boson production in bottom quark fusion at next-to-next-to leading order*, *Phys.Rev.* **D68** (2003) 013001, [arXiv:hep-ph/0304035](#) [hep-ph].
- [20] L. Reina and S. Dawson, *Next-to-leading order results for  $t$  anti- $t$   $h$  production at the Tevatron*, *Phys.Rev.Lett.* **87** (2001) 201804, [arXiv:hep-ph/0107101](#) [hep-ph].
- [21] W. Beenakker, S. Dittmaier, M. Kramer, B. Plumper, M. Spira, *et al.*, *Higgs radiation off top quarks at the Tevatron and the LHC*, *Phys.Rev.Lett.* **87** (2001) 201805, [arXiv:hep-ph/0107081](#) [hep-ph].
- [22] S. Dawson, L. Orr, L. Reina, and D. Wackerroth, *Associated top quark Higgs boson production at the LHC*, *Phys.Rev.* **D67** (2003) 071503, [arXiv:hep-ph/0211438](#) [hep-ph].
- [23] W. Beenakker, S. Dittmaier, M. Kramer, B. Plumper, M. Spira, *et al.*, *NLO QCD corrections to  $t$  anti- $t$   $H$  production in hadron collisions*, *Nucl.Phys.* **B653** (2003) 151–203, [arXiv:hep-ph/0211352](#) [hep-ph].
- [24] R. Raitio and W. W. Wada, *Higgs Boson Production at Large Transverse Momentum in QCD*, *Phys.Rev.* **D19** (1979) 941.
- [25] Z. Kunszt, *Associated Production of Heavy Higgs Boson with Top Quarks*, *Nucl.Phys.* **B247** (1984) 339.
- [26] S. Catani, D. de Florian, M. Grazzini, and P. Nason, *Soft gluon resummation for Higgs boson production at hadron colliders*, *JHEP* **0307** (2003) 028, [arXiv:hep-ph/0306211](#) [hep-ph].
- [27] U. Aglietti, R. Bonciani, G. Degrossi, and A. Vicini, *Two loop light fermion contribution to Higgs production and decays*, *Phys.Lett.* **B595** (2004) 432–441, [arXiv:hep-ph/0404071](#) [hep-ph].

- [28] S. Actis, G. Passarino, C. Sturm, and S. Uccirati, *NLO Electroweak Corrections to Higgs Boson Production at Hadron Colliders*, *Phys.Lett.* **B670** (2008) 12–17, [arXiv:0809.1301 \[hep-ph\]](#).
- [29] G. Altarelli, R. K. Ellis, and G. Martinelli, *Leptoproduction and Drell-Yan Processes Beyond the Leading Approximation in Chromodynamics*, *Nucl.Phys.* **B143** (1978) 521.
- [30] T. Matsuura and W. van Neerven, *Second Order Logarithmic Corrections to the Drell-Yan Cross-section*, *Z.Phys.* **C38** (1988) 623.
- [31] T. Matsuura, S. van der Marck, and W. van Neerven, *The Calculation of the Second Order Soft and Virtual Contributions to the Drell-Yan Cross-Section*, *Nucl.Phys.* **B319** (1989) 570.
- [32] R. Hamberg, W. van Neerven, and T. Matsuura, *A Complete calculation of the order  $\alpha - s^2$  correction to the Drell-Yan K factor*, *Nucl.Phys.* **B359** (1991) 343–405.
- [33] S. Moch, J. Vermaseren, and A. Vogt, *The Quark form-factor at higher orders*, *JHEP* **0508** (2005) 049, [arXiv:hep-ph/0507039 \[hep-ph\]](#).
- [34] S. Moch, J. Vermaseren, and A. Vogt, *Three-loop results for quark and gluon form-factors*, *Phys.Lett.* **B625** (2005) 245–252, [arXiv:hep-ph/0508055 \[hep-ph\]](#).
- [35] T. Gehrmann, T. Huber, and D. Maitre, *Two-loop quark and gluon form-factors in dimensional regularisation*, *Phys.Lett.* **B622** (2005) 295–302, [arXiv:hep-ph/0507061 \[hep-ph\]](#).
- [36] P. Baikov, K. Chetyrkin, A. Smirnov, V. Smirnov, and M. Steinhauser, *Quark and gluon form factors to three loops*, *Phys.Rev.Lett.* **102** (2009) 212002, [arXiv:0902.3519 \[hep-ph\]](#).
- [37] T. Gehrmann, E. Glover, T. Huber, N. Ikizlerli, and C. Studerus, *Calculation of the quark and gluon form factors to three loops in QCD*, *JHEP* **1006** (2010) 094, [arXiv:1004.3653 \[hep-ph\]](#).
- [38] S. Moch, J. Vermaseren, and A. Vogt, *The Three loop splitting functions in QCD: The Nonsinglet case*, *Nucl.Phys.* **B688** (2004) 101–134, [arXiv:hep-ph/0403192 \[hep-ph\]](#).
- [39] K. Chetyrkin, B. A. Kniehl, and M. Steinhauser, *Decoupling relations to  $O(\alpha-s^{**3})$  and their connection to low-energy theorems*, *Nucl.Phys.* **B510** (1998) 61–87, [arXiv:hep-ph/9708255 \[hep-ph\]](#).
- [40] D. de Florian and J. Mazzitelli, *A next-to-next-to-leading order calculation of soft-virtual cross sections*, *JHEP* **1212** (2012) 088, [arXiv:1209.0673 \[hep-ph\]](#).
- [41] S. Moch and A. Vogt, *Higher-order soft corrections to lepton pair and Higgs boson production*, *Phys.Lett.* **B631** (2005) 48–57, [arXiv:hep-ph/0508265 \[hep-ph\]](#).
- [42] E. Laenen and L. Magnea, *Threshold resummation for electroweak annihilation from DIS data*, *Phys.Lett.* **B632** (2006) 270–276, [arXiv:hep-ph/0508284 \[hep-ph\]](#).
- [43] A. Idilbi, X.-d. Ji, J.-P. Ma, and F. Yuan, *Threshold resummation for Higgs production in effective field theory*, *Phys.Rev.* **D73** (2006) 077501, [arXiv:hep-ph/0509294 \[hep-ph\]](#).
- [44] V. Ravindran, *On Sudakov and soft resummations in QCD*, *Nucl.Phys.* **B746** (2006) 58–76, [arXiv:hep-ph/0512249 \[hep-ph\]](#).
- [45] V. Ravindran, *Higher-order threshold effects to inclusive processes in QCD*, *Nucl.Phys.* **B752** (2006) 173–196, [arXiv:hep-ph/0603041 \[hep-ph\]](#).

- [46] W. B. Kilgore, *One-Loop Single-Real-Emission Contributions to  $pp \rightarrow H + X$  at Next-to-Next-to-Next-to-Leading Order*, *Phys.Rev.* **D89** (2014) 073008, [arXiv:1312.1296 \[hep-ph\]](#).
- [47] C. Anastasiou, C. Duhr, F. Dulat, F. Herzog, and B. Mistlberger, *Real-virtual contributions to the inclusive Higgs cross-section at  $N^3LO$* , *JHEP* **1312** (2013) 088, [arXiv:1311.1425 \[hep-ph\]](#).
- [48] C. Duhr, T. Gehrmann, and M. Jaquier, *Two-loop splitting amplitudes and the single-real contribution to inclusive Higgs production at  $N^3LO$* , [arXiv:1411.3587 \[hep-ph\]](#).
- [49] F. Dulat and B. Mistlberger, *Real-Virtual-Virtual contributions to the inclusive Higgs cross section at  $N^3LO$* , [arXiv:1411.3586 \[hep-ph\]](#).
- [50] C. Anastasiou, C. Duhr, F. Dulat, E. Furlan, T. Gehrmann, *et al.*, *Higgs boson gluonfusion production at threshold in  $N^3LO$  QCD*, *Phys.Lett.* **B737** (2014) 325–328, [arXiv:1403.4616 \[hep-ph\]](#).
- [51] T. Ahmed, M. Mahakhud, N. Rana, and V. Ravindran, *Drell-Yan production at threshold in  $N^3LO$  QCD*, *Phys.Rev.Lett.* **113** (2014) 112002, [arXiv:1404.0366 \[hep-ph\]](#).
- [52] Y. Li, A. von Manteuffel, R. M. Schabinger, and H. X. Zhu,  *$N^3LO$  Higgs and Drell-Yan production at threshold: the one-loop two-emission contribution*, *Phys.Rev.* **D90** (2014) 053006, [arXiv:1404.5839 \[hep-ph\]](#).
- [53] S. Catani, L. Cieri, D. de Florian, G. Ferrera, and M. Grazzini, *Threshold resummation at  $N^3LL$  accuracy and soft-virtual cross sections at  $N^3LO$* , *Nucl.Phys.* **B888** (2014) 75–91, [arXiv:1405.4827 \[hep-ph\]](#).
- [54] N. Lo Presti, A. Almasy, and A. Vogt, *Leading large- $x$  logarithms of the quarkgluon contributions to inclusive Higgs-boson and lepton-pair production*, *Phys.Lett.* **B737** (2014) 120–123, [arXiv:1407.1553 \[hep-ph\]](#).
- [55] D. de Florian, J. Mazzitelli, S. Moch, and A. Vogt, *Approximate  $N^3LO$  Higgs-boson production cross section using physical-kernel constraints*, *JHEP* **1410** (2014) 176, [arXiv:1408.6277 \[hep-ph\]](#).
- [56] C. Anastasiou, C. Duhr, F. Dulat, E. Furlan, T. Gehrmann, *et al.*, *Higgs boson gluon-fusion production beyond threshold in  $N^3LO$  QCD*, [arXiv:1411.3584 \[hep-ph\]](#).
- [57] T. Gehrmann and D. Kara, *The  $Hb\bar{b}$  form factor to three loops in QCD*, *JHEP* **09** (2014) 174, [arXiv:1407.8114 \[hep-ph\]](#).
- [58] N. Kidonakis, *Collinear and soft gluon corrections to Higgs production at NNNLO*, *Phys.Rev.* **D77** (2008) 053008, [arXiv:0711.0142 \[hep-ph\]](#).
- [59] T. Ahmed, N. Rana, and V. Ravindran, *Higgs boson production through  $b\bar{b}$  annihilation at threshold in  $N^3LO$  QCD*, *JHEP* **1410** (2014) 139, [arXiv:1408.0787 \[hep-ph\]](#).
- [60] C. Anastasiou, L. J. Dixon, K. Melnikov, and F. Petriello, *Dilepton rapidity distribution in the Drell-Yan process at NNLO in QCD*, *Phys.Rev.Lett.* **91** (2003) 182002, [arXiv:hep-ph/0306192 \[hep-ph\]](#).
- [61] C. Anastasiou, K. Melnikov, and F. Petriello, *Higgs boson production at hadron colliders: Differential cross sections through next-to-next-to-leading order*, *Phys.Rev.Lett.* **93** (2004) 262002, [arXiv:hep-ph/0409088 \[hep-ph\]](#).

- [62] C. Anastasiou, K. Melnikov, and F. Petriello, *Fully differential Higgs boson production and the di-photon signal through next-to-next-to-leading order*, *Nucl.Phys.* **B724** (2005) 197–246, [arXiv:hep-ph/0501130](#) [hep-ph].
- [63] S. Bhler, F. Herzog, A. Lazopoulos, and R. Miller, *The fully differential hadronic production of a Higgs boson via bottom quark fusion at NNLO*, *JHEP* **1207** (2012) 115, [arXiv:1204.4415](#) [hep-ph].
- [64] G. Ferrera, M. Grazzini, and F. Tramontano, *Associated WH production at hadron colliders: a fully exclusive QCD calculation at NNLO*, *Phys.Rev.Lett.* **107** (2011) 152003, [arXiv:1107.1164](#) [hep-ph].
- [65] G. Ferrera, M. Grazzini, and F. Tramontano, *Higher-order QCD effects for associated WH production and decay at the LHC*, *JHEP* **1404** (2014) 039, [arXiv:1312.1669](#) [hep-ph].
- [66] V. Ravindran, J. Smith, and W. van Neerven, *QCD threshold corrections to di-lepton and Higgs rapidity distributions beyond  $N^2$  LO*, *Nucl.Phys.* **B767** (2007) 100–129, [arXiv:hep-ph/0608308](#) [hep-ph].
- [67] T. Ahmed, M. Mandal, N. Rana, and V. Ravindran, *Rapidity distributions in Drell-Yan and Higgs productions at threshold in  $N^3$  LO QCD*, [arXiv:1404.6504](#) [hep-ph].
- [68] D. Choudhury, S. Majhi, and V. Ravindran, *NLO corrections to lepton pair production beyond the standard model at hadron colliders*, *JHEP* **0601** (2006) 027, [arXiv:hep-ph/0509057](#) [hep-ph].
- [69] O. Tarasov, A. Vladimirov, and A. Y. Zharkov, *The Gell-Mann-Low Function of QCD in the Three Loop Approximation*, *Phys.Lett.* **B93** (1980) 429–432.
- [70] O. Tarasov, *ANOMALOUS DIMENSIONS OF QUARK MASSES IN THREE LOOP APPROXIMATION*, .
- [71] T. van Ritbergen, J. Vermaseren, and S. Larin, *The Four loop beta function in quantum chromodynamics*, *Phys.Lett.* **B400** (1997) 379–384, [arXiv:hep-ph/9701390](#) [hep-ph].
- [72] M. Czakon, *The Four-loop QCD beta-function and anomalous dimensions*, *Nucl.Phys.* **B710** (2005) 485–498, [arXiv:hep-ph/0411261](#) [hep-ph].
- [73] V. Sudakov, *Vertex parts at very high-energies in quantum electrodynamics*, *Sov.Phys.JETP* **3** (1956) 65–71.
- [74] A. H. Mueller, *On the Asymptotic Behavior of the Sudakov Form-factor*, *Phys.Rev.* **D20** (1979) 2037.
- [75] J. C. Collins, *Algorithm to Compute Corrections to the Sudakov Form-factor*, *Phys.Rev.* **D22** (1980) 1478.
- [76] A. Sen, *Asymptotic Behavior of the Sudakov Form-Factor in QCD*, *Phys.Rev.* **D24** (1981) 3281.
- [77] A. Vogt, S. Moch, and J. Vermaseren, *The Three-loop splitting functions in QCD: The Singlet case*, *Nucl.Phys.* **B691** (2004) 129–181, [arXiv:hep-ph/0404111](#) [hep-ph].
- [78] S. Catani and L. Trentadue, *Resummation of the QCD Perturbative Series for Hard Processes*, *Nucl.Phys.* **B327** (1989) 323.
- [79] S. Catani and L. Trentadue, *Comment on QCD exponentiation at large  $x$* , *Nucl.Phys.* **B353** (1991) 183–186.

- [80] A. Vogt, *Next-to-next-to-leading logarithmic threshold resummation for deep inelastic scattering and the Drell-Yan process*, *Phys.Lett.* **B497** (2001) 228–234, [arXiv:hep-ph/0010146](#) [hep-ph].
- [81] V. Ravindran, J. Smith, and W. van Neerven, *Two-loop corrections to Higgs boson production*, *Nucl.Phys.* **B704** (2005) 332–348, [arXiv:hep-ph/0408315](#) [hep-ph].
- [82] T. Becher and M. Neubert, *Infrared singularities of scattering amplitudes in perturbative QCD*, *Phys.Rev.Lett.* **102** no. 19, (2009) 162001, [arXiv:0901.0722](#) [hep-ph].
- [83] E. Gardi and L. Magnea, *Factorization constraints for soft anomalous dimensions in QCD scattering amplitudes*, *JHEP* **0903** (2009) 079, [arXiv:0901.1091](#) [hep-ph].
- [84] J. Vermaseren, S. Larin, and T. van Ritbergen, *The four loop quark mass anomalous dimension and the invariant quark mass*, *Phys.Lett.* **B405** (1997) 327–333, [arXiv:hep-ph/9703284](#) [hep-ph].
- [85] A. Martin, W. Stirling, R. Thorne, and G. Watt, *Parton distributions for the LHC*, *Eur.Phys.J.* **C63** (2009) 189–285, [arXiv:0901.0002](#) [hep-ph].