

## Author's Proof

Before checking your proof, please read the instructions below

- Carefully read the entire proof and mark all corrections in the appropriate place, using the Adobe Reader commenting tools ([Adobe Help](#)).
- Provide your corrections in a single PDF file or post your comments in the Production Forum making sure to reference the relevant query/line number. Upload or post all your corrections directly in the Production Forum to avoid any comments being missed.
- We do not accept corrections in the form of edited manuscripts nor via email.
- Do not provide scanned or handwritten corrections.
- Before you submit your corrections, please make sure that you have checked your proof carefully as once you approve it, you won't be able to make any further corrections.
- To ensure the timely publication of your article, please submit the corrections within 48 hours. After submitting, do not email or query asking for confirmation of receipt.

Do you need help? Visit our [Production Help Center](#) for more information. If you can't find an answer to your question, contact your Production team directly by posting in the Production Forum.

### Quick Check-List

- Author names** - Complete, accurate and consistent with your previous publications
- Affiliations** - Complete and accurate. Follow this style when applicable: Department, Institute, University, City, Country
- Tables** - Make sure our formatting style did not change the meaning/alignment of your Tables.
- Figures** - Make sure we are using the latest versions.
- Funding and Acknowledgments** - List all relevant funders and acknowledgments.
- Conflict of Interest** - Ensure any relevant conflicts are declared.
- Supplementary files** - Ensure the latest files are published and that no line numbers and tracked changes are visible. Also, the supplementary files should be cited in the article body text.
- Queries** - Reply to all typesetters queries below
- Content** - Read all content carefully and ensure any necessary corrections are made.

## Author Queries Form

<b>Q1</b>	Confirm whether the insertion of the article title is correct.	
<b>Q2</b>	The emphases (colored text) from revisions were removed throughout the article. Confirm that this change is fine.	
<b>Q3</b>	Confirm that the short running title (top right corner starting from the 2nd page) is correct, making sure to keep it to a maximum of five words.	
<b>Q4</b>	The citation and surnames of all of the authors have been highlighted. Check that they are correct and consistent with the authors' previous publications, and correct if need be. Please note that this may affect the indexing of your article in repositories such as PubMed.	
<b>Q5</b>	Confirm that the email address in your correspondence section is accurate. Please note that any changes to the corresponding authorship would require individual confirmation from all original and added/removed corresponding authors.	

<b>Q6</b>	Please ask the following authors to register with Frontiers (at <a href="https://www.frontiersin.org/Registration/Register.aspx">https://www.frontiersin.org/Registration/Register.aspx</a> ) if they would like their LOOP profile to be linked to the final published version. Please ensure to provide us with the profile link(s) (not email addresses) when submitting the proof corrections. Non-registered authors and authors with profiles set to private mode will have the default profile image displayed. "Jingting Li"	
<b>Q7</b>	Confirm that all author affiliations are correctly listed. Note that affiliations are listed sequentially as per journal style and requests for non-sequential listing will not be applied. Note that affiliations should reflect those at the time during which the work was undertaken.	
<b>Q8</b>	Provide the expanded form of "IRCCS".	
<b>Q9</b>	Confirm that the keywords are correct and keep them to a maximum of eight and a minimum of five. (Note: a keyword can be comprised of one or more words.) Note that we have used the keywords provided at Submission. If this is not the latest version, please let us know.	
<b>Q10</b>	Check if the section headers (i.e., section leveling) were correctly captured.	
<b>Q11</b>	If you decide to use previously published, <a href="#">copyrighted figures</a> in your article, please keep in mind that it is your responsibility, as the author, to obtain the appropriate permissions and licenses and to follow any citation instructions requested by third-party rights holders. If obtaining the reproduction rights involves the payment of a fee, these charges are to be paid by the authors.	
<b>Q12</b>	Ensure that all the figures, tables and captions are correct, and that all figures are of the highest quality/resolution. Please note that Figures and Tables must be cited sequentially, as per section 2.2 of the <a href="#">author guidelines</a>	
<b>Q13</b>	Verify that all the equations and special characters are displayed correctly.	
<b>Q14</b>	Confirm that the Data Availability statement is accurate. Note that we have used the statement provided at Submission. If this is not the latest version, please let us know.	
<b>Q15</b>	Ensure to add all grant numbers and funding information, as after publication this will no longer be possible. All funders should be credited and all grant numbers should be correctly included in this section.	
<b>Q16</b>	Confirm whether the insertion of the Ethics Statement section is fine. Note that we have used the statement provided at Submission. If this is not the latest version, please let us know.	

<b>Q17</b>	<p>Ensure that any supplementary material is correctly published at this link:  <a href="https://www.frontiersin.org/articles/10.3389/fendo.2022.861869/full#supplementary-material">https://www.frontiersin.org/articles/10.3389/fendo.2022.861869/full#supplementary-material</a></p> <p>If the link does not work, you can check the file(s) directly in the production forum; the published supplementary files appear in green.</p> <p>Provide new files if you have any corrections and make sure all Supplementary files are cited. Please also provide captions for these files, if relevant.</p> <p>Frontiers will deposit ALL supplementary files to FigShare and they will receive a DOI.</p> <p>Notify us of any previously deposited material.</p> <p>If the Supplementary Material files contain identifiable images, please keep in mind that it is your responsibility, as the author, to ensure you have permission to use the images in the article. Please check this <a href="#">link</a> for author's responsibility for publication of identifiable images.</p>	
<b>Q18</b>	Confirm if the edit made in the "Author Contributions" section is correct.	
<b>Q19</b>	Frontiers guidelines require listing the first 6 authors + et al. for articles with more than 6 authors. Please provide the names of the other 3 authors for the following references. 1–8, 10–17, 19, 20, 22–24, 27, 28, 31.	
<b>Q20</b>	Provide the volume number for the following reference. 11.	
<b>Q21</b>	Confirm if the text included in the Conflict of Interest statement is correct.	



# Prognostic Impact of Inflammatory Markers PLR, LMR, PDW, MPV in Medullary Thyroid Carcinoma

Canxiao Li<sup>1</sup>, Han Zhang<sup>1</sup>, Shijie Li<sup>1</sup>, Daqi Zhang<sup>1</sup>, Jingting Li<sup>1</sup>, Gianlorenzo Dionigi<sup>2</sup>, Nan Liang<sup>1\*\*</sup> and Hui Sun<sup>1\*\*</sup>

<sup>1</sup> Division of Thyroid Surgery, The China-Japan Union Hospital of Jilin University, Jilin Provincial Key Laboratory of Surgical Translational Medicine, Jilin Provincial Precision Medicine Laboratory of Molecular Biology and Translational Medicine on Differentiated Thyroid Carcinoma, Changchun, China, <sup>2</sup> Division of General and Endocrine Surgery, Istituto Auxologico Italiano IRCCS, Department of Medical Biotechnology and Translational Medicine, University of Milan, Milan, Italy

## OPEN ACCESS

### Edited by:

Akira Sugawara,  
Tohoku University, Japan

### Reviewed by:

Erkan Cüre,  
Bagcilar Medilife Hospital, Turkey  
Toru Takano,  
Osaka University, Japan

### \*Correspondence:

Hui Sun  
s\_h@jlu.edu.cn  
Nan Liang  
liangnan2006@jlu.edu.cn

### †ORCID:

Hui Sun  
orcid.org/0000-0001-8348-4933  
Nan Liang  
orcid.org/0000-0002-7733-862X

### Specialty section:

This article was submitted to  
Thyroid Endocrinology,  
a section of the journal  
Frontiers in Endocrinology

Received: 25 January 2022

Accepted: 11 February 2022

Published: xx xx 2022

### Citation:

Li C, Zhang H, Li S, Zhang D, Li J, Dionigi G, Liang N and Sun H (2022) Prognostic Impact of Inflammatory Markers PLR, LMR, PDW, MPV in Medullary Thyroid Carcinoma. *Front. Endocrinol.* 13:861869. doi: 10.3389/fendo.2022.861869

**Background:** Neutrophil-to-lymphocyte ratio (NLR), platelet-to-lymphocyte ratio (PLR), lymphocyte-to-monocyte ratio (LMR), mean platelet volume (MPV), and platelet distribution width (PDW) have been used as prognostic biomarkers in various cancers. We aim to investigate the relationship between the above inflammatory indices, clinicopathological features, and postoperative calcitonin (Ctn) progression in medullary thyroid carcinoma (MTC).

**Methods:** Sixty-eight patients diagnosed with MTC who underwent surgery at our institution between 2009 and 2020 were retrospectively evaluated. Areas under the receiver operating characteristic curves (ROC) and logistic regression were applied to explore the potential risk factors.

**Results:** PDW was predictive of lymph node metastasis (LN) (AUC=0.645,  $P=0.044$ ), PLR, PDW, and MPV were predictive of capsule invasion (AUC=0.771,  $P=0.045$ ; AUC=0.857,  $P=0.008$ ; and AUC =0.914,  $P=0.002$ , respectively), and MPV and LMR were predictive of postoperative Ctn progression (AUC=0.728,  $P=0.003$ ; AUC=0.657,  $P=0.040$ ). Multivariate analysis revealed that  $PDW \leq 16.4$  (OR=7.8, 95% CI: 1.532-39.720,  $P=0.013$ ) and largest tumor size  $\geq 1$  cm (OR=4.833, 95% CI: 1.514-15.427,  $P=0.008$ ) were potential independent risk factors for lateral LN metastasis. We also found that,  $MPV \leq 8.2$  (OR=13.999, 95% CI: 2.842-68.965,  $P=0.001$ ),  $LMR \leq 4.7$  (OR=4.790, 95% CI: 1.034-22.187,  $P=0.045$ ), and N1 (OR=45.890, 95%CI:3.879-542.936,  $P=0.002$ ) were potential independent risk factors for postoperative Ctn progression. In addition, compared with the single indicator, the appropriate combination of MPV and LMR could improve the specificity and sensitivity of predicting postoperative Ctn progression.

**Conclusions:** PLR, LMR, PDW, and MPV were associated with clinicopathological features and postoperative Ctn progression in MTC, suggesting that those inflammatory indices might be potential biomarkers of MTC.

**Keywords:** inflammation-based score, medullary thyroid carcinoma, clinicopathological characteristics, calcitonin, biomarkers

**Q10** INTRODUCTION

Cancer-related inflammation plays a central role in the development and progression of malignancies (1). Specifically, in particular, an increasing number of studies have demonstrated the importance of host systemic inflammatory responses in clinical performance and tumor prognosis, and various inflammatory indices, including neutrophil-to-lymphocyte ratio (NLR), platelet-to-lymphocyte ratio (PLR), and lymphocyte-to-monocyte ratio (LMR), are now widely recognized as novel potential biomarkers for predicting cancer outcomes (2–4). Moreover, all inflammatory indices can be obtained as part of a routine complete blood count prior to surgery, so they do not impose an additional burden on patients. This suggests that an inflammation index has good applications in tumors.

Medullary thyroid carcinoma (MTC) originates from the parafollicular C-cells of the thyroid gland, which are derived from the neural crest (5). Although MTC accounts for only 1–2% of thyroid cancers, it is responsible for approximately 13% of all thyroid cancer-related deaths (6). However, there is still a lack of valid biomarkers to evaluate the clinicopathological features and prognosis of MTC. Given the potential of the aforementioned inflammatory indices, we aimed to investigate their potential applications in MTC.

Current American Thyroid Association guidelines recommend measuring serum calcitonin (Ctn) levels in patients diagnosed with histological MTC (5). However, the prognostic value of inflammatory index for postoperative Ctn progression has not been studied in MTC. Therefore, another aim of the present study was to clarify the value of inflammatory indices for postoperative progression of Ctn in MTC.

## MATERIALS AND METHODS

## Design of the Study

Retrospective observational study

## Setting

China-Japan Union Hospital of Jilin College, China

## Time Frame

2009–2020

## Eligibility

The inclusion criteria were as follows: consecutive enrollment and first-time diagnosis of primary MTC, and confirmation of diagnosis by histological analysis and imaging techniques. The following patients were excluded: Patients who had previously received antitumor treatment, including chemotherapy, radiotherapy, and immunotherapy; patients with other primary cancers; follow-up for < 6 months; coinfection or other inflammatory diseases (except autoimmune thyroid diseases such as chronic lymphocytic thyroiditis), not resectable at the time of referral, and lack of relevant information. All patients

who met the inclusion criteria during the study period were included in the study. In addition, follow-up was performed, and clinical features and outcomes were assessed. No intervention was performed during the treatment period

## Patients and Data Collection

A total of 101 patients had an initial diagnosis of MTC. Sixty-eight patients had complete data and follow-up, including preoperative complete blood count, Ctn serum levels, and clinical and histopathological data (**Figure 1**).

## Ethics

This study was approved by the institutional ethics committee of the China-Japan Union Hospital of Ji Lin College. Written informed consent was obtained from all the patients.

## MTC Diagnosis

Tumor tissue samples were obtained from all patients for histological analysis or surgical resection. MTC diagnosis was based on guidelines (7). Tumor size corresponded to the largest tumor diameter measured on pathological examination after surgery.

## Measurement of Biomarkers

NLR is the ratio of neutrophil count to lymphocyte count, PLR is the ratio of platelet count to lymphocyte count, and LMR is the ratio of lymphocyte count to monocyte count. The above inflammatory indices were calculated based on the preoperative complete blood count; Ctn was detected by chemiluminescence: Ctn (REF: LKCL1) using the Immulite 1000 system (Siemens Medical Diagnostics Solutions). The normal range of Ctn levels was 0.15–6.00 pg/mL in females and 0.15–9.20 pg/mL in males. The upper detection limit was 585.0 pg/mL.

## Follow-Up

By December 30, 2020, the follow-up time was 8–133 months, and the median follow-up time was 79 months.


## Outcomes Recorded

Considering the different tumor load of different patients, the assessments time of postoperative ctn can be divided into 1 week, 1 month, 3 months and half a year. If it is lower than the lower limit of detection or within the normal reference range, regular postoperative reexamination is adopted. The initial reexamination period is half a year, and if the condition is stable, it is gradually extended to once a year.

Patients were divided into three groups based on postoperative serum Ctn levels (5, 8). *Remission*: Postoperative Ctn levels decreased to normal and remained constant. *Stable*: Postoperative Ctn levels were stable but did not decrease to normal values. *Progression*: Postoperative Ctn level increased to 150 pg/mL or doubling time was <12 months (**Supplementary Table 1**).

## Statistical Analysis

The SPSS package version 23.0 (SPSS) was used to analyze all data. All results are presented as mean  $\pm$  standard deviation (SD),



**FIGURE 1** | Flow diagram of the patients included in the current study.

median, and interquartile range (IQR) or number (%). Fisher's exact test and rank sum test were used to analyze categorical variables. Continuous variables were analyzed using an ANOVA test or, in the case of non-normal distributions, the Kruskal-Wallis test. Binary logistic regression models were constructed to predict MTC, and the factors that contributed to the outcome in univariate analysis had a  $P < 0.05$ , and statistical significance was set at  $P < 0.05$ . The receiver-operating characteristic curve (ROC curve) is a graphical representation that illustrates the diagnostic ability of a binary classification system with a varying discrimination threshold.

## RESULTS

### Characteristics of the Patients

As shown in **Table 1**, a total of 68 patients were enrolled in this study, including 30 males (44.1%) and 38 females (55.9%), with a mean age of  $48.3 \pm 11.3$  years. Six of them (8.8%) underwent lobectomy thyroidectomy+ CND, 2(2.9%) underwent lobectomy thyroidectomy + CND+ ipsilateral LND, 4(5.9%) underwent subtotal thyroidectomy+ CND, 9(13.2%) underwent subtotal thyroidectomy+ CND+ ipsilateral LND, 10(14.7%) underwent total thyroidectomy+ CND, 28(41.2%) underwent total

Q11  
Q12



**TABLE 1 |** Baseline clinicopathological characteristics of MTC patients.

Features	N (%)	Features	N (%)
<b>Total</b>	68	Yes	3 (4.4%)
<b>Sex</b>		<b>Thyroid function</b>	
Female	38 (59.1%)	euthyroid	57 (83.8%)
Male	30 (44.1%)	hypothyroid	2 (2.9%)
<b>Age (years)</b>	48.3 ± 11.3	hyperthyroid	0 (0%)
<b>Largest tumor size (cm)</b>	1.2 (0.6-2.2)	subclinical hypothyroid	7 (10.3%)
<b>Multifocality</b>		subclinical hyperthyroid	2 (2.9%)
No	57 (83.2%)	<b>Type of operation</b>	
Yes	11 (16.2%)	Lobectomy thyroidectomy+ CND	6 (8.8%)
<b>Bilateral</b>		Lobectomy thyroidectomy+ CND+ Ipsilateral LND	2 (2.9%)
No	60 (88.2%)	Subtotal thyroidectomy+ CND	4 (5.9%)
Yes	8 (11.8%)	Subtotal thyroidectomy+ CND+ Ipsilateral LND	9 (13.2%)
<b>T stage</b>		Total thyroidectomy+ CND	10 (14.7%)
T1+T2	62 (91.2%)	Total thyroidectomy+ CND+ Ipsilateral LND	28 (41.2%)
T3+T4	6 (8.8%)	Total thyroidectomy+ CND+ Bilateral LND	9 (13.2%)
<b>N stage</b>		<b>Calcitonin changes in follow-up</b>	
NO	34 (50.0%)	Remission	39 (57.4%)
N1a	7 (10.3%)	Stable	8 (11.8%)
N1b	27 (39.7%)	Progress	21 (30.9%)
<b>TNM stage</b>		<b>Inflammation index</b>	
I+II	34 (50%)	LMR	5.2 ± 1.7
III+IV	34 (50%)	NLR	1.7 ± 0.6
<b>Capsule invasion</b>		PLR	115.1 ± 31.2
NO	63 (92.6%)	MPV	9 (8.1-10.0)
YES	5 (7.4%)	PDW	16.1 (15.5-16.5)
<b>Hashimoto's thyroiditis</b>		<b>Follow up time (months)</b>	72.0 ± 32.4
No	65 (95.6%)		

CND, central neck dissection; LND, lateral neck dissection; LMR, lymphocyte-to-monocyte ratio; NLR, neutrophil-to-lymphocyte ratio; PLR, platelet-to-lymphocyte ratio; MPV, average platelet volume; PDW, platelet distribution width.

thyroidectomy+ CND+ ipsilateral LND, 9(13.2%) underwent total thyroidectomy+ CND+ bilateral LND and postoperative pathology confirmed that 5 (7.4%) had capsule invasion. Lymph node metastasis occurred in 34 (50.0%), of whom 7 (10.3%) had central lymph node metastasis and 27 (39.7%) had lateral neck lymph node metastasis. After the follow-up period (median 79 months), 39 (57.4%) patients were classified as being in remission, 8 (11.8%) as stable, 21 (30.9%) as progressive, and five patients as recurrence.


## Predictive Performance of Inflammatory Markers

As shown in **Figure 2**, we used the ROC curve to evaluate the predictive power of each inflammatory index for capsule invasion, lymph node (LN) metastasis, postoperative Ctn progression, and recurrence. The corresponding AUC values are shown in **Table 2**. For predicting capsule invasion, the AUCs of PLR, PDW and MPV were 0.771 (95% CI: 0.546–0.996,  $P=0.045$ ), 0.857 (95% CI: 0.726–0.988,  $P=0.008$ ) and 0.914 (95% CI: 0.833–0.996,  $P=0.002$ ), respectively. However, LMR and NLR showed relatively low discriminative power. We then checked the discriminative value of each inflammatory indices in LN metastases, central LN metastases, and lateral LN metastases and found that all inflammatory indices had no discriminative power for LN metastases or central LN metastases. However, for predicting lateral LN metastasis, PDW was found to have significant discriminatory power, with an AUC measurement of 0.645 ( $P=0.044$ ), and a cut-off value of 16.4. Finally, we also

examined the predictive performance of inflammatory indices for postoperative Ctn progression and recurrence, although none of them predicted recurrence. LMR and MPV were found to be significantly predictive of postoperative Ctn progression, with AUC values of 0.657 ( $P=0.040$ ) and 0.728 ( $P=0.003$ ), respectively. With an LMR cut-off value of 4.7, the sensitivity and specificity were 61.9% and 66.0%, respectively. With an MPV cut-off value of 8.2, specificity was 85.1%, and sensitivity was 61.9%.

## Logistic Regression Analysis of Risk Factors for Lateral LN Metastasis and Postoperative Ctn Progression

We performed univariate and multivariate analyzes to further illustrate the risk factors for lateral LN metastasis and postoperative progression of Ctn (**Table 3**). Univariate analysis revealed that bilateral tumors ( $P=0.011$ ), multifocality ( $P=0.035$ ), largest tumor size  $\geq 1$  cm ( $P=0.003$ ) and PDW  $\leq 16.4$  ( $P=0.018$ ) were significantly related to lateral LN metastasis; further multivariate regression analysis showed that PDW  $\leq 16.4$  and largest tumor size  $\geq 1$  cm were potential independent risk factors of lateral LN metastasis. Specifically, the risk of lateral LN metastasis was 7.8 times higher in patients with a PDW  $\leq 16.4$  than in patients with a PDW  $> 16.4$  (OR=7.8, 95% CI: 1.532–39.720,  $P=0.013$ ) and the risk of lateral LN metastasis in patients with the largest tumor size  $\geq 1$  cm was 4.8 times higher than in patients with the largest tumor size  $< 1$  cm (OR=4.833, 95% CI: 1.514–15.427,  $P=0.008$ ).



514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570



**TABLE 2** | Area under the ROC curve.

Features	AUC	95%CI	P-value	Cut-off	sensitivity	specificity
<b>Capsule invasion</b>						
LMR	0.749	0.540-0.958	0.065	/	/	/
PLR	0.771	0.546-0.996	0.045	138.3	0.800	0.857
NLR	0.562	0.307-0.816	0.647	/	/	/
MPV	0.914	0.833-0.996	0.002	10.0	1.000	0.810
PDW	0.857	0.726-0.988	0.008	14.5	0.800	0.857
<b>LN metastasis</b>						
LMR	0.523	0.385-0.661	0.745	/	/	/
PLR	0.433	0.295-0.571	0.342	/	/	/
NLR	0.423	0.285-0.562	0.278	/	/	/
MPV	0.535	0.396-0.675	0.615	/	/	/
PDW	0.629	0.495-0.763	0.067	/	/	/
<b>Central LN metastasis</b>						
LMR	0.506	0.324-0.688	0.96	/	/	/
PLR	0.616	0.425-0.807	0.318	/	/	/
NLR	0.527	0.333-0.721	0.816	/	/	/
MPV	0.452	0.198-0.706	0.679	/	/	/
PDW	0.527	0.319-0.735	0.816	/	/	/
<b>Lateral LN metastasis</b>						
LMR	0.526	0.382-0.671	0.716	/	/	/
PLR	0.475	0.329-0.620	0.726	/	/	/
NLR	0.430	0.290-0.570	0.334	/	/	/
MPV	0.519	0.374-0.663	0.797	/	/	/
PDW	0.645	0.516-0.775	0.044	16.4	0.926	0.390
<b>Calcitonin progress</b>						
LMR	0.657	0.513-0.801	0.040	4.7	0.619	0.660
PLR	0.511	0.357-0.664	0.889	/	/	/
NLR	0.474	0.328-0.620	0.735	/	/	/
MPV	0.728	0.574-0.881	0.003	8.2	0.619	0.851
PDW	0.528	0.389-0.668	0.710	/	/	/
<b>Recurrence</b>						
LMR	0.598	0.402-0.795	0.466	/	/	/
PLR	0.667	0.467-0.866	0.217	/	/	/
NLR	0.597	0.440-0.754	0.474	/	/	/
MPV	0.568	0.205-0.931	0.613	/	/	/
PDW	0.589	0.370-0.808	0.511	/	/	/

LMR, lymphocyte-to-monocyte ratio; NLR, neutrophil-to-lymphocyte ratio; PLR, platelet-to-lymphocyte ratio; MPV, average platelet volume; PDW, platelet distribution width.  
For  $P \leq 0.05$ , cut-off, sensitivity, and specificity are not shown.

Similarly, we also examined the risk factors for postoperative Ctn progression (Table 4). Univariate analysis revealed that the largest tumor size  $\geq 1$  cm ( $P=0.024$ ), LN metastasis ( $P=0.001$ ),  $MPV \leq 8.2$  ( $P < 0.01$ ), and  $LMR \leq 4.7$  ( $P=0.032$ ) were significantly associated with postoperative Ctn progression; further multivariate analysis revealed that  $MPV \leq 8.2$ ,  $LMR \leq 4.7$ , and N1 were potential independent risk factors for postoperative Ctn progression. Specifically, the risk of postoperative Ctn progression was 14.0 times higher in patients with  $MPV \leq 8.2$  than in patients with  $MPV > 8.2$  (OR=13.999, 95% CI: 2.842-68.965,  $P=0.001$ ), and it was 4.8 times higher than in patients with  $LMR > 4.7$  (OR=4.790, 95% CI: 1.034-22.187,  $P=0.045$ ), and in patients with LN metastases it was 45.9 times higher than in patients without metastases (OR=45.890, 95% CI: 3.879-542.936,  $P=0.002$ ).

### Predictive Performance of MPV and LMR Together for Postoperative Ctn Progression

Based on the above, we further examined the predictive power of the combination of MPV and LMR for postoperative Ctn

progression. As shown in Figure 3 and Table 5, according to the cut-off value obtained from the ROC curve, we defined either  $MPV \leq 8.2$  or  $LMR \leq 4.7$  as MPV-LMR and  $MPV \leq 8.2$  and  $LMR \leq 4.7$  simultaneously as MPV+LMR simultaneously. The sensitivity of MPV-LMR in predicting postoperative Ctn progression increased from 61.9% to 81%, while the specificity of MPV+LMR in predicting postoperative Ctn progression increased from 66% to 95.7%.

## DISCUSSION

The inflammatory response in the tumor microenvironment is involved in the dual process of tumor progression and suppression (9). Inflammatory indices are newly discovered tumor markers based on host inflammatory response. Studies have shown that they correlate with clinicopathological features and prognosis in pancreatic cancer, lung cancer, and anaplastic thyroid carcinoma (2, 4, 10). However, there are few reports on MTC. Therefore, in our study, we sought to clarify and compare the applications of inflammatory indices as MTC tumor markers.

**TABLE 3 |** Univariate and multivariate analyses of the risk factors for lateral LN metastasis.

Features	Non-N1b	N1b	Univariate p-value	Multivariate	
				OR(95%CI)	p-value
<b>Age (year)</b>			0.421		
≤55	18	31			
>55	9	10			
<b>Sex</b>			0.649		
female	16	22			
male	11	19			
<b>Bilateral</b>			0.011		
No	20	40			
Yes	7	1			
<b>Multifocality</b>			0.035		
No	19	38			
Yes	8	3			
<b>Capsule invasion</b>			>0.05		
No	25	38			
Yes	2	3			
<b>Largest tumor size (cm)</b>			0.003		
≤1	6	24		1	
>1	21	17		4.833 (1.514-15.427)	0.008
<b>PDW</b>			0.018		
PDW ≤ 16.4	2	13		7.8 (1.532-39.720)	0.013
PDW>16.4	25	28		1	

**TABLE 4 |** Univariate and multivariate analyses of the risk factors for calcitonin progression.

Features	Non-progressive group	Progressive group	Univariate p-value	Multivariate	
				OR (95%CI)	P-value
<b>Age (year)</b>			0.938		
≤55	34	15			
>55	13	6			
<b>Sex</b>			0.359		
female	28	10			
male	19	11			
<b>Largest tumor size (cm)</b>			0.024		
≤1	25	5			
>1	22	16			
<b>Bilateral</b>			0.981		
No	42	18			
Yes	5	3			
<b>Multifocality</b>			0.942		
No	40	17			
Yes	7	4			
<b>Capsule invasion</b>			0.294		
No	42	21			
Yes	5	0			
<b>Coexisting thyroiditis</b>			1		
No	45	20			
Yes	2	1			
<b>T stage</b>			0.336		
I+II	45	18			
III+IV	2	3			
<b>N stage</b>			0.001		
N0	30	14		1	
N1	17	17		45.890(3.879-542.936)	0.002
<b>N stage</b>			0.049		
N0+N1a	32	9			
N1b	15	12			
<b>TNM stage</b>			0.001		

(Continued)


TABLE 4 | Continued

Features	Non-progressive group	Progressive group	Univariate <i>p</i> -value	Multivariate	
				OR (95%CI)	<i>P</i> -value
I+II	30	4			
III+IV	17	17			
<b>MPV</b>			<0.01		
>8.2	10	14		1	
≤8.2	37	7		13.999(2.842-68.965)	0.001
<b>LMR</b>	13/8	16/31	0.032		
>4.7	16	13		1	
≤4.7	31	8		4.790(1.034-22.187)	0.045

In addition, we investigated the relationship between inflammatory indices and postoperative Ctn progression for the first time.

The development of a scored-inflammatory marker system established here is useful to predict the invasiveness and progress of MTC patients. The inflammatory models established in this study can be used to predict prognosis.

The ROC analysis showed that PDW was significantly associated with lateral LN metastasis, preoperative PLR, PDW, and MPV were predictive of invasion, and MPV and LMR were predictive of postoperative Ctn progression. Further multivariate analysis revealed PDW ≤16.4 was a potential independent factor for predicting lateral LN metastasis, and LMR ≤4.7 and MPV ≤8.2 were potential independent factors for predicting postoperative Ctn progression. Notably, the combination of LMR and MPV has improved sensitivity and specificity for predicting postoperative Ctn progression.



**FIGURE 3** | ROC curves for the combination of preoperative MPV and LMR to predict calcitonin progression. LMR, lymphocyte-to-monocyte ratio; MPV, mean platelet volume. MPV-LMR, MPV ≤8.2 or LMR ≤4.7; MPV+LMR, MPV ≤8.2 and LMR ≤4.7.

Previous studies investigated the relationship between tumor invasion and inflammatory indices, but their conclusions were conflicting. After a retrospective analysis of data from 205 patients with papillary thyroid carcinoma (PTC) in Greece, Manatakis et al. found that the NLR was significantly higher in carcinomas with extrathyroidal invasion ( $2.74 \pm 1.24$  versus  $2.39 \pm 0.96$ ,  $P=0.04$ ) (11), but other studies reached different conclusions (12–16). Also in our study, invasion was found to be not significantly associated with preoperative NLR but significantly associated with high PLR, MPV and low PDW. The specific mechanism is not clear, although it may be possible that the release of various inflammatory mediators induces an increase in platelets, while activated platelets contribute to the release of platelet-derived growth factor, platelet-activating factor, and vascular endothelial growth factor, which accelerate tumor-related blood vessel formation and extracellular matrix degradation, promoting tumor growth and distant metastasis (17). Although lymphocytes are the smallest white blood cells, they can induce the production of various cytotoxins, such as perforin, which directly or indirectly exert anti-tumoral effects (18). To some extent, lymphopenia indicates a weakened antitumoral immune response, which lead to more invasion and poorer prognosis (19). Therefore, PLR could be a potential tumor marker for predicting invasion in MTC.

However, some studies have indicated that platelet activation is related to platelet size rather than platelet number (20). MPV and PDW are common indicators of platelet volume. MPV indicates the mean platelet size and reflects the platelet production rate and stimulation (21). Larger platelets have stronger metabolic and enzymatic activities than smaller platelets (22). After reviewing data from 45 patients with gastric cancer and 20 healthy controls, Osada et al. found that MPV was higher in patients with gastric cancer than in healthy controls (23). However, Inagaki et al. found that the MPV of patients with advanced non-small cell lung cancer was lower than that of the control group (24). Similarly, our study suggests that a high MPV was predictive of invasion (AUC=0.914, 95% CI: 0.833–0.996,  $P=0.002$ ), but an MPV ≤8.2 (OR=13.999, 95% CI: 2.842–68.965,  $P=0.001$ ) was a potential independent risk factor for postoperative Ctn progression. This seemingly contradictory phenomenon highlights the complicated interactions between inflammatory factors in the tumor

**TABLE 5** | Area under the ROC curve.

	AUC	95%CI	P-value	Cut-off	Sensitivity	Specificity
MPV	0.728	0.574-0.881	0.003	8.2	0.619	0.851
LMR	0.657	0.513-0.801	0.040	4.7	0.619	0.660
MPV-LMR	0.649	0.512-0.786	0.050	0.5	0.810	0.489
MPV+LMR	0.717	0.570-0.864	0.050	0.5	0.476	0.957

MPV-LMR,  $MPV \leq 8.2$  or  $LMR \leq 4.7$ ; MPV+LMR,  $MPV \leq 8.2$  and  $LMR \leq 4.7$ .

LMR, lymphocyte-to-monocyte ratio; MPV, mean platelet volume.

microenvironment. Moreover, it could also be explained by the indolent nature of DTC.

The PDW reflects the uniformity of platelet volume, and higher values indicate that abnormally small and large platelets are in circulation (25). Compared with MPV, PDW is more reliable in evaluating the causes of thrombocytopenia (26). Cheng et al. found a lower PDW in gastric cancer patients compared with the healthy control group and that a PDW  $\leq 11.95$  was an independent risk factor for disease-free survival of T1 gastric cancer (27). Similarly, we found that PDW had predictive power for invasion and lateral LN metastasis, and that PDW  $\leq 16.4$  (OR=0.128; 95% CI: 0.025–0.653;  $P=0.013$ ) was a potential independent risk factor for lateral LN metastasis. This suggests that PDW is a potential tumor marker for MTC, and the lower its value, the worse the clinicopathological features of the tumor.

Moreover, our study was the first to find that an LMR  $\leq 4.7$  (OR=4.790; 95% CI: 1.034–22.187;  $P=0.045$ ) was a potential independent risk factor for postoperative Ctn progression. Similarly, some studies have indicated that low LMR is associated with poor prognosis of PTC and anaplastic thyroid carcinoma (4, 28). This could be explained by the fact that circulating monocytes can transform into myeloid suppressor cells and tumor-associated macrophages, which are involved in tumor growth, invasion, and metastasis by accelerating the epithelial-mesenchymal transition (29, 30). Finally, we investigated the influence of the appropriate combination of inflammatory indices on diagnostic efficiency. The analysis showed that the sensitivity of MPV-LMR in predicting postoperative Ctn progression increased from 61.9% to 81%, while the specificity of MPV+LMR in predicting postoperative Ctn progression increased from 66% to 95.7%.

Limited by the small number of patients with recurrence ( $n=5$ ), no significant association between LMR and recurrence was found in our study, but 5 patients were classified as postoperative Ctn progression, and some previous studies found that postoperative Ctn is significantly related to MTC prognosis (31), suggesting that LMR has potential for evaluating MTC prognosis.

Our study has several limitations. First, this was a retrospective analysis. Second, although patients with diseases that could influence complete blood count were excluded, there may have been some unknown or undetectable factors that could potentially influence our results. Finally, the number of cases included in our study was relatively small. Further large-scale and prospective studies are needed to confirm our conclusions.

## CONCLUSIONS

In conclusion, this study preliminarily explored the potential applications of inflammatory indices in the clinicopathological features and prognosis of MTC. We found that inflammatory indices were associated with the clinicopathological characteristics and postoperative Ctn progression in MTC. Specifically, PDW was predictive of lymph node metastasis, PLR, PDW, and MPV were predictive of capsule invasion, and MPV and LMR were predictive of postoperative Ctn progression. Remarkably, the appropriate combination of MPV and LMR could improve the specificity and sensitivity of predicting postoperative Ctn progression. Overall, this suggests that inflammatory indices are potential biomarkers for predicting the clinicopathological features and prognosis of medullary thyroid carcinoma.

## DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

## ETHICS STATEMENT

The studies involving human participants were reviewed and approved by China-Japan Union Hospital Institutional Review Board. Written informed consent for participation was not required for this study in accordance with the national legislation and the institutional requirements. Written informed consent was obtained from the individual(s) for the publication of any potentially identifiable images or data included in this article.

## AUTHOR CONTRIBUTIONS

HS, CL, and NL contributed to the study conception and design. Material preparation and data collection were carried out by CL, HZ, SL, DZ, and JL. All authors proposed many professional suggestions when data analysis. The first draft of the manuscript was written by CL. GD and NL contributed to manuscript review and editing. HS was responsible for project administration and supervision. All authors contributed to the article and approved the submitted version.

## FUNDING

This study was supported by the National Nature Science Foundation of China [81972499]; Jilin Province educational program [JJKH20211142KJ], the Jilin Province Science and Technology Development Program [20210402011GH]; the Program of Jilin Provincial Finance Department [2020SCZ03]; Jilin University Bethune Project [2020B14].

## ACKNOWLEDGMENTS

The authors sincerely thank all the clinicians in the division for their support. They were Guang Zhang, Le Zhou, Xuehai Bian,

## REFERENCES

- Moore MM, Chua W, Charles KA, et al. Inflammation and Cancer: Causes and Consequences. *Clin Pharmacol Ther* (2010) 87(4):504–8. doi: 10.1038/clpt.2009.254
- Chen Y, Wang W, Zhang X, et al. Prognostic Significance of Combined Preoperative Platelet-to-Lymphocyte Ratio and Lymphocyte-to-Monocyte Ratio in Patients Undergoing Surgery With Stage IB non-Small-Cell Lung Cancer. *Cancer Manag Res* (2018) 10:5411–22. doi: 10.2147/cmar.s177320
- Proctor MJ, Morrison DS, Talwar D, et al. A Comparison of Inflammation-Based Prognostic Scores in Patients With Cancer. A Glasgow Inflammation Outcome Study. *Eur J Cancer* (2011) 47(17):2633–41. doi: 10.1016/j.ejca.2011.03.028
- Ahn J, Song E, Oh HS, et al. Low Lymphocyte-To-Monocyte Ratios Are Associated With Poor Overall Survival in Anaplastic Thyroid Carcinoma Patients. *Thyroid* (2019) 29(6):824–9. doi: 10.1089/thy.2018.0684
- Wells SA Jr., Asa SL, Dralle H, et al. Revised American Thyroid Association Guidelines for the Management of Medullary Thyroid Carcinoma. *Thyroid* (2015) 25(6):567–610. doi: 10.1089/thy.2014.0335
- Randle RW, Balentine CJ, Levenson GE, et al. Trends in the Presentation, Treatment, and Survival of Patients With Medullary Thyroid Cancer Over the Past 30 Years. *Surgery* (2017) 161(1):137–46. doi: 10.1016/j.surg.2016.04.053
- Haddad RI, Nasr C, Bischoff L, et al. NCCN Guidelines Insights: Thyroid Carcinoma, Version 2.2018. *J Natl Compr Canc Netw* (2018) 16(12):1429–40. doi: 10.6004/jnccn.2018.0089
- Saltiki K, Simeakis G, Anagnostou E, et al. Different Outcomes in Sporadic Versus Familial Medullary Thyroid Cancer. *Head Neck* (2019) 41(1):154–61. doi: 10.1002/hed.25463
- Grivennikov SI, Greten FR, Karin M. Immunity, Inflammation, and Cancer. *Cell* (2010) 140(6):883–99. doi: 10.1016/j.cell.2010.01.025
- Lee BM, Chung SY, Chang JS, et al. The Neutrophil-Lymphocyte Ratio and Platelet-Lymphocyte Ratio Are Prognostic Factors in Patients With Locally Advanced Pancreatic Cancer Treated With Chemoradiotherapy. *Gut Liver* (2018) 12(3):342–52. doi: 10.5009/gnl17216
- Manatakis DK, Tseleni-Balafouta S, Balalis D, et al. Association of Baseline Neutrophil-To-Lymphocyte Ratio With Clinicopathological Characteristics of Papillary Thyroid Carcinoma. *Int J Endocrinol* (2017):8471235. doi: 10.1155/2017/8471235
- Liu CL, Lee JJ, Liu TP, et al. Blood Neutrophil-to-Lymphocyte Ratio Correlates With Tumor Size in Patients With Differentiated Thyroid Cancer. *J Surg Oncol* (2013) 107(5):493–7. doi: 10.1002/jso.23270
- Liu J, Du J, Fan J, et al. The Neutrophil-To-Lymphocyte Ratio Correlates With Age in Patients With Papillary Thyroid Carcinoma. *ORL J Otorhinolaryngol Relat Spec* (2015) 77(2):109–16. doi: 10.1159/000375534
- Kim JY, Park T, Jeong SH, et al. Prognostic Importance of Baseline Neutrophil to Lymphocyte Ratio in Patients With Advanced Papillary Thyroid Carcinomas. *Endocrine* (2014) 46(3):526–31. doi: 10.1007/s12020-013-0089-6
- Lang BH, Ng CP, Au KB, et al. Does Preoperative Neutrophil Lymphocyte Ratio Predict Risk of Recurrence and Occult Central Nodal Metastasis in

Xiaoli Liu, SL, Lina Zhao, JL, Qingfeng Fu, Hongbo Li, Yishen Zhao, Gaofeng Xue, Xueyan Zhang, Tie Wang, Changlin Li, Jiao Zhang, Jingwei Xin, and Tianyu Yu.

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fendo.2022.861869/full#supplementary-material>

**Supplementary Table 1** | Classification based on postoperative calcitonin.

- Papillary Thyroid Carcinoma? *World J Surg* (2014) 38(10):2605–12. doi: 10.1007/s00268-014-2630-z
- Kim SM, Kim EH, Kim BH, et al. Association of the Preoperative Neutrophil-To-Lymphocyte Count Ratio and Platelet-To-Lymphocyte Count Ratio With Clinicopathological Characteristics in Patients With Papillary Thyroid Cancer. *Endocrinol Metab (Seoul)* (2015) 30(4):494–501. doi: 10.3803/EnM.2015.30.4.494
- Schumacher D, Strilic B, Sivaraj KK, et al. Platelet-Derived Nucleotides Promote Tumor-Cell Transendothelial Migration and Metastasis via P2Y2 Receptor. *Cancer Cell* (2013) 24(1):130–7. doi: 10.1016/j.ccr.2013.05.008
- Dunn GP, Old LJ, Schreiber RD. The Immunobiology of Cancer Immunosurveillance and Immunoeediting. *Immunity* (2004) 21(2):137–48. doi: 10.1016/j.immuni.2004.07.017
- Ray-Coquard I, Cropet C, Van Glabbeke M, et al. Lymphopenia as a Prognostic Factor for Overall Survival in Advanced Carcinomas, Sarcomas, and Lymphomas. *Cancer Res* (2009) 69(13):5383–91. doi: 10.1158/0008-5472.can-08-3845
- Wen W, Wu P, Li J, et al. Predictive Values of the Selected Inflammatory Index in Elderly Patients With Papillary Thyroid Cancer. *J Transl Med* (2018) 16(1):261. doi: 10.1186/s12967-018-1636-y
- Threatte GA. Usefulness of the Mean Platelet Volume. *Clin Lab Med* (1993) 13(4):937–50. doi: 10.1016/S0272-2712(18)30418-9
- Mangalpally KK, Siqueiros-Garcia A, Vaduganathan M, et al. Platelet Activation Patterns in Platelet Size Sub-Populations: Differential Responses to Aspirin *In Vitro*. *J Thromb Thrombolysis* (2010) 30(3):251–62. doi: 10.1007/s1239-010-0489-x
- Osada J, Rusak M, Kamocki Z, et al. Platelet Activation in Patients With Advanced Gastric Cancer. *Neoplasma* (2010) 57(2):145–50. doi: 10.4149/neo\_2010\_02\_145
- Inagaki N, Kibata K, Tamaki T, et al. Prognostic Impact of the Mean Platelet Volume/Platelet Count Ratio in Terms of Survival in Advanced non-Small Cell Lung Cancer. *Lung Cancer* (2014) 83(1):97–101. doi: 10.1016/j.lungcan.2013.08.020
- Dincel O, Bayraktar C. Evaluation of Platelet Indices as a Useful Marker in Papillary Thyroid Carcinoma. *Bratisl Lek Listy* (2017) 118(3):153–5. doi: 10.4149/bl\_2017\_030
- Alsweedan SA, Al-Shurman A, Mahmoud AS. Diagnostic Value of Platelet Indices in Children With Leukemia. *J Pediatr Hematol Oncol* (2008) 30(12):953–5. doi: 10.1097/MPH.0b013e318182e7a9
- Cheng S, Han F, Wang Y, et al. The Red Distribution Width and the Platelet Distribution Width as Prognostic Predictors in Gastric Cancer. *BMC Gastroenterol* (2017) 17(1):163. doi: 10.1186/s12876-017-0685-7
- Yokota M, Katoh H, Nishimiya H, et al. Lymphocyte-Monocyte Ratio Significantly Predicts Recurrence in Papillary Thyroid Cancer. *J Surg Res* (2020) 246:535–43. doi: 10.1016/j.jss.2019.09.034
- Katoh H, Watanabe M. Myeloid-Derived Suppressor Cells and Therapeutic Strategies in Cancer. *Mediators Inflammation* (2015) 2015:159269. doi: 10.1155/2015/159269
- Marvel D, Gabrilovich DI. Myeloid-Derived Suppressor Cells in the Tumor Microenvironment: Expect the Unexpected. *J Clin Invest* (2015) 125(9):3356–64. doi: 10.1172/jci80005



- 1141 31. Pellegriti G, Leboulleux S, Baudin E, et al. Long-Term Outcome of Medullary  
 1142 Thyroid Carcinoma in Patients With Normal Postoperative Medical Imaging.  
 1143 *Br J Cancer* (2003) 88(10):1537–42. doi: 10.1038/sj.bjc.6600930

1144 **Q21** **Conflict of Interest:** The authors declare that the research was conducted in the  
 1145 absence of any commercial or financial relationships that could be construed as a  
 1146 potential conflict of interest.

1147 **Publisher's Note:** All claims expressed in this article are solely those of the authors  
 1148 and do not necessarily represent those of their affiliated organizations, or those of  
 1149

the publisher, the editors and the reviewers. Any product that may be evaluated in  
 this article, or claim that may be made by its manufacturer, is not guaranteed or  
 endorsed by the publisher.

*Copyright © 2022 Li, Zhang, Li, Zhang, Li, Dionigi, Liang and Sun. This is an open-  
 access article distributed under the terms of the Creative Commons Attribution License  
 (CC BY). The use, distribution or reproduction in other forums is permitted, provided the  
 original author(s) and the copyright owner(s) are credited and that the original  
 publication in this journal is cited, in accordance with accepted academic practice. No  
 use, distribution or reproduction is permitted which does not comply with these terms.*