## Daemonic ergotropy in continuously-monitored open quantum batteries

Daniele Morrone,<sup>1,2,\*</sup> Matteo A. C. Rossi,<sup>2,3,4</sup> and Marco G. Genoni<sup>1,†</sup>

<sup>1</sup>Quantum Technology Lab, Dipartimento di Fisica Aldo Pontremoli,

<sup>2</sup>Algorithmiq Ltd., Kanavakatu 3C, FI-00160 Helsinki, Finland

<sup>3</sup>InstituteQ - the Finnish Quantum Institute, Aalto University, Finland

<sup>4</sup>QTF Centre of Excellence, Department of Applied Physics, Aalto University, FI-00076 Aalto, Finland

(Dated: October 31, 2023)

The amount of work that can be extracted from a quantum system can be increased by exploiting the information obtained from a measurement performed on a correlated ancillary system. The concept of daemonic ergotropy has been introduced to properly describe and quantify this work extraction enhancement in the quantum regime. We here explore the application of this idea in the context of continuously-monitored open quantum systems, where information is gained by measuring the environment interacting with the energy-storing quantum device. We first show that the corresponding daemonic ergotropy takes values between the ergotropy and the energy of the corresponding unconditional state. The upper bound is achieved by assuming an initial pure state and a perfectly efficient projective measurement on the environment, independently of the kind of measurement performed. On the other hand, if the measurement is inefficient or the initial state is mixed, the daemonic ergotropy is generally dependent on the measurement strategy. This scenario is investigated via a paradigmatic example of an open quantum battery: a two-level atom driven by a classical field and whose spontaneously emitted photons are continuously monitored via either homodyne, heterodyne, or photo-detection.

### I. INTRODUCTION

The field of quantum thermodynamics aims to extend the laws of classical thermodynamics to the quantum regime [1]. One of its main goals is to understand the limits of work extraction from a quantum system, which has both fundamental and practical implications. Allahverdyan et al. [2] introduced the concept of ergotropy, as the maximum amount of work that can be extracted from a quantum state through unitary dynamics.

Extracting work from a quantum state is extremely useful in the context of quantum batteries (QBs), energystoring devices that follow the laws of quantum mechanics during charging and discharging processes. Research on QBs has then focused on exploring quantum enhancements in the charging process [3–23] and various implementations have been proposed, including single qubits, collective spins, and quantum harmonic oscillators [6, 12, 24–26]. Recent experiments have shown promising results in realizing a quantum-enhanced QB [27–30].

The analysis of open quantum batteries (OQBs), that is QBs in an open quantum system scenario, is crucial to discuss and guarantee their real-world implementation. Research on OQBs has focused so far on studying the effect of different environmental models [31–35] and devising quantum control strategies to counteract the impact of noise [19, 36–41]. We here want to address the situation where some information leaking to the environment can be recovered via continuous measurements [42, 43],



FIG. 1. Pictorial representation of a continuously-monitored quantum battery: a coherent drive supplies energy with intensity  $\alpha$  to a two-level atom (green) which is spontaneously emitting into its environment with rate  $\kappa$ . The emitted field is continuously monitored via a detector with efficiency  $\eta$ .

and then exploited for enhancing the work extraction protocol (see Fig. 1). Since the paradigmatic example of Maxwell's demon, it is well known that acquiring information via a measurement effectively brings a system out of equilibrium and allows to extract useful work via conditional operations [44]. This idea was then brought to the quantum realm, also highlighting the relationship between extractable work and quantum correlations [45– 56]. Francica et al. introduced the concept of *daemonic ergotropy* [52], as the maximum average work that can be extracted from a quantum system via unitary operations by exploiting the information obtained by measuring a correlated ancilla.

Continuously-monitored open quantum systems [42, 43] have been extensively studied from a theoretical point of view, mainly for feedback-assisted quantum state engineering protocols [57–73] and for quantum estimation purposes [74–91]; more recently their quantum thermodynamics properties have been also analyzed, being

Università degli Studi di Milano, I-20133 Milano, Italy

<sup>\*</sup> daniele.morrone@unimi.it

<sup>&</sup>lt;sup>†</sup> marco.genoni@fisica.unimi.it

a paradigmatic example of out-of-equilibrium quantum systems [89, 92–103]. Remarkably, the possibility of observing single trajectories has been now experimentally shown in different platforms, such as superconducting circuits [104–107], optomechanical [108, 109] and hybrid [110] systems. Recently, feedback protocols able to cool mechanical oscillators have also been demonstrated [111–113].

From a fundamental point of view, during its evolution, an open quantum system is correlated to the environment; for this reason, being able to measure the environment makes this scenario the ideal playground where to investigate the properties of the daemonic ergotropy both for its fundamental aspects, but also for its possible practical implementation in OQBs.

In this paper we derive the general properties of the daemonic ergotropy in open quantum systems and we apply them to the simplest example of an OQB, that is a two-level atom driven via a classical field and spontaneously emitting photons into its electromagnetic environment. We show that the daemonic ergotropy of a continuously-monitored system surpasses the ergotropy of the unconditional state and, with perfectly efficient measurements, can even reach the energy of the unconditional state. We then discuss the performance of different types of measurements with non-unit efficiency.

The manuscript is organized as follows: in Sec. II we review the concept of daemonic ergotropy and present our first results. In Sec. III we extend the concept of daemonic ergotropy to continuously-monitored open quantum systems, along with some general results that apply in this scenario. In Sec. IV we discuss these in a minimal example of an open quantum battery: a two level system driven by a classical field and whose spontaneous emitted photons are continuously monitored. Finally, in Sec. V, we give our conclusions and propose further outlooks.

### **II. DAEMONIC ERGOTROPY**

We start by recalling the definition and main properties of the ergotropy of a quantum state. Let us consider a quantum system described by a Hamiltonian  $\hat{H}_0$ , where, without losing generality, we fix its smallest eigenvalue equal to zero. We define the ergotropy  $\mathcal{E}(\varrho)$  of a given quantum state  $\varrho$  describing such quantum system as the maximum amount of work that can be extracted via unitary dynamics [2]:

$$\mathcal{E}(\varrho) = \max_{\hat{U}} \left[ E(\varrho) - E(\hat{U}\varrho\hat{U}^{\dagger}) \right], \qquad (1)$$

where  $E(\varrho) = \text{Tr}\left[\hat{H}_0 \varrho\right]$  denotes the average energy of the quantum state  $\varrho$ . A closed formula for  $\mathcal{E}(\varrho)$  in terms of eigenvalues and eigenvectors of  $\varrho$  and  $\hat{H}_0$  can be straightforwardly derived and the following main properties can be demonstrated: i) the ergotropy is upper bounded by the energy, and this upper bound is saturated for pure

quantum states, i.e  $\mathcal{E}(|\psi\rangle\langle\psi|) = E(|\psi\rangle\langle\psi|)$ ; ii) ergotropy is equal to zero  $\mathcal{E}(\varrho) = 0$  if and only if  $\varrho$  is a *passive state*, i.e. it is diagonal in the Hamiltonian  $\hat{H}_0$  eigenbasis, and its eigenvalues do not admit energy inversion [2].

Let us now consider a bipartite quantum state  $\rho^{SA}$ , where the quantum system S represents our energy storing device, while A is an ancillary system. If the ancilla is discarded, the maximum amount of extractable work is simply equal to  $\mathcal{E}(\rho^S)$ , with  $\rho^S = \text{Tr}_A[\rho^{SA}]$ . We now assume that a measurement is performed on the ancilla, described by a positive operator-valued measurement (POVM) { $\hat{\Pi}_a^A$ }. We can thus define the *daemonic ergotropy* as the average ergotropy of the corresponding conditional states [52]

$$\overline{\mathcal{E}}_{\{\hat{\Pi}_a^A\}} = \sum_a p_a \mathcal{E}(\varrho_a^S) \,, \tag{2}$$

where

$$p_a = \operatorname{Tr}_{SA}[\varrho^{SA}(\hat{\mathbb{1}}^S \otimes \hat{\Pi}_a^A)] \tag{3}$$

$$\varrho_a^S = \text{Tr}_A[\varrho^{SA}(\hat{\mathbb{1}}^S \otimes \hat{\Pi}_a^A)]/p_a \tag{4}$$

denote respectively the probability and the conditional state corresponding to the measurement outcome a. The ergotropy is a convex quantity in the quantum state  $\varrho$  [52, 55]; as a consequence, since  $\varrho^S = \sum_a p_a \varrho_a^S$ , one obtains that

$$\overline{\mathcal{E}}_{\{\hat{\Pi}_{\alpha}^{A}\}} \ge \mathcal{E}(\varrho^{S}), \qquad (5)$$

that is the average work that may be extracted is increased thanks to the information obtained from the ancilla. Furthermore, the daemonic ergotropy can be rewritten as

$$\overline{\mathcal{E}}_{\{\hat{\Pi}_a^A\}} = E(\varrho^S) - \sum_a p_a \min_{\hat{U}_a} E(\hat{U}_a \varrho_a^S \hat{U}_a^\dagger) \,. \tag{6}$$

From this formula it is then clear that, in order to achieve the daemonic enhancement, one needs to implement a conditional unitary evolution  $\hat{U}_a$  that depends on the conditional state  $\varrho_a^S$ . Furthermore,  $\overline{\mathcal{E}}_{\{\hat{\Pi}_a^A\}}$  generally depends on the specific POVM  $\{\hat{\Pi}_a^A\}$  implemented (see [55] for a recipe to obtain the optimal POVM maximizing  $\overline{\mathcal{E}}_{\{\hat{\Pi}_a^A\}}$ ). Before addressing the scenario of continuoslymonitored quantum systems, we here present the first result of our work via the following proposition.

**Proposition 1.** Given a bipartite system system+ancilla prepared in an initial pure state  $|\Psi\rangle^{SA}$ , and assuming a projective (rank-one) measurement on the ancilla  $\{\Pi_a^A = |\phi_a\rangle\langle\phi_a|\}$ , then

$$\overline{\mathcal{E}}_{\{\hat{\Pi}_a^A\}} = E(\varrho^S), \qquad (7)$$

that is, the daemonic ergotropy is equal to the energy of the reduced state of the system  $\varrho^S = \text{Tr}_A[\varrho^{SA}]$ , independently of the measurement performed. *Proof.* In order to prove this theorem, we first observe that the conditional state remains pure for any measurement outcome, i.e.  $\varrho_a^S = |\xi_a\rangle\langle\xi_a|$ , with

$$|\xi_a\rangle = \frac{1}{\sqrt{p_a}} \langle \phi_a | \Psi \rangle^{SA} \tag{8}$$

As a consequence one has

$$\overline{\mathcal{E}}_{\{\hat{\Pi}_{a}^{A}\}} = \sum_{a} p_{a} \mathcal{E}(|\xi_{a}\rangle\langle\xi_{a}|)$$
$$= \sum_{a} p_{a} E(|\xi_{a}\rangle\langle\xi_{a}|) = E(\varrho^{S}), \qquad (9)$$

where we have exploited: i) the fact that the ergotropy of pure states is equal to their energy, ii) the linearity of the trace, and iii) the relationship  $\rho^S = \sum_a p_a |\xi_a\rangle \langle \xi_a|$ .  $\Box$ 

### III. DAEMONIC ERGOTROPY IN CONTINUOUSLY-MONITORED OPEN QUANTUM SYSTEMS

We start by briefly introducing the basic notions on continuously-monitored quantum systems and quantum trajectories [42, 43]. We assume that our quantum system is interacting with a Markovian environment such that its unconditional evolution is described by a master equation in Lindblad form

$$d\varrho_{\rm unc}(t)/dt = -i[\hat{H}_s(t), \varrho_{\rm unc}(t)] + \mathcal{D}[\hat{c}]\varrho_{\rm unc}(t), \quad (10)$$

where  $\hat{H}_s(t)$  denotes the Hamiltonian ruling the evolution of the system and  $\mathcal{D}[\hat{c}]\varrho = \hat{c}\varrho\hat{c} - (\hat{c}^{\dagger}\hat{c}\varrho + \varrho\hat{c}^{\dagger}\hat{c})/2$  is the Lindbladian superoperator [114].

If one assumes that the environment is continuously monitored, the system evolution will be described by a stochastic master equation (SME) for the conditional state  $\rho_{c}(t)$ , which is typically referred to as a quantum trajectory. In general it is always true that

$$\varrho_{\rm unc}(t) = \sum_{\rm traj} p_{\rm traj} \, \varrho_{\rm c}(t), \tag{11}$$

where  $p_{\text{traj}}$  denotes the probability of each quantum trajectory  $\rho_{c}(t)$ . Different measurement strategies will correspond to different unravelling of the unconditional master equation (10), that is to different SMEs for the conditional state  $\rho_{c}(t)$  and, mathematically speaking, to different convex combinations for the unconditional state  $\rho_{unc}(t)$ . We will later consider the most paradigmatic examples of such unravellings, corresponding to the scenarios where the environment is continuously-monitored via either photo-detection (PD), homodyne-detection (HoD) or heterodyne-detection (HeD). The explicit formulas for the corresponding SMEs can be found in appendix A. Remarkably, while we here focus on Markovian open quantum systems described by a Lindblad master equation (10), all the results that follow apply whenever one can write the unconditional state as a mixture of trajectories as of (11), including monitored quantum systems exhibiting non-Markovian behaviour.

Besides the kind of detection performed on the environment, such unravellings are characterized by their measurement efficiency  $\eta$ , that comprehensively quantifies the portion of the environment that is accessible and the efficiency of the detector. In particular, we recall that for  $\eta = 1$ , that is when one assumes that the environment is fully accessible and measurable via a projective (rank-one) measurement, and for an initial pure state for the system, one can prove that the evolution can be described via a stochastic Schrödinger equation, and the conditional state remains pure during the whole dynamics [42, 43]. We can now present one of the main results via the following proposition:

**Proposition 2.** Given an open quantum system, whose charging process is described by a (possibly time-dependent) Hamiltonian  $\hat{H}_s(t)$ , and whose environment can be continuously monitored with efficiency  $\eta$ , the corresponding daemonic ergotropy

$$\overline{\mathcal{E}}_{\mathsf{unr},\eta}(t) = \sum_{a} p_{\mathsf{traj}} \mathcal{E}(\varrho_{\mathsf{c}}(t)) \tag{12}$$

is bounded as

$$\mathcal{E}(\varrho_{\mathsf{unc}}(t)) \le \overline{\mathcal{E}}_{\mathsf{unr},\eta}(t) \le E(\varrho_{\mathsf{unc}}(t)) \,. \tag{13}$$

The upper bound can be achieved in the presence of Markovian environment, whenever the system is initially prepared in a pure state and the monitoring is performed with unit efficiency  $\eta = 1$ , independently of the kind of unravelling, i.e.

$$\overline{\mathcal{E}}_{\mathsf{unr},\eta=1}(t) = E(\varrho_{\mathsf{unc}}(t)). \,. \tag{14}$$

*Proof.* The first inequality in Eq. (13) is a generalization of Eq. (5), applied to (11). Similarly the upper bound is a consequence of Eq. (6), while the fact that the upper bound can be achieved for unravellings of pure states follows straightforwardly from Prop. 1.  $\Box$ 

In the following, we show that as a corollary, the inequality above can be trivially extended to the case where ergotropies and energies are rescaled by the evolution time t, and thus in terms of figure of merits characterizing the charging power of the protocol.

# A. Average power of continuously-monitored quantum batteries

Let us assume that the state of the quantum battery at the beginning of the charging process is initially in a pure state  $(\varrho_0 = |\psi_0\rangle\langle\psi_0|)$ , and a charging protocol evolve the system from a time t = 0 to  $t = \tau$ . We can use the definitions of energy ( $\mathcal{E}$ ), ergotropy ( $\mathcal{E}$ ) and daemonic ergotropy ( $\overline{\mathcal{E}}_{\{\hat{\Pi}_{\tau}\}}$ ) introduced in the manuscript, to define respectively the average power (P), the average ergotropic power  $(\mathcal{P})$  and the average daemonic power  $(\overline{\mathcal{P}}_{\{\hat{\Pi}^A\}})$  as follows:

$$P(\varrho(\tau)) = \frac{E(\varrho(\tau)) - E(\varrho_0)}{\tau}$$
(15)

$$\mathcal{P}(\varrho(\tau)) = \frac{\mathcal{E}(\varrho(\tau)) - E(\varrho_0)}{\tau}$$
(16)

$$\overline{\mathcal{P}}_{\{\hat{\Pi}_a^A\}}(\tau) = \frac{\overline{\mathcal{E}}_{\{\hat{\Pi}_a^A\}}(\varrho(\tau)) - E(\varrho_0)}{\tau}$$
(17)

We can thus formulate the following corollary, that can be easily proven starting from Prop. 2 of the manuscript.

**Corollary 1.** Given an open quantum system, whose charging process is described by a (possibly time-dependent) Hamiltonian  $\hat{H}_s(t)$ , and whose environment can be continuously monitored with efficiency  $\eta$ , the corresponding average daemonic power  $\overline{\mathcal{P}}_{unr,\eta}(t) = \sum_a p_{traj} \mathcal{P}(\varrho_c(t))$  is bounded as

$$\mathcal{P}(\varrho_{\mathsf{unc}}(t)) \le \overline{\mathcal{P}}_{\mathsf{unr},\eta}(t) \le P(\varrho_{\mathsf{unc}}(t)) \,. \tag{18}$$

The upper bound can be achieved in the presence of Markovian environment, whenever the system is initially prepared in a pure state and the monitoring is performed with unit efficiency  $\eta = 1$ , independently of the kind of unravelling, i.e.

$$\overline{\mathcal{P}}_{\mathsf{unr},\eta=1}(t) = P(\varrho_{\mathsf{unc}}(t)).$$
(19)

We have thus proved that, as expected, the extractable work and power can be increased in an open quantum system if one obtains some information by monitoring the environment. Remarkably, the maximum daemonic ergotropy is equal to the unconditional energy and can be achieved via unit efficiency monitoring, independently of the measurement strategy. In the following we will rather investigate what happens in the more practical and experimentally relevant scenario of monitoring with nonunit efficiency, where a hierarchy between the different unravellings is established.

### IV. A CONTINUOUSLY-MONITORED OPEN QUANTUM BATTERY

We now consider the paradigmatic example of an OQB, that is a two-level atom characterized by a Hamiltonian  $\hat{H}_0 = (\omega_0/2)(\hat{\sigma}_z + 1)$ , driven by a resonant classical field of intensity  $\alpha$ , which acts as a charger, and spontaneously emitting with rate  $\kappa$  (see Fig. 1). In interaction picture with respect to  $\hat{H}_0$ , the evolution of the system is described by a Markovian master equation of the form (10), where the Hamiltonian ruling the evolution and the jump operator respectively read  $\hat{H}_s = \alpha \hat{\sigma}_x$  and  $\hat{c} = \sqrt{\kappa} \sigma_-$ , i.e.

$$\frac{d\varrho_{\mathsf{unc}}(t)}{dt} = -i\alpha[\hat{\sigma}_x, \varrho_{\mathsf{unc}}(t)] + \kappa \mathcal{D}[\hat{\sigma}_-]\varrho_{\mathsf{unc}}(t) \,. \tag{20}$$



FIG. 2. Daemonic ergotropies  $\overline{\mathcal{E}}_{unr,\eta}(t)$  as a function of time for different unravellings (HoD with photocurrents monitoring either  $\hat{\sigma}_x$  or  $\hat{\sigma}_y$ , HeD and PD) with  $\alpha/\kappa = 1$ ,  $\eta = 0.4$  and averaged over  $n = 5 \times 10^4$  trajectories. Black and gray dashed lines correspond respectively to the ergotropy and energy of the unconditional state. The different daemonic ergotropies lie as expected between these two lines, with HeD and  $\hat{\sigma}_x$ -HoD giving the larger values of extractable work. The upper and the lower bounds would be saturated for all the unravellings in the case of respectively perfect monitoring  $(\eta = 1)$  and nomonitoring  $(\eta = 0)$ .

An analytical solution can be obtained for  $\rho_{unc}(t)$  and we report here the corresponding steady-state values of energy and ergotropy

$$E_{\rm unc}^{\rm ss}/\omega_0 = \frac{4\alpha^2}{8\alpha^2 + \kappa}\,,\tag{21}$$

$$\mathcal{E}_{\text{unc}}^{\text{ss}}/\omega_0 = \frac{\kappa}{2} \left( \frac{\sqrt{16\alpha^2 + \kappa^2} - \kappa}{8\alpha^2 + \kappa^2} \right), \qquad (22)$$

where we have exploited the following formula for the ergotropy of a qubit state

$$\mathcal{E}(\varrho) = E(\varrho) + \frac{\omega_0}{2}\sqrt{2\mu(\varrho) - 1},$$
(23)

in terms of energy and purity  $\mu(\varrho) = \text{Tr}[\varrho^2]$ . One observes that the steady state energy  $E_{\text{unc}}^{\text{ss}}$  grows monotonically as a function of  $\alpha/\kappa$  and asymptotically reaches the value  $E_{\text{unc,max}}^{\text{ss}} = \omega_0/2$  in the limit of large driving. Oppositely, the steady-state ergotropy  $\mathcal{E}_{\text{unc}}^{\text{ss}}$  presents a maximum at

$$\alpha/\kappa = \sqrt{(1+\sqrt{2})/8}, \qquad (24)$$

where it reaches its peak value

$$\mathcal{E}_{unc,max}^{ss} = \omega_0(\sqrt{2} - 1)/2. \qquad (25)$$

If we now assume that a photo-counting detector is able to measure the spontaneously emitted photons with efficiency  $\eta$ , the dynamics of the conditional states  $\rho_{c}(t)$ is described by a SME of the form (A2); in particular the statistics of the corresponding Poissonian increment is univocally identified by its average value

$$\mathbb{E}[dN_t] = \eta \kappa \langle \hat{\sigma}_+ \hat{\sigma}_- \rangle_t dt \,, \tag{26}$$

and thus depends on the average value of  $\hat{\sigma}_z$  (we remind that  $\hat{\sigma}_+ \hat{\sigma}_- = (\hat{\sigma}_z + \hat{\mathbb{1}}_2)/2$ ).

If one rather considers a homodyne detection on the emitted field, one has a SME of the form (A3), where in particular the continuous homodyne photo-current reads

$$dy_t = \sqrt{\eta\kappa} \langle \hat{\sigma}_- e^{i\phi} + \hat{\sigma}_+ e^{-i\phi} \rangle_t \, dt + dW_t \,, \tag{27}$$

$$= \sqrt{\eta \kappa} \langle \cos \phi \hat{\sigma}_x + \sin \phi \hat{\sigma}_y \rangle_t \, dt + dW_t \,, \qquad (28)$$

where  $\phi$  corresponds to the homodyne phase. In particular for  $\phi = 0$  and  $\phi = \pi/2$  one has photocurrents depending respectively to the average values of  $\hat{\sigma}_x$  and  $\hat{\sigma}_y$ .

Heterodyne detection on the environment leads to a similar SME, as can be indeed thought and implemented as a double homodyne, measuring orthogonal quadratures with half efficiency. As a consequence one gets a SME of the form (A4) where the two photocurrents depend on  $\hat{\sigma}_x$  and  $\hat{\sigma}_y$  as

$$dy_t^{(1)} = \sqrt{\frac{\eta\kappa}{2}} \langle \hat{\sigma}_x \rangle_t \, dt + dW_t^{(1)} \,, \tag{29}$$

$$dy_t^{(2)} = \sqrt{\frac{\eta\kappa}{2}} \langle \hat{\sigma}_y \rangle_t \, dt + dW_t^{(2)} \,. \tag{30}$$

We will denote with  $\overline{\mathcal{E}}_{PD,\eta}$ ,  $\overline{\mathcal{E}}_{HoD,\eta}$  and  $\overline{\mathcal{E}}_{HeD,\eta}$  the daemonic ergotropies corresponding to continuous monitoring of the fluorescence field due to the atomic spontaneous emission via respectively PD, HoD and HeD with efficiency  $\eta$ .

According to Prop 2, we know that for unravellings of pure states with  $\eta = 1$ , one obtains  $\overline{\mathcal{E}}_{unr,\eta=1}(t) = E(\varrho_{unc}(t))$ , that is, the energy of the unconditional state can be fully extracted via conditional unitary operations for all the possible detection strategies.

We first consider the situation where the system is initially prepared in the ground state  $|0\rangle$ , while the environment is not fully accessible and it is thus monitored with non-unit efficiency  $\eta$ . We have numerically solved the corresponding SMEs [115, 116] and evaluated the corresponding daemonic ergotropy by averaging over a large number of trajectories. In this case we find that there is a hierarchy between the different unravellings: as one can observe in Fig. 2, for  $\eta = 0.4$  and  $\alpha = \kappa$ , homodyne detection with photocurrent proportional to  $\langle \hat{\sigma}_x \rangle_t$ (corresponding to  $\phi = 0$ ) and heterodyne detection lead to the largest values of daemonic ergotropy, while photodetection is the least performing strategy. The results presented in this plot are quite general: in all our numerical simulations we find that, as regards HoD,  $\overline{\mathcal{E}}_{\text{HoD},\eta=1}(t)$ 



FIG. 3. Steady-state daemonic ergotropy  $\overline{\mathcal{E}}_{unr,\eta}^{ss}$  for the different unravellings (HoD with photocurrents monitoring either  $\hat{\sigma}_x$  or  $\hat{\sigma}_y$ , HeD, PD) as a function of  $\alpha$  and for  $\eta = 0.1$  (dashed-dotted) and  $\eta = 0.7$  (dashed) (number of trajectories  $n = 5 \times 10^4$ ). Also in this case HeD and  $\hat{\sigma}_x$ -HoD yield the largest values. Black and gray solid lines correspond respectively to ergotropy and energy of the unconditional steady-state, that is to the lower and upper bound for the daemonic ergotropies, that would be obtained for respectively no-monitoring ( $\eta = 0$ ) or perfect monitoring ( $\eta = 1$ ).

is maximized (minimized) for  $\phi = 0$  ( $\phi = \pi/2$ ), that is for a  $\hat{\sigma}_x$ -dependent ( $\hat{\sigma}_y$ -dependent) photocurrent. Furthermore, in the whole range of parameters we have explored,  $\hat{\sigma}_x$ -HoD and HeD generally yield very similar values of daemonic ergotropy. The same behavior can be indeed observed if we focus on the steady-state properties as shown in Fig. 3, where steady-state daemonic ergotropies  $\overline{\mathcal{E}}_{unr,\eta}^{ss}$  for the different unravellings are plotted as a function of  $\alpha$  and for two different values of  $\eta$ . From the figure one can clearly observe the hierarchy existing between the different strategies, and how, by increasing the monitoring efficiency, the daemonic ergotropy can reach values, approaching the unconditional energy  $E_{unc}^{ss}$  in the limit of  $\eta = 1$ . Finally, in Fig. 4 we consider the other scenario where different unravellings lead to different daemonic ergotropies, that is when the environment is monitored with unit efficiency, but the initial state is mixed. In particular, we consider the maximally mixed state  $\rho_0 = \hat{1}/2$  as initial state and we plot  $\overline{\mathcal{E}}_{unr,\eta}(t)$  as a function of time. We observe that at steady state all the unravellings lead to the same value of daemonic ergotropy, equal to the corresponding unconditional energy: the monitoring will eventually purify all the trajectories and thus one falls back into the scenario described in Prop. 2. However, different unravellings lead to a different purification speed, an effect that has been widely discussed in the literature [117–122]. With respect to the scenarios described in these works, we here have



FIG. 4. Daemonic ergotropy  $\overline{\mathcal{E}}_{unr,\eta}(t)$  as a function of time for different unravellings (HoD with photocurrents monitoring either  $\hat{\sigma}_x$  or  $\hat{\sigma}_y$ , HeD, PD) with  $\alpha/\kappa = 0.4$ ,  $\eta = 1$  and by considering the maximally mixed state  $\rho_0 = \hat{1}/2$  as initial state (number of trajectories  $n = 5 \times 10^4$ ). Black and gray lines correspond respectively to ergotropy and energy of the unconditional state when initial states are the maximally mixed state (dashed) and the ground state (dotted).

a non-trivial system Hamiltonian  $\hat{H}_s$  ruling the dynamics, and a fixed jump operator  $\hat{c} = \sqrt{\kappa}\hat{\sigma}_{-}$ , representing the interaction between system and environment. Remarkably, at small times we find that the purification speed of HeD and, in the second instance, HoD (almost independently on the phase  $\phi$ ) allows to achieve values of daemonic ergotropy evidently larger than the maximum daemonic ergotropy obtainable by starting from a ground state, whose upper bound (the unconditional energy represented as a dotted-gray line in Fig. 4) can be interpreted as the amount of energy injected into the system by the driving laser; we can thus conclude that in monitoring-enhanced battery charging protocols, the effects of purification may be more efficient than pure energy injection at small times. On the other hand we also observe that PD still yields the lowest values of daemonic ergotropy in the transient leading to steady-state, and for small times the enhancement due to the monitoring respect to the unconditional ergotropy is almost negligible.

#### V. CONCLUSIONS

We have extended the concept of daemonic ergotropy to the open quantum system scenario, where some information leaking into the environment can be continuously monitored and exploited in order to enhance the work extraction protocol. Our findings reveal that the daemonic ergotropy not only surpasses the unconditional state ergotropy, but can even reach its energy in the ideal scenario of unit efficiency detectors. We have then discussed the simplest, but practically relevant, example of an OQB, that is a two-level atom classically driven by an external field acting as a charger.

Our main results (Props.1,2) require very few assumptions compared to the ones implied in the model of OQB considered. In particular, they hold even for charging protocols with time-dependent Hamiltonian [9, 10], allowing one to further assess the performances of these protocols in the case of continuous monitoring and to envisage more efficient quantum feedback protocols. Moreover, one can also consider trajectories describing continuously monitored open quantum systems in the presence of a non-Markovian environment; while the interpretation of unravellings of non-Markovian master equation in terms of monitoring is in general not guaranteed [123–127], our approach can be directly pursued in the non-Markovian setting by describing such quantum conditional dynamics via continuously measured collisional models [35, 100, 128].

A proof-of-principle experimental demonstration of our results can be readily pursued in a circuit-QED platform, where quantum trajectories corresponding to HeD of atom fluorescence have been recently observed [105, 106]. In general our results pave the way to further investigation on the relationship between measurement energy cost and work extraction in continuously-monitored quantum systems [129–137], and on the design of a new generation of monitoring-enhanced and noise-resilient quantum batteries.

### ACKNOWLEDGMENTS

The authors acknowledge helpful discussions with F. Albarelli, A. Serafini and A. Smirne. DM acknowledges financial support from MUR under the "PON Ricerca e Innovazione 2014-2020". MACR acknowledges financial support from the Academy of Finland via the Centre of Excellence program (Project No. 336810). MGG acknowledges support from UniMi via PSR-2 2021. The authors wish to acknowledge CSC – IT Center for Science, Finland, for computational resources.

 F. Binder, L. A. Correa, C. Gogolin, J. Anders, and G. Adesso, eds., *Thermodynamics in the Quantum* Regime, Fundamental Theories of Physics, Vol. 195

(Springer International Publishing, Cham, 2018).

- [2] A. E. Allahverdyan, R. Balian, and T. M. Nieuwenhuizen, Maximal work extraction from finite quantum systems, Europhysics Letters 67, 565 (2004).
- [3] R. Alicki and M. Fannes, Entanglement boost for extractable work from ensembles of quantum batteries, Physical Review E 87, 042123 (2013).
- [4] F. C. Binder, S. Vinjanampathy, K. Modi, and J. Goold, Quantacell: Powerful charging of quantum batteries, New Journal of Physics 17, 075015 (2015).
- [5] F. Campaioli, F. A. Pollock, F. C. Binder, L. Céleri, J. Goold, S. Vinjanampathy, and K. Modi, Enhancing the Charging Power of Quantum Batteries, Physical Review Letters 118, 150601 (2017).
- [6] F. Campaioli, F. A. Pollock, and S. Vinjanampathy, Quantum Batteries - Review Chapter (2018).
- [7] D. Ferraro, M. Campisi, G. M. Andolina, V. Pellegrini, and M. Polini, High-Power Collective Charging of a Solid-State Quantum Battery, Physical Review Letters 120, 117702 (2018).
- [8] G. M. Andolina, M. Keck, A. Mari, M. Campisi, V. Giovannetti, and M. Polini, Extractable Work, the Role of Correlations, and Asymptotic Freedom in Quantum Batteries, Physical Review Letters 122, 047702 (2019).
- [9] Y.-Y. Zhang, T.-R. Yang, L. Fu, and X. Wang, Powerful harmonic charging in a quantum battery, Physical Review E 99, 052106 (2019).
- [10] A. Crescente, M. Carrega, M. Sassetti, and D. Ferraro, Charging and energy fluctuations of a driven quantum battery, New Journal of Physics 22, 063057 (2020).
- [11] S. Julià-Farré, T. Salamon, A. Riera, M. N. Bera, and M. Lewenstein, Bounds on the capacity and power of quantum batteries, Physical Review Research 2, 023113 (2020).
- [12] D. Rossini, G. M. Andolina, D. Rosa, M. Carrega, and M. Polini, Quantum Advantage in the Charging Process of Sachdev-Ye-Kitaev Batteries, Physical Review Letters 125, 236402 (2020).
- [13] J.-Y. Gyhm, D. Šafránek, and D. Rosa, Quantum Charging Advantage Cannot Be Extensive without Global Operations, Physical Review Letters 128, 140501 (2022).
- [14] S. Seah, M. Perarnau-Llobet, G. Haack, N. Brunner, and S. Nimmrichter, Quantum Speed-Up in Collisional Battery Charging, Physical Review Letters 127, 100601 (2021).
- [15] R. Salvia, M. Perarnau-Llobet, G. Haack, N. Brunner, and S. Nimmrichter, Quantum advantage in charging cavity and spin batteries by repeated interactions, arxiv:2205.00026 [cond-mat, physics:quant-ph] (2022).
- [16] G. T. Landi, Battery Charging in Collision Models with Bayesian Risk Strategies, Entropy 23, 1627 (2021).
- [17] F. Mayo and A. J. Roncaglia, Collective effects and quantum coherence in dissipative charging of quantum batteries, Physical Review A 105, 062203 (2022).
- [18] F. Barra, Efficiency fluctuations in a quantum battery charged by a repeated interaction process, Entropy 24, 820 (2022).
- [19] R. R. Rodriguez, B. Ahmadi, G. Suarez, P. Mazurek, S. Barzanjeh, and P. Horodecki, Optimal quantum control of charging quantum batteries, arxiv:2207.00094 [quant-ph] (2022).
- [20] S.-f. Qi and J. Jing, Magnon-mediated quantum battery under systematic errors, Physical Review A 104, 032606

(2021).

- [21] A. Crescente, D. Ferraro, M. Carrega, and M. Sassetti, Enhancing coherent energy transfer between quantum devices via a mediator, Physical Review Research 4, 033216 (2022).
- [22] F. Mazzoncini, V. Cavina, G. M. Andolina, P. A. Erdman, and V. Giovannetti, Optimal Control Methods for Quantum Batteries, Physical Review A 107, 032218 (2023).
- [23] C. Rodríguez, D. Rosa, and J. Olle, Ai-discovery of a new charging protocol in a micromaser quantum battery, arxiv:2301.09408 [quant-ph] (2023).
- [24] G. M. Andolina, D. Farina, A. Mari, V. Pellegrini, V. Giovannetti, and M. Polini, Charger-mediated energy transfer in exactly-solvable models for quantum batteries, Physical Review B 98, 205423 (2018).
- [25] D. Rosa, D. Rossini, G. M. Andolina, M. Polini, and M. Carrega, Ultra-stable charging of fast-scrambling SYK quantum batteries, Journal of High Energy Physics 2020, 67 (2020).
- [26] V. Shaghaghi, V. Singh, G. Benenti, and D. Rosa, Micromasers as quantum batteries, Quantum Science and Technology 7, 04LT01 (2022).
- [27] C.-K. Hu, J. Qiu, P. J. P. Souza, J. Yuan, Y. Zhou, L. Zhang, J. Chu, X. Pan, L. Hu, J. Li, Y. Xu, Y. Zhong, S. Liu, F. Yan, D. Tan, R. Bachelard, C. J. Villas-Boas, A. C. Santos, and D. Yu, Optimal charging of a superconducting quantum battery, Quantum Science and Technology 7, 045018 (2022).
- [28] J. Q. Quach, K. E. McGhee, L. Ganzer, D. M. Rouse, B. W. Lovett, E. M. Gauger, J. Keeling, G. Cerullo, D. G. Lidzey, and T. Virgili, Superabsorption in an organic microcavity: Toward a quantum battery, Science Advances 8, eabk3160 (2022).
- [29] J. Joshi and T. S. Mahesh, Experimental investigation of a quantum battery using star-topology nmr spin systems, Phys. Rev. A 106, 042601 (2022).
- [30] I. M. de Buy Wenniger, S. E. Thomas, M. Maffei, S. C. Wein, M. Pont, N. Belabas, S. Prasad, A. Harouri, A. Lemaître, I. Sagnes, N. Somaschi, A. Auffèves, and P. Senellart, Experimental analysis of energy transfers between a quantum emitter and light fields, 2202.01109 [quant-ph] (2023).
- [31] D. Farina, G. M. Andolina, A. Mari, M. Polini, and V. Giovannetti, Charger-mediated energy transfer for quantum batteries: An open system approach, Physical Review B 99, 035421 (2019).
- [32] S. Zakavati, F. T. Tabesh, and S. Salimi, Bounds on charging power of open quantum batteries, Physical Review E 104, 054117 (2021).
- [33] M. Carrega, A. Crescente, D. Ferraro, and M. Sassetti, Dissipative dynamics of an open quantum battery, New Journal of Physics 22, 083085 (2020).
- [34] F. H. Kamin, F. T. Tabesh, S. Salimi, F. Kheirandish, and A. C. Santos, Non-Markovian effects on charging and self-discharging process of quantum batteries, New Journal of Physics 22, 083007 (2020).
- [35] D. Morrone, M. A. C. Rossi, A. Smirne, and M. G. Genoni, Charging a quantum battery in a non-Markovian environment: A collisional model approach, Quantum Science and Technology 8, 035007 (2023).
- [36] A. C. Santos, B. Çakmak, S. Campbell, and N. T. Zinner, Stable adiabatic quantum batteries, Physical Review E 100, 032107 (2019).

- [37] J. Q. Quach and W. J. Munro, Using Dark States to Charge and Stabilize Open Quantum Batteries, Physical Review Applied 14, 024092 (2020).
- [38] S. Gherardini, F. Campaioli, F. Caruso, and F. C. Binder, Stabilizing open quantum batteries by sequential measurements, Physical Review Research 2, 013095 (2020).
- [39] M. T. Mitchison, J. Goold, and J. Prior, Charging a quantum battery with linear feedback control, Quantum 5, 500 (2021).
- [40] Y. Yao and X. Q. Shao, Optimal charging of open spinchain quantum batteries via homodyne-based feedback control, Physical Review E 106, 014138 (2022).
- [41] K. Koshihara and K. Yuasa, Quantum ergotropy and quantum feedback control, Physical Review E 107, 064109 (2023).
- [42] H. M. Wiseman and G. J. Milburn, *Quantum Measure*ment and Control (Cambridge University Press, Cambridge, 2009).
- [43] K. Jacobs, Quantum Measurement Theory and its Applications (Cambridge University Press, Cambridge, 2014).
- [44] K. Maruyama, F. Nori, and V. Vedral, Colloquium: The physics of Maxwell's demon and information, Reviews of Modern Physics 81, 1 (2009).
- [45] S. Lloyd, Quantum-mechanical Maxwell's demon, Physical Review A 56, 3374 (1997).
- [46] J. Oppenheim, M. Horodecki, P. Horodecki, and R. Horodecki, Thermodynamical Approach to Quantifying Quantum Correlations, Physical Review Letters 89, 180402 (2002).
- [47] W. H. Zurek, Quantum discord and Maxwell's demons, Physical Review A 67, 012320 (2003).
- [48] K. Maruyama, F. Morikoshi, and V. Vedral, Thermodynamical detection of entanglement by Maxwell's demons, Physical Review A 71, 012108 (2005).
- [49] S. W. Kim, T. Sagawa, S. De Liberato, and M. Ueda, Quantum Szilard Engine, Physical Review Letters 106, 070401 (2011).
- [50] M. Perarnau-Llobet, K. V. Hovhannisyan, M. Huber, P. Skrzypczyk, N. Brunner, and A. Acín, Extractable Work from Correlations, Physical Review X 5, 041011 (2015).
- [51] E. G. Brown, N. Friis, and M. Huber, Passivity and practical work extraction using Gaussian operations, New Journal of Physics 18, 113028 (2016).
- [52] G. Francica, J. Goold, F. Plastina, and M. Paternostro, Daemonic ergotropy: Enhanced work extraction from quantum correlations, npj Quantum Information 3, 1 (2017).
- [53] M. A. Ciampini, L. Mancino, A. Orieux, C. Vigliar, P. Mataloni, M. Paternostro, and M. Barbieri, Experimental extractable work-based multipartite separability criteria, npj Quantum Information 3, 1 (2017).
- [54] M. Brunelli, M. G. Genoni, M. Barbieri, and M. Paternostro, Detecting Gaussian entanglement via extractable work, Physical Review A 96, 062311 (2017).
- [55] F. Bernards, M. Kleinmann, O. Gühne, and M. Paternostro, Daemonic Ergotropy: Generalised Measurements and Multipartite Settings, Entropy 21, 771 (2019).
- [56] D. Šafránek, D. Rosa, and F. Binder, Work extraction from unknown quantum sources, Physical Review Letters 130, 210401 (2023).

- [57] H. M. Wiseman and G. J. Milburn, Quantum theory of optical feedback via homodyne detection, Physical Review Letters 70, 548 (1993).
- [58] H. M. Wiseman, Quantum theory of continuous feedback, Physical Review A 49, 2133 (1994).
- [59] A. C. Doherty and K. Jacobs, Feedback control of quantum systems using continuous state estimation, Physical Review A 60, 2700 (1999).
- [60] L. K. Thomsen, S. Mancini, and H. M. Wiseman, Spin squeezing via quantum feedback, Physical Review A 65, 061801 (2002).
- [61] A. Serafini and S. Mancini, Determination of Maximal Gaussian Entanglement Achievable by Feedback-Controlled Dynamics, Physical Review Letters 104, 220501 (2010).
- [62] A. Szorkovszky, A. C. Doherty, G. I. Harris, and W. P. Bowen, Mechanical Squeezing via Parametric Amplification and Weak Measurement, Physical Review Letters 107, 213603 (2011).
- [63] M. G. Genoni, S. Mancini, and A. Serafini, Optimal feedback control of linear quantum systems in the presence of thermal noise, Physical Review A 87, 042333 (2013).
- [64] M. G. Genoni, J. Zhang, J. Millen, P. F. Barker, and A. Serafini, Quantum cooling and squeezing of a levitating nanosphere via time-continuous measurements, New Journal of Physics 17, 073019 (2015).
- [65] S. G. Hofer and K. Hammerer, Entanglement-enhanced time-continuous quantum control in optomechanics, Physical Review A 91, 033822 (2015).
- [66] L. Martin, F. Motzoi, H. Li, M. Sarovar, and K. B. Whaley, Deterministic generation of remote entanglement with active quantum feedback, Physical Review A 92, 062321 (2015).
- [67] M. Brunelli, D. Malz, and A. Nunnenkamp, Conditional Dynamics of Optomechanical Two-Tone Backaction-Evading Measurements, Physical Review Letters 123, 093602 (2019).
- [68] L. S. Martin, Quantum feedback for measurement and control, arxiv:2004.09766 [quant-ph] (2020).
- [69] Y. Jiang, X. Wang, L. Martin, and K. B. Whaley, Optimality of feedback control for qubit purification under inefficient measurement, Physical Review A 102, 022612 (2020).
- [70] S. Zhang, L. S. Martin, and K. B. Whaley, Locally optimal measurement-based quantum feedback with application to multiqubit entanglement generation, Physical Review A 102, 062418 (2020).
- [71] A. Di Giovanni, M. Brunelli, and M. G. Genoni, Unconditional mechanical squeezing via backaction-evading measurements and nonoptimal feedback control, Physical Review A 103, 022614 (2021).
- [72] A. Candeloro, C. Benedetti, M. G. Genoni, and M. G. A. Paris, Feedback-Assisted Quantum Search by Continuous-Time Quantum Walks, Advanced Quantum Technologies 6, 2200093 (2023).
- [73] F. W. Isaksen and U. L. Andersen, Mechanical cooling and squeezing using optimal control, Physical Review A 107, 023512 (2023).
- [74] H. Mabuchi, Dynamical identification of open quantum systems, Quantum and Semiclassical Optics: Journal of the European Optical Society Part B 8, 1103 (1996).
- [75] J. Gambetta and H. M. Wiseman, State and dynamical parameter estimation for open quantum systems,

Physical Review A 64, 042105 (2001).

- [76] JM. Geremia, J. K. Stockton, A. C. Doherty, and H. Mabuchi, Quantum Kalman Filtering and the Heisenberg Limit in Atomic Magnetometry, Physical Review Letters 91, 250801 (2003).
- [77] M. Guţă, B. Janssens, and J. Kahn, Optimal Estimation of Qubit States with Continuous Time Measurements, Communications in Mathematical Physics 277, 127 (2008).
- [78] M. Tsang, Optimal waveform estimation for classical and quantum systems via time-symmetric smoothing. ii. applications to atomic magnetometry and hardy's paradox, Phys. Rev. A 81, 013824 (2010).
- [79] M. Tsang, H. M. Wiseman, and C. M. Caves, Fundamental Quantum Limit to Waveform Estimation, Physical Review Letters 106, 090401 (2011).
- [80] M. Tsang, Quantum metrology with open dynamical systems, New Journal of Physics 15, 073005 (2013).
- [81] S. Gammelmark and K. Mølmer, Bayesian parameter inference from continuously monitored quantum systems, Physical Review A 87, 032115 (2013).
- [82] S. Gammelmark and K. Mølmer, Fisher Information and the Quantum Cram\'er-Rao Sensitivity Limit of Continuous Measurements, Physical Review Letters 112, 170401 (2014).
- [83] P. Six, Ph. Campagne-Ibarcq, L. Bretheau, B. Huard, and P. Rouchon, Parameter estimation from measurements along quantum trajectories, in 2015 54th IEEE Conference on Decision and Control (CDC) (2015) pp. 7742-7748.
- [84] C. Catana, L. Bouten, and M. Guţă, Fisher informations and local asymptotic normality for continuoustime quantum Markov processes, Journal of Physics A: Mathematical and Theoretical 48, 365301 (2015).
- [85] A. H. Kiilerich and K. Mølmer, Bayesian parameter estimation by continuous homodyne detection, Physical Review A 94, 032103 (2016).
- [86] J. F. Ralph, S. Maskell, and K. Jacobs, Multiparameter estimation along quantum trajectories with sequential Monte Carlo methods, Physical Review A 96, 052306 (2017).
- [87] M. G. Genoni, Cram\'er-Rao bound for time-continuous measurements in linear Gaussian quantum systems, Physical Review A 95, 012116 (2017).
- [88] F. Albarelli, M. A. C. Rossi, M. G. A. Paris, and M. G. Genoni, Ultimate limits for quantum magnetometry via time-continuous measurements, New Journal of Physics 19, 123011 (2017).
- [89] M. Rossi, L. Mancino, G. T. Landi, M. Paternostro, A. Schliesser, and A. Belenchia, Experimental Assessment of Entropy Production in a Continuously Measured Mechanical Resonator, Physical Review Letters 125, 080601 (2020).
- [90] T. Ilias, D. Yang, S. F. Huelga, and M. B. Plenio, Criticality-Enhanced Quantum Sensing via Continuous Measurement, PRX Quantum 3, 010354 (2022).
- [91] D. Yang, S. F. Huelga, and M. B. Plenio, Efficient information retrieval for sensing via continuous measurement, arxiv:2209.08777 [quant-ph] (2022).
- [92] C. Elouard and M. H. Mohammady, Work, Heat and Entropy Production Along Quantum Trajectories, in *Thermodynamics in the Quantum Regime: Fundamental Aspects and New Directions*, Fundamental Theories of Physics, edited by F. Binder, L. A. Correa,

C. Gogolin, J. Anders, and G. Adesso (Springer International Publishing, Cham, 2018) pp. 363–393.

- [93] G. Manzano and R. Zambrini, Quantum thermodynamics under continuous monitoring: A general framework, AVS Quantum Science 4, 025302 (2022).
- [94] J. P. Garrahan and I. Lesanovsky, Thermodynamics of Quantum Jump Trajectories, Physical Review Letters 104, 160601 (2010).
- [95] J. P. Garrahan, A. D. Armour, and I. Lesanovsky, Quantum trajectory phase transitions in the micromaser, Physical Review E 84, 021115 (2011).
- [96] J. P. Garrahan and M. Guţă, Catching and reversing quantum jumps and thermodynamics of quantum trajectories, Physical Review A 98, 052137 (2018).
- [97] D. Cilluffo, S. Lorenzo, G. M. Palma, and F. Ciccarello, Quantum jump statistics with a shifted jump operator in a chiral waveguide, Journal of Statistical Mechanics: Theory and Experiment **2019**, 104004 (2019).
- [98] A. Belenchia, L. Mancino, G. T. Landi, and M. Paternostro, Entropy production in continuously measured Gaussian quantum systems, npj Quantum Information 6, 1 (2020).
- [99] G. T. Landi and M. Paternostro, Irreversible entropy production: From classical to quantum, Reviews of Modern Physics 93, 035008 (2021).
- [100] G. T. Landi, M. Paternostro, and A. Belenchia, Informational Steady States and Conditional Entropy Production in Continuously Monitored Systems, PRX Quantum 3, 010303 (2022).
- [101] B. Bhandari and A. N. Jordan, Continuous measurement boosted adiabatic quantum thermal machines, Physical Review Research 4, 033103 (2022).
- [102] K. Yanik, B. Bhandari, S. K. Manikandan, and A. N. Jordan, Thermodynamics of quantum measurement and Maxwell's demon's arrow of time, Physical Review A 106, 042221 (2022).
- [103] B. Bhandari, R. Czupryniak, P. A. Erdman, and A. N. Jordan, Measurement-Based Quantum Thermal Machines with Feedback Control, Entropy 25, 204 (2023).
- [104] K. W. Murch, S. J. Weber, C. Macklin, and I. Siddiqi, Observing single quantum trajectories of a superconducting quantum bit, Nature 502, 211 (2013).
- [105] P. Campagne-Ibarcq, P. Six, L. Bretheau, A. Sarlette, M. Mirrahimi, P. Rouchon, and B. Huard, Observing Quantum State Diffusion by Heterodyne Detection of Fluorescence, Physical Review X 6, 011002 (2016).
- [106] Q. Ficheux, S. Jezouin, Z. Leghtas, and B. Huard, Dynamics of a qubit while simultaneously monitoring its relaxation and dephasing, Nature Communications 9, 1926 (2018).
- [107] Z. K. Minev, S. O. Mundhada, S. Shankar, P. Reinhold, R. Gutiérrez-Jáuregui, R. J. Schoelkopf, M. Mirrahimi, H. J. Carmichael, and M. H. Devoret, To catch and reverse a quantum jump mid-flight, Nature 570, 200 (2019).
- [108] W. Wieczorek, S. G. Hofer, J. Hoelscher-Obermaier, R. Riedinger, K. Hammerer, and M. Aspelmeyer, Optimal State Estimation for Cavity Optomechanical Systems, Physical Review Letters 114, 223601 (2015).
- [109] M. Rossi, D. Mason, J. Chen, and A. Schliesser, Observing and Verifying the Quantum Trajectory of a Mechanical Resonator, Physical Review Letters **123**, 163601 (2019).
- [110] R. A. Thomas, M. Parniak, C. Østfeldt, C. B. Møller,

C. Bærentsen, Y. Tsaturyan, A. Schliesser, J. Appel, E. Zeuthen, and E. S. Polzik, Entanglement between distant macroscopic mechanical and spin systems, Nature Physics **17**, 228 (2021).

- [111] M. Rossi, D. Mason, J. Chen, Y. Tsaturyan, and A. Schliesser, Measurement-based quantum control of mechanical motion, Nature 563, 53 (2018).
- [112] L. Magrini, P. Rosenzweig, C. Bach, A. Deutschmann-Olek, S. G. Hofer, S. Hong, N. Kiesel, A. Kugi, and M. Aspelmeyer, Real-time optimal quantum control of mechanical motion at room temperature, Nature 595, 373 (2021).
- [113] F. Tebbenjohanns, M. L. Mattana, M. Rossi, M. Frimmer, and L. Novotny, Quantum control of a nanoparticle optically levitated in cryogenic free space, Nature 595, 378 (2021).
- [114] We are here considering a single jump operator  $\hat{c}$  describing the interaction between the system and a zero-temperature environment, but the formalism can be straightforwardly generalized to multiple jump operators and to generic thermal environments.
- [115] J. R. Johansson, P. D. Nation, and F. Nori, QuTiP: An open-source Python framework for the dynamics of open quantum systems, Computer Physics Communications 183, 1760 (2012).
- [116] J. R. Johansson, P. D. Nation, and F. Nori, QuTiP 2: A Python framework for the dynamics of open quantum systems, Computer Physics Communications 184, 1234 (2013).
- [117] K. Jacobs, How to project qubits faster using quantum feedback, Physical Review A 67, 030301 (2003).
- [118] J. Combes and K. Jacobs, Rapid State Reduction of Quantum Systems Using Feedback Control, Physical Review Letters 96, 010504 (2006).
- [119] H. M. Wiseman and J. F. Ralph, Reconsidering rapid qubit purification by feedback, New Journal of Physics 8, 90 (2006).
- [120] A. N. Jordan and A. N. Korotkov, Qubit feedback and control with kicked quantum nondemolition measurements: A quantum Bayesian analysis, Physical Review B 74, 085307 (2006).
- [121] H. M. Wiseman and L. Bouten, Optimality of Feedback Control Strategies for Qubit Purification, Quantum Information Processing 7, 71 (2008).
- [122] R. Ruskov, J. Combes, K. Mølmer, and H. M. Wiseman, Qubit purification speed-up for three complementary continuous measurements, Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences **370**, 5291 (2012).
- [123] L. Diósi, Non-markovian continuous quantum measure-

ment of retarded observables, Physical Review Letters **100**, 080401 (2008).

- [124] H. M. Wiseman and J. M. Gambetta, Pure-state quantum trajectories for general non-markovian systems do not exist, Physical Review Letters 101, 140401 (2008).
- [125] J. Piilo, S. Maniscalco, K. Härkönen, and K.-A. Suominen, Non-markovian quantum jumps, Physical Review Letters 100, 180402 (2008).
- [126] A. Barchielli, C. Pellegrini, and F. Petruccione, Stochastic schrödinger equations with coloured noise, Europhysics Letters **91**, 24001 (2010).
- [127] A. Smirne, M. Caiaffa, and J. Piilo, Rate operator unraveling for open quantum system dynamics, Physical Review Letters 124, 190402 (2020).
- [128] F. Ciccarello, S. Lorenzo, V. Giovannetti, and G. M. Palma, Quantum collision models: Open system dynamics from repeated interactions, Physics Reports 954, 1 (2022).
- [129] R. Alicki, M. Horodecki, P. Horodecki, and R. Horodecki, Thermodynamics of Quantum Information Systems — Hamiltonian Description, Open Systems & Information Dynamics 11, 205 (2004).
- [130] T. Sagawa and M. Ueda, Second Law of Thermodynamics with Discrete Quantum Feedback Control, Physical Review Letters 100, 080403 (2008).
- [131] T. Sagawa and M. Ueda, Minimal Energy Cost for Thermodynamic Information Processing: Measurement and Information Erasure, Physical Review Letters 102, 250602 (2009).
- [132] K. Jacobs, Second law of thermodynamics and quantum feedback control: Maxwell's demon with weak measurements, Physical Review A 80, 012322 (2009).
- [133] J. A. Vaccaro and S. M. Barnett, Information erasure without an energy cost, Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences 467, 1770 (2011).
- [134] K. Jacobs, Quantum measurement and the first law of thermodynamics: The energy cost of measurement is the work value of the acquired information, Physical Review E 86, 040106 (2012).
- [135] S. Deffner, J. P. Paz, and W. H. Zurek, Quantum work and the thermodynamic cost of quantum measurements, Physical Review E 94, 010103 (2016).
- [136] G. Manzano, F. Plastina, and R. Zambrini, Optimal Work Extraction and Thermodynamics of Quantum Measurements and Correlations, Physical Review Letters 121, 120602 (2018).
- [137] K. Abdelkhalek, Y. Nakata, and D. Reeb, Fundamental energy cost for quantum measurement, arxiv:1609.06981 [cond-mat, physics:quant-ph] (2018).

### Appendix A: Continuously-monitored quantum systems

In this appendix we briefly introduce the formalism behind continuously monitored quantum systems, presenting the stochastic master equations (SMEs) corresponding to continuous photo-detection, homodyne-detection and heterodyne-detection.

We assume that our quantum system is interacting with a Markovian environment such that its unconditional evolution is described by a master equation in Lindblad form

$$\frac{d\varrho_{\rm unc}(t)}{dt} = -i[\hat{H}_s, \varrho_{\rm unc}(t)] + \mathcal{D}[\hat{c}]\varrho_{\rm unc}(t), \qquad (A1)$$

where  $\hat{H}_s$  denotes the Hamiltonian ruling the evolution of the system and  $\mathcal{D}[\hat{c}]\varrho = \hat{c}\varrho\hat{c} - (\hat{c}^{\dagger}\hat{c}\varrho + \varrho\hat{c}^{\dagger}\hat{c})/2$  is the Lindbladian superoperator. We remark that we will here consider a single jump operator  $\hat{c}$  describing the interaction between the system and a zero-temperature environment, but the formalism can be straightforwardly generalized to multiple jump operators and to generic thermal environments.

We will now assess the scenario where the environment is continuously monitored. Different measurement strategies lead to different possible evolutions of the corresponding conditional states  $\rho_{c}(t)$ , i.e. to different unravellings of the unconditional master equation (A1). We will here consider three different measurements: photo-detection (PD), homodyne-detection (HoD) and heterodyne-detection (HeD).

In the case of PD the evolution is described by a stochastic master equation

$$d\varrho_{\mathsf{c}}(t) = -i[\hat{H}_{s}, \varrho_{\mathsf{c}}(t)] dt + (1 - \eta) \mathcal{D}[\hat{c}]\varrho_{\mathsf{c}}(t) dt - \frac{\eta}{2} (\hat{c}^{\dagger} \hat{c} \varrho_{\mathsf{c}}(t) + \varrho_{\mathsf{c}}(t) \hat{c}^{\dagger} \hat{c}) + \eta \langle \hat{c}^{\dagger} \hat{c} \rangle_{t} \varrho_{c}(t) dt + \left( \frac{\hat{c} \varrho_{\mathsf{c}}(t) \hat{c}^{\dagger}}{\langle \hat{c}^{\dagger} \hat{c} \rangle_{t}} - \varrho_{\mathsf{c}}(t) \right) dN_{t},$$
(A2)

where  $\langle \hat{A} \rangle_t = \text{Tr} \Big[ \rho_{\mathsf{c}}(t) \hat{A} \Big]$ ,  $\eta$  is the efficiency of the detector and  $dN_t$  is a Poisson increment taking value 0 (no-click event) or 1 (detector click event), and having average value  $\mathbb{E}[dN_t] = \eta \langle \hat{c}^{\dagger} \hat{c} \rangle_t dt$ .

For HoD one has the following diffusive SME

$$d\varrho_{\mathsf{c}}(t) = -i[H_s, \varrho_{\mathsf{c}}(t)] dt + \mathcal{D}[\hat{c}]\varrho_{\mathsf{c}}(t) dt + \sqrt{\eta} \mathcal{H}[\hat{c}e^{i\phi}]\varrho_{\mathsf{c}} dW_t$$
(A3)

where  $\mathcal{H}[\hat{c}]\varrho = \hat{c}\varrho + \varrho\hat{c}^{\dagger} - \langle \hat{c} + \hat{c}^{\dagger} \rangle_t \varrho$ ,  $\phi$  is the phase of the quadrature monitored via the homodyne, and  $dW_t$  is a Wiener increment related to the measured photocurrent  $dy_t = \sqrt{\eta} \langle \hat{c}e^{i\phi} + \hat{c}^{\dagger}e^{-i\phi} \rangle_t dt + dW_t$ . The unravelling for HeD is a generalization of the HoD case, corresponding to a double-homodyne scheme leading to the SME

$$d\varrho_{\mathsf{c}}(t) = -i[\hat{H}_s, \varrho_{\mathsf{c}}(t)] dt + \mathcal{D}[\hat{c}]\varrho_{\mathsf{c}}(t) dt + \sqrt{\eta/2} \mathcal{H}[\hat{c}]\varrho_{\mathsf{c}} dW_t^{(1)} + \sqrt{\eta/2} \mathcal{H}[i\hat{c}]\varrho_{\mathsf{c}} dW_t^{(2)}$$
(A4)

where  $dW_t^{(1)}$  and  $dW_t^{(2)}$  are uncorrelated Wiener increments corresponding to the two photocurrents  $dy_t^{(1)} = \sqrt{\eta/2} \langle \hat{c} + \hat{c}^{\dagger} \rangle_t dt + dW_t^{(1)}$  and  $dy_t^{(2)} = \sqrt{\eta/2} \langle i\hat{c} - i\hat{c}^{\dagger} \rangle_t dt + dW_t^{(2)}$ .