

ABSTRACT BOOK



ICAR2020+2

Bologna, Italy

19th International Congress
on Animal Reproduction
BOLOGNA (ITALY), 26th-30th JUNE 2022

Monday, 27th June 2022

- Avian species reproduction M01 - M05
- Bovine reproduction M06 - M107
- Canine and feline reproduction M109 - M127
- Control of estrous cycle M128 - M133
- Cryobiology of gametes and embryos M134 - M156
- Reproductive pathologies M157 - M165

Tuesday, 28th June 2022

- Buffalo reproduction T01 - T09
- Male reproductive physiology T10 - T18
- Metabolism and reproduction T19 - T28
- Ovary and oocyte T29 - T64
- Pregnancy, placental function and parturition T66 - T76 and T168
- Reproduction exotic anim and wild species T77 - T99
- Reproduction in fish T100 - T101
- Reproduction in sheep and goats T102 - T139
- Testis T141
- Uterus T143 - T146
- Oviduct T147 - T153
- Embryogenesis in vitro T154 - T167

Wednesday, 29th June 2022

- Camelid reproduction W01 - W10
- Embryo development and differentiation W11 - W28
- Female reproductive physiology W29 - W34 and W171
- Horse reproduction W35 - W81
- Imaging methods in reproduction W82 - W90
- Neuroendocrine control of reproduction W91 - W93
- Nutrition and reproduction W94 -W102 and W170
- Other W106 - W111
- Pig reproduction W112 - W135
- Reproductive system diseases W136 - W141
- Spermatology and sperm quality W142 - W169

ABSTRACTS WORKSHOPS

SPONSORS & PARTNERS

W87

EFFECT OF PATERNAL EXPOSURE TO HIGH TEMPERATURE-HUMIDITY INDEX ON SUBSEQUENT IN VITRO EMBRYO MORPHOKINETICS IN BOS TAURUS

*M. Melean*¹, *E. Malama*¹, *H. Bollwein*¹, *C. Herrera*¹

¹Clinic of Reproductive Medicine, Vetsuisse Faculty, University of Zurich

BACKGROUND-AIM

Bulls exposed to high temperature-humidity index (THI) experience adverse effects on sperm quality and in vitro embryo production (IVEP) outcomes. However, detailed information about the influence of paternal high-THI exposure on in vitro embryo morphokinetics is missing. The aim of this study was to evaluate the effect of bull exposure to high THI on the subsequent in vitro embryo morphokinetics by means of time-lapse image monitoring (TLM).

METHODS

Twelve ejaculates from six mature Simmental bulls that were exposed to high (n=6) and low (n=6) THI during the epididymal maturation phase were used for IVEP. Briefly, 96 in vitro matured cumulus-oocyte complexes retrieved from slaughterhouse ovaries were inseminated with high (n=48) and low (n=48) THI sperm samples during six IVEP sessions. A total of 73 fertilized embryos were cultured in vitro (IVC) in a micro well group culture dish (Vitrolife, Denmark) and monitored with a TLM system (PrimoVision, Vitrolife, Denmark). Images were recorded every 10 minutes until day 9 of IVC. Insemination time was considered as time 0 and used to calculate the developmental events. The time of first (T1) and second (T2) cleavage, last cleavage before entering the lag-phase (T9), cleavage resumption after lag phase (RCI), length of lag-phase (LP), start time of blastocyst expansion (tSB) and time of hatching (tHB) were recorded.

RESULTS

Both groups of embryos were similar regarding T1, T2, T9, RCI, LP and tHB (P>0.05). However, the in vitro derived embryos from bulls exposed to high-THI presented a faster blastocyst expansion (P<0.05) when compared to low-THI (159 ± 11 and 169 ± 21, respectively).

CONCLUSIONS

Our results described for the first time the effect of paternal high-THI exposure on the morphokinetic parameters of in vitro produced bovine embryos and suggest a relationship between paternal high-THI exposure and the time of blastocyst expansion.

W88

THERMAL COMFORT AND THERMOGRAPHIC SCROTAL GRADIENTS OF BULLS RAISED ON SHADED TROPICAL PASTURES: PRELIMINARY RESULTS

*A. Rossetto Garcia*¹, *N. Romanello*², *A. Nascimento Barreto*³, *M.A. Chagas Jacintho*¹, *A.C. Campos Bernardi*¹, *J.R. Macedo Pezzopane*¹, *F. Luzi*⁶, *V. Redaelli*⁶, *L. Nanni Costa*⁵, *M. Zappaterra*⁵, *F. Tonato*¹, *C. Righetti Marcondes*¹, *A. Faria Pedroso*¹, *F. Zandonadi Brandão*⁴, *W. Barioni Junior*¹

¹Embrapa - Brazilian Agricultural Research Corporation, São Carlos-SP, Brazil

²FMVZ/USP, Universidade de São Paulo, Pirassununga-SP, Brazil

³PPGCAN, Universidade Federal do Pará, Castanhal-PA, Brazil

⁴UFF, Universidade Federal Fluminense, Niterói-RJ, Brazil

⁵UNIBO, Università di Bologna, Bologna-BO, Italy

⁶UNIMI, Università degli Studi di Milano, Milano-MI, Italy

BACKGROUND-AIM

This study aimed to evaluate the thermal comfort and the scrotal temperature gradients of bulls of two breeds raised on natural shaded tropical pastures.

METHODS

Nelore (NEL, n=16; *Bos indicus*; 30 months; 478kg) and Canchim bulls (CAN, n=16; 5/8 *Bos taurus* x 3/8 *Bos indicus*; 32 months; 488kg) were raised on pastures with a 20% shaded area (*Eucalyptus urograndis*) in São Carlos, Brazil (21°57'S, 47°50'W). Monthly, respiratory rate (RR, breaths/min), rectal temperature (RT, °C), and infrared thermographic scrotal surface patterns were evaluated (Testo 890-2), during the day's greatest thermal challenge period (11:00 am to 2:30 pm). Thermal gradients (°C) were determined by the difference between the RT and the temperatures of the scrotum (G1), spermatic cord (G2), proximal testicular pole (G3), distal testicular pole (G4), tail of epididymis (G5), and between the temperatures of both testicular poles (G6). The data comprised the period from April to June (Mean air temperature: 20.2°C; relative humidity: 69.2%; THI: 66.8). Means were compared by ANOVA and Tukey's post-hoc test.

RESULTS

The RR (34.1±1.1 vs 34.5±1.6) and RT (39.37±0.08 vs 39.43±0.07) of NEL and CAN did not differ and were within the physiological range, indicating no thermal stress. NEL and CAN bulls did not differ regarding G1 (5.75±0.14 vs 5.70±0.18), G3 (5.07±0.15 vs 4.70±0.16), G4 (6.64±0.12 vs 6.96±0.19) and G5 (8.14±0.18 vs 8.72±0.26). However, NEL bulls presented higher G2 (3.78±0.18 vs 3.18±0.17, P<0.05) probably due to the longer and narrower testes and a greater number of scrotal sweat glands, which benefit both sensible heat and latent heat transfers. CAN bulls presented higher G6 (2.26 ± 0.14 vs 1.57 ± 0.12, P <0.01) probably because of the volume of scrotal sweat glands, that increases from the proximal to the distal testicular poles both in *Bos taurus* and their crossbreeds.

CONCLUSIONS

Shaded pastures promoted thermal comfort to Nelore and Canchim bulls. Despite the breed, thermal gradients particularities of the spermatic cord and the testicular poles indicated that these anatomical regions play a critical role in scrotal thermoregulation. The use of infrared thermography may be an interesting complementary tool to the breeding soundness evaluation of zebu and composite bulls. (FAPESP Process 2019/04528-6).