

MEASURE RIGIDITY OF RICCI CURVATURE LOWER BOUNDS

FABIO CAVALLETTI AND ANDREA MONDINO

ABSTRACT. The measure contraction property, MCP for short, is a weak Ricci curvature lower bound conditions for metric measure spaces. The goal of this paper is to understand which structural properties such assumption (or even weaker modifications) implies on the measure, on its support and on the geodesics of the space.

We start our investigation from the euclidean case by proving that if a positive Radon measure m over \mathbb{R}^d is such that $(\mathbb{R}^d, |\cdot|, m)$ verifies a weaker variant of MCP, then its support $\text{spt}(m)$ must be convex and m has to be absolutely continuous with respect to the relevant Hausdorff measure of $\text{spt}(m)$. This result is then used as a starting point to investigate the rigidity of MCP in the metric framework.

We introduce the new notion of *reference measure* for a metric space and prove that if (X, d, m) is essentially non-branching and verifies MCP, and μ is an essentially non-branching MCP reference measure for $(\text{spt}(m), d)$, then m is absolutely continuous with respect to μ , on the set of points where an inversion plan exists. As a consequence, an essentially non-branching MCP reference measure enjoys a weak type of uniqueness, up to densities. We also prove a stability property for reference measures under measured Gromov-Hausdorff convergence, provided an additional uniform bound holds.

In the final part we present concrete examples of metric spaces with reference measures, both in smooth and non-smooth setting. The main example will be the Hausdorff measure over an Alexandrov space. Then we prove that the following are reference measures over smooth spaces: the volume measure of a Riemannian manifold, the Hausdorff measure of an Alexandrov space with bounded curvature, and the Haar measure of the subRiemannian Heisenberg group.

1. INTRODUCTION

The notion of curvature for a smooth space, i.e. a Riemannian n -dimensional manifold (M, g) , is one of most basic geometrical concepts and goes back to the work of Gauss and Riemann; the idea being to consider suitable combinations of second derivatives of the metric g . If the space M is non-smooth this approach has few chances to be carried out, so one has to understand what are the fundamental geometric consequences of the curvature and then set these as defining properties. This was the approach by Alexandrov who, in last century, defined what it means for a metric space (X, d) to satisfy an upper or lower bound on the sectional curvature. Such non-smooth spaces are now called Alexandrov spaces and there is a huge literature about their properties (we do not even attempt to give a selection of papers; we just recall that Alexandrov, Gromov and Perelman among others gave major contributions to this theory and we refer to the textbook [8] for more references).

While, as just recalled, the notion of lower bound on the sectional curvature makes perfect sense for a metric space (X, d) , i.e. a set endowed with a distance function, for defining what it means for a non-smooth space to satisfy a lower Ricci curvature bound one also has to fix a measure m on (X, d) , thus getting a so called metric measure space (X, d, m) , m.m.s. for short. In the last ten years there has been a flourishing of literature about different notions and properties of lower Ricci curvature bounds for m.m.s.: on one hand Bakry-Émery-Ledoux [6, 7] developed the so called Gamma-calculus based on the fact that, roughly speaking, the Bochner inequality characterizes

lower Ricci bounds; on the other hand there is a parallel theory via optimal transport based on the idea that one can detect Ricci curvature by examining how the volume changes when optimally transported along geodesics. A third way to study the non-smooth spaces with lower Ricci bounds, having its origins in the work of Gromov and mainly due to Cheeger-Colding [13, 14, 15] (see also the more recent developments by Colding-Naber [16]), is to concentrate the attention on those m.m.s. arising as limits (in the measured Gromov-Hausdorff sense) of smooth Riemannian manifolds satisfying Ricci curvature lower bounds; such a point of view is very powerful if one is interested in the structure of these limit spaces.

In this paper we will focus on the second approach via optimal transportation, but let us mention that there is a precise correspondence between the first two (see [2] for the infinite dimensional case and [18, 4] for the finite dimensional one) and there are some recent results on the local structure of such spaces even in the abstract framework (see [22, 23, 30]).

The weakest Ricci curvature condition for a m.m.s. is the so called measure contraction property, MCP for short, which keeps track of the distortion of the volume of a set when it is transported to a Dirac delta. The quantitative formalization of having Ricci bounded from below by $K \in \mathbb{R}$ and dimension bounded from above by $N > 1$ via this approach, the so called $\text{MCP}(K, N)$ condition, is due to Ohta [34] and has its roots in earlier works by Grigor'yan [24], Sturm [36] and Kuwae-Shioya [27]. Such condition is compatible with the smooth counterpart (i.e. a smooth Riemannian manifold of dimension N has Ricci curvature bounded below by K if and only if it satisfies $\text{MCP}(K, N)$) and is stable under pointed measured Gromov-Hausdorff convergence, $pmGH$ for short, so that the limit spaces of Cheeger-Colding are included in this theory. Also finite dimensional Alexandrov spaces with lower curvature bounds endowed with the corresponding Hausdorff measure satisfy the measure contraction property [34] (see also [28] for a subsequent independent proof) as well as the Heisenberg group endowed with the Haar measure and the Carnot-Carathéodory distance [25].

Stronger notions of “Ricci bounded from below by $K \in \mathbb{R}$ and dimension bounded above by $N \in (1, \infty]$ ” for m.m.s. are the so called $\text{CD}(K, N)$ -conditions introduced independently by Lott-Villani [29] and by Sturm [37, 38], and the even stronger notions of $\text{RCD}(K, \infty)$ and $\text{RCD}^*(K, N)$ -spaces (the first were introduced in [3] and further investigated in [1, 2], for the second ones see [4, 18, 22, 23, 30, 39]). All these classes include the smooth Riemannian manifolds satisfying the corresponding curvature-dimension bounds and their $pmGH$ -limits, as well as finite dimensional Alexandrov spaces with curvature bounded from below. On the other hand let us recall that the Heisenberg group does not satisfy any $\text{CD}(K, N)$ condition but it does verify MCP, see [25]. So MCP is a strictly weaker notion of curvature than CD.

The goal of the present paper is to investigate the structural properties (in particular on the measure, on its support, and on the geodesics) forced by MCP.

In order to introduce the problem let us start by analyzing the behavior of the euclidean space \mathbb{R}^d , endowed with the d -dimensional Lebesgue measure \mathcal{L}^d and the Euclidean distance $|\cdot|$. It is almost trivial to check that the triple $(\mathbb{R}^d, |\cdot|, \mathcal{L}^d)$ verifies the measure contraction property with curvature $K = 0$ and dimension $N = d$, i.e. $\text{MCP}(0, d)$. Moreover it is well known that, at the price of changing the lower bound on the curvature and the upper bound on the dimension, in the triple $(\mathbb{R}^d, |\cdot|, \mathcal{L}^d)$ it is possible to replace the Lebesgue measure with some equivalent measure and still obtain MCP, provided the density verifies some concavity estimates.

A first goal of this work is to investigate, and give affirmative answer to, the reverse question:

- Let \mathfrak{m} be a positive Radon measure on \mathbb{R}^d such that the triple $(\mathbb{R}^d, |\cdot|, \mathfrak{m})$ verifies MCP (or even a weaker condition). Can we deduce that \mathfrak{m} is absolutely continuous with respect to \mathcal{L}^k , for some $k \leq d$?
- Moreover, does the support of \mathfrak{m} have some nice geometric properties?

The next theorem contains an affirmative answer to these natural questions; before stating it let us remark that the *non-degeneracy condition* for a positive Radon measure \mathfrak{m} is given in Definition 3.1 and it is ensured by the measure contraction property (it is indeed much weaker since no quantitative or uniform lower bound on the transported measure is assumed).

Theorem 1.1. *Let \mathfrak{m} be a positive Radon measure over \mathbb{R}^d and denote with Ω its support. If the metric measure space $(\Omega, |\cdot|, \mathfrak{m})$ verifies the non-degeneracy condition then there exists a natural number $k \leq d$ so that:*

- Ω is convex and contained in a k -dimensional affine subspace of \mathbb{R}^d , say V^k ;
- the measure \mathfrak{m} is absolutely continuous with respect to $\mathcal{L}^k \llcorner V^k$.

The same question can be reformulated in the metric framework only once the choice of a favorite measure is done. More precisely, given a m.m.s. (X, \mathfrak{d}, μ) satisfying MCP and maybe some other structural assumption, one can ask:

- Is the support of μ convex?
- If \mathfrak{m} is a positive Radon measure on X also satisfying MCP, can we deduce that \mathfrak{m} is absolutely continuous with respect to μ ?

This kind of properties were proved by Cheeger-Colding [15] in the framework of *pmGH*-limits of Riemannian manifolds satisfying lower Ricci curvature bounds: more precisely they showed that in the limit space the “favorite measure” is the Hausdorff measure of the relevant dimension, and any other possible limit measure has to be absolutely continuous with respect to it. Therefore this paper can be seen as an intrinsic-non smooth analogue of the Cheeger-Colding result.

Regarding the first question, in Proposition 4.5, we will prove the affirmative answer: the support of any Radon measure satisfying MCP has to be weakly convex (more precisely the support of any measure satisfying the strong qualitative MCP condition, defined in (4.3), must be weakly convex; i.e. every couple of points of $\text{spt}(\mathfrak{m})$ is joined by a geodesic entirely contained in $\text{spt}(\mathfrak{m})$).

In order to solve the second problem, we will give a precise meaning of what is for us a “favorite measure”: this is what we call *reference measure* (see Definition 5.1) for a complete and separable metric space (X, \mathfrak{d}) . The crucial property being to behave nicely under geodesic inversion with respect to almost every point of the space.

Definition 1.2 (Reference measure). A positive Radon measure $\mu \in \mathcal{M}^+(X)$ is a *reference measure* for (X, \mathfrak{d}) provided it is non-zero, and for μ -a.e. $z \in X$ there exists $\pi^z \in \mathcal{M}^+(X \times X)$ so that

$$(P_1)_\# \pi^z = \mu, \quad \pi^z(X \times X \setminus H(z)) = 0, \quad (P_2)_\# \pi^z \ll \mu,$$

where $P_i : X \times X \rightarrow X$ is the projection on the i -th component, for $i = 1, 2$ and

$$H(z) := \{(x, y) \in X \times X : \mathfrak{d}(x, y) = \mathfrak{d}(x, z) + \mathfrak{d}(z, y)\}.$$

The measure π^z will be called *inversion plan* and those points where an inversion plan exists will be called *inversion points* and denoted by $\text{lp}(\mu)$.

Let us stress that this concept is completely new, to best of our knowledge. In Remark 5.2 we will also point out that existence of an inversion plan in a point z is closely related to the regularity of the ambient space (X, \mathfrak{d}) at z . In particular if a conical singularity happens in z then no inversion plan at z exists.

Another ingredient in the next theorem will be the essentially non-branching condition (re-called in Section 2.1): it is an important structural assumption on a m.m.s. and it is fulfilled by a large class of geometrically relevant examples, for instance Riemannian manifolds, Alexandrov spaces with lower curvature bounds, *pmGH*-limits of Riemannian manifolds with lower Ricci curvature bounds, metric measure spaces verifying $\text{RCD}(K, \infty)$ or $\text{RCD}^*(K, N)$, the Heisenberg Group endowed with the Carnot-Carathéodory metric, etc.. Moreover, since in the proof of the next result we do not need the quantitative controls assumed in $\text{MCP}(K, N)$, we just assume a weaker qualitative MCP condition (for the precise notion see Definition 4.1).

Theorem 1.3. *Let (X, d, m) be an essentially non-branching m.m.s. that verifies the qualitative MCP condition (4.1). Assume the existence of a reference measure μ for (Ω, d) , where $\Omega = \text{spt}(m)$, so that (Ω, d, μ) verifies the qualitative MCP condition (4.1) and it is essentially non-branching. If $m(\Omega \setminus \text{lp}(\mu)) = 0$, then*

$$m \ll \mu.$$

In particular, if also m is a reference measure and $\text{spt}(m) = \text{spt}(\mu)$ with $\mu(\Omega \setminus \text{lp}(m)) = 0$, then μ and m are equivalent measures, i.e.

$$\mu \ll m \quad \text{and} \quad m \ll \mu.$$

In other words, once the support is fixed, a reference measure satisfying MCP is uniquely determined up to densities.

Once observed that metric measure spaces verifying RCD are all essentially non-branching, Theorem 1.3 can be used straightforwardly to obtain the next uniqueness result.

Theorem 1.4. *Let (X, d, μ) be a m.m.s. that verifies $\text{RCD}^*(K, N)$, with $N < \infty$ and $X = \text{spt}(\mu)$. Assume that μ is a reference measure for (X, d) . If (X, d, m) is a m.m.s. verifying $\text{RCD}^*(K', N')$, possibly for different K' and N' , with $X = \text{spt}(m)$ and $m(X \setminus \text{lp}(\mu)) = 0$, then*

$$m \ll \mu.$$

In particular, if also m is a reference measure with $\mu(X \setminus \text{lp}(m)) = 0$, then μ and m are equivalent measures, i.e. $\mu \ll m$ and $m \ll \mu$.

The second part of Theorem 1.4 therefore says that once the support X and a metric d are fixed, there is a unique reference measure, up to densities, so that the triple (X, d, μ) verifies RCD^* .

Both Theorem 1.3 and Theorem 1.4 can be restated dropping the assumption of $m(\Omega \setminus \text{lp}(\mu)) = 0$ (and the analogous one in the uniqueness part) at the price of assuming regularity properties on the geometry of $\text{lp}(\mu)$ (and on $\text{lp}(m)$): one way (see Theorem 5.5) is to assume that $\text{lp}(\mu)$ is μ -connected. For the precise meaning of this assumption see Definition 5.4, here we only mention that it is inspired by, and resemble, the convexity property of regular points in Alexandrov geometry. Another way is to assume that $\text{lp}(\mu)$ forms an open set in X , see Theorem 5.6.

Section 6 is devoted to the investigation of general properties of reference measures: we will establish the local-to-global property of reference measures under the non-branching assumption, we will show that the multiplication by a density does not affect the property of being a reference measure and finally we will address the problem of stability of reference measures with respect to measured Gromov-Hausdorff convergence: it will be proved that if the reference measures are probabilities and if a uniform bound on the second marginal of the inversion plan holds, the notion of reference measure is stable. See Theorem 6.6 for the precise statement and Theorem 6.9 for its application to uniqueness of limit reference measures. This last result should be compared with the aforementioned analogous statement for Ricci limits spaces obtained by Cheeger-Colding [15].

The final part of the paper, that is formed by Section 7 and Section 8, is devoted to discuss examples and applications of the introduced techniques.

The main result will be the existence of an inversion plan at almost every regular point of an Alexandrov space with curvature bounded from below, where the almost everywhere and the inversion plan are referred to the Hausdorff measure of the right dimension, see Theorem 7.3. This shows that the Hausdorff measure is a reference measure and permits to apply all the general theorems stated so far to Alexandrov spaces.

In the last section we then present smooth examples. The term smooth here is used to emphasize that in those spaces every point admits an inversion plan. The smooth examples will be: the standard volume measure on a smooth Riemannian manifold, the Hausdorff measure on an Alexandrov space with curvature bounded from above and below (see Corollary 8.3) and the Heisenberg group endowed with the Carnot-Carathéodory distance and the Haar measure (see Corollary 8.9).

As a final comment we want to stress that, while the proof of the existence of an inversion plan in a smooth context relies mainly on the bi-Lipschitz regularity of the exponential map, such ingredient on a general Alexandrov space fails to hold. Hence a finer analysis is needed. Combining the Lipschitz regularity of the exponential map together with the fact that it maps sets of positive measure to sets of positive measure, using also Disintegration Theorem, one can construct a “local” inversion plan. Suitably iterating this construction one can obtain a “global” inversion plan and prove Theorem 7.3. The existence of an inversion plan can be therefore understood as a measure-theoretic reformulation of a bi-Lipschitz control on the behavior of geodesic hinges and, as for a general Alexandrov space not so much can be proved on the regularity of the exponential map, Theorem 7.3 seems to be a novelty also in this direction.

We end the introduction with some comments regarding possible future applications of the techniques here introduced.

- In the class of those metric measure spaces admitting a bi-Lipschitz map with a subset of a Euclidean space, one of the relevant questions to ask is whether the image of such a metric space has positive Lebesgue measure or not. To this aim, let (X, d, μ) satisfy MCP, $\phi : X \rightarrow \mathbb{R}^d$ be a bi-Lipschitz map for some positive d and call $\eta := (\phi)_\# \mu$. Proving that η verifies the non-degeneracy condition (see Definition 3.1) would imply, thanks to Theorem 1.1, that $\mathcal{L}^d(\phi(X)) > 0$; yielding the rectifiability property of (X, d, μ) .
- In the recent [17], Csoranyi and Jones proved that if a Radon measure μ over \mathbb{R}^d makes the triple $(\mathbb{R}^d, |\cdot|, \mu)$ a Lipschitz differentiability space (see [12] for the definition of Lipschitz differentiability space), then μ is absolutely continuous with respect to the d -dimensional Lebesgue measure \mathcal{L}^d . Then Theorem 1.1 could be compared with [17]. In particular a relevant question could be to check if a metric measure space verifying the *non-degeneracy condition* is also a Lipschitz differentiability space.
- Understanding and characterizing the singular part of a non smooth space is an important and challenging issue. As explained above the set of inversion points is strongly linked with the regular set, i.e. the complementary of the singular set. It would be interesting to further investigate this link for instance in Alexandrov, or more generally in $\text{RCD}^*(K, N)$ -spaces.
- Given an $\text{RCD}^*(K, N)$ -space (X, d, \mathfrak{m}) , it is an interesting open problem whether the measure \mathfrak{m} is absolutely continuous with respect to the Hausdorff measure of the relevant dimension. We believe that the techniques here introduced could be useful to attach this question which would be the natural generalization to $\text{RCD}^*(K, N)$ -spaces to the aforementioned result of Cheeger-Colding [15] in the framework of *pmGH*-limits of Riemannian manifolds satisfying lower Ricci curvature bounds.

- In Section 6 we address the problem of stability of a reference measure with respect to the *pmGH*-convergence. As a next step one would like to study measured tangent spaces to pointed m.m.s. endowed with a reference measure. Recall that the set $\text{Tan}(X, \mathbf{d}, \mu, \bar{x})$ of tangent spaces to (X, \mathbf{d}, μ) at \bar{x} is the collection of all *pmGH* limits of sequences $(X, 1/r_i \cdot \mathbf{d}, \mu_{\bar{x}}^{r_i}, \bar{x})$ where $\mu_{\bar{x}}^{r_i}$ is a properly rescaled version of μ .

Acknowledgment. A.M. acknowledges the support of the ETH Fellowship. The project started during the conference YEP XI at EURANDOM (Eindhoven, The Netherlands) and continued during the “ERC Research Period on Calculus of Variations and Analysis in Metric Spaces” at Scuola Normale Superiore (Pisa-Italy), the authors wish to thank the institutions for the excellent working conditions and the organizers of the events for the stimulating atmosphere. Many thanks also to Luigi Ambrosio and Tapio Rajala for carefully reading a preliminary version of the paper, to Anton Petrunin for a fruitful conversation on Alexandrov spaces and to the two anonymous referees for the valuable comments and suggestions.

2. SETTING

We now recall some terminology and general notation. In what follows we say that a triple $(X, \mathbf{d}, \mathbf{m})$ is a metric measure space, m.m.s. for short, if and only if:

- (X, \mathbf{d}) is a complete and separable metric space;
- the measure \mathbf{m} belongs to $\mathcal{M}^+(X)$,

where $\mathcal{M}^+(X)$ denotes the space of positive Radon measure over X . We will also sometimes assume \mathbf{m} to be a probability measure, that is $\mathbf{m}(X) = 1$, but this will be specified at the beginning of each section. Moreover a metric space is a geodesic space if and only if for each $x, y \in X$ there exists $\gamma \in \text{Geo}$ so that $\gamma_0 = x, \gamma_1 = y$. Here we are using the following notation for the space of geodesics:

$$\text{Geo} := \{\gamma \in C([0, 1], X) : \mathbf{d}(\gamma_s, \gamma_t) = (s - t)\mathbf{d}(\gamma_0, \gamma_1), s, t \in [0, 1]\}.$$

The metric ball is denoted with $B_r(o) := \{z \in X : \mathbf{d}(z, o) < r\}$. Recall that for complete geodesic spaces local compactness is equivalent to properness (a metric space is proper if every closed ball is compact). As we will always deal with spaces enjoying some type of measure contraction property implying an upper bound on the dimension of the space, we directly assume the ambient space (X, \mathbf{d}) to be proper. Hence from now on we assume the following: the ambient metric space (X, \mathbf{d}) is geodesic, complete, separable and proper.

We will also use quite often the set of geodesics ending in a fixed point of the space: for $z \in X$

$$\text{Geo}(z) := \{\gamma \in \text{Geo} : \gamma_1 = z\} = \text{Geo} \cap e_1^{-1}(z),$$

where for any $t \in [0, 1]$, e_t denotes the evaluation map:

$$e_t : \text{Geo} \rightarrow X, \quad e_t(\gamma) := \gamma_t.$$

A geodesic metric space (X, \mathbf{d}) is said to be non-branching if and only if for any $\gamma^1, \gamma^2 \in \text{Geo}$, it holds:

$$\gamma_0^1 = \gamma_0^2, \gamma_{\bar{t}}^1 = \gamma_{\bar{t}}^2, \bar{t} \in (0, 1) \implies \gamma_s^1 = \gamma_s^2, \quad \forall s \in [0, 1].$$

We denote with $\mathcal{P}_2(X)$ the space of probability measures with finite second moment endowed with the L^2 -Wasserstein distance W_2 defined as follows: for $\mu_0, \mu_1 \in \mathcal{P}_2(X)$ we set

$$(2.1) \quad W_2^2(\mu_0, \mu_1) = \inf_{\pi} \int_X \mathbf{d}^2(x, y) \pi(dx dy),$$

where the infimum is taken over all $\pi \in \mathcal{P}(X \times X)$ with μ_0 and μ_1 as the first and the second marginal. Assuming the space (X, d) to be geodesic, also the space $(\mathcal{P}_2(X), W_2)$ is geodesic. It turns out that any geodesic $(\mu_t)_{t \in [0,1]}$ in $(\mathcal{P}_2(X), W_2)$ can be lifted to a measure $\nu \in \mathcal{P}(\text{Geo})$, so that $(e_t)_\# \nu = \mu_t$ for all $t \in [0,1]$. Given $\mu_0, \mu_1 \in \mathcal{P}_2(X)$, we denote by $\text{OptGeo}(\mu_0, \mu_1)$ the space of all $\nu \in \mathcal{P}(\text{Geo})$ for which $(e_0, e_1)_\# \nu$ realizes the minimum in (2.1). If (X, d) is geodesic, then the set $\text{OptGeo}(\mu_0, \mu_1)$ is non-empty for any $\mu_0, \mu_1 \in \mathcal{P}_2(X)$.

It is also convenient to introduce the following closed sets:

$$(2.2) \quad \begin{aligned} H &:= \{(x, y, z) \in X^3 : d(x, y) = d(x, z) + d(z, y)\}, \\ H(z) &:= P_{12}(H \cap (X^2 \times \{z\})) = \{(x, y) \in X^2 : d(x, y) = d(x, z) + d(z, y)\}, \end{aligned}$$

where for any $n \in \mathbb{N}$, $P_{ij} : X^n \rightarrow X^2$ is the projection on the ij -th component, for $i, j = 1, \dots, n$.

2.1. The essential non-branching property. We recall the following definition.

Definition 2.1. A metric measure space (X, d, m) is *essentially non-branching* if and only if for any $\mu_0, \mu_1 \in \mathcal{P}_2(X)$ which are absolutely continuous with respect to m any element of $\text{OptGeo}(\mu_0, \mu_1)$ is concentrated on a set of non-branching geodesics.

If (X, d, m) is essentially non-branching, one can deduce some information on the branching points as follows. Consider the set

$$(2.3) \quad D := \{(z, w) \in X \times X : \gamma^1, \gamma^2 \in \text{Geo}, \gamma_0^1 = \gamma_0^2 = w, \gamma_1^1 = \gamma_1^2 = z, \gamma^1 \neq \gamma^2\},$$

and observe that thanks to the properness assumption, it is σ -compact: it can be written as countable union of compact sets. Observe that also $X^2 \setminus D$ is σ -compact and D is symmetric: if $(z, w) \in D$ also $(w, z) \in D$. From essential non-branching we can deduce that the σ -compact set

$$D(z) := P_2(D \cap (\{z\} \times X)),$$

has m -measure zero (see for instance [39], in any case this statement will follow by the arguments below which give a more detailed description of the branching set).

We need a more refined property of branching structures that can be obtained observing that any L^2 optimal transportation between any measure and a Dirac delta, say in $z \in X$, is also an L^1 -optimal transportation. What follows is contained in [10] and is proved for RCD spaces but the same result holds for essentially non-branching spaces.

Consider the following closed sets:

$$\Gamma := \{(x, y, z) \in X^3 : d(x, z) - d(y, z) = d(x, y)\}, \quad \Gamma^{-1} := \{(x, y, z) \in X^3 : (y, x, z) \in \Gamma\},$$

and $R := \Gamma \cup \Gamma^{-1}$. We want to analyze possible branching structures inside $X \times X$. We therefore consider the set of forward branching:

$$(2.4) \quad A_+ := \{(x, z) \in X \times X : \exists y, w \in X, (x, y, z), (x, w, z) \in \Gamma, (y, w, z) \notin R\},$$

and the set of backward branching:

$$(2.5) \quad A_- := \{(x, z) \in X \times X : \exists y, w \in X, (x, y, z), (x, w, z) \in \Gamma^{-1}, (y, w, z) \notin R\}.$$

As the space X is proper by assumption, it is easily seen that both A_+ and A_- are σ -compact sets. The information we will use is the following: define

$$A_+(z) := P_1((X \times \{z\}) \cap A_+), \quad A_-(z) := P_1((X \times \{z\}) \cap A_-),$$

then it follows from Proposition 4.5 of [10] (notice that what is denoted with \mathcal{T} in [10], in the present framework coincides with the whole $X \times X$) that

$$(2.6) \quad m(A_+(z)) = m(A_-(z)) = 0,$$

provided (X, d, \mathbf{m}) is essentially non-branching. In particular the σ -compact set

$$\mathcal{T}_{nb}(z) := X \setminus (A_+(z) \cup A_-(z)),$$

contains all the points, up to a set of \mathbf{m} -measure zero, moved via optimal transportation towards the end point z and the whole set $\mathcal{T}_{nb}(z)$ is covered by a family of non-branching geodesics. It will also be convenient to introduce the set $\mathcal{T}_{nb} := X \times X \setminus (A_+ \cup A_-)$ and to observe that trivially

$$\mathcal{T}_{nb}(z) = P_1((X \times \{z\}) \cap \mathcal{T}_{nb}).$$

2.2. Disintegration of measures. We include here a version of the Disintegration Theorem (for a comprehensive treatment see for instance [19]).

Given a measurable space (R, \mathcal{R}) , i.e. \mathcal{R} is a σ -algebra of subsets of R , and a function $r : R \rightarrow S$, with S general set, we can endow S with the *push forward σ -algebra* \mathcal{S} of \mathcal{R} :

$$Q \in \mathcal{S} \iff r^{-1}(Q) \in \mathcal{R},$$

which could be also defined as the biggest σ -algebra on S such that r is measurable. Moreover given a probability measure ρ on (R, \mathcal{R}) , define a probability measure η on (S, \mathcal{S}) by push forward via r , i.e. $\eta := r_{\#}\rho$.

Definition 2.2. A *disintegration* of ρ consistent with r is a map (with slight abuse of notation still denoted with) $\rho : \mathcal{R} \times S \rightarrow [0, 1]$ such that, set $\rho_s(B) := \rho(B, s)$, the following hold:

- (1) $\rho_s(\cdot)$ is a probability measure on (R, \mathcal{R}) for all $s \in S$,
- (2) $\rho_s(B)$ is η -measurable for all $B \in \mathcal{R}$,

and satisfies for all $B \in \mathcal{R}, C \in \mathcal{S}$ the consistency condition

$$\rho(B \cap r^{-1}(C)) = \int_C \rho_s(B) \eta(ds).$$

A disintegration is *strongly consistent with respect to r* if for all s we have $\rho_s(r^{-1}(s)) = 1$. The measures ρ_s are called *conditional probabilities*.

We recall the following version of the disintegration theorem that can be found in [19, Section 452]. Recall that a σ -algebra \mathcal{J} is *countably generated* if there exists a countable family of sets so that \mathcal{J} coincide with the smallest σ -algebra containing them.

Theorem 2.3 (Disintegration of measures). *Assume that (R, \mathcal{R}, ρ) is a countably generated probability space and $R = \cup_{s \in S} R_s$ is a partition of R . Denote with $r : R \rightarrow S$ the quotient map:*

$$s = r(x) \iff x \in R_s,$$

and with (S, \mathcal{S}, η) the quotient measure space. Assume $(S, \mathcal{S}) = (X, \mathcal{B}(X))$ with X Polish space, where $\mathcal{B}(X)$ denotes the Borel σ -algebra. Then there exists a unique strongly consistent disintegration $s \mapsto \rho_s$ w.r.t. r , where uniqueness is understood in the following sense: if ρ_1, ρ_2 are two consistent disintegrations then $\rho_{1,s}(\cdot) = \rho_{2,s}(\cdot)$ for η -a.e. $s \in S$.

3. NON-DEGENERATE REFERENCE MEASURE ON \mathbb{R}^d

Let \mathfrak{m} be a positive Radon measure over \mathbb{R}^d that we will consider to be equipped with the Euclidean distance. Denote with Ω the support of \mathfrak{m} : $\Omega := \text{spt}(\mathfrak{m})$.

Definition 3.1 (Non-Degenerate measure). Consider the m.m.s. $(\Omega, |\cdot|, \mathfrak{m})$. We say that the measure \mathfrak{m} is *non-degenerate* if and only if for any compact set $A \subset \Omega$ of positive \mathfrak{m} -measure it holds

$$(3.1) \quad \mathfrak{m}(A_{t,o}) > 0, \quad \forall t \in [0, 1),$$

where o is any element of Ω and

$$(3.2) \quad A_{t,o} := \{z \in \mathbb{R}^d : z = (1-t)y + to, y \in A\}.$$

Remark 3.2. The non degeneracy condition may be regarded as a very weak curvature condition on the m.m.s. $(\Omega, |\cdot|, \mathfrak{m})$. Indeed it is implied by $\text{MCP}(K, n)$, for any K and n .

For the rest of this section we tacitly assume that \mathfrak{m} is a non-degenerate measure. We start by proving a geometric property of Ω .

Lemma 3.3. *The set Ω is a convex subset of \mathbb{R}^d .*

Proof. Consider $x, y \in \text{spt}(\mathfrak{m})$. Then for any positive radius δ it holds $\mathfrak{m}(B_\delta(x) \cap \Omega) > 0$. Then (3.1) implies that

$$\mathfrak{m}((\bar{B}_\delta(x) \cap \Omega)_{t,y}) > 0, \quad \forall t \in [0, 1).$$

Now observe that for any $t \in [0, 1)$,

$$(\bar{B}_\delta(x) \cap \Omega)_{t,y} \subset B_{(1-t)\delta}(x_t), \quad \text{with } x_t := (1-t)x + ty.$$

It follows that $\mathfrak{m}(B_{(1-t)\delta}(x_t)) > 0$ for any positive δ , therefore $x_t \in \text{spt}(\mathfrak{m}) = \Omega$ and the claim follows. \square

We deduce from Lemma 3.3 that Ω has a well defined dimension: there exists a minimal $k \in \mathbb{N}$ with $k \leq d$ so that Ω is contained in a k -dimensional affine subspace of \mathbb{R}^d that we identify with \mathbb{R}^k .

We can therefore consider the relative interior and the relative boundary of Ω in \mathbb{R}^k : there exist $U, C \subset \mathbb{R}^k$ so that $\Omega = U \cup C$ and $U \cap C = \emptyset$ with U maximal open subset in \mathbb{R}^k contained in Ω .

Proposition 3.4. *If k is the dimension of Ω , the measure \mathfrak{m}_U is absolutely continuous with respect to the k -dimensional Lebesgue measure \mathcal{L}^k .*

Proof. Assume by contradiction the existence of a compact set $K \subset U$ so that

$$\mathfrak{m}(K) > 0, \quad \mathcal{L}^k(K) = 0.$$

The statement we are proving is local, therefore we can consider $x_0 \in K$ and $\delta > 0$ so that $B_\delta(x_0) \subset U$ and $K \subset B_\delta(x_0)$. Since the measure \mathfrak{m} is non-degenerate in $(\Omega, |\cdot|, \mathfrak{m})$, for every

$o \in \partial B_\delta(x_0)$ and $t \in [0, 1]$ it holds $\mathfrak{m}(K_{t,o}) > 0$. Therefore we have:

$$\begin{aligned}
0 &< \int_{[0,1]} \int_{\partial B_\delta(x_0)} \mathfrak{m}(K_{t,o}) \mathcal{H}^{k-1}(do) \mathcal{L}^1(dt) \\
&= \mathfrak{m} \otimes \mathcal{H}^{k-1} \otimes \mathcal{L}^1(\{(z, o, t) \in B_\delta(x_0) \times \partial B_\delta(x_0) \times [0, 1] : z = (1-t)x + to, x \in K\}) \\
&= \int_{B_\delta(x_0)} \mathcal{H}^{k-1} \llcorner_{\partial B_\delta(x_0)} \otimes \mathcal{L}^1 \llcorner_{[0,1]}(\{(o, t) : f_z(o, t) \in K\}) \mathfrak{m}(dz) \\
(3.3) \quad &= \int_{B_\delta(x_0)} (f_z)_\# \left(\mathcal{H}^{k-1} \llcorner_{\partial B_\delta(x_0)} \otimes \mathcal{L}^1 \llcorner_{[0,1]} \right) (K) \mathfrak{m}(dz),
\end{aligned}$$

where

$$f_z(o, t) := \frac{z - to}{1 - t}, \quad \forall t \in [0, 1].$$

Since $\mathcal{H}^{k-1} \llcorner_{\partial B_\delta(x_0)} \otimes \mathcal{L}^1 \llcorner_{[0,1]}$ is equivalent to $\mathcal{L}^k \llcorner_{B_\delta(x_0)}$, the function f_z can be defined directly on $B_\delta(x_0)$. For ease of notation we also assume $x_0 = 0$, then:

$$f_z(w) := \frac{z - w}{1 - |w|/\delta} = \delta \cdot \frac{z - w}{\delta - |w|}.$$

Then

$$\begin{aligned}
df_z(w) &= \delta(z - w) \nabla \left(\frac{1}{\delta - |w|} \right)^T - \frac{\delta}{\delta - |w|} Id \\
&= \delta(z - w) \frac{1}{(\delta - |w|)^2} \left(\frac{w}{|w|} \right)^T - \frac{\delta}{\delta - |w|} Id.
\end{aligned}$$

Using the following Lemma 3.5, we get

$$\begin{aligned}
\det \left(-\frac{(\delta - |w|)}{\delta} df_z(w) \right) &= \det \left(Id - \frac{1}{\delta - |w|} (z - w) \left(\frac{w}{|w|} \right)^T \right) \\
&= 1 - \frac{1}{(\delta - |w|)|w|} \langle z - w, w \rangle.
\end{aligned}$$

Thus $\det \left(-\frac{(\delta - |w|)}{\delta} df_z(w) \right) = 0$ only when $\delta|w| = \langle z, w \rangle$. Since $z, w \in B_\delta(0)$, it follows that $\det(df_z(w)) \neq 0$ for all w and z . Therefore, $\mathcal{L}^k(K) = 0$ implies $(f_z)_\# \left(\mathcal{H}^{k-1} \llcorner_{\partial B_\delta(x_0)} \otimes \mathcal{L}^1 \llcorner_{[0,1]} \right) (K) = 0$ contradicting (3.3) and the claim is proved. \square

Lemma 3.5. *Let Id be the $n \times n$ identity matrix and let a, b be $n \times 1$ column vectors. Then*

$$\det(Id + ab^T) = 1 + b^T a.$$

Proof. Observe that one can write

$$\begin{bmatrix} Id & 0 \\ b^T & 1 \end{bmatrix} \cdot \begin{bmatrix} Id + ab^T & a \\ 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} Id & 0 \\ -b^T & 1 \end{bmatrix} = \begin{bmatrix} Id & a \\ 0 & 1 + b^T a \end{bmatrix}$$

The proof follows then by the standard product rules of the determinant and of matrices in block forms. \square

To conclude we analyze the boundary. Putting together the next Proposition and the previous results, Theorem 1.1 is proved.

Proposition 3.6. *If C is the boundary of U for the trace topology of \mathbb{R}^k , then $\mathfrak{m}(C) = 0$.*

Proof. Since Ω is convex, there exists $o \in U$ so that $C_{t,o} \subset U$ for all $t \in (0, 1)$, where $C_{t,o}$ is defined as in (3.2). Suppose now by contradiction that $\mathfrak{m}(C) > 0$. Then by non-degeneracy

$$\mathfrak{m}(C_{t,o}) > 0, \quad \forall t \in (0, 1),$$

and therefore from Proposition 3.4, $\mathcal{L}^k(C_{t,o}) > 0$ for all $t \in (0, 1)$. But on the other hand, C is the boundary of a convex set U in \mathbb{R}^k , thus $\mathcal{L}^k(C) = 0$; therefore, since the map

$$C \ni x \mapsto T_t(x) := (1-t)x + to$$

is a diffeomorphism, it follows that $\mathcal{L}^k(C_{t,o}) = 0$. This gives a contradiction and the claim follows. \square

4. GENERAL PROPERTIES OF MCP

This Section serves as a general picture of the properties of m.m.s. verifying MCP and it is divided into two parts. In the first one we analyze the geodesic convexity of the support of the measure, in the second one we investigate the implications of MCP on the extendability of geodesics.

So again we recall that (X, d, \mathfrak{m}) is a metric measure space; i.e. (X, d) is a geodesic, proper, complete and separable metric space and \mathfrak{m} is a positive Radon measure on X .

The quantitative dependance of MCP with respect to lower curvature bound K and the dimension upper bound N will not play any role in our proofs, we prefer therefore to consider the following *qualitative* form of MCP.

Definition 4.1. Let (X, d, \mathfrak{m}) be a m.m.s. as before, denote with $\Omega := \text{spt}(\mathfrak{m})$ and let $\bar{x} \in X$. The triple (X, d, \mathfrak{m}) verifies the *qualitative* MCP if and only if there exist a sequence $R_j \uparrow +\infty$ and continuous functions $f_j : [0, 1] \rightarrow \mathbb{R}^+$ with $f_j(0) = 1$, $f_j(1) = 0$ and $f_j(t) > 0$ for any $t \in [0, 1)$, $j \in \mathbb{N}$, so that for every $o \in \Omega \cap B_{R_j}(\bar{x})$:

$$(4.1) \quad \mathfrak{m}(A_{t,o}) \geq f_j(t) \mathfrak{m}(A), \quad \forall t \in [0, 1),$$

where

$$(4.2) \quad A_{t,o} := \{\gamma_t : \gamma \in \text{Geo}(o), \gamma_0 \in A\},$$

and $A \subset \Omega \cap B_{R_j}(\bar{x})$ is any Borel set.

Clearly the property of satisfying the qualitative MCP condition above is independent of the center \bar{x} , just by changing the sequence of radii $R_j \uparrow +\infty$.

At this level of generality not so much can be said on the geometry of Ω . We therefore introduce the next stronger version of the qualitative measure contraction property.

Definition 4.2 (Strong Qualitative MCP). A triple $(\Omega, d, \mathfrak{m})$ as above satisfies the *strong qualitative* MCP provided there exist a sequence $R_j \uparrow +\infty$ and continuous functions $f_j : [0, 1] \rightarrow \mathbb{R}^+$ with $f_j(0) = 1$, $f_j(1) = 0$ and $f_j(t) > 0$ for any $t \in [0, 1)$, $j \in \mathbb{N}$, so that for every $o \in \Omega \cap B_{R_j}(\bar{x})$:

$$(4.3) \quad \mathfrak{m}(A) \geq f_j(t) \mathfrak{m}(e_0(e_t^{-1}(A) \cap \text{Geo}(o))), \quad \forall t \in [0, 1),$$

where $A \subset \Omega \cap B_{R_j}(\bar{x})$ is any Borel set.

Remark 4.3. Few comments on Definition 4.2 are in order. The adjective “strong” is due to the fact that Definition 4.2 is equivalent to impose (4.1) replacing $A_{t,o}$ with $e_t(H)$, for all the $H \subset \text{Geo}(o)$ so that $e_0(H) = A$. Hence Definition 4.2 imposes the evolution estimate along the support of any W_2 -geodesic connecting $\mathfrak{m}(A)^{-1}\mathfrak{m}_A$ to δ_o . It follows therefore that strong qualitative MCP implies qualitative MCP.

On the other hand, if (X, d, m) satisfies the qualitative MCP and it is essentially non-branching (see (2.3)), then it verifies the strong qualitative MCP:

Fix any $o \in \Omega$ and $A \in B_{R_j}(\bar{x})$, and let $H = e_t^{-1}(A) \cap \text{Geo}(o)$. Denote with \hat{X} the set $X \setminus D(o)$ and recall that, by the essentially non-branching condition, one has $m(D(o)) = 0$. The qualitative MCP implies that

$$\begin{aligned} m(A) &\geq m(e_t(H)) \geq m\left(\left(e_0(H) \cap \hat{X}\right)_{t,o}\right) \geq f_j(t) m(e_0(H) \cap \hat{X}) = f_j(t) m(e_0(H)) \\ &= f_j(t) m(e_0(e_t^{-1}(A) \cap \text{Geo}(o))), \end{aligned}$$

which is (4.3).

All the constructions that we will consider will be then used for spaces verifying the essentially non-branching property. Hence we prefer to not pursue the most general result and to prove the next two statements under the strong version of the qualitative MCP. At the moment we do not know if they hold under the general qualitative MCP.

The terminology “strong” is justified by the next result.

Lemma 4.4. *Let (Ω, d, m) be a metric measure space verifying the strong qualitative MCP condition (4.3). Then for any $A \subset \Omega$ measurable set with $m(A) \in (0, \infty)$, $o \in \Omega$ and for any geodesic $(\mu_t)_{t \in [0,1]} \subset \mathcal{P}_2(X)$ so that*

$$\mu_0 = \frac{1}{m(A)} m_{\perp A}, \quad \mu_1 = \delta_o,$$

it holds $\mu_t \ll m$ for all $t \in [0, 1]$.

Proof. Fix $A \subset \Omega$ measurable with $m(A) \in (0, \infty)$, a point $o \in \Omega$ and let $(\mu_t)_{t \in [0,1]} \subset \mathcal{P}_2(X)$ be an arbitrary W_2 -geodesic from $\mu_0 := \frac{1}{m(A)} m_{\perp A}$ to $\mu_1 := \delta_o$. Since (X, d) is a geodesic space, we can associate a measure $\nu \in \mathcal{P}(\text{Geo})$ so that for any $t \in [0, 1]$ it holds $(e_t)_\# \nu = \mu_t$; finally denote with $H \subset \text{Geo}(o)$ a set where ν is concentrated.

Fix $t \in [0, 1)$ and $K \subset \Omega$ compact subset so that $m(K) = 0$. Then, by (4.3), we have

$$m(e_0(e_t^{-1}(K) \cap \text{Geo}(o))) = 0,$$

and consequently

$$\begin{aligned} \mu_t(K) &= (e_t)_\# \nu(K) = \nu(e_t^{-1}(K) \cap \text{Geo}(o)) \\ &\leq \nu(e_0^{-1}(e_0(e_t^{-1}(K) \cap \text{Geo}(o)))) \\ &= (e_0)_\# \nu(e_0(e_t^{-1}(K) \cap \text{Geo}(o))) \\ &= \frac{1}{m(A)} m_{\perp A}(e_0(e_t^{-1}(K) \cap \text{Geo}(o))) \\ &= 0. \end{aligned}$$

□

In analogy of what we proved in \mathbb{R}^d , also in this framework one has that the support Ω of m is convex.

Proposition 4.5. *Let (X, d, m) be a m.m.s. and $\Omega := \text{spt}(m)$ is so that the triple (Ω, d, m) satisfies the strong qualitative MCP condition (4.3). Then Ω is a weakly geodesically convex subset, i.e for every $x_0, x_1 \in \Omega$ there exists $\hat{\gamma} \in \text{Geo}$ with $\hat{\gamma}_i = x_i$, $i = 0, 1$, such that $\hat{\gamma}([0, 1]) \subset \Omega$.*

Proof. Let $x, y \in \Omega$ and $\delta > 0$. We consider the d^2 -optimal transportation problem between the marginal measures

$$\mu_0 := \frac{1}{\mathfrak{m}(B_\delta(x))} \mathfrak{m} \llcorner_{B_\delta(x)}, \quad \mu_1 := \delta_y.$$

The associated measure $\nu^\delta \in \mathcal{P}(\text{Geo})$ is so that $(e_t)_\#(\nu^\delta)_{t \in [0,1]}$ is a W_2 -geodesic connecting μ_0 to μ_1 . We now consider a sequence $\delta_n > 0$ converging to 0 as $n \rightarrow \infty$. The associated measure ν^{δ_n} gives measure one to the following set

$$G_n := \{\gamma \in \text{Geo} : \gamma_0 \in \overline{B_{\delta_n}}(x), \gamma_1 = y\}.$$

By compactness, there exists a subsequence, still denoted with δ_n , so that $G_n \rightarrow H$ in the Hausdorff distance, with H compact subset of

$$G := \{\gamma \in \text{Geo} : \gamma_0 = x, \gamma_1 = y\}.$$

Also the set $\{\nu^{\delta_n}\}_{n \in \mathbb{N}} \subset \mathcal{P}(\text{Geo})$ is W_2 -precompact, and extracting another subsequence we deduce the existence of $\nu \in \mathcal{P}(\text{Geo})$ so that $\nu^{\delta_n} \rightarrow \nu$ in the W_2 -metric. Moreover $\nu(H) = 1$ and there exists $\hat{\gamma} \in H$ so that $\nu(B_\alpha^{\text{Geo}}(\hat{\gamma})) > 0$ for all $\alpha > 0$, where

$$B_\alpha^{\text{Geo}}(\hat{\gamma}) := \{\gamma \in \text{Geo} : \sup_{t \in [0,1]} d(\gamma_t, \hat{\gamma}_t) < \alpha\}.$$

By lower-semicontinuity over open sets,

$$\liminf_{n \rightarrow \infty} \nu^{\delta_n}(B_\alpha^{\text{Geo}}(\hat{\gamma})) \geq \nu(B_\alpha^{\text{Geo}}(\hat{\gamma})) > 0.$$

Hence for every $\alpha > 0$ there exists δ_n so that

$$0 < \nu^{\delta_n}(B_\alpha^{\text{Geo}}(\hat{\gamma})) \leq ((e_t)_\# \nu^{\delta_n})(e_t(B_\alpha^{\text{Geo}}(\hat{\gamma}))) \leq ((e_t)_\# \nu^{\delta_n})(B_\alpha(\hat{\gamma}_t)),$$

for all $t \in [0, 1]$. By Lemma 4.4 we also deduce that $((e_t)_\# \nu^{\delta_n}) \ll \mathfrak{m} \llcorner_{e_t(G_n)}$, for all $n \in \mathbb{N}$ and $t \in [0, 1]$. We can therefore summarize what we obtained as follows: there exists $\hat{\gamma} \in \text{Geo}(X)$ so that

$$\mathfrak{m}(B_\alpha(\hat{\gamma}_t)) > 0, \quad \text{for every } \alpha > 0 \text{ and } t \in [0, 1].$$

It follows that $\hat{\gamma}_t \in \text{spt}(\mathfrak{m}) = \Omega$ for every $t \in [0, 1]$, as desired. \square

Thanks to Proposition 4.5, $(\Omega, d, \mathfrak{m})$ is a geodesic metric measure space. We will identify with no loss of generality the whole space X with the support Ω of the measure \mathfrak{m} . Therefore any future assumption on the smoothness or geometry of the m.m.s. (X, d, \mathfrak{m}) is indeed an assumption that should be verified only on Ω .

4.1. Structure of geodesics under MCP. In this section we establish some structural properties of the geodesics implied by the qualitative MCP condition (4.1) under the essentially non-branching condition. The next is a well-known property of m.m.s. enjoying MCP(K, N) and it is valid as well under the qualitative MCP (see for instance the proof of [23, Lemma 3.1]).

Let the m.m.s. (X, d, \mathfrak{m}) satisfy the qualitative MCP. Then for any $z \in X$, \mathfrak{m} -a.e. point of X is an interior point of some geodesic ending in z :

$$\mathfrak{m}(\{\gamma_t : \gamma \in \text{Geo}(z), t \in (0, 1)\}) = 1.$$

The goal of this section is to improve such statement. For any point z of the space, consider the set of geodesics having z as an interior point, called *rays through z* and the evaluation of them, called *points through z* . They are defined in the following way:

$$R(z) := \{\gamma \in \text{Geo} : \gamma_s = z, \text{ for some } s \in (0, 1)\}, \quad e_{(0,1)}(R(z)) := \{\gamma_t : \gamma \in R(z), t \in (0, 1)\}.$$

We can now state the result.

Proposition 4.6. *Let the m.m.s. (X, d, m) be essentially non-branching and satisfy the qualitative MCP condition (4.1). Then for m-a.e. $z \in X$ the following holds:*

$$(4.4) \quad m(X \setminus e_{(0,1)}(R(z))) = 0.$$

Proof. We will prove the following: for any $x \in X$ and $r > 0$, for m-a.e. $z \in B_r(x)$, it holds

$$m(B_r(x) \setminus e_{(0,1)}(R(z))) = 0.$$

So let us fix $x \in X$ and $r > 0$. We also consider the function $f = f_j$ given by the qualitative MCP condition associated to an open ball containing $B_r(x)$. Then we have

$$\int_{B_r(x)} m\left((B_r(x))_{t,z}\right) m(dz) \geq f(t) m(B_r(x))^2.$$

In equivalent terms,

$$\begin{aligned} m(B_r(x))^2 f(t) &\leq m \otimes m(\{(z, w) \in B_r(x) \times X : w = \gamma_t, \gamma \in \text{Geo}(z), \gamma_0 \in B_r(x)\}) \\ &= \int_X m(\{z \in B_r(x) : w = \gamma_t, \gamma \in \text{Geo}(z), \gamma_0 \in B_r(x)\}) m(dw) \\ &= \int_{B_{(1+2t)r}(x)} m(\{z \in B_r(x) : w = \gamma_t, \gamma \in \text{Geo}(z), \gamma_0 \in B_r(x)\}) m(dw) \\ &\leq \int_{B_{(1+2t)r}(x)} m(\{z \in B_r(x) : w \in \gamma_{(0,1)}, \gamma \in \text{Geo}(z), \gamma_0 \in B_r(x)\}) m(dw). \end{aligned}$$

Taking the limit as $t \rightarrow 0$, since f is continuous and $f(0) = 1$, we get that for m-a.e. $w \in B_r(x)$

$$m(\{z \in B_r(x) : w \in \gamma_{(0,1)}, \gamma \in \text{Geo}(z), \gamma_0 \in B_r(x)\}) = m(B_r(x)).$$

To conclude we prove that set in the left hand side of the previous identity coincides with $e_{(0,1)}(R(z))$, up to a set of m-measure zero. To this purpose it is enough to show that the set of initial points of maximal geodesics has m-measure zero; but this follows from Proposition 4.8 below, since $m(D(z)) = 0$ by the essential non-branching assumption. \square

Proposition 4.6 has a nice consequence on the symmetric cut locus. Let us recall that the *symmetric cut locus* $SC \subset X \times X$ is defined by

$$(4.5) \quad \begin{aligned} SC &:= X \times X \setminus \{(x, y) \in X \times X : \exists \gamma \in \text{Geo}, \exists s, t \in (0, 1) \text{ such that } \gamma_s = x, \gamma_t = y\} \\ &= \{(x, y) \in X \times X : x \in C(y) \text{ or } y \in C(x)\}, \end{aligned}$$

where, given $z \in X$, the *cut locus* of z denoted with $C(z) \subset X$ is defined by

$$(4.6) \quad C(z) := X \setminus \{x \in X : \exists \gamma \in \text{Geo}, \exists t \in [0, 1) \text{ such that } \gamma_0 = z, \gamma_t = x\}.$$

It was already well-known that qualitative MCP implies that $m(C(z)) = 0$, see for instance (2.6). From Proposition 4.6, once we observe that for every $z \in X$ it holds

$$\{z\} \times e_{(0,1)}(R(z)) \subset X \times X \setminus SC,$$

it is then possible to deduce the following property of SC .

Corollary 4.7. *Let the m.m.s. (X, d, m) be essentially non-branching and satisfy the qualitative MCP. Then*

$$m \otimes m(SC) = 0.$$

The next result is already present in literature, see for instance [9] where a more general case is considered. To keep presentation as self contained as possible, we include it here in a form adjusted to our framework.

Proposition 4.8 (Disintegration with MCP). *Let the m.m.s. (X, d, m) with $m(X) = 1$ be essentially non-branching, and satisfy the qualitative MCP condition (4.1). Fix any $z \in X$ and define $\hat{X} := X \setminus \{D(z) \cup \{z\}\}$. Consider in \hat{X} the equivalence relation T :*

$$(x, y) \in T \iff d(y, z) = d(y, x) + d(x, z), \text{ or } d(x, z) = d(x, y) + d(y, z),$$

whose equivalence classes are maximal geodesics in $\text{Geo}(z)$. Then there exists an m -measurable map $Q : \hat{X} \rightarrow \hat{X}$ and a Borel subset $K \subset \hat{X}$ with $m(K) = 1$ so that

$$(x, y) \in T \iff Q(x) = Q(y) \quad \text{and} \quad (x, Q(x)) \in T, \quad \forall x, y \in K.$$

Moreover

$$m = m_{\hat{X}} = \int_{Q(\hat{X})} m^\alpha q(d\alpha), \quad m^\alpha \ll \mathcal{H}^1_{\perp \gamma^\alpha}, \quad q\text{-a.e. } \alpha \in Q(\hat{X}),$$

where $q = (Q)_\# m_{\hat{X}}$ is the quotient measure and $\gamma^\alpha \in \text{Geo}(z)$ is the maximal geodesic containing $\alpha \in Q(\hat{X})$.

Proof. Step 1. Since $m(\hat{X}) = 1$, we can assume the existence of $\hat{K} \subset \hat{X}$, σ -compact, so that $m(\hat{K}) = 1$; observe that the equivalence relation restricted to \hat{K}

$$T = \{(x, y) \in \hat{K} \times \hat{K} : d(y, z) = d(y, x) + d(x, z)\} \cup \{(x, y) \in K \times K : d(x, z) = d(x, y) + d(y, z)\},$$

is σ -compact as well. With a slight abuse of notation we set $T(\omega)$ to be the class of ω , i.e. the maximal geodesic through ω and z . Fix now a countable dense family of points $\{x_i\}_{i \in \mathbb{N}} \subset \hat{K}$ and consider for $i, j, k \in \mathbb{N}$

$$W_{ijk} := \{w \in \bar{B}_{2^{-j}}(x_i) \cap \hat{K} : L(T(w) \cap B_{2^{1-j}}(x_i)) \leq 2^{-k}, L(T(w)) \geq 2^{2-k}\},$$

where $L(\gamma)$ denotes the length of the geodesic γ . It can be proved that W_{ijk} form a countable cover of \hat{K} of class \mathcal{A} , the σ -algebra generated by the analytic sets, see [9, Lemma 4.1] for the details. As $W_{ijk} \in \mathcal{A}$, there exists $N_{ijk} \subset W_{ijk}$ with $m(N_{ijk}) = 0$ so that $W_{ijk} \setminus N_{ijk}$ is Borel. Then

$$H_{ijk} := T^{-1}(W_{ijk} \setminus N_{ijk}) = P_1(\{(v, w) \in T : w \in W_{ijk} \setminus N_{ijk}\})$$

is a countable covering of $\hat{K} \setminus \cup N_{ijk}$ into *saturated* (sets of the form $T^{-1}(A)$) analytic sets. Observing that the difference of two saturated sets is still saturated, we can find a countable disjoint family of sets $\{K_{ijk}\}$ made of \mathcal{A} saturated sets so that

$$K_{ijk} \subset H_{ijk}, \quad \cup_{ijk} K_{ijk} = \cup_{ijk} H_{ijk}.$$

For ease of notation we fix a bijection from $\mathbb{N} \rightarrow \mathbb{N}^3$ and denote with K_n the set $K_{i(n)j(n)k(n)}$. We can now define the following multivalued map:

$$K_n \ni x \mapsto F(x) := T(x) \cap \bar{B}_{2^{-(j(n))}}(x_{i(n)}),$$

that is easily seen to be \mathcal{A} -measurable. Observe that $F(x)$ is a closed subset of \hat{K} . Then by [9, Corollary 2.7] there exists $f_n : K_n \rightarrow \bar{B}_{2^{-(j(n))}}(x_{i(n)})$ \mathcal{A} -measurable so that

$$(x, f_n(x)) \in T, \quad (x, y) \in T \iff f_n(x) = f_n(y).$$

We can repeat the same procedure for any $m \in \mathbb{N}$ and then define $Q(x) := f_n(x)$ for all $x \in K_n$ and arbitrarily outside of $K := \cup_n K_n$. As all K_n are disjoint and saturated, the definition of Q

is well posed and as $\hat{X} \setminus K$ has \mathbf{m} -measure 0, Q is \mathbf{m} -measurable and the first part of the claim follows.

Step 2. The existence of an \mathbf{m} -measurable map Q obtained before is in fact equivalent to the strong consistency of the disintegration, see Proposition 4.4 of [9] and subsequent discussions. Hence

$$\mathbf{m} = \mathbf{m}_{\llcorner \hat{X}} = \int_{Q(\hat{X})} \mathbf{m}^\alpha q(d\alpha), \quad \mathbf{m}^\alpha(\gamma^\alpha) = 1, \quad q\text{-a.e. } \alpha \in Q(\hat{X}),$$

where $q = (Q)_\# \mathbf{m}_{\llcorner \hat{X}}$ is the quotient measure and $\gamma^\alpha \in \text{Geo}(z)$ is the unique maximal geodesic containing α . To obtain that q -a.e. \mathbf{m}^α is absolutely continuous with respect to $\mathcal{H}^1_{\llcorner \gamma^\alpha}$ it is enough to write the qualitative MCP between any Borel set and the fixed point $z \in X$. Then the equivalence classes are invariant sets for the evolution and, for q -a.e. α , each measure \mathbf{m}^α has to satisfy a non-degeneracy property of the same type of the one introduced in Section 3. Hence the claim follows (for the details of this second part of the proof, see [9, Theorem 5.7]). \square

5. REFERENCE MEASURES

We are now ready to define what is for us a reference measure. Recall that (X, \mathbf{d}) is a geodesic, proper, complete and separable metric space even if the next definition will make sense in a general complete and separable metric space. During this Section all the assumptions regarding finiteness of measures are dropped. Denote with $\mathcal{M}^+(X)$ the space of positive Radon measures.

Definition 5.1. A positive Radon measure $\mu \in \mathcal{M}^+(X)$ is a *reference measure* for (X, \mathbf{d}) provided it is non-zero, and for μ -a.e. $z \in X$ there exists $\pi^z \in \mathcal{M}^+(X \times X)$ so that

$$(5.1) \quad (P_1)_\# \pi^z = \mu, \quad \pi^z(X \times X \setminus H(z)) = 0, \quad (P_2)_\# \pi^z \ll \mu,$$

where $P_i : X \times X \rightarrow X$ is the projection on the i -th component, for $i = 1, 2$ and $H(z)$ is defined in (2.2). The measure π^z will be called *inversion plan*.

Let us briefly observe that, since $H(z) \subset X \times X$ is closed, clearly the requirement $\pi^z(X \times X \setminus H(z)) = 0$ is equivalent to $\text{spt}(\pi^z) \subset H(z)$. Moreover we will use the following notation

$$(5.2) \quad \text{Ip}(\mu) := \{z \in X : \exists \pi^z \text{ inversion plan}\},$$

to denote the set of *Inversion points* of μ , those point where an inversion plan exists.

Remark 5.2 (Inversion points are regular points). Let us remark that the condition for a point $p \in X$ to be an inversion point is strictly related to the regularity of the space (X, \mathbf{d}) at p . Indeed if (X, \mathbf{d}) has a conical singularity at p then clearly p cannot be an inversion point (unless $\mathbf{m}(\{p\}) > 0$) since all geodesics end to minimize length once they cross p . Indeed, since $H(p) \subset X \times \{p\} \cup \{p\} \cup X$, if $\pi \in \mathcal{M}^+(X)$ with $\pi(X \times X \setminus H(p)) = 0$ and $(P_1)_\#(\pi) = \mathbf{m}$, then $(P_2)_\#(\pi) = \mathbf{m}(X) \delta_p$.

On the other hand in Section 7 we will prove that \mathcal{H}^n -a.e. regular point in an Alexandrov space with lower curvature bound is an inversion point (for the precise statement see Theorem 7.3).

In order to have a clear statement for the main result of this section, we drop the identification between the ambient space X and the support of \mathbf{m} , Ω . Notice that, once we assume $(X, \mathbf{d}, \mathbf{m})$ to be essentially non-branching and satisfy the qualitative MCP condition (4.1), then by Proposition 4.5 we have that $\Omega = \text{spt}(\mathbf{m})$ is a geodesic space.

Theorem 5.3. *Let (X, d, m) be an essentially non-branching m.m.s. that verifies the qualitative MCP condition (4.1). Assume the existence of a reference measure μ for (Ω, d) , where $\Omega = \text{spt}(m)$, so that (Ω, d, μ) verifies the qualitative MCP condition (4.1) and it is essentially non-branching. If $m(X \setminus \text{lp}(\mu)) = 0$, then*

$$m \ll \mu.$$

Proof. We start by observing that from Proposition 4.5, the triples (Ω, d, m) and (Ω, d, μ) are geodesic, essentially non-branching m.m.s. verifying the qualitative MCP. Once the measures will be restricted to sets of finite measure, we will use the result of Proposition 4.8

Step 1. Assume by contradiction the claim is false. Then there exists $A \subset \Omega$, that we can assume to be a subset of $B_r(o) \subset \Omega$ for some $o \in \Omega$ and $r > 0$, so that

$$m(A) > 0 \quad \text{and} \quad \mu(A) = 0.$$

Since m verifies the qualitative MCP, we have the following strict inequality

$$\int_{B_r(o)} m(A_{1/2,x} \cap \mathcal{T}_{nb}(x)) \mu(dx) > 0,$$

where $\mathcal{T}_{nb}(x)$ has been introduced in Section 2.1 and guarantees that the transportation towards x moves along non-branching geodesics (recall that $\mathcal{T}_{nb}(x)$ has full measure thanks to (2.6)). The measurability of $A_{1/2,x} \cap \mathcal{T}_{nb}(x)$ follows from the fact that \mathcal{T}_{nb} is σ -compact; note we also used that $\mathcal{T}_{nb}(x)$ is of full m -measure thanks to the essential non-branching assumption. Using Fubini's Theorem to change order of integration, we get

$$(5.3) \quad 0 < \int_{B_{2r}(o)} \mu(\{x \in B_r(o) : \exists \gamma \in \text{Geo}(x) \text{ s.t. } z = \gamma_{1/2}, \gamma_0 \in A, \gamma_0, z \in \mathcal{T}_{nb}(x)\}) m(dz).$$

As $m(X \setminus \text{lp}(\mu)) = 0$, the set of integration can be substituted by $B_{2r}(o) \cap \text{lp}(\mu)$ without changing anything. Fix any $z \in B_{2r}(o) \cap \text{lp}(\mu)$. Since (Ω, d, m) is a geodesic m.m.s., from the qualitative MCP and Proposition 4.8 applied to z and $\mu_{\perp B_{2r}(z)}$ it follows that ^(a)

$$(5.4) \quad \mu_{\perp B_{2r}(z)} = \int_{Q(B_{2r}(z))} \mu^\alpha q(d\alpha), \quad \mu^\alpha \ll \mathcal{H}^1_{\perp \gamma^\alpha}, \quad q\text{-a.e. } \alpha \in Q(B_{2r}(z)),$$

where $q = Q_{\#}(\mu_{\perp B_{2r}(z)})$ is the quotient measure and γ^α is the maximal geodesic in $\text{Geo}(z)$ containing α . Moreover for q -a.e. $\alpha \in Q(B_{2r}(z))$, the density of μ^α with respect to $\mathcal{H}^1_{\perp \gamma^\alpha}$ is strictly positive at any point and so $\mathcal{H}^1_{\perp \gamma^\alpha} \ll \mu^\alpha$ for q -a.e. $\alpha \in Q(B_{2r}(z))$. Hence, $\mu(A) = 0$ implies that

$$\int_{Q(B_{2r}(z))} \mathcal{H}^1(\gamma^\alpha \cap A) q(d\alpha) = 0.$$

There exists therefore a q -measurable set $B \subset Q(B_{2r}(z))$ with $q(B) = 0$ so that

$$(5.5) \quad \mathcal{H}^1(\gamma^\alpha \cap A) = 0, \quad \forall \alpha \in Q(B_{2r}(z)) \setminus B.$$

Step 2. We now write the set

$$E = E(z) := \{x \in B_r(o) : \exists \gamma \in \text{Geo}(x) \text{ s.t. } z = \gamma_{1/2}, \gamma_0 \in A, \gamma_0, z \in \mathcal{T}_{nb}(x)\}$$

^(a)Observe that convexity of $B_{2r}(z)$ is not needed to apply Disintegration theorem, we only use that for each point in $x \in B_{2r}(z)$, any intermediate point between x and z belongs to $B_{2r}(z)$. The same property permits to use MCP to obtain the absolute continuity of the conditional measures with respect to the Hausdorff measure of dimension 1.

as union of the following two sets:

$$\begin{aligned} E_1 &:= \{x \in B_r(o) : \exists \gamma \in \text{Geo}(x) \text{ s.t } z = \gamma_{1/2}, \gamma_0 \in A, \gamma_0, z \in \mathcal{T}_{nb}(x), Q(\gamma_0) \in B\}, \\ E_2 &:= \{x \in B_r(o) : \exists \gamma \in \text{Geo}(x) \text{ s.t } z = \gamma_{1/2}, \gamma_0 \in A, \gamma_0, z \in \mathcal{T}_{nb}(x), Q(\gamma_0) \in Q(B_{2r}(z)) \setminus B\}. \end{aligned}$$

Note that by the definition of E , the map Q is well defined over E and the definitions of E_1 and E_2 are well posed. In order to get the thesis it is enough to show that both $\mu(E_1) = 0$ and $\mu(E_2) = 0$; this will give $\mu(E_z) = 0$, for every $z \in B_{2r}(o) \cap \text{lp}(\mu)$, contradicting (5.3).

In order to prove $\mu(E_1) = 0$, observe that thanks to the non-branching property of \mathcal{T}_{nb} it holds

$$(5.6) \quad (X \times Q^{-1}(B)) \cap H(z) \cap \mathcal{T}_{nb}^{-1} = (P_1((X \times Q^{-1}(B)) \cap H(z) \cap \mathcal{T}_{nb}^{-1}) \times X) \cap H(z) \cap \mathcal{T}_{nb}^{-1},$$

where we have used the following notation: $\mathcal{T}_{nb}^{-1} := \{(z, x) \in X \times X : (x, z) \in \mathcal{T}_{nb}\}$. Indeed it is easily checked that

$$\begin{aligned} &\{(x, y) \in X \times Q^{-1}(B) : (x, y) \in H(z) \cap \mathcal{T}_{nb}^{-1}\} \\ &= (\{x \in X : \exists y \in Q^{-1}(B), (x, y) \in H(z) \cap \mathcal{T}_{nb}^{-1}\} \times X) \cap H(z) \cap \mathcal{T}_{nb}^{-1}. \end{aligned}$$

The inclusion \subset is trivial, and the reverse inclusion follows directly by the non branching property of \mathcal{T}_{nb} . As the right hand side term can be rewritten as

$$(P_1((X \times Q^{-1}(B)) \cap H(z) \cap \mathcal{T}_{nb}^{-1}) \times X) \cap H(z) \cap \mathcal{T}_{nb}^{-1},$$

the identity (5.6) is proved.

Since $\mu(Q^{-1}(B)) = 0$, the definition (5.1) of inversion plan π^z implies that

$$\begin{aligned} 0 &= \pi^z((X \times Q^{-1}(B)) \cap H(z) \cap \mathcal{T}_{nb}^{-1}) \\ &= \pi^z((P_1((X \times Q^{-1}(B)) \cap H(z) \cap \mathcal{T}_{nb}^{-1}) \times X) \cap H(z) \cap \mathcal{T}_{nb}^{-1}) \\ &= \mu(P_1((X \times Q^{-1}(B)) \cap H(z) \cap \mathcal{T}_{nb}^{-1})). \end{aligned}$$

As $E_1 \subset P_1((X \times Q^{-1}(B)) \cap H(z) \cap \mathcal{T}_{nb}^{-1})$, we have that $\mu(E_1) = 0$.

The fact that $\mu(E_2) = 0$ follows easily by the disintegration (5.4). Indeed by (5.5), we infer $\mathcal{H}^1(\gamma^\alpha \cap A) = 0$ for every $\alpha \in Q(B_{2r}(z)) \setminus B$. Therefore, after the inversion with respect to z , we have

$$\mathcal{H}^1(\gamma^\alpha \cap E_2) = 0, \quad \forall \alpha \in Q(E_2).$$

Recalling that $\mu^\alpha \ll \mathcal{H}^1 \llcorner \gamma^\alpha$, for q -a.e. $\alpha \in Q(B_{2r}(z))$, the disintegration (5.4) then implies

$$\mu(E_2) = \int_{Q(E_2)} \mu^\alpha(E_2) q(d\alpha) = 0.$$

□

One can drop the hypothesis that \mathfrak{m} is concentrated on $\text{lp}(\mu)$, and still obtain the same claim at the price of adding regularity properties on the set of inversion points of the reference measure μ . We see two different cases; but, before that, let us give the following definition.

Definition 5.4. A set $C \subset X$ is said to be μ -connected if for any compact set $A \subset C$ there exists a set $U \subset C$ with $\mu(U) > 0$ and $\bar{t} \in (0, 1)$ so that

$$(A, U)_{\bar{t}} := \{\gamma_{\bar{t}} : \gamma \in \text{Geo}, \gamma_0 \in A, \gamma_1 \in U\} \subset C.$$

Let us mention that Definition 5.4 resembles the convexity property of regular points in Alexandrov spaces.

Theorem 5.5. *Let $(X, \mathbf{d}, \mathbf{m})$ be an essentially non-branching m.m.s. that verifies the qualitative MCP condition (4.1). Assume the existence of a reference measure μ for (Ω, \mathbf{d}) , where $\Omega = \text{spt}(\mathbf{m})$, so that $(\Omega, \mathbf{d}, \mu)$ verifies the qualitative MCP condition (4.1) and it is essentially non-branching. If $\text{lp}(\mu)$ is μ -connected, then*

$$\mathbf{m} \llcorner_{\text{lp}(\mu)} \ll \mu.$$

Proof. The proof follows the same lines of the proof of Theorem 5.3 and we only sketch the first part of it. Assume by contradiction the claim is false. Then there exists a compact set $A \subset \Omega \cap \text{lp}(\mu)$, that we can assume to be a subset of $B_r(o) \subset \Omega$ for some $o \in \Omega$ and $r > 0$, so that

$$\mathbf{m}(A) > 0 \quad \text{and} \quad \mu(A) = 0.$$

Since \mathbf{m} verifies the qualitative MCP, we have the following strict inequality

$$\int_U \mathbf{m}(A_{\bar{t}, x} \cap \mathcal{T}_{nb}(x)) \mu(dx) > 0,$$

where $U \subset \text{lp}(\mu)$ and $\bar{t} \in (0, 1)$ are given by the μ -connectedness of $\text{lp}(\mu)$. Using Fubini's Theorem to change order of integration, we get

$$0 < \int_{(A, U)_{\bar{t}}} \mu(\{x \in B_r(o) : \exists \gamma \in \text{Geo}(x) \text{ s.t. } z = \gamma_{\bar{t}}, \gamma_0 \in A, \gamma_0, z \in \mathcal{T}_{nb}(x)\}) \mathbf{m}(dz).$$

Since $(A, U)_{\bar{t}} \subset \text{lp}(\mu)$, the proof continues as in Theorem 5.3, obtaining a contradiction and therefore the claim. \square

We can also assume $\text{lp}(\mu)$ to be open and prove the next result.

Theorem 5.6. *Let $(X, \mathbf{d}, \mathbf{m})$ be an essentially non-branching m.m.s. that verifies the qualitative MCP condition (4.1). Assume the existence of a reference measure μ for (Ω, \mathbf{d}) , where $\Omega = \text{spt}(\mathbf{m})$, so that $(\Omega, \mathbf{d}, \mu)$ verifies the qualitative MCP condition (4.1) and it is essentially non-branching. If $\text{lp}(\mu)$ is open in Ω , then*

$$\mathbf{m} \llcorner_{\text{lp}(\mu)} \ll \mu.$$

Proof. Assume by contradiction the claim is false. Then there exists $A \subset \Omega \cap \text{lp}(\mu)$, that we can assume to be a subset of $B_R(o) \subset \Omega$ for some $o \in \Omega$ and $R > 0$, so that

$$\mathbf{m}(A) > 0 \quad \text{and} \quad \mu(A) = 0.$$

By inner regularity we can also assume A to be compact in Ω . As $\text{lp}(\mu)$ is open, for any $w \in A$ there exists r_w so that $B_{2r_w}(w) \subset \text{lp}(\mu)$. Then by compactness there exists $w \in A$ so that

$$\mathbf{m}(A \cap B_{r_w}(w)) > 0,$$

and therefore we can assume with no loss of generality that $A \subset B_{r_w}(w)$. Since \mathbf{m} verifies the qualitative MCP, we have the following strict inequality

$$\int_{B_{r_w}(w)} \mathbf{m}(A_{\frac{1}{2}, x} \cap \mathcal{T}_{nb}(x)) \mu(dx) > 0.$$

Using Fubini's Theorem to change order of integration, we get

$$0 < \int_{B_{2r_w}(w)} \mu(\{x \in B_r(o) : \exists \gamma \in \text{Geo}(x) \text{ s.t. } z = \gamma_{\frac{1}{2}}, \gamma_0 \in A, \gamma_0, z \in \mathcal{T}_{nb}(x)\}) \mathbf{m}(dz).$$

Since $B_{2r_w}(w) \subset \text{lp}(\mu)$, the proof continues as in Theorem 5.3, obtaining a contradiction and therefore the claim. \square

6. PROPERTIES OF REFERENCE MEASURES

In this section we study few natural properties of reference measures.

As one would expect, the property of being a reference measure is invariant under multiplication by a positive function.

Proposition 6.1 (Invariance). *Let (X, d, μ) be a m.m.s. with μ reference measure for (X, d) . Let $f : X \rightarrow \mathbb{R}$ be a Borel function such that $\mathfrak{m} := f \cdot \mu$ is still a Radon measure. If*

$$\mu(\{x \in X : f(x) = 0\}) = 0,$$

then \mathfrak{m} is a reference measure for (X, d) .

Proof. Consider the set of inversion points of μ , $\mathfrak{lp}(\mu)$, and observe that $\mathfrak{m}(X \setminus \mathfrak{lp}(\mu)) = 0$. For $z \in \mathfrak{lp}(\mu)$ let π^z be the associated inversion plan and define the rescaled plan

$$\hat{\pi}^z := f \circ P_1 \cdot \pi^z.$$

Since clearly $(P_1)_\# \hat{\pi}^z = \mathfrak{m}$, it only remains to prove that $(P_2)_\# \hat{\pi}^z \ll \mathfrak{m}$. To this aim let $A \subset X$ be a Borel set with $\mathfrak{m}(A) = 0$ and observe that, by the assumption on f , also $\mu(A) = 0$. By using that π^z is an inversion plan we infer

$$\pi^z(X \times A) = (P_2)_\# \pi^z(A) = 0.$$

Hence $(P_2)_\# \hat{\pi}^z(A) = f \circ P_1 \cdot \pi^z(X \times A) = 0$ and the claim follows. \square

The next property we would like to investigate is the locality, that means to obtain an inversion plan starting from a *local inversion plan*. Let us define the latter object:

Given a m.m.s. (X, d, μ) and $z \in X$ we say that $\pi^z \in \mathcal{M}_+(X \times X)$ is a *local inversion plan* for μ provided

$$(6.1) \quad (P_1)_\# \pi^z = \mu \llcorner_{B_r(z)}, \quad \pi^z(X \times X \setminus H(z)) = 0, \quad (P_2)_\# \pi^z \ll \mu,$$

for some $r > 0$. The set of those points admitting a local inversion plan will be denoted by $\mathfrak{lp}_{loc}(\mu)$ and will be called the set of points of local inversion.

Trivially restricting the inversion plan to the set $B_r(z) \times X$, any inversion point is a local inversion point, that is $\mathfrak{lp}_{loc}(\mu) \subset \mathfrak{lp}(\mu)$. Also the converse inclusion holds, provided (X, d) is non-branching.

Proposition 6.2 (Locality). *Let (X, d, μ) be a non-branching m.m.s. that verifies the qualitative MCP condition. Then*

$$\mathfrak{lp}_{loc}(\mu) = \mathfrak{lp}(\mu).$$

Proof. Consider $z \in \mathfrak{lp}_{loc}(\mu)$, the corresponding local inversion plan π^z and the corresponding open ball $B_r(z)$. We will now construct a measure $\eta \in \mathcal{M}_+(X \times X)$ such that

$$(P_1)_\# \eta = \mu \llcorner_{C_{r,2r}(z)}, \quad \eta(X \times X \setminus H(z)) = 0, \quad (P_1)_\# \eta \ll \mu,$$

where $C_{r,2r}(z) = B_{2r}(z) \setminus B_r(z)$. Once such a η is obtained, one can repeat the same argument for $C_{nr,(n+1)r}(z)$ and obtain the measure η^n , for any $n \in \mathbb{N}$ with $n \geq 2$. Then

$$\hat{\pi}^z := \pi^z + \sum_{n \geq 1} \eta^n,$$

will be an inversion plan.

Reasoning as in (5.4) we get

$$\mu_{\perp B_{2r}(z)} = \int_{Q(B_{2r}(z))} \mu^\alpha q(d\alpha), \quad \mu^\alpha \sim \mathcal{H}^1_{\perp \gamma^\alpha}, \quad q\text{-a.e. } \alpha \in Q(B_{2r}(z)),$$

where $q = Q_{\#}(\mu_{\perp B_{2r}(z)})$ is the quotient measure and γ^α is the maximal geodesic in $\text{Geo}(z)$ containing α . We can also assume each γ^α is parametrized with speed 1. From the non-branching assumption, each $\gamma^\alpha \cap B_r(z)$ is sent by π^z to a unique $\gamma^{\beta(\alpha)}$, that is

$$\pi^z \left(X \times X \setminus \{(x, y) \in B_r(z) \times X : x \in \gamma^\alpha, y \in \gamma^{\beta(\alpha)}\} \right) = 0, \quad \beta : Q \rightarrow Q,$$

with β μ -measurable map. Moreover from Proposition 4.6, for q -a.e. $\alpha \in Q$

$$\mathcal{L}^1 \left(\{s \in [r, 2r] : \gamma_r^\alpha \notin e_{(0,1)}R(z)\} \right) = 0.$$

In particular for q -a.e. $\alpha \in Q$

$$\sup\{\tau : (\gamma_{2r}^\alpha, \gamma_\tau^{\beta(\alpha)}) \in H(z)\} = \max\{\tau : (\gamma_{2r}^\alpha, \gamma_\tau^{\beta(\alpha)}) \in H(z)\} > 0.$$

Denote the maximal τ with τ_α and consider the map

$$T^\alpha : \{\gamma_s^\alpha : s \in [r, 2r]\} \rightarrow \{\gamma_s^{\beta(\alpha)} : s \in [\tau_\alpha/2, \tau_\alpha]\}, \quad T^\alpha(\gamma_{(1-\varepsilon)r+\varepsilon 2r}^\alpha) := \gamma_{(1-\varepsilon)\frac{\tau_\alpha}{2}+\varepsilon\tau_\alpha}^{\beta(\alpha)}.$$

Then the map $T : C_{r,2r}(z) \rightarrow X$, defined by $T(x) = T^\alpha(x)$ for $x \in \gamma^\alpha$, is μ -measurable and satisfies

$$(x, T(x)) \subset H(z),$$

for μ -a.e. $x \in C_{r,2r}(z)$. Now define $\eta := (id, T)_{\#} \mu_{\perp C_{r,2r}(z)}$ and observe that in order to have the claim it only remains to show that $(P_2)_{\#} \eta \ll \mu$.

To this aim consider a Borel set $A \subset X$ such that $\mu(A) = 0$. The claim is equivalent to prove that $\mu(T^{-1}(A)) = 0$. Consider the set $Q(A)$ and write it as the union of

$$Q_1(A) := \{\alpha \in Q(A) : \mu^\alpha(A) > 0\}, \quad Q_2(A) := \{\alpha \in Q(A) : \mu^\alpha(A) = 0\}.$$

For each $\alpha \in Q_2(A)$, since $\mu^\alpha \sim \mathcal{H}^1_{\perp \gamma^\alpha}$,

$$\mu^{\beta^{-1}(\alpha)} \left(T^{\beta^{-1}(\alpha)}(A) \right) = 0.$$

Moreover, since $\mu(A) = 0$, it follows that $q(Q_1(A)) = 0$. Then since β was also associated with the local inversion plan, $q(\beta^{-1}(Q_1(A))) = 0$. Hence $\mu(T^{-1}(A)) = 0$ and the claim follows. \square

Remark 6.3. It seems to be a challenging problem to weaken the non-branching assumption in Proposition 6.2 with the essentially non-branching one. Regarding this issue, a relevant role is played by the following phenomenon.

Consider a point $z \in X$ of local inversion for μ and $\gamma \in \text{Geo}(z)$, i.e. $\gamma_1 = z$. Suppose that γ can be extended through z with a geodesic η^1 of length $\varepsilon > 0$ with $\eta_0^1 = z$, so that

$$d(\gamma_0, \eta_1^1) = d(\gamma_0, \gamma_1) + d(\gamma_1, \eta_1^1).$$

Now consider the constant speed reparametrization $\hat{\gamma}$ of γ restricted to $[1/2, 1]$. With no violation of essentially non-branching, it could happen that $\hat{\gamma}$ admits also a different extension through z , that is there exists a geodesic $\eta^2 \neq \eta^1$ again such that $\eta_0^2 = z$ and

$$d(\hat{\gamma}_0, \eta_1^2) = d(\hat{\gamma}_0, \hat{\gamma}_1) + d(\hat{\gamma}_1, \eta_1^2).$$

Clearly also η^1 is a good extension of $\hat{\gamma}$.

Since z is a point of local inversion, we know that the local (multivalued) inverting map sends sets of positive μ -measure to sets of positive μ -measure. It may be the case that it sends $\hat{\gamma}_0$ to η_1^2 but γ_0 is forced to be moved to a point belonging to η^1 , if for instance

$$d(\gamma_0, \eta_1^2) > d(\gamma_0, z) + d(z, \eta_1^2).$$

In this situation no information can be obtained from the existence of the local inversion plan. Notice that the non-branching assumption indeed forces η^1 to coincide with η^2 .

6.1. Stability. Only for this section the reference measures will be assumed to be a probability measure.

We consider a sequence of pointed metric measures spaces $(X_k, d_k, \mu_k, \bar{x}_k)$ with μ_k *almost uniform* reference measure for (X_k, d_k) . We will show that if $(X_k, d_k, \mu_k, \bar{x}_k)$ converges to (X, d, μ, \bar{x}) in the pointed measured Gromov-Hausdorff topology, then μ is an *almost uniform* reference measure for (X, d) .

Definition 6.4. Let (X, d, μ) be a m.m.s. with $\mu \in \mathcal{P}(X)$ reference measure for (X, d) .

- We say that μ is an *almost uniform* reference measure provided that for μ -a.e. $z \in X$ there exist a constant $C_z > 0$ and an inversion plan π^z satisfying

$$(6.2) \quad (P_2)_\# \pi^z \leq C_z \mu.$$

- We say that μ is a *uniform* reference measure provided that there exists a constant $C > 0$ such that for μ -a.e. $z \in X$ there exists an inversion plan π^z satisfying (6.2) with $C_z \equiv C$.

Let us briefly recall the notion of pointed measured Gromov-Hausdorff convergence. A map $f : (X, d_X) \rightarrow (Y, d_Y)$ between metric spaces is an ε -isometry if and only if:

- it almost preserves distances: for all $z, w \in X$,

$$|d_X(z, w) - d_Y(f(z), f(w))| \leq \varepsilon;$$

- it is almost surjective:

$$\forall y \in Y, \exists x \in X \text{ such that } d_Y(f(x), y) \leq \varepsilon.$$

The following is a well-known notion of convergence that we will use for pointed m.m.s..

Definition 6.5. Let $(X_k, d_k, \mu_k, \bar{x}_k), (X, d, \mu, \bar{x})$ be pointed m.m.s. with $\mu_k \in \mathcal{P}(X_k)$, for all $k \in \mathbb{N}$, and $\mu \in \mathcal{P}(X)$. We say that $(X_k, d_k, \mu_k, \bar{x}_k) \rightarrow (X, d, \mu, \bar{x})$ in the *pointed measured Gromov-Hausdorff* topology, (*pmGH* for short) provided there exist sequences $R_k \rightarrow +\infty$ and $\varepsilon_k \rightarrow 0$ and Borel maps $f_k : X_k \rightarrow X$ so that

- i) $f_k(\bar{x}_k) = \bar{x}$;
- ii) $\sup \{ |d_k(x, y) - d(f_k(x), f_k(y))| : x, y \in B_{R_k}(\bar{x}_k) \} \leq \varepsilon_k$;
- iii) the ε_k -neighborhood of $f_k(B_{R_k}(x_k))$ contains $B_{R_k - \varepsilon_k}(\bar{x})$;
- iv) the sequence of probability measures $(f_k)_\# \mu_k$ converges to μ narrowly; i.e calling $C_b(X)$ the space of bounded continuous functions on X it holds

$$\lim_{k \rightarrow \infty} \int_X \varphi d[(f_k)_\# \mu_k] = \int_X \varphi d\mu, \quad \forall \varphi \in C_b(X).$$

In Definition 6.5 one can consider also sequences of pointed m.m.s. with measures not necessarily of total mass one. In the case of measures with possibly infinite total mass, i.e. Radon measures, one asks the weak convergence of *iv)* to hold for any $f \in C_b(X)$ with bounded support. We can state the stability result. Recall that during this section we will always assume all the metric measure spaces to be geodesic, proper, complete and separable.

Theorem 6.6. *Let $(X_k, \mathbf{d}_k, \mu_k, \bar{x}_k)$ be a sequence of pointed m.m.s. which converges in pmGH-sense to a limit pointed m.m.s. $(X, \mathbf{d}, \mu, \bar{x})$. Assume that, for infinitely many k , μ_k is an almost uniform reference measure for X_k and that for μ -a.e. $z \in X$ there exists a sequence $\{z_k\}_{k \in \mathbb{N}}$ with $z_k \in X_k$, such that*

$$(6.3) \quad \mathbf{d}(z, f_k(z_k)) \rightarrow 0 \quad \text{and} \quad \liminf_{k \in \mathbb{N}} C_{z_k} < \infty.$$

Then μ is an almost uniform reference measure for (X, \mathbf{d}) .

Proof. Let $z \in \text{spt}(\mu)$ be as in the assumption and consider a sequence of points $z_k \in X_k$ so that $\mathbf{d}(z, f_k(z_k)) \rightarrow 0$ and, for infinitely many k 's, z_k are inversion points for μ_k .

Up to extracting a suitable subsequence, for each $k \in \mathbb{N}$ we can consider the inversion plan $\pi^k \in \mathcal{P}(X_k \times X_k)$ around z_k ; moreover, by assumption, there exists $C = C_z$ such that

$$(P_1)_\# \pi^k = \mu_k, \quad (P_2)_\# \pi^k \leq C\mu_k, \quad \text{spt}(\pi^k) \subset H_k(z_k).$$

It is natural to consider the push-forward of the inversion plan via the ε_k -isometries as follows:

$$\eta^k := (f_k, f_k)_\# \pi^k \in \mathcal{P}(X \times X).$$

Since by assumption $(f_k)_\# \mu_k \rightarrow \mu$, it follows that both $(P_1)_\# \eta^k = (f_k)_\# \mu_k$ and $(P_2)_\# \eta^k \leq C \cdot (f_k)_\# \mu_k$ are tight. This in turn implies that $\eta^k \in \mathcal{P}(X \times X)$ are tight and then, by Prokhorov Theorem, they converge narrowly to some $\eta \in \mathcal{P}(X \times X)$, up to subsequences:

$$\int_{X \times X} \psi(x, y) d\eta_k \rightarrow \int_{X \times X} \psi(x, y) d\eta, \quad \text{as } k \rightarrow \infty, \quad \forall \psi \in C_b(X \times X).$$

In the rest of the proof we show that η is an inversion plan for μ at z .

Claim 1: $(P_1)_\# \eta = \mu$ and $(P_2)_\# \eta \leq C\mu$.

First of all observe that for $\varphi \in C_b(X)$ one has $\varphi \circ P_i \in C_b(X \times X)$, for $i = 1, 2$. It follows that, for every $\varphi \in C_b(X)$, it holds

$$\begin{aligned} \int_X \varphi d\mu &= \lim_{k \rightarrow \infty} \int_X \varphi d[(f_k)_\# \mu_k] = \lim_{k \rightarrow \infty} \int_X \varphi d[(f_k)_\# (P_1)_\# \pi_k] \\ &= \lim_{k \rightarrow \infty} \int_{X \times X} (\varphi \circ P_1) d\eta_k = \int_{X \times X} (\varphi \circ P_1) d\eta = \int_X \varphi d[(P_1)_\# \eta]. \end{aligned}$$

Since $\varphi \in C_b(X)$ is arbitrary, we infer that $(P_1)_\# \eta = \mu$. By analogous computations, using this time that $(P_2)_\# (\pi_k) \leq C\mu_k$, we get that, for every $\varphi \in C_b(X)$ with $\varphi \geq 0$, it holds

$$\int_X \varphi d[(P_2)_\# \eta] = \lim_{k \rightarrow \infty} \int_{X \times X} (\varphi \circ P_2) d\eta_k = \lim_{k \rightarrow \infty} \int_X \varphi d[(f_k)_\# (P_2)_\# \pi_k] \leq C \int_X \varphi d\mu.$$

Since $\varphi \in C_b(X)$ with $\varphi \geq 0$ is arbitrary, we infer that $(P_2)_\# \eta \leq C\mu$ and thus the proof of claim 1 is complete.

Claim 2: $\text{spt}(\eta) \subset H(z)$.

Fix a point $y \in X \times X$ and an increasing sequence of real numbers $R_i \rightarrow \infty$ so that

$$\eta(\partial B_{R_i}(y)) = 0, \quad \forall i, k \in \mathbb{N},$$

where the ball is taken in $X \times X$. Thus for each $i \in \mathbb{N}$ it holds

$$\lim_{k \rightarrow \infty} \eta^k(B_{R_i}(y)) = \eta(B_{R_i}(y)) > 0.$$

Consider then, for each $i \in \mathbb{N}$, the following probability measures:

$$\xi^k := (\eta^k(B_{R_i}(y)))^{-1} \eta^k \llcorner_{B_{R_i}(y)}, \quad \xi := (\eta(B_{R_i}(y)))^{-1} \eta \llcorner_{B_{R_i}(y)},$$

and observe that $\xi^k \rightharpoonup \xi$ narrowly. For ease of notation we have ignored the dependence on i . Thanks to the inner regularity of probability measures, for each $k \in \mathbb{N}$ there exists a compact set $C_k \subset B_{R_i}(y)$ so that

$$\xi^k(C_k) \geq 1 - \frac{1}{k}, \quad C_k \subset (f_k, f_k)(H_k(z_k)) \cap B_{R_i}(y).$$

Up to subsequences, there exists $C \subset \bar{B}_{R_i}(y)$ so that $C_k \rightarrow C$ in the Hausdorff distance. Therefore for any $\delta > 0$ there exists $k_\delta \in \mathbb{N}$ so that $C_k \subset C^\delta$, for all $k \geq k_\delta$ where C^δ denotes the closed tubular neighborhood of C of radius δ . Using upper-semicontinuity over compact sets of weakly converging measures, we infer

$$\xi(C^\delta) \geq \limsup_{k \rightarrow \infty} \xi^k(C^\delta) \geq \limsup_{k \rightarrow \infty} \xi^k(C_k) = 1.$$

Sending $\delta \rightarrow 0$, one obtains $\xi(C) = 1$. It remains to show that $C \subset H(z)$. As $C_k \rightarrow C$, for each $(x, y) \in C$ there exists a sequence $\{(x_k, y_k)\}_{k \in \mathbb{N}}$ with

$$(x_k, y_k) \in C_k \subset (f_k, f_k)(H_k(z_k)) \cap B_{R_i}(y), \quad (x_k, y_k) \rightarrow (x, y).$$

Hence $(x_k, y_k) = (f_k(u_k), f_k(w_k))$, with $(u_k, w_k) \in H_k(z_k)$. We can deduce from *ii*) and *iii*) of Definition 6.5 that

$$\begin{aligned} \mathbf{d}(x, y) - \mathbf{d}(x, z) - \mathbf{d}(z, y) &= \lim_{k \rightarrow \infty} \mathbf{d}(x_k, y_k) - \mathbf{d}(x_k, f_k(z_k)) - \mathbf{d}(f_k(z_k), y_k) \\ &= \lim_{k \rightarrow \infty} \mathbf{d}_k(u_k, w_k) - \mathbf{d}_k(u_k, z_k) - \mathbf{d}_k(z_k, w_k) \\ &= 0. \end{aligned}$$

It follows that $(x, y) \in H(z)$ and therefore $\xi(H(z)) = 1$. Hence $\text{spt}(\eta) \cap B_{R_i}(y) \subset H(z)$. Letting $R_i \rightarrow \infty$ yields $\text{spt}(\eta) \subset H(z)$, as desired. \square

The first part of the next corollary follows directly from the statement of Theorem 6.6; the second claim is instead a consequence of the proof above.

Corollary 6.7. *• Let $(X_k, \mathbf{d}_k, \mu_k, \bar{x}_k)$ be a sequence of pointed m.m.s. with $\mu_k \in \mathcal{P}(X_k)$ uniform reference measure for (X_k, \mathbf{d}_k) with the constant in (6.2) uniform in $k \in \mathbb{N}$. Assume the existence of a pointed m.m.s. $(X, \mathbf{d}, \mu, \bar{x})$ so that $(X_k, \mathbf{d}_k, \mu_k, \bar{x}_k) \rightarrow (X, \mathbf{d}, \mu, \bar{x})$ in the pmGH-sense, where $\mu \in \mathcal{P}(X)$. Then μ is a uniform reference measure for (X, \mathbf{d}) .*

• If μ is a uniform reference measure for (X, \mathbf{d}) then every point $z \in \text{spt}(\mu)$ is an inversion point.

Remark 6.8. It is a challenging open problem to replace in Theorem 6.6 the assumption that μ_k are almost uniform reference measures satisfying (6.3) by a uniform lower bound on the Ricci curvature of $(X_k, \mathbf{d}_k, \mu_k)$ (in MCP or CD or RCD sense). This would have as remarkable consequence that the limit measure on a Ricci limit space is unique up to multiplication by an L^1 function (note that this last fact was already established by Cheeger-Colding [15] via a completely different argument).

6.2. Uniqueness of the limit. In order to state and prove the uniqueness result, let us introduce a last small piece of notation: a sequence of pointed m.m.s. $(X_k, \mathbf{d}_k, \mathbf{m}_k, \bar{x}_k)$ is said to *uniformly* satisfy the qualitative MCP condition if there exist a sequence of radii $R_j \uparrow +\infty$ and continuous functions $f_j : [0, 1] \rightarrow \mathbb{R}^+$ with $f_j(0) = 1$, $f_j(1) = 0$ and $f_j(t) > 0$ for any $t \in [0, 1]$, $j \in \mathbb{N}$, such that the qualitative MCP condition (4.1) holds for every $(X_k, \mathbf{d}_k, \mathbf{m}_k, \bar{x}_k)$, $k \in \mathbb{N}$, with these fixed sequence $\{R_j, f_j\}_{j \in \mathbb{N}}$.

Theorem 6.9. *Let $(X_k, \mathbf{d}_k, \mu_k, \bar{x}_k)$ be a sequence of essentially non-branching pointed m.m.s. that uniformly verify the qualitative MCP condition in the above sense. Assume also that, for every $k \in \mathbb{N}$, $\mu_k \in \mathcal{P}(X_k)$ is a uniform reference measure for (X_k, \mathbf{d}_k) , with the constant in (6.2) uniform in $k \in \mathbb{N}$.*

Let $(X, \mathbf{d}, \mu, \bar{x})$ and $(X, \mathbf{d}, \xi, \bar{y})$ be two limit points in the pointed measured Gromov Hausdorff convergence of the two sequences $(X_k, \mathbf{d}_k, \mu_k, \bar{x}_k)$ and $(X_k, \mathbf{d}_k, \mu_k, \bar{y}_k)$, respectively. If (X, \mathbf{d}, μ) and (X, \mathbf{d}, ξ) are both essentially non-branching and

$$\text{spt}(\xi) = \text{spt}(\mu),$$

then $\mu \sim \xi$, i.e. $\mu \ll \xi$ and $\xi \ll \mu$.

Proof. The proof of the stability of the uniform qualitative MCP condition is completely analogous to the proof of the stability of $\text{MCP}(K, N)$, see [34, Theorem 6.8]. It follows that both (X, \mathbf{d}, μ) and (X, \mathbf{d}, ξ) verify the qualitative MCP and both are proper. Hence if $\Omega = \text{spt}(\xi) = \text{spt}(\mu)$, then by Corollary 6.7 both μ and ξ are uniform reference measures for (Ω, \mathbf{d}) , in particular $\text{lp}(\mu) = \text{lp}(\xi) = \Omega$. Then by Theorem 5.3 it follows that $\xi \ll \mu$ and $\mu \ll \xi$, i.e. $\xi \sim \mu$. \square

7. INVERSION OVER ALEXANDROV SPACES WITH CURVATURE BOUNDED FROM BELOW

The goal of this section is to show that, given a compact n -dimensional Alexandrov space (X, \mathbf{d}) , the n -dimensional Hausdorff measure \mathcal{H}^n is a reference measure. The idea is to combine the Lipschitz regularity of the exponential map together with the fact that it maps sets of positive measure to sets of positive measure (see Lemma 7.2), and then use Disintegration Theorem in order to construct a “local” inversion plan. Suitably iterating this construction one can obtain a “global” inversion plan and prove Theorem 7.3.

We first recall few definitions and notations that will be needed only during this section. Let (X, \mathbf{d}) be an Alexandrov space with curvature bounded below by $k \in \mathbb{R}$, that is (X, \mathbf{d}) is a complete, locally compact, geodesic space of curvature $\geq -k$ and of Hausdorff dimension $n < \infty$.

7.1. Exponential map. Fix any $p \in X$ and consider the set of segments emanating from p

$$\tilde{W}_p := \{pq : q \in X\},$$

where $pq : [0, 1] \rightarrow X$ with $pq \in \text{Geo}$, $pq_0 = p$ and $pq_1 = q$. Denote then with $\tilde{\Sigma}_p$ the set of equivalence classes of minimal segments emanating from p where pq is equivalent to pr if and only if one of pq and pr is contained in the other. The space $\tilde{\Sigma}_p$ has distance naturally induced by the angle \angle between minimal segments from p (for the definition of \angle see for instance [8] or [35]). The completion of $\tilde{\Sigma}_p$ is the *space of directions* and is denoted with Σ_p . For any minimal segment pq in X , the symbol v_{pq} denotes the direction at p corresponding to pq . Then for any pq and pr , it holds

$$\mathbf{d}_{\Sigma_p}(v_{pq}, v_{pr}) := \angle qpr.$$

For ease of notation we also denote \mathbf{d}_{Σ_p} with \angle . We say that p is a *regular point* if $(\Sigma_p, \mathbf{d}_{\Sigma_p})$ is isometric to the standard $(n-1)$ sphere, S^{n-1} .

The *tangent cone* K_p is obtained from $[0, \infty) \times \Sigma_p$ by identifying together all elements of the form $(0, u_0)$, with $u_0 \in \Sigma_p$. Its elements are denoted by tu_0 for $t \geq 0$ and $u_0 \in \Sigma_p$. The natural cone distance on K_p is given by

$$\mathbf{d}_{K_p}(tu_0, sv_0) := \sqrt{t^2 + s^2 - 2st \cos \angle u_0 v_0}.$$

Equivalently, a point $p \in X$ is regular if and only if K_p is isometric to \mathbb{R}^n . By considering $\tilde{W}_p \subset K_p$ one can define the exponential map at p as follows

$$\text{Exp}_p : \tilde{W}_p \rightarrow X, \quad \text{Exp}_p(|pq|v_{pq}) := q.$$

By Toponogov convexity it follows that Exp is Lipschitz on balls: there exists $L = L(k) > 0$ such that

$$(7.1) \quad \mathbf{d}(\text{Exp}_p(v), \text{Exp}_p(u)) \leq L \mathbf{d}_{K_p}(v, u), \quad \forall u, v \in B_1^{K_p}(0),$$

with $L = 1$ when $k = 0$. Denote now with $D(\text{Exp}_p)$ the identification of \tilde{W}_p in $K_p \equiv \mathbb{R}^n$.

Lemma 7.1. *Let $p \in X$ be a regular point. Then*

$$(7.2) \quad \mathcal{L}^n \left(B_1^{K_p}(0) \setminus D(\text{Exp}_p) \right) = 0.$$

Proof. Consider the cut locus at p , $C(p)$, see (4.6) for its definition. One can prove that $\mathcal{H}^n(C(p)) = 0$. Denote then with $E := X \setminus C(p)$ and define

$$\log_p : E \rightarrow D(\text{Exp}_p) \subset K_p,$$

so that $\text{Exp}_p \circ \log_p \llcorner E = \text{Id} \llcorner E$. As geodesics do not branch, the definition is well-posed. Moreover, recalling (7.1), for any $x, y \in E$ it trivially holds

$$\mathbf{d}(x, y) \leq L \mathbf{d}_{K_p}(\log_p(x), \log_p(y)),$$

with L depending on k and n , such that $L \rightarrow 1$ as $k \rightarrow 0$. For any $r \in (0, 1)$ consider X_r , the Alexandrov space X endowed with the rescaled distance $r^{-1}\mathbf{d}$; note that $B_1^{X_r}(p) := B_r(p)$ and that

$$\mathcal{H}_{X_r}^n = \frac{1}{r^n} \mathcal{H}^n, \quad \text{curv}(X_r) \geq r^2 k.$$

Then we have:

$$\begin{aligned} \frac{1}{r^n} \mathcal{H}^n(B_r(p)) &= \frac{1}{r^n} \mathcal{H}^n(E \cap B_r(p)) = \frac{1}{r^n} \mathcal{H}^n(E \cap B_1^{X_r}(p)) \\ &= \mathcal{H}_{X_r}^n(E \cap B_1^{X_r}(p)) \\ &\leq L_r^n \cdot \mathcal{L}^n(\log_p(E) \cap B_1(0)) \\ &\leq L_r^n \cdot \mathcal{L}^n(D(\text{Exp}_p) \cap B_1(0)). \end{aligned}$$

Since p is a regular point, $r^{-n} \mathcal{H}^n(B_r(p)) \rightarrow \omega_n$, where ω_n is the volume of the n -dimensional Euclidean unit ball. Since, as $r \rightarrow 0$ we have that $\text{curv}(X_r) \rightarrow 0$, we get that $L_r^n \rightarrow 1$ and the claim follows. \square

We also need the following crucial property of the Exponential map. Since we don't have a reference for it, we include our proof.

Lemma 7.2. *Let (X, \mathbf{d}) be an Alexandrov space with curvature bounded from below and dimension n . Fix $p \in X$ a regular point. Then for any $A \subset K_p$ of positive n -dimensional Lebesgue measure, it holds*

$$\mathcal{H}^n(\text{Exp}_p(A)) > 0.$$

Proof. We present here only the proof in the case of non-negative curvature.

Step 1. Suppose by contradiction the claim is false and pick $A \subset K_p$ contradicting the claim. For ease of notation we identify K_p with \mathbb{R}^n and for any $v \in S^{n-1}$ we denote the half line in the direction v , that is the set $\{tv : t \geq 0\}$, with $\text{Span}_+\{v\}$. Without loss of generality we can also assume that

$$\mathcal{L}^1(A \cap \text{Span}_+\{v\}) > 0,$$

for all those $v \in S^{n-1}$ so that $A \cap \text{Span}_+\{v\} \neq \emptyset$.

Observing that on each half line $\text{Span}_+\{v\}$ the map Exp_p is an isometry, and from Disintegration Theorem

$$0 = \mathcal{H}^n(\text{Exp}_p(A)) = \int_Q (h_\alpha \mathcal{H}^1)(\text{Exp}_p(A) \cap \gamma^\alpha) q(d\alpha),$$

where q is the quotient measure of \mathcal{H}^n associated to the rays decomposition and $h_\alpha \mathcal{H}^1$ the conditional measures associated to the disintegration, we obtain that

$$q(\{\alpha \in Q : \text{Exp}_p(A) \cap \gamma^\alpha \neq \emptyset\}) = 0.$$

This implies that we can consider the following set:

$$\text{Span}(A) := \{t \cdot v : t \in [0, 1], v \in S^{n-1} \text{ s.t. } \exists s > 0, s \cdot v \in A\} \subset \mathbb{R}^n,$$

and still obtain that

$$\mathcal{L}^n(\text{Span}(A)) > 0, \quad \mathcal{H}^n(\text{Exp}_p(\text{Span}(A))) = 0.$$

Step 2. We now obtain a contradiction using the non-expanding property of Exp_p . Since p is a regular point the volume of the $B_r^X(p)$ has the same infinitesimal behavior of r^n . So we have

$$\begin{aligned} 1 &= \lim_{r \rightarrow 0} \frac{\mathcal{H}^n(B_r^X(p))}{\omega_n r^n} = \lim_{r \rightarrow 0} \frac{\mathcal{H}^n(B_r^X(p) \setminus \text{Exp}_p(\text{Span}(A)))}{\omega_n r^n} \\ &\leq \lim_{r \rightarrow 0} \frac{\mathcal{L}^n(B_r(0) \setminus \text{Span}(A))}{\omega_n r^n}, \end{aligned}$$

where for the last inequality we used Lemma 7.1, the non-expanding of Exp_p and ω_n is the volume of the unitary ball in \mathbb{R}^n . It follows that $0 \in \mathbb{R}^n$ is a point of density 0 for $\text{Span}(A)$. From the definition of $\text{Span}(A)$ we conclude that this is possible only if

$$\mathcal{H}^{n-1}(v \in S^{n-1} : s \cdot v \in A, s > 0) = 0.$$

Since the last identity implies that $\mathcal{L}^n(A) = 0$ we have a contradiction and the claim is proved. \square

7.2. Inversion plan. We now show that \mathcal{H}^n is a reference measure for an n -dimensional compact Alexandrov space (X, d) . More precisely we will construct an inversion plan at \mathcal{H}^n -a.e. regular point $p \in X$.

Theorem 7.3. *Let (X, d) be a compact Alexandrov space with curvature bounded from below by k and Hausdorff dimension $n \in \mathbb{N}$. Then \mathcal{H}^n is a reference measure for (X, d) . More precisely we show that an inversion plan exists at all regular point $p \in X$ so that (4.4) holds.*

Proof. Since (X, d) is compact and it is well known that (X, d, \mathcal{H}^n) is non-branching and verifies MCP(k, n). We will construct an inversion plan for any regular point p such that (7.3) holds true.

Step 1. From Proposition 4.6 it follows that for \mathcal{H}^n -a.e. $p \in X$

$$(7.3) \quad \mathcal{H}^n(e_{(0,1)}(R(p))) = \mathcal{H}^n(X),$$

where $R(p) = \{\gamma \in \text{Geo} : \gamma_t = p, \text{ for some } t \in (0, 1)\}$. So we fix once and for all a regular point $p \in X$ so that (7.3) holds true. For ease of notation we denote with R_p the set $e_{(0,1)}(R(p))$. We perform the disintegration of $\mathcal{H}^n \llcorner R_p$ as in Proposition 4.8:

$$\mathcal{H}^n \llcorner R_p = \int_S \eta^\alpha q(d\alpha), \quad \eta^\alpha(\gamma^\alpha) = 1, \quad q - a.e. \alpha \in S,$$

where $\gamma^\alpha \in \text{Geo}(p)$ is the unique geodesic in $\text{Geo}(p)$ through α , $S = Q(R_p)$ is the quotient set and Q the quotient map. Observe that convexity of R_p is not needed to apply Disintegration theorem, we only use that for each point in $z \in R_p$, any intermediate point between p and z belongs to R_p . The same property permits to use MCP to obtain the absolute continuity of the conditional measures with respect to the Hausdorff measure of dimension 1.

We can write R_p in the following way:

$$R_p = \bigcup_{m=2}^{\infty} e_{(0,1)}(\{\gamma \in \text{Geo} : \gamma_t = p, t \leq 1 - 1/m\}) = \bigcup_{m=2}^{\infty} e_{(0,1)}(R_m(p)) = \bigcup_{m=2}^{\infty} R_{p,m}.$$

Note that if $\gamma \in \text{Geo}(p)$ is so that $\gamma_0 \in R_{p,m}$, then $\gamma_s \in R_{p,m}$ for all $s \in [0, 1)$. Indeed $\gamma_0 \in R_{p,m}$ implies the existence of $\xi \in \text{Geo}$ so that

$$\xi_0 = \gamma_0, \quad \xi_t = p.$$

with $t \leq 1 - 1/m$. Then for any $\bar{s} \in (0, 1)$ there exists $\tau < t$ so that $\xi_\tau = \gamma_{\bar{s}}$. Consider $\sigma(\tau) = t + (1-t)(t-\tau)t^{-1}$ and define the new curve

$$\xi_s^\tau := \xi_{\tau(1-s) + \sigma(\tau)s}.$$

Being a subset of ξ , it follows that $\xi^\tau \in \text{Geo}$, moreover $\xi_0^\tau = \gamma_{\bar{s}}$ and $\xi_t^\tau = p$. Hence $\gamma_{\bar{s}} \in R_{p,m}$. It follows that for each $\alpha \in S$ either $\gamma_s^\alpha \notin R_{p,m}$ for all $s \in [0, 1)$ (but this will never happen because of the definition of R_p) or there exists a minimal $s(\alpha) \in [0, 1)$ so that

$$\gamma_s^\alpha \in R_{p,m}, \quad \forall s \in [s(\alpha), 1), \quad \gamma_s^\alpha \notin R_{p,m}, \quad \forall s < s(\alpha).$$

As $R_{m,p}$ can be written as countable union of closed sets, $\alpha \mapsto s(\alpha)$ is measurable. Therefore we have the following representation:

$$\mathcal{H}^n \llcorner R_{p,m} = \int_S \eta_m^\alpha q_m(d\alpha), \quad \eta_m^\alpha(\gamma^{\alpha,m}) = 1, \quad q_m - a.e. \alpha \in S,$$

where $\gamma^{\alpha,m}$ is the reparametrization of γ^α such that $\gamma_0^{\alpha,m} = \gamma_{s(\alpha)}^\alpha$ and $\gamma_1^{\alpha,m} = p$. To keep notation simple, we will omit the dependence on m of the geodesic, so $\gamma^{\alpha,m} = \gamma^\alpha$.

Step 2. From the definition of $R_{p,m}$ and the non-branching property of (X, d) , for q_m -a.e. $\alpha \in S$ there exists a unique $\beta \in S$, that we will denote with $\beta(\alpha)$, such that

$$(7.4) \quad \{(\gamma_s^\alpha, \gamma_t^\beta) \in X \times X : s \in [0, 1], t \in I(\alpha)\} \subset H(p),$$

with $I(\alpha) := \left\{s \in [0, 1] : s \geq 1 - L(\gamma^\alpha) \left((m-1)L(\gamma^{\beta(\alpha)}) \right)^{-1}\right\}$, where L denotes the length of the geodesic. Indeed notice that for q_m -a.e. $\alpha \in S$ there exists, as before, $\xi \in \text{Geo}$ so that

$$\xi_0 = \gamma_0^\alpha, \quad \xi_t = p, \quad \xi_1 \in \gamma_{(0,1)}^{\beta(\alpha)}, \quad d(p, \xi_1) \geq \frac{1}{m-1}L(\gamma^\alpha)$$

and moreover $\{(\xi_s, \xi_\tau) : s \leq t \leq \tau\} \subset H(p)$. From the bound $t \leq 1 - 1/m$ it is immediate to check that if $s \in I(\alpha)$ then

$$\gamma_s^{\beta(\alpha)} \in \xi_{(t,1)}$$

and therefore (7.4) is proved. For the next step we find convenient the following abuse of notation: we denote with $\gamma^{\beta(\alpha)}$ the reparametrization of the whole geodesic $\gamma^{\beta(\alpha)} \in \text{Geo}(p)$ to the interval $I(\alpha)$.

Step 3. We construct the inversion plan. First of all observe that $R_{p,1} = \emptyset$; for any $m \geq 2$ and for q_m -a.e. $\alpha \in S$, we consider the coupling $\pi_m \in \mathcal{P}(X \times X)$ defined by

$$\pi_m := (\text{Id}, F_m)_\# \mathcal{H}^n \llcorner_{(R_{p,m} \setminus \bigcup_{l=1}^{m-1} R_{p,l})},$$

where

$$F_m : R_{p,m} \mapsto R_{p,m}, \quad F_m(\gamma_t^\alpha) := \gamma_{1-t}^{\beta(\alpha)}.$$

From Step 2, π_m is well defined and by construction we have

$$\pi_m(X \times X \setminus H(p)) = 0 \quad \text{and} \quad (P_1)_\# \pi_m = \mathcal{H}^n \llcorner_{(R_{p,m} \setminus \bigcup_{l=1}^{m-1} R_{p,l})}.$$

The fact that $(P_2)_\# \pi_m \ll \mathcal{H}^n$ follows easily from Lemma 7.2: indeed if by contradiction it was false, then there would exist a subset $A \subset R_{p,m}$ so that

$$(P_2)_\# \pi_m(A) > 0, \quad \mathcal{H}^n(A) = 0,$$

that can be restated as $\mathcal{H}^n(F_m^{-1}(A)) > 0$ and $\mathcal{H}^n(A) = 0$. But since Exp_p is locally Lipschitz we have on one hand $\mathcal{L}^n(\log_p(F_m^{-1}(A))) > 0$ and on the other hand, from Lemma 7.2, $\mathcal{L}^n(\log_p(A)) = 0$. Now we can map $\log_p(A)$ to $\log_p(F_m^{-1}(A))$ using the inversion around the origin composed with a rescaling on each line, therefore either both have Lebesgue measure 0 or Lebesgue measure strictly positive. Hence we have a contradiction and $(P_2)_\# \pi_m \ll \mathcal{H}^n$.

The desired inversion plan is then given by summation:

$$\pi := \sum_{m=2}^{\infty} \pi_m.$$

Indeed, since R_p has full measure by construction, we have

$$(P_1)_\#(\pi) = \sum_{m=2}^{\infty} \mathcal{H}^n \llcorner_{(R_{p,m} \setminus \bigcup_{l=1}^{m-1} R_{p,l})} = \mathcal{H}^n \llcorner_{\bigcup_{m=2}^{\infty} R_{p,m}} = \mathcal{H}^n \llcorner_{R_p} = \mathcal{H}^n;$$

it is also clear that $(P_2)_\#(\pi) \ll \mathcal{H}^n$ and $\pi(X \times X \setminus H(p)) = \sum_{m=2}^{\infty} \pi_m(X \times X \setminus H(p)) = 0$. \square

The following is a straightforward consequence of Theorem 5.3 and Theorem 7.3.

Corollary 7.4. *Let (X, \mathbf{d}) be a compact Alexandrov space with curvature bounded from below by k and with Hausdorff dimension $n \in \mathbb{N}$. Let \mathbf{m} be a Borel finite measure on X so that $(X, \mathbf{d}, \mathbf{m})$ verifies the qualitative MCP condition (4.1). If \mathbf{m} is concentrated on those regular points of X such that \mathcal{H}^n verifies (4.4), then*

$$\mathbf{m} \ll \mathcal{H}^n.$$

8. EXAMPLES AND APPLICATIONS

In this final section, we give "smooth" examples where the inversion plan exists at every point; as a consequence, the application of Theorem 5.3 will be very neat. We will consider the following examples: smooth Riemannian manifolds, Alexandrov spaces with bounded curvature, and the Heisenberg group endowed with the Carnot-Carathéodory distance and the Haar measure.

8.1. Example 1: Riemannian manifolds.

Proposition 8.1 (The volume measure is reference). *Let (M, g) be a complete (non necessarily compact but connected and without boundary) n -dimensional smooth Riemannian manifold and denote respectively with d_g, μ_g the Riemannian distance and the volume measure induced by g . Then μ_g is a reference measure for (M, d_g) , in the sense of Definition 5.1. More precisely, we show that there exists an inversion plan at every point, i.e. $\text{lp}(\mu_g) = M$.*

Proof. For every $z \in M$ we have to construct an inversion plan π^z satisfying (5.1).

Step 1. First of all we claim that there exist compact subsets $K_n \subset M$, $n \in \mathbb{N}$, satisfying the following:

- $\mu_g(M \setminus \bigcup_{n \in \mathbb{N}} K_n) = 0$;
- for every $x \in K_n$, there exists a unique minimizing geodesic $\gamma_{xz} : [0, 1] \rightarrow M$ from x to z and it is extendable to $[0, 1 + \frac{2}{n}]$ as minimizing geodesic;
- the map $\Phi_n : K_n \rightarrow M$ defined by $\Phi_n(x) := \gamma_{xz}(1 + \frac{1}{n})$ is bi-Lipschitz onto its image.

In order to prove the above properties, first of all recall that the cut locus of z has null measure, i.e. $\mu_g(C(z)) = 0$. Moreover for every $x \in M \setminus C(z)$ it is known that $z \notin C(x)$ (see for instance [26, Lemma 2.1.11]), so that for every $x \in M \setminus C(z)$ there exists a unique minimizing geodesic $\gamma_{xz} : [0, 1] \rightarrow M$ from x to z and it is extendable (still as minimizing geodesic) to a bigger interval $[0, 1 + \varepsilon]$ for some $\varepsilon = \varepsilon(x) > 0$. Therefore, called

$$(8.1) \quad \tilde{K}_n := \left\{ x \in M : \exists! \gamma_{xz} \in \text{Geo}, \gamma_{xz} : [0, 1] \rightarrow M \text{ geodesic from } x \text{ to } z, \right. \\ \left. \text{and it is extendable to } \left[0, 1 + \frac{2}{n}\right] \right\},$$

we get that $M \setminus C(z) = \bigcup_{n \in \mathbb{N}} \tilde{K}_n$, so $\mu_g(M \setminus \bigcup_{n \in \mathbb{N}} \tilde{K}_n) = 0$; notice also that $K_n \subset K_{n+1}$, for every $n \in \mathbb{N}$. By continuous dependence on initial data in the geodesic equation (see for instance the proofs of [26, Lemma 2.1.5, Proposition 2.1.10]), one has that $\tilde{K}_n \subset M$ is a closed subset. Hence, called

$$K_n := \tilde{K}_n \cap \overline{B_n(z)},$$

we get that $K_n \subset M$ is a compact subset for every $n \in \mathbb{N}$, and $\mu_g(M \setminus \bigcup_{n \in \mathbb{N}} K_n) = 0$. The second claim is clear from the definition (8.1) and the third claim follows by the compactness of K_n and the fact that the exponential map centered at z is a diffeomorphism once restricted to $M \setminus C(z)$.

Step 2 Define a map $\Phi : M \rightarrow M$ in the following way:

- if $x \in \bigcup_{n \in \mathbb{N}} K_n$, called $n_0 := \min\{n \in \mathbb{N} : x \in K_n\}$, set $\Phi(x) := \Phi_{n_0}(x)$;
- if $x \in M \setminus \bigcup_{n \in \mathbb{N}} K_n$, set $\Phi(x) := z$.

Now let $\pi^z := (Id, \Phi)_\#(\mu_g)$. Clearly we have $(P_1)_\#(\pi^z) = \mu_g$ and, by construction, π^z is concentrated on $H(z)$. Finally we also infer that $(P_2)_\#(\pi^z) \ll \mu_g$, indeed for every $E \subset M$ with $\mu_g(E) = 0$ we have

$$\begin{aligned} \left[(P_2)_\#(\pi^z) \right](E) &= \pi^z(P_2^{-1}(E)) = \mu_g(\Phi^{-1}(E)) = \mu_g\left(\Phi^{-1}(E) \cap \left(\bigcup_{n \in \mathbb{N}} K_n\right)\right) \\ &= \mu_g\left(\bigcup_{n \in \mathbb{N}} \Phi^{-1}(E) \cap (K_n \setminus \cup_{1 \leq j \leq n-1} K_j)\right) \\ &\leq \sum_{n \in \mathbb{N}} \mu_g(\Phi_n^{-1}(E)) = 0, \end{aligned}$$

where in the last equality we used that Φ_n are bi-Lipschitz by construction. We conclude that π^z is an inversion plan satisfying (5.1) hence, since $z \in M$ was arbitrary, μ_g is a reference measure. \square

Corollary 8.2 (MCP measures on complete Riemannian manifolds). *Let (M, g) be a (possibly non compact) complete n -dimensional smooth Riemannian manifold with Ricci curvature bounded from below:*

$$\text{Ric}_g \geq Kg, \quad \text{for some } K \in \mathbb{R}.$$

Let $\mathfrak{m} \in \mathcal{M}^+(M)$ be a positive Radon measure with $\text{spt}(\mathfrak{m}) = M$, such that $(M, \mathbf{d}_g, \mathfrak{m})$ verifies the qualitative MCP condition (4.1). Then, called μ_g the Riemannian volume measure of (M, g) , we have

$$(8.2) \quad \mathfrak{m} \ll \mu_g.$$

If moreover \mathfrak{m} is a reference measure for (M, \mathbf{d}_g) and $\mu_g(M \setminus \text{lp}(\mathfrak{m})) = 0$, then $\mathfrak{m} \sim \mu_g$.

Proof. By Proposition 8.1, we know that μ_g is a reference measure with $\text{lp}(\mu_g) = M$, moreover it is clear that (M, \mathbf{d}_g, μ_g) is a non-branching metric space satisfying the qualitative MCP (notice that, more precisely it satisfies the $\text{MCP}(K, n)$ condition). The first claim (8.2) is then a direct consequence of Theorem 5.3. If \mathfrak{m} is a reference measure, we can reverse the role of \mathfrak{m} and μ_g and conclude that, if $\mu_g(M \setminus \text{lp}(\mathfrak{m})) = 0$, then $\mu_g \ll \mathfrak{m}$ and hence $\mathfrak{m} \sim \mu_g$. \square

8.2. Example 2: Alexandrov spaces with bounded curvature.

Corollary 8.3 (MCP measures on Alexandrov spaces with curv. bounded above and below). *Let (X, \mathbf{d}) be an n -dimensional geodesically complete Alexandrov space with curvature bounded from above and below, and let $\mu := \mathcal{H}^n$ be the n -dimensional Hausdorff measure. Let $\mathfrak{m} \in \mathcal{M}^+(X)$ be a positive Radon measure with $\text{spt}(\mathfrak{m}) = X$, such that $(X, \mathbf{d}, \mathfrak{m})$ verifies the qualitative MCP condition (4.1). Then*

$$(8.3) \quad \mathfrak{m} \ll \mu.$$

Proof. Thanks to the structural properties of Alexandrov spaces, the proof is not too far from the Riemannian one above. Indeed it is well known that the curvature bound from below implies non-branching, so that trivially any measure is essentially non-branching. On the other hand, the geodesic completeness combined with the upper curvature bound ensures that every point $p \in X$ has a neighborhood U with the following properties (see for instance [8, Sections 9.1.3, 9.1.7]):

- the closure \bar{U} is compact with Lipschitz boundary ∂U ;
- for every $x, y \in U$ there exists a unique geodesic γ of X joining them which is contained in U , i.e. $\gamma(0) = x, \gamma(1) = y, \gamma([0, 1]) \subset U$; moreover such geodesic is extendable to a minimizing geodesic until it intersects ∂U ;
- every triple of points $xyz \in U$ satisfies the triangle comparison conditions with the two model spaces corresponding to the upper and lower curvature bounds.

In particular, since (U, \mathbf{d}) is itself an Alexandrov space with curvature bounded from below, it is non-branching and its n -dimensional Hausdorff measure $\mu \llcorner U$ satisfies the qualitative MCP condition (4.1) (more precisely it satisfies $\text{MCP}((n-1)K, n)$ where $K \in \mathbb{R}$ stands for the lower curvature bound, see [34, Proposition 2.8] which is based on the previous work [27]).

In order to apply Theorem 5.3 it is then enough to show that $\mu \llcorner U$ is a reference measure for (U, \mathbf{d}) in the sense of Definition 5.1. To this aim, for every $z \in U$, define the following geodesic inversion map $\Phi_z : U \rightarrow U$ with center z

$$(8.4) \quad \Phi_z(x) = \text{Exp}_z(-\alpha_z(x) \cdot \text{Exp}_z^{-1}(x)), \quad \forall x \in U,$$

where Exp_z is the exponential map of the Alexandrov space centered at z and

$$\alpha_z(x) := \frac{\sup\{t : Exp_z(-t Exp_z^{-1}(x)) \in U\}}{\sup\{t : Exp_z(t Exp_z^{-1}(x)) \in U\}},$$

notice that $\alpha : U \rightarrow (0, +\infty)$ is Lipschitz since U is geodesically convex and ∂U is Lipschitz. Hence, by comparison with geodesic triangles in the two model spaces corresponding to the upper and lower curvature bounds we get that the above geodesic inversion $\Phi_z : U \rightarrow U$ with center z is a bi-Lipschitz map. Therefore $(\Phi_z)_\#(\mu \llcorner U) \ll \mu \llcorner U$ and, setting

$$\pi^z := (Id|_U, \Phi_z)_\#(\mu \llcorner U),$$

we get that $\mu_g \llcorner U$ is a reference measure for (U, \mathbf{d}) . The thesis follows then by applying Theorem 5.3 locally to $\mu_g \llcorner U$ and $\mathbf{m} \llcorner U$, since U was a suitable neighborhood of an arbitrary point $p \in X$. \square

Remark 8.4. Let us remark that from the work of Nikolaev [33], it is known that a space with curvature bounded from above and below in the Alexandrov sense has the structure of a $C^{3,\alpha}$ -manifold with a $C^{1,\alpha}$ -metric tensor. But, since the geodesic equations involve first order derivatives of the metric which a priori may not be locally Lipschitz, this remarkable structural result is not so useful to our purpose, that is to show that there is bi-Lipschitz geodesic inversion at every point. This is the reason why we used the argument above using classical comparison, and why we believe that the notion of inversion plan is not trivial also for such spaces.

8.3. Example 3: the Heisenberg group. Let us start by recalling some basic facts about the n -dimensional Heisenberg group \mathbb{H}^n . Such space is a non-commutative stratified nilpotent Lie group; as a set it can and will be identified with its Lie algebra $\mathbb{R}^{2n+1} \simeq \mathbb{C}^n \times \mathbb{R}$. We denote a point in \mathbb{H}^n indifferently by $x = (\xi, \eta, t) = [\zeta, t]$, where $\xi = (\xi_1, \dots, \xi_n), \eta = (\eta_1, \dots, \eta_n) \in \mathbb{R}^n$, $t \in \mathbb{R}$ and $\zeta = \xi + i\eta \in \mathbb{C}^n$. The group law in this system of coordinates is given by

$$[\zeta, t] \cdot [\zeta', t'] = \left[\zeta + \zeta', t + t' + 2 \sum_{j=1}^n \text{Im} \zeta_j \bar{\zeta}'_j \right].$$

The center of the group is

$$(8.5) \quad L := \{[0, s] \in \mathbb{H}^n : s \in \mathbb{R}\};$$

define also L^*, L^+, L^- restricting s to belong to, respectively, $\mathbb{R} \setminus \{0\}, [0, +\infty), (-\infty, 0]$. The Haar measure of the group is given by the Lebesgue measure \mathcal{L}^{n+1} on $\mathbb{H}^n \simeq \mathbb{R}^{2n+1}$. The left invariant vector fields

$$(8.6) \quad X_j := \partial_{\xi_j} + 2\eta_j \partial_t, \quad Y_j := \partial_{\eta_j} - 2\xi_j \partial_t$$

Lie generate the Lie algebra of the group (which is \mathbb{R}^{2n+1}), the only nontrivial bracket relation being $[X_j, Y_j] = -4Z$, where $Z := \partial_t$. We endow \mathbb{H}^n with the standard Carnot-Carathéodory distance \mathbf{d}_c defined as follows. An absolutely continuous curve $\gamma : [0, T] \rightarrow \mathbb{R}^{2n+1}$ is said to be *horizontal* if there exist measurable functions $h_j : [0, T] \rightarrow \mathbb{R}$ such that

$$(8.7) \quad \dot{\gamma}(s) = \sum_{j=1}^n h_j(s) X_j(\gamma(s)) + h_{n+j}(s) Y_j(\gamma(s)).$$

We say that γ is sub-unit if moreover $\sum_{j=1}^{2n} h_j(s)^2 \leq 1$ for a.e. $s \in [0, T]$. Since $(X_1, \dots, X_n, Y_1, \dots, Y_n)$ Lie generate the Lie algebra of the group it follows from Chow's Theorem that any couple of points

x, y in \mathbb{H}^n can be joined by a sub-unit curve. Then the Carnot-Carathéodory distance d_c between x and y is defined as

$$(8.8) \quad d_c(x, y) := \inf\{T > 0 : \exists \text{ a sub-unit curve } \gamma : [0, T] \rightarrow \mathbb{H}^n \text{ such that } \gamma(0) = x, \gamma(T) = y\}.$$

It follows easily from the left invariance and homogeneity with respect to the dilations of the vector fields X_j and Y_j that d_c is a left invariant and homogeneous distance on \mathbb{H}^n . We say that a curve $\gamma : [0, T] \rightarrow \mathbb{H}^n$ is a *minimal geodesic with unit speed* if

$$d_c(\gamma(t), \gamma(t')) = |t - t'|, \quad \forall t, t' \in [0, T].$$

By the work of Pansu [40] on differentiability of Lipschitz functions between stratified and nilpotent Lie groups (see also [5, Theorem 3.2]), it follows that for any $x, y \in \mathbb{H}^n$ there exists a sub-unit (a fortiori unit speed) minimal geodesic joining x to y . The equations for the minimal geodesics can be explicitly computed and can be found in the literature (see for instance [20, 31, 32]), here we follow the presentation of [5] by recalling just the main results that we will need to prove Corollary 8.9.

Set $\mathbb{S} := \{a + ib \in \mathbb{C}^n : |a + ib| = 1\}$. For any $a + ib \in \mathbb{S}, v \in \mathbb{R}$ and $r > 0$, we say that a curve $\gamma : [0, r] \rightarrow \mathbb{H}^n$ is a *curve with parameter* $(a + ib, v, r)$, if $\gamma(s) = (\xi(s), \eta(s), t(s))$, where

$$(8.9) \quad \begin{aligned} \xi_j(s) &:= \frac{b_j(1 - \cos \frac{vs}{r}) + a_j \sin \frac{vs}{r}}{v} r, & \eta_j &:= \frac{-a_j(1 - \cos \frac{vs}{r}) + b_j \sin \frac{vs}{r}}{v} r, & t &= 2 \frac{v - \sin \frac{vs}{r}}{v^2} r^2, & \text{if } v \neq 0; \\ \xi_j(s) &:= a_j s, & \eta_j &:= b_j s, & t &\equiv 0, & \text{if } v = 0. \end{aligned}$$

As the next theorem states (for the proof see for instance [5, Theorem 3.4]), these are the only geodesics in \mathbb{H}^n starting from the origin.

Theorem 8.5 (Geodesics in \mathbb{H}^n). *Sub-unit minimal geodesics starting from the origin 0 are curves γ with parameter $(a + ib, v, r)$, for some $(a + ib, v, r) \in \mathbb{S} \times [-2\pi, 2\pi] \times (0, +\infty)$. In particular, curves γ with parameter $(a + ib, v, r) \in \mathbb{S} \times \mathbb{R} \times (0, +\infty)$ with $|v| > 2\pi$ are not minimal geodesics. Conversely, any curve γ with parameter $(a + ib, v, r) \in \mathbb{S} \times [-2\pi, 2\pi] \times (0, +\infty)$ is a sub-unit minimal geodesic starting from 0. More precisely the following holds:*

- For any $x = [0, t] \in L^*$ with $t > 0$, resp. $t < 0$, sub-unit minimal geodesics from 0 to x form a family of curves parametrized on \mathbb{S} , each one being obtained from one another by a rotation around the axis L . More precisely this family is composed by all curves γ with parameter $(a + ib, 2\pi, \sqrt{\pi t})$, resp. $(a + ib, -2\pi, \sqrt{\pi |t|})$, where $a + ib \in \mathbb{S}$.
- For any $x \in \mathbb{H}^n \setminus L$ there exists a unique $(a + ib, v, r) \in \mathbb{S} \times (-2\pi, 2\pi) \times (0, +\infty)$ so that the curve γ with parameter $(a + ib, v, r)$ joins 0 to x and in particular $\gamma : [0, r] \rightarrow \mathbb{H}^n$ is the unique sub-unit minimal geodesic from 0 to x . Moreover such γ is extendable to a minimizing geodesic (via the same formulas (8.9)) defined on the maximal interval $[0, 2\pi r/|v|] \supset [0, r]$.

Consider now the endpoint map $\Phi : \mathbb{S} \times [-2\pi, 2\pi] \times (0, +\infty) \rightarrow \mathbb{H}^n$, $(a + ib, v, r) \mapsto (\xi, \eta, t)$, which associates to each geodesic γ with parameter $(a + ib, v, r)$ its endpoint (ξ, η, t) given by

$$(8.10) \quad \begin{aligned} \xi_j &:= \frac{b_j(1 - \cos v) + a_j \sin v}{v} r, & \eta_j &:= \frac{-a_j(1 - \cos v) + b_j \sin v}{v} r, & t &= 2 \frac{v - \sin v}{v^2} r^2, & \text{if } v \neq 0; \\ \xi_j &:= a_j r, & \eta_j &:= b_j r, & t &\equiv 0, & \text{if } v = 0. \end{aligned}$$

Theorem 8.5 directly implies that

- the range of Φ is \mathbb{H}^n ;
- $d_c(0, \Phi(a + ib, v, r)) = r$, for all $(a + ib, v, r) \in \mathbb{S} \times [-2\pi, 2\pi] \times (0, +\infty)$;

- the map Φ is bijective from D to $\mathbb{H}^n \setminus L$, where

$$(8.11) \quad D := \mathbb{S} \times (-2\pi, 2\pi) \times (0, +\infty);$$

- the map Φ is not injective on $\{(a + ib, v, r) : a + ib \in \mathbb{S}, |v| = 2\pi, r > 0\}$ whose image corresponds to the center L of the group. More precisely

$$\begin{aligned} L^+ \setminus \{0\} &= \{\Phi(a + ib, v, r) \in \mathbb{H}^n : a + ib \in \mathbb{S}, v = 2\pi, r > 0\}; \\ L^- \setminus \{0\} &= \{\Phi(a + ib, v, r) \in \mathbb{H}^n : a + ib \in \mathbb{S}, v = -2\pi, r > 0\}. \end{aligned}$$

Moreover, an explicit computation of the Jacobian of Φ (see [32, Page 161] or [25, Propositions 1.7, 1.12]) shows that it does not vanish on D . Therefore we have the next

Proposition 8.6. *The map Φ is a diffeomorphism from D onto $\mathbb{H}^n \setminus L$.*

The next lemma of linear algebra will be useful in the sequel.

Lemma 8.7. *Let $v \neq 0$ and consider the linear map $L_v : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by*

$$(x_1, x_2) \mapsto L_v(x_1, x_2) := (x_1 \sin v + x_2 (1 - \cos v), -x_1 (1 - \cos v) + x_2 \sin v).$$

Then the operator $A_v : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ given by $A_v := (L_{-v})^{-1} L_v$ is an orthogonal transformation of \mathbb{R}^2 .

Proof. First of all observe that $\det(L_v) = \det(L_{-v}) = 2(1 - \cos v) \neq 0$, since by assumption $v \neq 0$. Moreover the inverse map is given by

$$(8.12) \quad (L_v)^{-1} = \frac{1}{\det(L_v)} L_v^T,$$

where L_v^T is the transpose operator of L_v . We conclude that

$$A_v A_v^T = \frac{1}{\det(L_v)^2} (L_{-v}^T L_v) (L_{-v}^T L_v)^T = L_{-v}^{-1} L_v L_v^{-1} L_{-v} = Id.$$

□

Let us consider the following map

$$(8.13) \quad \begin{aligned} \Psi : \mathbb{S} \times \left((-2\pi, 0) \cup (0, 2\pi) \right) \times (0, +\infty) &\rightarrow \mathbb{S} \times \left((-2\pi, 0) \cup (0, 2\pi) \right) \times (0, +\infty) \\ (a + ib, v, r) &\mapsto \left(A_v(a + ib), -\frac{v(2\pi - |v|)}{2|v|}, \frac{2\pi - |v|}{2|v|} r \right), \end{aligned}$$

where A_v was defined in Lemma 8.7, and $A_v(a + ib)$ is a short notation for the $2n$ -vector $(A_v(a_j + ib_j))_{j=1, \dots, n}$. Notice that A_v maps \mathbb{S} to \mathbb{S} thanks to Lemma 8.7. The next lemma is the key to show that the Heisenberg group \mathbb{H}^n enters into the framework of Theorem 5.3.

Lemma 8.8. *The following hold:*

- (1) *The map $\Psi : \mathbb{S} \times \left((-2\pi, 0) \cup (0, 2\pi) \right) \times (0, +\infty) \rightarrow \mathbb{S} \times \left((-2\pi, 0) \cup (0, 2\pi) \right) \times (0, +\infty)$ defined in (8.13) is a diffeomorphism.*
- (2) *The map*

$$(8.14) \quad \Lambda : \mathbb{H}^n \setminus \left(L \cup \{[\zeta, t] \in \mathbb{H}^n : t = 0\} \right) \rightarrow \mathbb{H}^n \setminus \left(L \cup \{[\zeta, t] \in \mathbb{H}^n : t = 0\} \right), \quad \Lambda := \Phi \circ \Psi \circ \Phi^{-1}$$

is a diffeomorphism. Moreover

$$(8.15) \quad \mathbf{d}_c(x, \Lambda(x)) = \mathbf{d}_c(x, 0) + \mathbf{d}_c(0, \Lambda(x)), \quad \forall x \in \mathbb{H}^n \setminus \left(L \cup \{[\zeta, t] \in \mathbb{H}^n : t = 0\} \right).$$

In other words $(x, \Lambda(x)) \in H(0)$, for every $x \in \mathbb{H}^n \setminus \left(L \cup \{[\zeta, t] \in \mathbb{H}^n : t = 0\} \right)$, where $H(0)$ was defined in (2.2).

Proof. The first claim directly follows from the formula (8.13) and Lemma 8.7, we pass then to the proof of the second claim. Recalling the definition (8.10) of Φ , Theorem 8.5 and Proposition 8.6, we have that

$\Phi : \mathbb{S} \times \left((-2\pi, 0) \cup (0, 2\pi) \right) \times (0, +\infty) \rightarrow \mathbb{H}^n \setminus \left(L \cup \{[\zeta, t] \in \mathbb{H}^n : t = 0\} \right)$ is a diffeomorphism.

In particular, for every $x \in \mathbb{H}^n \setminus \left(L \cup \{[\zeta, t] \in \mathbb{H}^n : t = 0\} \right)$ we can set

$$(8.16) \quad (a + ib, v, r) := \Phi^{-1}(x), \quad \text{with } (a + ib, v, r) \in \mathbb{S} \times \left((-2\pi, 0) \cup (0, 2\pi) \right) \times (0, +\infty).$$

Let us denote with $\gamma_{a+ib, v, r}$ the curve with parameter $(a + ib, v, r)$ defined in (8.9). By Theorem 8.5, we know that $\gamma_{a+ib, v, r} : [0, 2\pi r/|v|] \rightarrow \mathbb{H}^n$ is a minimizing geodesic and, by construction, $\gamma_{a+ib, v, r}(0) = 0$, $\gamma_{a+ib, v, r}(r) = x$. Clearly the reverse-parametrized curve

$$s \in [0, r] \mapsto \gamma_{a+ib, v, r}(r - s)$$

is a geodesic from x to 0 . Since $x^{-1} = -x$ and the left translation in \mathbb{H}^n is an isometry, we infer that

$$s \in [0, r] \mapsto -x \cdot \gamma_{a+ib, v, r}(r - s)$$

is a geodesic from 0 to $-x = x^{-1}$. On the other hand, from the definition of A_v in Lemma 8.7 and the explicit expression of the endpoint map Φ given in (8.10), it is readily checked that

$$\Phi(A_v(a + ib), -v, r) = -\Phi(a + ib, v, r) = -x.$$

In other words $\gamma_{A_v(a+ib), -v, r} : [0, r] \rightarrow \mathbb{H}^n$ is a geodesic from 0 to $-x$. Since $x \notin L$, the geodesic from 0 to $-x$ is unique thanks to Theorem 8.5, so in particular we have that

$$\gamma_{A_v(a+ib), -v, r}(s) = -x \cdot \gamma_{a+ib, v, r}(r - s), \quad \forall s \in [0, r].$$

Recalling that actually $\gamma_{A_v(a+ib), -v, r}$ is length minimizing on the larger interval $[0, 2\pi r/|v|]$, it follows that we can extend $s \mapsto \gamma_{a+ib, v, r}(r - s)$ up to $[0, 2\pi r/|v|]$ to a length minimizing geodesic. By observing that $\frac{|v|+2\pi}{2|v|}r \in (r, 2\pi r/|v|)$, in particular we get

$$(8.17) \quad \begin{aligned} \gamma_{A_v(a+ib), -v, r}\left(\frac{2\pi - |v|}{2|v|}r\right) &= \gamma_{a+ib, v, r}\left(\frac{|v| - 2\pi}{2|v|}r\right) \\ \mathbf{d}_c\left(x, \gamma_{A_v(a+ib), -v, r}\left(\frac{2\pi - |v|}{2|v|}r\right)\right) &= \mathbf{d}_c(x, 0) + \mathbf{d}_c\left(0, \gamma_{A_v(a+ib), -v, r}\left(\frac{2\pi - |v|}{2|v|}r\right)\right). \end{aligned}$$

The explicit parametrizations given in (8.9) and (8.10) imply that

$$(8.18) \quad \gamma_{A_v(a+ib), -v, r}\left(\frac{2\pi - |v|}{2|v|}r\right) = \Phi\left(A_v(a + ib), -\frac{v(2\pi - |v|)}{2|v|}, \frac{2\pi - |v|}{2|v|}r\right) = \Phi \circ \Psi \circ \Phi^{-1}(x) = \Lambda(x),$$

where we used (8.16), (8.13) and (8.14). The claim (8.15) follows then by the combination of (8.17) and (8.18); finally, Λ is a diffeomorphism since composition of diffeomorphisms. \square

We are now ready to prove the main result of this subsection.

Corollary 8.9. *Let (\mathbb{H}^n, d_c) be the n -dimensional Heisenberg group endowed with the Carnot-Carathéodory distance and let $\mathfrak{m} \in \mathcal{M}^+(\mathbb{H}^n)$ be a positive Radon measure with $\text{spt}(\mathfrak{m}) = \mathbb{H}^n$, such that $(\mathbb{H}^n, d_c, \mathfrak{m})$ satisfies the qualitative MCP condition (4.1). Then \mathfrak{m} is absolutely continuous with respect to the Haar measure on (\mathbb{H}^n, d_c) , which coincides with the $(2n+1)$ -Lebesgue measure \mathcal{L}^{2n+1} under the identification $\mathbb{H}^n \simeq \mathbb{R}^{2n+1}$.*

Proof. First of all it is well known (for example it follows from Theorem 8.5) that (\mathbb{H}^n, d_c) is a non-branching geodesic metric space, moreover the Heisenberg group (\mathbb{H}^n, d_c) endowed with the Haar measure \mathcal{L}^{2n+1} satisfies the qualitative MCP condition (4.1) (more precisely it satisfies $\text{MCP}(0, N)$ if and only if $N \geq 2n + 3$ by the work of Juillet [25]). In order to apply Theorem 5.3 it is then enough to show that \mathcal{L}^{2n+1} is a reference measure for (\mathbb{H}^n, d_c) .

To this aim, observe since that the map

$$\Lambda : \mathbb{H}^n \setminus \left(L \cup \{[\zeta, t] \in \mathbb{H}^n : t = 0\} \right) \rightarrow \mathbb{H}^n \setminus \left(L \cup \{[\zeta, t] \in \mathbb{H}^n : t = 0\} \right)$$

defined in (8.14) is a diffeomorphism, and since

$$\mathcal{L}^{2n+1}(L \cup \{[\zeta, t] \in \mathbb{H}^n : t = 0\}) = 0,$$

then

$$(8.19) \quad \Lambda_{\#}(\mathcal{L}^{2n+1}) \ll \mathcal{L}^{2n+1}.$$

Defining then $\pi^0 := (\mathcal{L}^{2n+1}, \Lambda_{\#}(\mathcal{L}^{2n+1}))$, we get that π^0 is concentrated on the graph of Λ , which is contained in $H(0)$ thanks to Lemma 8.8. Therefore $\pi^0(X \times X \setminus H(0)) = 0$ and (5.1) holds at 0. The construction of π^x for $x \in \mathbb{H}^n$, satisfying (5.1), can be reduced to the one of π^0 just by conjugation with x . We conclude that $\text{lp}(\mathcal{L}^{2n+1}) = \mathbb{H}^n$ and the thesis follows directly by Theorem 5.3. \square

REFERENCES

- [1] L. Ambrosio, N. Gigli, A. Mondino and T. Rajala, Riemannian Ricci curvature lower bounds in metric spaces with σ -finite measure, arxiv:1207.4924, (2012), to appear on Trans. Amer. Math. Soc.
- [2] L. Ambrosio, N. Gigli, G. Savaré Bakry-Émery curvature-dimension condition and Riemannian Ricci curvature bounds. *Annals of Probab.*, to appear.
- [3] ———, Metric measure spaces with Riemannian Ricci curvature bounded from below, *Duke Math. J.*, 163, 1405–1490, (2014).
- [4] L. Ambrosio, A. Mondino, G. Savaré, Nonlinear diffusion equations and curvature conditions in metric measure spaces, *in preparation*.
- [5] L. Ambrosio, S. Rigot, Optimal mass transportation in the Heisenberg group. *J. Funct. Anal.*, 208, 261–301, (2004).
- [6] D. Bakry and M. Émery, Diffusions hypercontractives *Seminaire de Probabilités XIX, Lecture Notes in Math.*, Springer-Verlag, New York. 1123 (1985), 177–206.
- [7] D. Bakry and M. Ledoux, A logarithmic Sobolev form of the Li-Yau parabolic inequality. *Rev. Mat. Iberoam.* 22, (2006), 683–702.
- [8] D. Burago, Y. Burago and S. Ivanov, A course in metric geometry. *Grad. Studies in Math.*, Vol.33, Americ. Math. Soc., (2001).
- [9] S. Bianchini and F. Cavalletti, The Monge problem for distance cost in geodesic spaces. *Commun. Math. Phys.*, 318, 615 – 673 (2013).
- [10] F. Cavalletti, Monge problem in metric measure spaces with Riemannian curvature-dimension condition. *Nonlinear Anal.*, 99 (2014), 136 – 151.
- [11] F. Cavalletti and M. Huesmann, Existence and uniqueness of optimal transport maps. *Ann. I. H. Poincaré AN*, (2014) <http://dx.doi.org/10.1016/j.anihpc.2014.09.006>.
- [12] G. Cheeger Differentiability of Lipschitz functions on metric measure spaces. *Geom. Funct. Anal.*, 9 (1999), no. 3, 428 – 517.

- [13] G. Cheeger and T.H. Colding, On the structure of spaces with Ricci curvature bounded below. I. *J. Diff. Geom.*, 45 (1997), 406 – 480.
- [14] ———, On the structure of spaces with Ricci curvature bounded below. II. *J. Diff. Geom.*, 54, (2000), 13–35.
- [15] ———, On the structure of spaces with Ricci curvature bounded below. III. *J. Diff. Geom.*, 54 (2000), 37 – 74.
- [16] T.H. Colding and A. Naber, Sharp Hölder continuity of tangent cones for spaces with a lower Ricci curvature bound and applications. *Ann. of Math. (2)*, 176 (2012), 1173–1229.
- [17] M. Csornyei and P. Jones, Product Formulas for Measures and Applications to Analysis and Geometry. www.math.sunysb.edu/Videos/dfest/PDFs/38-Jones.pdf.
- [18] M Erbar, Kuwada, K.T. Sturm, On the Equivalence of the Entropic Curvature-Dimension Condition and Bochner’s Inequality on Metric Measure Space, Preprint arXiv:1303.4382, (2013). To appear in *Invent. Math.*
- [19] D. H. Fremlin, *Measure Theory*, volume 4. Torres Fremlin, 2002.
- [20] B. Gaveau, Principe de moindre action, propagation de la chaleur et estimées sous elliptiques sur certains groupes nilpotents. *Acta Mathematica* 139, 1– 2, 95–153, (1977).
- [21] N. Gigli, Optimal maps in non branching spaces with Ricci curvature bounded from below, *Geom. Funct. Anal.*, 22 (2012) no. 4, 990–999.
- [22] N. Gigli, The splitting theorem in non-smooth context, *preprint arXiv:1302.5555*, (2013).
- [23] N. Gigli, A. Mondino and T. Rajala, Euclidean spaces as weak tangents of infinitesimally Hilbertian metric measure spaces with Ricci curvature bounded below, *Journal für die Reine und Angewandte Mathematik (Crelle’s journal)* (in press). DOI 10.1515/crelle-2013-0052.
- [24] A. Grigor’yan, Heat equation on a non-compact Riemannian manifold, *Math USSR Sb.* 72 (1992) no. 1, 47–77. (Translated from Russian *Matem. Sbornik*, 182 (1991), no. 1, 55–87).
- [25] N. Juillet, Geometric Inequalities and Generalized Ricci Bounds in the Heisenberg Group. *Int. Math. Res. Not.*, 2009, 13, 2347–2373, (2009).
- [26] W.P. Klingenberg, Riemannian geometry. Second edition. *de Gruyter Studies in Mathematics*, 1. Walter de Gruyter & Co., Berlin, (1995), x+409 pp.
- [27] K. Kuwae and T. Shioya, On generalized measure contraction property and energy functionals over Lipschitz maps. *Potential Anal.*, 15, 105–121, (2001).
- [28] ———, Infinitesimal Bishop-Gromov condition for Alexandrov spaces. Probabilistic approach to geometry. *Adv. Stud. Pure Math.* 57, Math. Soc. Japan, Tokyo, (2010)
- [29] J. Lott and C. Villani, Ricci curvature for metric-measure spaces via optimal transport, *Ann. of Math.*, 169, 903–991 (2009).
- [30] A. Mondino and A. Naber, Structure Theory of Metric-Measure Spaces with Lower Ricci Curvature Bounds I, *preprint arXiv:1405.2222*, (2014).
- [31] R. Montgomery, A tour of subriemannian geometries, their geodesics and applications. *Mathematical Surveys and Monographs*, Vol. 91, Amer. Math. Soc., Providence, RI, (2002), xx+259pp.
- [32] R. Monti, Some properties of Carnot- Carathéodory balls in the Heisenberg group, *Rend. Mat. Acc. Lincei*, 11, 155–167, (2000).
- [33] I.G. Nikolaev, Smoothness of the metric of spaces with two-sided bounded Alexandrov curvature, *Siberian Math. Journ.*, 24, 247–263, (1983).
- [34] S.I. Ohta, On the measure contraction property of metric measure spaces *Comment. Math. Helv.*, 82, 805–828, (2007).
- [35] Y. Otsu and T. Shioya, Riemannian structure of Alexandrov spaces. *J. Diff. Geom.*, 39, 629–658, (1994).
- [36] K.T. Sturm, Diffusion processes and heat kernels on metric spaces. *Annals of Probab.*, 26, 1–55, (1998).
- [37] ———, On the geometry of metric measure spaces. I, *Acta Math.* 196, 65–131, (2006) .
- [38] ———, On the geometry of metric measure spaces. II, *Acta Math.*, 196, 133–177, (2006)
- [39] T. Rajala and K.T. Sturm, Non-branching geodesics and optimal maps in strong $CD(K, \infty)$ -spaces. *Calc. Var. Partial Differential Equations*, 50, 3-4, 831–846, (2014).
- [40] P. Pansu, Métriques de Carnot-Carathéodory et quasiisométries des espaces symétriques de rang un. *Ann. of Math.*, 129, 1, 1–60, (1989).

CRM DE GIORGI - SCUOLA NORMALE SUPERIORE, PIAZZA DEI CAVALIERI 3, I-56126 PISA (ITALY)
E-mail address: fabio.cavalletti@sns.it

ETH, RÄMISTRASSE 101, 8092 ZÜRICH (SWITZERLAND)
E-mail address: andrea.mondino@math.ethz.ch