

## VIII. REFERENCES

- [1] Rubin BP., Tucker RP., Martin D., Chiquet-Ehrismann, R. (1999). Teneurins: a novel family of neuronal cell surface proteins in vertebrates, homologous to the *Drosophila* pair-rule gene product Ten-m. *Dev. Biol.* 216, 195-209.
- [2] Baumgartner S., Chiquet-Ehrismann R. (1993). Tena, a *Drosophila* gene related to tenascin, shows selective transcript localization. *Mech. Dev.* 40, 165 – 176.
- [3] Baumgartner S., Martin D., Hagios C., Chiquet-Ehrismann, R. (1994). Ten-m, a *Drosophila* gene related to tenascin, is a new pair-rule gene. *EMBO J.* 13, 3728 – 3740.
- [4] Levine A., Bashan-Ahrend A., Budai-Hadrian O., Gartenberg D., Mena-sherow S., Wides R. (1994). Odd Oz: a novel *Drosophila* pair rule gene. *Cell* 77, 587–598.
- [5] Tucker RP., Chiquet-Ehrismann R. (2006). Teneurins: A conserved family of transmembrane proteins involved in intercellular signalling during development. *Dev. Biol* 209, 237-245.
- [6] Brellier F., Martina E., Chiquet M., Ferralli J., van der Heyden M., Orend G., Schittny J.C., Chiquet-Ehrismann R., Tucker RP. (2012). The adhesion modulating properties of tenascin-w. *Int J Biol Sci.* 8, 187-94.
- [7] Chavez M. and Réka A. (2008). Studying the effect of cell division on expression patterns of the segment polarity genes. *J. R. Soc. Interface* 5. Suppl 1 S71-S84
- [8] Jürgens G., Wieschaus E., Niisslein-Volhard, C and Kluding H. (1984). Mutation affecting the pattern of the larval cuticle in *Drosophila Melanogaster*. *Roux's Arch Dev Biol.* 193, 283-295.
- [9] Minet AD., Rubin BP., Tucker R.P., Baumgartner S., Chiquet-Ehrismann R. (1999). Teneurin-1, a vertebrate homologue of the *Drosophila* pair-rule gene ten-m, is a neuronal protein with a novel type of heparin-binding domain. *J. Cell Sci.* 112, 2019–2032.
- [10] Levine A., Weiss C., Wides R. (1997). Expression of the pair-rule gene odd Oz (odz) in imaginal tissues. *Dev. Dyn.* 209, 1–14.
- [11] Jasper H., Benes V., Atzberger A., Sauer S., Ansorge W., Bohmann D., (2002). A genomic switch at the transition from cell proliferation to terminal differentiation in the *Drosophila* eye. *Dev. Cell* 3, 511–521.
- [12] Fascetti N., Baumgartner S. (2002). Expression of *Drosophila* Ten-a, a dimeric receptor during embryonic development. *Mech. Dev.* 114, 197.
- [13] Drabikowski K., Trzebiatowska A., Chiquet-Ehrismann R. (2005). ten-1, an essential gene for germ cell development, epidermal morphogenesis, gonad migration, and neuronal pathfinding in *Caenorhabditis elegans*. *Dev. Biol.* 282, 27–38.
- [14] Trzebiatowska A., Topf U., Sauder U., Drabikowski K., Chiquet-Ehrismann R. (2008) *Caenorhabditis elegans* teneurin, *ten-1*, is required for gonadal and pharyngeal basement membrane integrity and acts redundantly with integrin *ina-1* and dystroglycan *dgn-1*. *Mol Biol Cell*, 19 3898-3908.
- [15] Bagutti C., Forro G., Ferralli J., Rubin BP., Chiquet-Ehrismann R. (2003). The intracellular domain of teneurin-2 has a nuclear function and represses zic-1-mediated transcription. *J. Cell Sci.* 116, 2957–2966.
- [16] Wang XZ., Kuroda M., Sok J., Batchvarova N., Kimmel R., Chung P., Zinszner H., Ron D. (1998). Identification of novel stress-induced genes downstream of chop. *EMBO J.* 17, 3619–3630.
- [17] Oohashi T., Zhou XH., Feng K., Richter B., Morgelin M., Perez MT., Su WD.,

- Chiquet-Ehrismann R., Rauch U., Fassler R. (1999). Mouse ten- m/Odz is a new family of dimeric type II transmembrane proteins expressed in many tissues. *J. Cell Biol.* 145, 563–577.
- [18] Ben-Zur T., Feige E., Motro B., Wides R. (2000). The mammalian Odz gene family: homologs of a *Drosophila* pair-rule gene with expression implying distinct yet overlapping developmental roles. *Dev. Biol.* 217, 107–120.
- [19] Ben-Zur T. and Wides R. (1999). Mapping homologs of *Drosophila* odd Oz (odz): Doc4/Odz4 to mouse chromosome 7, Odz1 to mouse chromosome 11, and ODZ3 to human chromosome Xq25. *Genomics* 58, 107–120.
- [20] Zhou X.H., Brandau O., Feng K., Oohashi T., Ninomiya Y., Rauch U., Fassler R. (2003). The murine Ten-m/Odz genes show distinct but overlapping expression patterns during development and in adult brain. *Gene Expression Patterns* 3, 397–405.
- [21] Brandau O., Schuster V., Weiss M., Hellebrand H., Fink FM., Kreczy A., Friedrich W., Strahm B., Niemeyer C., Belohradsky BH., Meindl A. (1999). Epstein-Barr virus-negative boys with non-Hodgkin lymphoma are mutated in the SH2D1A gene, as are patients with X-linked lymphoproliferative disease (XLP). *Hum. Mol. Genet.* 13, 2407–2413.
- [22] Mieda M., Kikuchi Y., Hirate Y., Aoki M., Okamoto H. (1999). Compartmentalized expression of zebrafish ten-m3 and ten-m4, homologues of the *Drosophila* ten (m)/odd Oz gene in the central nervous system. *Mech. Dev.* 87, 223–227.
- [23] Otaki JM. and Firestein S. (1999). Neurestin: putative transmembrane molecule implicated in neuronal development. *Dev. Biol.* 212, 165–181.
- [24] Feng K., Zhou XH., Oohashi T., Morgelin M., Lustig A., Hirakawa S., Ninomiya Y., Engel J., Rauch U., Fassler R. (2002). All four members of the Ten-m/Odz family of transmembrane proteins form dimers. *J. Biol. Chem.* 277, 26128–26135.
- [25] Rubin BP., Tucker RP., Brown-Luedi M., Martin D., Chiquet-Ehrismann R. (2002). Teneurin 2 is expressed by the neurons of the thalamofugal visual system in situ and promotes homophilic cell-cell adhesion in vitro. *Development* 129, 4697–4705.
- [26] Feulner G., Gray J.A., Kirschman J.A., Lehner AF., Sadosky AB., Vlazny DA., Zhang J., Zhao S., Hill C.W. (1990). Structure of the rhsA locus from *Escherichia coli* K-12 and comparison of rhsA with other members of the rhs multigene family. *J. Bacteriol.* 172, 446–456.
- [27] Minet AD. and Chiquet-Ehrismann R. (2000). Phylogenetic analysis of teneurin genes and comparison to the rearrangement hot spot elements of *E. coli*. *Gene* 257, 87–97.
- [28] Wang L., Rotzinger S., Al Chawaf A., Elias CF., Barsyte-Lovejoy D., Qian X., Wang NC., De Cristofaro A., Belsham D., Bittencourt J.C., Vaccarino F., Lovejoy DA., (2005). Teneurin proteins possess a carboxy terminal sequence with neuromodulatory activity. *Mol. Brain Res.* 133, 253 – 265.
- [29] Lovejoy DA., Chawaf A., Cadinouche A. (2006). Teneurin C-terminal associated peptides: An enigmatic family of neuropeptides with structural similarity to the corticotropin-releasing factor and calcitonin families of peptides. *Gen Comp Endocrinol.* 15;(3), 299-305.
- [30] Tucker RP, Drabikowski K, Hess JF, Ferralli J, Chiquet-Ehrismann R, Adams JC. (2006). Phylogenetic analysis of the tenascin gene family: evidence of origin early in the chordate lineage. *BMC Evol Biol.* 7;6, 60-77.
- [31] Stanley P. (2007). Regulation of Notch signaling by glycosylation. *Curr Opin Struct Biol.* 17(5): 530-535.
- [32] Rampal R, Luther KB, Haltiwanger RS.(2007). Notch signaling in normal and disease States: possible therapies related to glycosylation. *Curr Mol Med.* 7(4), 427-445.
- [33] Moloney, DJ., Panin, VM., Johnston, SH., Chen, J., Shao, L., Wilson, R., Wang, Y.,

- Stanley, P., Irvine, KD., Haltiwanger, RS., Vogt TF. (2000). Fringe is a glycosyltransferase that modifies Notch. *Nature* 27;406, 369-375.
- [34] Haines N., Irvine KD. (2003). Glycosylation regulates Notch signalling. *Nat Rev Mol Cell Biol.* 4(10), 786-797.
- [35] Young TR., Leamey CA. (2009). Teneurins: Important regulators of neural circuitry. *Int. J. Bioch Cell Biol.* 41, 990-993.
- [36] Kenzelmann D., Chiquet-Ehrismann R., Leachman NT., Tucker RP. (2008). Teneurin-1 is expressed in interconnected regions of the developing brain and is processed in vivo. *BMC Dev. Biol* 8:30.
- [37] Tucker RP., Martin D., Kos R., Chiquet-Ehrismann R. (2000). The expression of teneurin-4 in the avian embryo. *Mech. Dev.* 98, 187-191.
- [38] Tucker RP., Chiquet-Ehrismann R., Chevron MP., Martin D., Hall RJ., Rubin BP. (2001). Teneurin-2 is expressed in tissues that regulate limb and somite pattern formation and is induced in vitro and in situ by FGF8. *Dev. Dyn.* 220, 27-39.
- [39] Kenzelmann D., Tucker RP., Leachman NT., Chiquet-Ehrismann R. (2010). The expression of teneurin-4 in the avian embryo: potential roles in patterning of the limb and nervous system. *Int. J. Dev. Biol.* 54(10):1509-1516.
- [40] Kenzelmann D. (2008). Teneurins in Development and Disease. Doctoral Thesis. Unpublished observation.
- [41] Robert Marcus., David Feldman., Dorothy Nelson., Clifford J. Rosen. *Fundamental of Osteoporosis.* (2009). Accademic Press. Chapter 8, 209-215.
- [42] Panman L and Zeller R. (2003). Patterning the limb before and after SHH signalling. *J Anat.* 202(1): 3-12.
- [43] Leamey CA, Glendining KA, Kreiman G, Kang ND, Wang KH, Fassler R, Sawatari A, Tonegawa S, Sur M. (2008). Differential gene expression between sensory neocortical areas: potential roles for Ten\_m3 and Bcl6 in patterning visual and somatosensory pathways. (2008). *Cereb Cortex.* 18(1):53-66.
- [44] Leamey CA., Merlin S., Lattouf P., Sawatari A., Zhou X., Demel N., Glendining KE., Oohashi T., Sur M., Fassler R. (2007). Ten\_m3 Regulates Eyes-Specific Patterning in the Mammalian Visual Pathway and Is Required for binocular Vision. *PLoS Biol* 5, 2077-2092
- [45] Getchell TV, Liu H., Vaishnav RA., Kwong K., Stromberg AJ., Getchell ML. (2005). Temporal profiling of gene expression during neurogenesis and remodeling in the olfactory epithelium at short intervals after target ablation. *Journal of Neuroscience Research* 80(3), 309-329,
- [46] Li L., Bishop KM., O'Leary DM. (2006). Potential target genes of EMX2 include Odz/Ten-M and other gene families with implications for cortical patterning. *Mol Cell Neurosci* 33(2), 136-149.
- [47] Lossie AC., Nakamura H., Thomas SE., Justice MJ. (2004). Mutation of 17Rn3 shows that Odz4 is required for mouse gastrulation. *Genetics* 169(1):285-99.
- [48] Patent application title: TENEURIN AND CANCER (2011). Patent application number: 20110229494. Inventors: Ruth Chiquet-Ehrismann and Daniela Kenzelmann Broz
- [49] [www.genecards.org](http://www.genecards.org)
- [50] Nunes SM., Ferralli J., Choi K., Brown-Luedi M., Minet AD., Chiquet-Ehrismann R. (2005). The intracellular domain of teneurin-1 interacts with MBD1 and CAP/ponsin resulting in subcellular codistribution and translocation to the nuclear matrix. *Exp. Cell Res.* 305, 122-132.
- [51] Ching RW., Dellaire G., Eskiw CH., Bazett-Jones DP. (2005). PML bodies: a meeting

- place for genomic loci?. *J. Cell Sci.* 118, 847–854.
- [52] Mandai K., Nakanishi H., Satoh A., Takahashi K., Satoh K., Nishioka H., Mizoguchi A., Takai Y., (1999). Ponsin/SH3P12: an I-afadin- and vinculin- binding protein localized at cell – cell and cell – matrix adherens junctions. *J. Cell Biol.* 144, 1001–1017.
- [53] Urban S. and Freeman M. (2002). Intramembrane proteolysis controls diverse signalling pathways throughout evolution. *Curr. Opin. Genet. Dev.* 5, 512 – 518.
- [54] <http://www.ncbi.nlm.nih.gov>
- [55] [http://www.ncbi.nlm.nih.gov/nucleotide/NM\\_001098816.2](http://www.ncbi.nlm.nih.gov/nucleotide/NM_001098816.2)
- [56] Qian X., Baryte-Lovejoy D., Wang L., Chewpoy B., Gautam N., Al Chawaf A., Lovejoy D.A. (2004). Cloning and characterization of teneurin C-terminus associated peptide (TCAP)-3 from the hypothalamus of an adult rainbow trout (*Oncorhynchus mykiss*). *Gen. Comp. Endocrinol.* 137, 205 – 216.
- [57] Tan L., Al Chawafa A., Vaccarino FJ., Boutros PC., Lovejoy DA. (2011) Teneurin C-terminal associated peptide (TCAP)-1 modulates dendritic morphology in hippocampal neurons and decreases anxiety-like behaviors in rats. *Physiol Behav.* 104(2): 199-204.
- [58] Al Chawafa A., Xuc K., Tan L., Vaccarino FJ., Lovejoy DA., Rotzinger S. (2007). Corticotropin-releasing factor (CRF)-induced behaviors are modulated by intravenous administration of teneurin C-terminal associated peptide-1 (TCAP-1). *Peptides* 28(7):1406-1415.
- [59] Trubiani G., Al Chawaf A., Belsham DD, Baryte-Lovejoy D, Lovejoy DA. (2007). Teneurin carboxy (C)-terminal associated peptide-1 inhibits alkalosis-associated necrotic neuronal death by stimulating superoxide dismutase and catalase activity in immortalized mouse hypothalamic cells. *Brain Res.* 1176:27-36.
- [60] Ng T., Chand D., Song L., Al Chawaf A., Watson JD., Boutros PC, Belsham DD, Lovejoy DA. (2011). Identification of a novel Brain Derived Neurotrophic Factor (BDNF)-inhibitory factor: Regulation of BDNF by Teneurin C-terminal Associated Peptide (TCAP)-1 in immortalized embryonic mouse hypothalamic cells. *Regul Pept.* In press.
- [61] <http://www.protagenic.com/publications.html>
- [62] Luckasson R and Reeve A. (2001). Naming, defining, and classifying in mental retardation. *Ment Retard.* 39(1):47-52.
- [63] Attree O, Olivos IM, Okabe I, Bailey LC, Nelson DL, Lewis RA, McInnes RR, Nussbaum RL. (1992). The Lowe's oculocerebrorenal syndrome gene encodes a protein highly homologous to inositol polyphosphate-5-phosphatase. *Nature* 358(6383):239-242.
- [64] Garcia CC., Blair HJ., Seager M., Coulthard A., Tennant S., Buddles M., Curtis A., Goodship JA. (2004). Identification of a mutation in synapsin I, a synaptic vesicle protein, in a family with epilepsy. *J Med Genet.* 41(3):183-186.
- [65] Amir RE., Van den Veyver IB., Wan M., Tran CQ., Francke U., Zoghbi HY.(1999). Rett syndrome is caused by mutations in X-linked MECP2, encoding methyl-CpG-binding protein 2. *Nat Genet.* 23(2):185-188.
- [66] Li H., Yamagata T., Mori M., Yasuhara A., Momoi MY. (2005). Mutation analysis of methyl-CpG binding protein family genes in autistic patients. *Brain Dev.* 27(5):321-325.
- [67] Beckmann J., Vitobello A., Ferralli J., Brož DK., Rijli FM., Chiquet-Ehrismann R. (2011). Human teneurin-1 is a direct target of the homeobox transcription factor EMX2 at a novel alternate promoter. *BMC Dev Biol.* 8;11:35.
- [68] Brown SA., Warburton D., Brown LY., Yu CY., Roeder ER., Stengel-Rutkowski S., Hennekam RC., Muenke M. (1998). Holoprosencephaly due to mutations in ZIC2, a homologue of *Drosophila* odd-paired. *Nat Genet.* 20(2):180-3.

- [69] Nathanson KL., Shugart YY., Omaruddin R., Szabo C., Goldgar D., Rebbeck TR., Weber BL. (2002). CGH-targeted linkage analysis reveals a possible BRCA1 modifier locus on chromosome 5q. *Hum Mol Genet* 11:1327-1332.
- [70] Minami M., Daimon Y., Mori K., Takashima H., Nakajima T., Itoh Y., Okanou T. (2005). Hepatitis B virus-related insertional mutagenesis in chronic hepatitis B patients as an early drastic genetic change leading to hepatocarcinogenesis. *Oncogene* 24:4340-4348.
- [71] Lee S., Medina D., Tsimelzon A., Mohsin SK., Mao S., Wu Y., Allred DC. (2007). Alterations of gene expression in the development of early hyperplastic precursors of breast cancer. *Am J Pathol* 171:252-262.
- [72] Vinatzer U., Gollinger M., Mullauer L., Raderer M., Chott A., Streubel B. (2008). Mucosa-associated lymphoid tissue lymphoma: novel translocations including rearrangements of ODZ2, JMJD2C, and CNN3. *Clin Cancer Res* 14:6426-6431.
- [73] Ziegler A., Cerciello F., Bigosch C., Bausch-Fluck D., Felley-Bosco E., Ossola R., Soltermann A., Stahel RA., Wollscheid B. (2011). Proteomic surfaceome analysis of mesothelioma. *Lung Cancer* 75(2):189-196.
- [74] Carbone M., Kratzke RA., Testa JR. (2002). The pathogenesis of mesothelioma. *Semin. Oncol.* 29:2-17
- [75] Husain AN., Colby TV., Ordonez NG., Krausz T., Borczuk A., Cagle PT, et al. (2009). Guidelines for pathologic diagnosis of malignant mesothelioma: a consensus statement from the International Mesothelioma Interest Group. *Arch Pathol Lab Med* 133:1317-1331.
- [76] Roberts F., McCall AE., Burnett RA. (2001). Malignant mesothelioma: a comparison of biopsy and postmortem material by light microscopy and immunohistochemistry. *J Clin Pathol* 54:766-70.
- [77] Jemal A., Siegel R., Ward E., Murray T., Xu J., Thun MJ. (2007). Cancer statistics, 2007, *CA Cancer. J Clin* 57:43-466.
- [78] Duffy MJ., Bonfrer JM., Kulpa J., Rustin GJ., Soletormos G., Torre GC., Tuxen MK., Zwirner M. (2005). CA125 in ovarian cancer: European Group on Tumor Markers guidelines for clinical use. *Int J Gynecol Cancer* 15(5):679-691.
- [79] Landen CN., Birrer MJ., Sood AK. (2008). Early events in the pathogenesis of epithelial ovarian cancer, *J Clin Oncol* 26:995-1005.
- [80] King MC., Marks JH., Mandell JB. (2003). Breast and ovarian cancer risks due to inherited mutations in BRCA1 and BRCA2. *Science* 302:643-646 57.
- [81] Stirling D., Evans DG., Pichert G., Shenton A., Kirk EN., Rimmer S., Steel CM., Lawson S., Busby-Earle RM., Walker J., Lalloo FI., Eccles DM., Lucassen AM, Porteous ME. (2005). Screening for familial ovarian cancer: failure of current protocols to detect ovarian cancer at an early stage according to the international Federation of gynecology and obstetrics system. *J Clin Oncol* 23:5588-5596.
- [82] Aarnio M., Sankila R., Pukkala E., Salovaara R., Aaltonen LA., de la Chapelle A., Peltomaki P., Mecklin JP., Jarvinen HJ. (1999). Cancer risk in mutation carriers of DNA-mismatch-repair genes. *Int J Cancer* 81:214-218
- [83] Kaku T., Ogawa S., Kawano Y., Ohishi Y., Kobayashi H., Hirakawa T., Nakano H. (2003). Histological classification of ovarian cancer. *Med Electron Microsc* 36:9-17
- [84] Scully RE. (1987). Classification of human ovarian tumors. *Environ Health Perspect* 73:15-24
- [85] Chang K., Pastan I., Willingham MC.(1992). Isolation and characterization of a monoclonal antibody, K1, reactive with ovarian cancers and normal mesothelium. *Int J Cancer* 50:373-381

- [86] Langseth H., Hankinson SE., Siemiatycki J., Weiderpass E. (2008). Perineal use of talc and risk of ovarian cancer. *J Epidemiol Community Health* 62:358-360
- [87] Robinson BW., Creaney J., Lake R., Nowak A., Musk AW., de Klerk N., Winzell P., Hellstrom KE., Hellstrom I. (2005). Soluble mesothelin-related protein—a blood test for mesothelioma. *Lung Cancer* 49 Suppl 1:S109-111
- [88] Bao LH., Sakaguchi H., Fujimoto J., Tamaya T. (2007). Osteopontin in metastatic lesions as a prognostic marker in ovarian cancers. *J Biomed Sci* 14:373-381
- [89] Visintin I., Feng Z., Longton G., Ward DC., Alvero AB., Lai Y., Tenthorey J., Leiser A., Flores-Saaib R., Yu H, Azori M et al. (2008). Diagnostic Markers for Early Detection of Ovarian Cancer. *Clin Cancer Res* 15;(16):5308.
- [90] Cheng JQ., Godwin AK., Bellacosa A., Taguchi T., Franke TF., Hamilton TC., Tschlis PN., Testa JR. (1992). AKT2, a putative oncogene encoding a member of a subfamily of protein-serine/threonine kinases, is amplified in human ovarian carcinomas. *Proc Natl Acad Sci USA* 89:9267-9271
- [91] Ramos-Nino ME., Vianale G., Sabo-Attwood T., Mutti L., Porta C., Heintz N., Mossman BT. (2005). Human mesothelioma cells exhibit tumor cell-specific differences in phosphatidylinositol 3-kinase/AKT activity that predict the efficacy of Onconase. *Mol Cancer* 4:835-842
- [92] Naylor MS., Stamp GW., Davies BD., Balkwill FR. (1994). Expression and activity of MMPS and their regulators in ovarian cancer. *Int J Cancer* 58:50-56
- [93] Paley PJ., Staskus KA., Gebhard K., Mohanraj D., Twigg LB., Carson LF., Ramakrishnan S. (1997). Vascular endothelial growth factor expression in early stage ovarian carcinoma. *Cancer* 1997, 80:98-106
- [94] Xu L., Fidler IJ. (2000). Interleukin 8: an autocrine growth factor for human ovarian cancer. *Oncol Res* 12:97-106 103.
- [95] Kurose K., Zhou XP., Araki T., Cannistra SA., Maher ER., Eng C. (2001). Frequent loss of PTEN expression is linked to elevated phosphorylated Akt levels, but not associated with p27 and cyclin D1 expression, in primary epithelial ovarian carcinomas. *Am J Pathol* 158:2097-2106
- [96] Havrilesky LJ., Alvarez AA., Whitaker RS., Marks JR., Berchuck A. (2001). Loss of expression of the p16 tumor suppressor gene is more frequent in advanced ovarian cancers lacking p53 mutations. *Gynecol Oncol* 83:491-500
- [97] Gatcliffe TA., Monk BJ., Planutis K., Holcombe RF. (2008). Wnt signaling in ovarian tumorigenesis. *Int J Gynecol Cancer* (5):954-62
- [98] Fox S., Dharmarajan A. (2006). WNT signaling in malignant mesothelioma. *Front Biosci* 11:2106-2112
- [99] Davidson B., Zhang Z., Kleinberg L., Li M., Florenes VA., Wang TL., Shih Ie M. (2006). Gene expression signatures differentiate ovarian/peritoneal serous carcinoma from diffuse malignant peritoneal mesothelioma. *Clin Cancer Res* 12:5944-5950.
- [100] <http://www.ncbi.nlm.nih.gov/geo/>
- [101] <https://www.oncomine.org>
- [102] ATCC Global Bioresource Center. [www.atcc.org](http://www.atcc.org)
- [103] (<http://www.ebi.ac.uk/Tools/msa/clustalw2/>; default alignment parameters were used).
- [104] NCBI Tool. BLAST 2.2.26. Zheng Zhang, Scott Schwartz, Lukas Wagner, and Webb Miller (2000), "A greedy algorithm for aligning DNA sequences", *J Comput Biol* 2000; 7(1-2):203-14.
- [105] UCSC Genome Browser. BLAT Tool. Kent Informatics, Inc.

<http://genome.ucsc.edu/cgi-bin/hgBlat>

[106] <http://www.premierbiosoft.com/netprimer/>

[107] <http://www.idtdna.com/Home/Home.aspx>

[108] Wang XZ., Jolicoeur EM., Conte N., Chaffanet M., Zhang Y., Mozziconacci M., Feiner H., Birnbaum D., Pébusque MJ and Ron D. (1999). Heregulin is the product of a chromosomal translocation fusing the *DOC4* and *HGL/NRG1* genes in the MDA-MB-175 breast cancer cell line. *Oncogene* 18, 5718-5721

[109] [http://www.macrogen.com/eng/macrogen/macrogen\\_main.jsp](http://www.macrogen.com/eng/macrogen/macrogen_main.jsp)

[110] UCSC Genome Bioinformatics Browser

[111] Ensembl Genome Browser

[112] <http://www.ncbi.nlm.nih.gov/Sequin/index.html>

[113] Frohman MA. (1994). On beyond classic RACE (rapid amplification of cDNA ends). *Genome Res.* 4: S40-S58

[114] Molecular Cloning (2001). A Laboratory Manual (eds. Sambrook, J. & Russell, D.W.) Chapters 6,7,8-10. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, USA.

[115] Taylor TD., et al. (2004). Human chromosome 11 DNA sequence and analysis including novel gene identification. *Nature* 440 (23), 497-500

[119] Ota T., et al. (2003). Complete sequencing and characterization of 21243 full-length human cDNA. *Nature Genetics* 36, 41-45

[120] Nagase T., Kukino R., Ishikawa K., Hirosawa M. and Ohara O. (2000). Prediction of the Coding Sequences of Unidentified Human Genes. XVI. The complete sequences of 150 new cDNA clones from brain which code for large proteins in vitro. *DNA Research* 7, 65-73

[121] GenBank Accession n. AK309124. (2008). Wakamatsu A., Yamamoto J., Kimura K., Ishii S., Watanabe K., Sugiyama A., Murakawa K., Kaida T., Tsuchiya K., Fukuzumi Y., et al. Direct submission

[122] Tucker RP., Beckmann J., Leachman NT., SchoLer J. and Chiquet-Ehrismann R. (2011). Phylogenetic Analysis of the Teneurins: Conserved Features and Premetazoan Ancestry. *Mol Bio Evol.* MBE Advances Access December 2011. doi:10.1093/molbev/msr271