

**ENHANCED OXIDATION OF BIS(3,5-DIBROMOSALICYL) FUMARATE
 α - α CROSS LINKED HEMOGLOBIN BY FREE RADICALS
GENERATED BY XANTHINE/XANTHINE OXIDASE**

M. Samaja, R. Motterlini and E. Rovida
Department of Biomedical Science and Technology
Scientific Institute San Raffaele
University of Milano
via Olgettina 60, 20132 Milano, ITALY

ABSTRACT

The xanthine/xanthine oxidase reaction produces reproducible amounts of oxygen-derived free radicals that oxidize human oxyhemoglobin (Hb). We monitored the kinetics of the oxidation of stripped Hb (sHb), purified HbA₀ and α - α cross-linked Hb (HbXL99 α) at [Hb] in the 5 to 150 μ M (heme) range. For increasing [Hb], the oxidation half-time ($t_{1/2}$) increased for all Hbs, but $t_{1/2}$ was always less for HbXL99 α than for HbA₀ and sHb. Such feature was attributed to the lower affinity for O₂ of HbXL99 α and may represent a serious problem for use of this Hb as blood substitute.

INTRODUCTION

The cytotoxicity of free plasma hemoglobin (Hb) involves free radicals [1,2]. Hb is known to generate free radicals during autooxidation [3,4] and to increase through the Fenton reaction the peroxidation of fatty acids initiated by free radicals generated by external sources [5-7]. Here, we focus on Hb as *target* of oxygen-derived free radicals (ODFR) generated by xanthine (XAN) and xanthine oxidase (XO). The main

features of this approach are: 1) The amount of ODFR is modulated and reproducible because they are produced stoichiometrically by an enzymatic system; 2) The primary target of ODFR is distinguished from secondary targets such as peroxidation of lipids; 3) The time course of this reaction is faster than that of Hb autooxidation; 4) The effects of ODFR are easily monitored with spectrophotometric methods.

We characterized this model monitoring the spectral changes of HbO₂ when mixed with XAN/XO assuming the mechanism drawn in Fig.1 (unbalanced equations). We show that ODFR oxidize HbO₂ in a concentration-dependent fashion with an action likely directed towards the heme. The Hb derivative cross-linked between the α chains by bis(3,5-dibromosalicyl) fumarate (HbXL99 α) appeared more susceptible than stripped Hb (sHb) and HbA₀ to the oxidation by ODFR perhaps for its lower O₂ affinity.

MATERIALS AND METHODS

Materials. The medium contained 50 mM K₂HPO₄, 1 mM ethylenediaminetetraacetate (EDTA), 0.5 mM XAN, 20 mM KCN, pH 7.3 at 37°C. Dimethylthiourea (DMTU, Aldrich), XO (20 U/mL, cow milk), superoxide dismutase (SOD, 5,000 U/mg, bovine erythrocytes), catalase (260,000 U/mL, beef liver), and cytochrome c (Boehringer Biochemia) were used without further treatment.

We prepared sHb from freshly drawn blood of non-smoker donors (HbCO<2%) [8]. Total [Hb] was determined by the met-cyanide method [9]. MetHb was obtained mixing HbO₂ with 20% molar excess solid K₃Fe(CN)₆ for 15 min at room temperature [8]. HbA₀ was purified by anion exchange high-performance liquid chromatography [10]. HbXL99 α was obtained reacting HbA₀ with 3,5-bis-dibromosalicyl-fumarate [11].

Methods. Visible and UV spectra were obtained with a DU-70 (Beckman Instruments Inc., Fullerton, CA). Kinetics were recorded at 37°C by a DW-2a (American Instrumentation Co., Silver Spring, MD) dual wavelength spectrophotometer operating at 577-600 nm (3 nm slit opening). The 1-cm path length quartz cuvette was filled with 1 mL of air-saturated buffer ([O₂]₀ 0.2 mM [12]) and appropriate amounts of HbO₂ were added to it. The reaction was started by adding 100 mU/mL XO to




FIGURE 1. Scheme of the mechanisms assumed in this work (unbalanced reactions). Abbreviations: OEC, oxygen equilibrium curve; SOD, superoxide dismutase; XO, xanthine oxidase.

the mixture. The rate of $\text{O}_2^{\cdot-}$ production by XAN/XO was measured mixing 100 μM cytochrome c ($E_{550 \text{ nm}}=24.3 \text{ mM}^{-1} \text{ cm}^{-1}$ in the reduced form) with 100 mU/mL XO and 0.5 mM XAN [13]. The Hb- O_2 affinity was measured at pH 7.4, 37°C (Hemox oxygen equilibrium curve analyzer).

RESULTS

Generation of ODFRs and Hb spectra. Under the selected conditions, $\text{O}_2^{\cdot-}$ was generated steadily at 0.045 mM/min for 2 min without initial lag phases. The oxidation of HbO₂ (all types) by 0.5 mM XAN or urate in the absence of XO was <0.25 $\mu\text{M}/\text{min}$ (0.8%/min) and <0.6 $\mu\text{M}/\text{min}$ (2%/min), respectively. The oxidation rate was not limited by shortage of XAN nor O_2 , which was regenerated during the reaction (Fig.1). KCN was added to the medium to convert metHb to CNmetHb.

Spectra analysis in the visible and UV regions (not shown) indicated that HbO₂ was oxidized to CNmetHb. The absorbance change at 700 nm, index of aspecific protein denaturation [14], was <0.003 absorbance units/min under all conditions.

Kinetics. The oxidation of 50 $\mu\text{M}/\text{L}$ sHb (Fig.2) was typically sigmoidal with maximal rate (V_{max})=22.7±2.1 $\mu\text{M}/\text{min}$ (mean±SD, n=10).




FIGURE 2. Kinetics of the oxidation of $50 \mu\text{M}$ stripped HbO_2 by pre-incubated mixtures of XAN/XO (from right to left, 0, 15, 30, 60, 120, 180 and 300 s). Each point represents the mean of 4 to 6 experiments (2% average S.D., not shown for clarity). $\Delta\text{Abs} = \text{Abs}_{577} - \text{Abs}_{600}$. Insert: Half-time ($t_{1/2}$) of the reaction vs the corresponding incubation time.

In the presence of CO, V_{max} was $0.7 \pm 0.3 \mu\text{M}/\text{min}$ ($n=5$, $p < 0.0001$). When sHb was added to pre-incubated XAN/XO (0 to 300 s) the sigmoidal shape disappeared and the oxidation half-time ($t_{1/2}$) decreased as function of the incubation time (Fig.2).

The oxidation of sHb was indistinguishable from that of HbA_0 . Fig.3 shows the increase of $t_{1/2}$ at increasing $[\text{Hb}]$ in the 5 to $150 \mu\text{M}$ range. At all concentrations, $t_{1/2}$ for HbXL99 α was significantly less than for HbA_0 .

In the presence of 50 mM DMTU, scavenger of $\cdot\text{OH}$, V_{max} reduced to $15.1 \pm 1.2 \mu\text{M}/\text{min}$ ($n=6$, $p < 0.0001$). In the presence of 2000 U/ml SOD, scavenger of $\text{O}_2^{\cdot-}$, V_{max} increased by 7-9% ($p = \text{NS}$). The effect of catalase, which quickly decomposes H_2O_2 , was not tested for the inhibition of this enzyme by KCN, but experiments in the absence of KCN demonstrated that catalase prevents the oxidation of HbO_2 (not shown).




FIGURE 3. Halftime ($t_{1/2}$) of the oxidation of HbXL99 α and HbA₀ (virtually the same as stripped Hb) as function of [Hb]. Dashed lines are the 95% confidence limits. The differences were significant at the $p=0.03$, 0.002, 0.0005 and 0.0005 levels (Student's t-test for unpaired data) at the four [Hb]'s.

The P_{50} 's of stripped Hb, HbA₀ and HbXL99 α were 10.0, 11.5 and 30.0 mmHg, respectively, pH 7.4.

DISCUSSION

The XAN/XO system generates reproducible and accurate amounts of ODFRs that oxidize HbO₂ by specific mechanisms which are faster than HbO₂ autooxidation [6]. The oxidation rate depended on [Hb] for all Hb's, but HbXL99 α was always oxidized faster than sHb and HbA₀.

Kinetic data fit a model where the first product of XAN/XO (O₂^{•-}) is unable to oxidize HbO₂, but slowly dismutates to H₂O₂ which is a strong oxidant. Several observations support this view: 1) When HbO₂ was added to pre-incubated mixtures of XAN/XO, the sigmoidal shape of the oxidation curve disappeared and the oxidation was faster; 2) SOD slightly

increased the oxidation rate presumably accelerating the formation of H_2O_2 [15-17]; 3) Catalase prevented the oxidation; 4) DMTU slowed significantly the oxidation rate scavenging the $\cdot OH$ radical.

The dependence of the oxidation rate on $[Hb]$ could be related to the Hb tetramer splitting into dimers and to different reactivities of tetramers and dimers. For a dissociation constant of 0.001 mM [18], the tetrameric form accounts for 92.2% and 64.2% of total Hb at $[Hb]=150$ and $5 \mu M$, respectively. However, the presence of α - α covalent linkage in HbXL99 α should have prevented splitting in this Hb, but nevertheless the kinetics of HbXL99 α depended on $[Hb]$ in the same way as HbA $_0$ and sHb, indicating that it is unlikely that the Hb tetramers and dimers have different reactivities in this model.

The heme appeared the elective site of action of ODFR because the oxidation was nearly abolished when O_2 was replaced by CO, consistently with the slow rate of CO dissociation from HbCO [19]. The faster oxidation of HbXL99 α with respect to that of HbA $_0$ and sHb may thus be explained hypothesizing that the overall reaction is limited by the release of O_2 from HbO $_2$ reflecting different Hb- O_2 affinities of the various tested Hbs.

These data are in essential agreement with the faster autooxidation rate of HbXL99 α with respect to HbXL82 β and HbA [20]. However, we suggest that it is not only during storage that HbXL99 α must be protected from oxidation, but also when infused into a patient, because this Hb is particularly susceptible to oxidation by circulating oxidative factors, especially those arising from reperfusion of ischemic tissues.

ACKNOWLEDGEMENTS

We thank Dr. R.M. Winslow for kindly providing samples of HbA $_0$ and HbXL99 α which have been used for this work. This study was supported by the Target Project Biotechnology and Bioinstrumentation, Rome, Italy.

REFERENCES

- 1] C.T. White, A.J. Murray, J.R. Greene, D.J. Smith, F. Medina, G.T. Makovec, E.J. Martin and R.B. Bolin, Toxicity of human hemoglobin solution infused into rabbits. *J Labor Clin Med* 108:121-131 (1986)

- 2] S.M.H. Sadrzadeh and J.W. Eaton, Hemoglobin-mediated oxidant damage to the central nervous system requires endogenous ascorbate. *J Clin Invest* 82:1510-1515 (1988)
- 3] W.J. Wallace, R.A. Houtchens, J.C. Maxwell and W.S. Caughey, Mechanism of autooxidation for hemoglobins and myoglobins. Promotion of superoxide production by protons and anions. *J Biol Chem* 257:4966-4977 (1982)
- 4] J.A. Watkins, S. Kawanishi and W.S. Caughey, Autoxidation reactions of hemoglobin a free from other red cell components: a minimal mechanism. *Biochem Biophys Res Comm* 132:742-748 (1985)
- 5] S.M.H. Sadrzadeh, E. Graf, S.S. Panter, P.E. Hallaway and J.W. Eaton, Hemoglobin: a biological Fenton reagent. *J Biol Chem* 259:14354-14356 (1984)
- 6] A. Puppo and B. Halliwell, Formation of hydroxyl radicals from hydrogen peroxide in the presence of iron. *Biochem J* 249:185-190 (1988)
- 7] J.M.C. Gutteridge, Iron promoters of the fenton reaction and lipid peroxidation can be released from hemoglobin by peroxidase. *FEBS Letters* 201:291-295 (1986)
- 8] M. Samaja, E. Rovida, M. Niggeler, M. Perrella and L. Rossi Bernardi, The dissociation of carbon monoxide from hemoglobin intermediates. *J Biol Chem* 262:4528-4533 (1987)
- 9] O.W. Van Assendelft, Spectrophotometry of haemoglobin derivatives (Royal Vangorcum Ltd., Assen, NL, 1970).
- 10] S.M. Christensen, F. Medina, R.M. Winslow, S.M. Snell, A. Zegna and M.A. Marini, Preparation of human hemoglobin A₀ for possible use as a substitute. *J Biochem Biophys Meth* 17:143-154 (1988)
- 11] S.R. Snyder, E.V. Welty, R.Y. Walder, L.A. Williams and J.A. Walder, HbxL99 α : a hemoglobin derivative that is cross-linked between the α subunits is useful as a blood substitute. *Proc Natl Acad Sci USA* 84:7280-7284 (1987)
- 12] F.J.W. Roughton and J.W. Severinghaus, Accurate determination of O₂ dissociation curve of human above 98.7% saturation with data on O₂ solubility in unmodified human blood from 0° to 37°C. *J Appl Physiol* 35:861-869 (1973)

- 13] J.M. McCord and I. Fridovich, The reduction of cytochrome c by milk xanthine oxidase. *J Biol Chem* 243:5753-5760 (1968)
- 14] E.A. Rachmilewitz, J. Peisach and W.E. Blumberg, Studies on the stability of oxyhemoglobin A and its constituent chains and their derivatives. *J Biol Chem* 246:3356-3366 (1971)
- 15] K. Yusa and K. Shikama, Oxidation of oxymyoglobin to metmyoglobin with hydrogen peroxide: involvement of ferryl intermediate. *Biochemistry* 26:6684-6688 (1987)
- 16] K.D. Whitburn, The interaction of oxymyoglobin with hydrogen peroxide: the formation of ferrylmyoglobin at moderate excesses of hydrogen peroxide. *Arch Biochem Biophys* 253:419-430 (1987)
- 17] M.R. Prasad, R.M. Engelman, R.M. Jones and D.K. Das, Effects of oxyradicals on oxymyoglobin. Deoxygenation, haem removal and iron release. *Biochem J* 263:731-736 (1989)
- 18] A.D. Barksdale and A. Rosenberg, Thermodynamic characterization of subunit association in liganded ferrohemeoglobin. *J Biol Chem* 253:4881-4885 (1978)
- 19] M. Samaja and E. Rovida, The dissociation of carbon monoxide from the alpha and the subunits of human carbonmonoxy hemoglobin. *Biochem Biophys Res Comm* 148:1196-1201 (1987)
- 20] T. Yang and K.W. Olsen, The effect of crosslinking by bis(3-5-dibromosalicyl) on the autoxidation of hemoglobin. *Biochem Biophys Res Comm* 163:733-738 (1989)