

## Contribute 1

# Multivariate functional data depth measure based on variance-covariance operators

Rachele Biasi, Francesca Ieva, Anna Maria Paganoni, Nicholas Tarabelloni

**Abstract** We introduce a generalization of the simplicial depth measure to multivariate functional data, exploiting the role of the variance-covariance operators in weighting the components that define the depth. We propose the use of this nonparametric method for supervised classification purpose.

## Introduction

In recent times, more and more data coming from various real-life applications can be studied with the paradigm of functional data. This framework requires the development and use of ad-hoc statistical instruments (for an introduction, see [1]). Here we consider multivariate functional data, that means data where each observation is a set of possibly correlated functions. These functions can be viewed as trajectories of stochastic processes defined on a given infinite dimensional functional space, and characterized by a variance covariance operator (see the monograph [3] for more theoretical details). We consider the notion of depth measure for functional data introduced in [7, 8] for univariate functional data, and extended to multivariate functional framework in [6]. Depth measure belongs to the field of non-parametric functional data analysis (see, for instance, [2]) and seems to be promising in classifying curves belonging to different generating processes. According to definition in [6], it is necessary to make a choice of the weights averaging the contribution of each component of the multivariate

---

Rachele Biasi  
Politecnico di Milano, Italy, e-mail: rachele.biasi@mail.polimi.it

Francesca Ieva  
Università degli Studi di Milano, Italy, e-mail: francesca.ieva@unimi.it

Anna Maria Paganoni  
Politecnico di Milano, Italy, e-mail: anna.paganoni@polimi.it

Nicholas Tarabelloni  
Politecnico di Milano, Italy, e-mail: nicholas.tarabelloni@polimi.it

signal to the depth itself. This choice is usually problem-driven, and in general no gold rules have been given so far. We develop a general method for defining such weights. In particular, we propose to choose them taking into account the distance between the estimated covariance operators of the two groups. In fact, the covariance structure of the multivariate functional signals contains information about the reciprocal role of the signal components with respect one to each other. This should be taken into account in measuring the depth of a signal, and in general when comparing signal features with reference traces. In the following, we consider many different distances between covariance operators in the infinite dimensional setting, as discussed in [9], and by means of a simulation study we discuss the behavior and the robustness of the weight choice with respect to different distances and as long as the correlation between components changes. In [4] a wider study of this problem as well as an application to a case study for disease probability prediction from electrocardiographic signals are presented. All the analyses are carried out using R statistical software [10] and the ad-hoc C++/MPI parallel library for computational statistics HPCS (for further details see the website: <https://github.com/ntarabelloni/HPCS>, Code is available upon request) [5].

## 1.1 Multivariate depth measure with weights based on variance covariance operators

Let  $X$  be stochastic process taking values in the space  $\mathcal{C}(I; \mathbb{R}^h)$  of continuous functions  $\mathbf{f} = (f_1, \dots, f_h) : I \rightarrow \mathbb{R}^h$ , where  $I$  is a compact interval of  $\mathbb{R}$ . The multivariate depth measure is defined as

$$MBD_n^J(\mathbf{f}) = \sum_{k=1}^h p_k MBD_{n,k}^J(f_k), \quad p_k > 0 \quad \forall k = 1, \dots, h, \quad \sum_{k=1}^h p_k = 1 \quad (1.1)$$

where for each function  $f_k \in F \subset \mathcal{C}(I; \mathbb{R})$ ,  $k = 1, \dots, h$ , the  $MBD_{n,k}^J(f_k)$  measures the proportion of time interval  $I$  where the graph of  $f_k$  belongs to the envelopes of the  $j$ -tuples  $(f_{i_1;k}, \dots, f_{i_j;k})$ ,  $j = 1, \dots, J$ , extracted from  $F$ . In other words, measuring that the curve  $f_k$  is in the band determined by the  $j$  curves  $(f_{i_1;k}, \dots, f_{i_j;k})$ , means computing

$$MBD_{n,k}^J(f_k) = \sum_{j=2}^J \binom{n}{j}^{-1} \sum_{1 \leq i_1 < i_2 < \dots < i_j \leq n} \tilde{\lambda}\{E(f_k; f_{i_1;k}, \dots, f_{i_j;k})\},$$

where  $E(f_k; f_{i_1;k}, \dots, f_{i_j;k}) = \{t \in I, \min_{r=i_1, \dots, i_j} f_{r;k}(t) \leq f_k(t) \leq \max_{r=i_1, \dots, i_j} f_{r;k}(t)\}$  and  $\tilde{\lambda}(A) = \lambda(A)/\lambda(I)$ ,  $\forall A \subseteq I$ , with  $\lambda$  the Lebesgue measure on  $I$ . Statistical properties of the depth measure defined in (1.1) as well as inferential tools based on this concept are detailed in [6]. For subsequent analyses we fixed  $J = 2$  in (1.1) once and for all.

Let us consider two different multivariate stochastic processes  $\mathbf{X}$  and  $\mathbf{Y}$ . Without loss of generality we assume that their means are equal to zero. We indicate their covariance operators with  $\mathcal{V}_{\mathbf{X}}$  and  $\mathcal{V}_{\mathbf{Y}}$ . The kernel of the covariance operator of  $\mathbf{X}$ ,  $V_{\mathbf{X}}(s, t)$ , is defined by

$$V_{\mathbf{X}}(s, t) = \mathbb{E}[\mathbf{X}(s) \otimes \mathbf{X}(t)], \quad s, t \in I$$

where  $\otimes$  is an outer product in  $\mathbb{R}^h$ . So  $V_{\mathbf{X}}(s, t)$  is a  $h \times h$  matrix, whose elements will be denoted as  $V_{\mathbf{X}}^{kq}(s, t)$ , for  $k, q = 1, \dots, h$ . Analogously for  $V_{\mathbf{Y}}(s, t)$ .

We consider the distances introduced in [9], generalizing them to the case of non necessarily positive definite operators, in fact we are interested in quantifying the distance also between the inter-component blocks, i.e.  $V_{\mathbf{X}}^{kq}(s, t)$  and  $V_{\mathbf{Y}}^{kq}(s, t)$  when  $k \neq q$ . Let  $d(V, W)$  denote a distance between two operators. We compute for each  $k = 1, \dots, h$  the quantity  $d_k = \sum_{q=1}^h d(V_{\mathbf{X}}^{kq}(s, t), V_{\mathbf{Y}}^{kq}(s, t))$ , considering the following distances:

- –  **$L^2$  distance**

$$d_L(V, W) = \sqrt{\int_I \int_I (v(s, t) - w(s, t))^2 ds dt}, \quad (1.2)$$

where  $v(s, t)$  and  $w(s, t)$  are the kernels of the operators  $V$  and  $W$  respectively.

- – **Spectral distance**

$$d_S(V, W) = |\lambda_1|, \quad (1.3)$$

where  $|\lambda_1|$  is the maximum eigenvalue of the difference operator  $V - W$ .

- – **Square root pseudo distance**

$$d_R(V, W) = \left\| |V|^{\frac{1}{2}} - |W|^{\frac{1}{2}} \right\|_{HS}, \quad (1.4)$$

where the Hilbert-Schmidt norm of an Hilbert-Schmidt compact operator  $T$  is  $\|T\|_{HS} = \sqrt{\text{trace} T^* T}$ ,  $T^*$  is the adjoint operator of  $T$ ,  $|T|^{\frac{1}{2}}$  is such that  $|T|^{\frac{1}{2}} v_k = |\lambda_k|^{\frac{1}{2}} v_k$ ,  $\{v_k\}_k$  is the orthonormal basis of  $L^2$  of the eigenfunctions of  $T$  and  $\{\lambda_k\}_k$  is the sequence of the related eigenvalues.

- – **Frobenius distance**

$$d_F(V, W) = \|V - W\|_{HS} = \sqrt{\text{trace}(V - W)^*(V - W)}. \quad (1.5)$$

- – **Procrustes pseudo distance**

$$d_P(V, W) = d_P(|V|, |W|) = \inf_{R \in O(L^2(I))} \|L_1 - L_2 R\|_{HS}, \quad (1.6)$$

where  $O(L^2(I))$  is the space of all unitary operators on  $L^2(I)$  and  $L_1$  and  $L_2$  are such that  $V = L_1 L_1^*$  and  $W = L_2 L_2^*$ .






Figure 1.1: Weights of MBD in the simulation study, determined through the distances between variance-covariance operators previously defined.

## Bibliography

- [1] Silverman, B.W., Ramsay, J.O. (2005), *Functional Data Analysis*, Springer.
- [2] Ferraty, F., Vieu, P. (2006), *Nonparametric functional data analysis: theory and practice*, Springer.
- [3] Horváth, L., Kokoszka, P. (2012), *Inference for functional data with applications*, Springer.
- [4] Biasi, R., Ieva, F., Paganoni, A.M., Tarabelloni, N. (2013), Use of depth measure for multivariate functional data in disease prediction: an application to electrocardiographic signals *Tech. Rep. MOX, Math. Dept.*, Politecnico di Milano. [Online] <http://mox.polimi.it/it/progetti/pubblicazioni/quaderni/54-2013.pdf>
- [5] Tarabelloni, N. (2013) Tools for computational statistics coded in C++, [online] <https://github.com/ntarabelloni/HPCS>
- [6] Ieva, F., Paganoni, A.M. (2013a), Depth Measures for Multivariate Functional Data, *Communication in Statistics - Theory and Methods*, **42**, 7, 1265–1276.
- [7] Lopez-Pintado, S., Romo, J. (2007), Depth-based inference for functional data, *Computational Statistics & Data Analysis*, **51**, 10, 4957–4968.
- [8] Lopez-Pintado, S., Romo, J. (2009), On the Concept of Depth for Functional Data, *Journal of the American Statistical Association*, **104**, 486, 718–734.
- [9] Pigoli, D., Aston, J.A.D., Dryden, I.L., Secchi, P. (2012), Distances and Inference for Covariance Functions *Tech. Rep. MOX, Math. Dept.*, Politecnico di Milano. [Online] <http://mox.polimi.it/it/progetti/pubblicazioni/quaderni/35-2012.pdf>
- [10] R Development Core Team (2009), R: A language and environment for statistical computing, R Foundation for Statistical Computing, Vienna, Austria. [online] <http://www.R-project.org>