

A condition for a perfect-fluid space-time to be a generalized Robertson-Walker space-time

Carlo Alberto Mantica, Luca Guido Molinari, and Uday Chand De

Citation: *Journal of Mathematical Physics* **57**, 022508 (2016); doi: 10.1063/1.4941942

View online: <http://dx.doi.org/10.1063/1.4941942>

View Table of Contents: <http://scitation.aip.org/content/aip/journal/jmp/57/2?ver=pdfcov>

Published by the [AIP Publishing](#)

Articles you may be interested in

[Geometrization conditions for perfect fluids, scalar fields, and electromagnetic fields](#)

J. Math. Phys. **56**, 072503 (2015); 10.1063/1.4926952

[A fully covariant information-theoretic ultraviolet cutoff for scalar fields in expanding Friedmann Robertson Walker spacetimes](#)

J. Math. Phys. **54**, 022301 (2013); 10.1063/1.4790482

[Brownian motion in Robertson–Walker spacetimes from electromagnetic vacuum fluctuations](#)

J. Math. Phys. **50**, 062501 (2009); 10.1063/1.3133946

[Variational approach to Robertson–Walker spacetimes with homogeneous scalar fields](#)

J. Math. Phys. **47**, 112505 (2006); 10.1063/1.2383068

[A space–time in toroidal coordinates](#)

J. Math. Phys. **44**, 3046 (2003); 10.1063/1.1580999



Computational Science and Engineering

Computing
SCIENCE ENGINEERING

Scientific Software Communities

PHYSICS
AIP
JOURNAL OF MATHEMATICAL PHYSICS

Broaden your impact to scientists and engineers in 50+ societies. Submit your computational article to *CISE*.

A condition for a perfect-fluid space-time to be a generalized Robertson-Walker space-time

Carlo Alberto Mantica,^{1,2,a)} Luca Guido Molinari,^{1,3,b)} and Uday Chand De^{4,a)}

¹Physics Department, Università degli Studi di Milano, Via Celoria 16, 20133 Milano, Italy

²I.I.S. Lagrange, Via L. Modignani 65, 20161 Milano, Italy

³I.N.F.N. sez. Milano, Via Celoria 16, 20133 Milano, Italy

⁴Department of Pure Mathematics, University of Calcutta, 35 Ballygaunge Circular Road, Kolkata 700019, West Bengal, India

(Received 28 September 2015; accepted 2 February 2016; published online 19 February 2016)

A perfect-fluid space-time of dimension $n \geq 4$, with (1) irrotational velocity vector field and (2) null divergence of the Weyl tensor, is a generalised Robertson-Walker space-time with an Einstein fiber. Condition (1) is verified whenever pressure and energy density are related by an equation of state. The contraction of the Weyl tensor with the velocity vector field is zero. Conversely, a generalized Robertson-Walker space-time with null divergence of the Weyl tensor is a perfect-fluid space-time. © 2016 AIP Publishing LLC. [<http://dx.doi.org/10.1063/1.4941942>]

I. INTRODUCTION

Standard cosmology is modelled on Robertson-Walker metrics for the high symmetry imposed on space-time by the cosmological principle (spatial homogeneity and isotropy). A wide generalization is the “generalized Robertson-Walker spacetimes,” introduced in 1995 by Alías, Romero, and Sánchez.^{1,2}

Definition 1.1. An n -dimensional Lorentzian manifold is a generalized Robertson-Walker (GRW) space-time if locally the metric may take the following form:

$$ds^2 = -dt^2 + q(t)^2 g_{\alpha\beta}^*(x_2, \dots, x_n) dx^\alpha dx^\beta, \quad \alpha, \beta = 2 \dots n, \quad (1)$$

that is, it is the warped product $(-1) \times q^2 \mathcal{M}^*$, where \mathcal{M}^* is an $(n-1)$ -dimensional Riemannian manifold. If \mathcal{M}^* has dimension 3 and has constant curvature, the space-time is a Robertson-Walker space-time.

Such spaces include the Einstein-de Sitter space-time, the Friedmann cosmological models, the static Einstein space-time, and the de Sitter space-time. They are the stage for treatment of small perturbations of the Robertson-Walker metric. We refer to the works by Romero *et al.*,²⁹ Sánchez,^{30,31} and Gutiérrez and Olea²⁰ for a comprehensive presentation of geometric properties and physical motivations.

Recently, Bang-Yen Chen proved the following deep result:⁸ A Lorentzian manifold of dimension $n \geq 4$ is a GRW space-time if and only if it admits a time-like vector, $X^j X_j < 0$, such that

$$\nabla_k X_j = \rho g_{kj}. \quad (2)$$

According to Yano,³⁷ a vector field v is *torse-forming* if $\nabla_k v_j = \omega_k v_j + f g_{jk}$, where f is a scalar function and ω_k is a 1-form. Its properties in pseudo-Riemannian manifolds were studied by Mikeš and Rachůnek.^{26,28} The vector is named *concircular* if ω_k is a gradient (or locally a gradient); in this case, v can be rescaled to a vector X with property (2).²⁶

Mantica *et al.*²⁵ proved two sufficient conditions for a Lorentzian manifold of dimension $n \geq 4$ to be a GRW space-time: the first one is the existence of a concircular vector such that $u^i u_i = -1$. The other sufficient condition restricts the Weyl and Ricci tensors: $\nabla_m C_{jkl}{}^m = 0$ and $R_{ij} = B u_i u_j$, where B is a scalar field and u is a time-like vector field.

^{a)}E-mail addresses: carloalberto.mantica@libero.it and uc_de@yahoo.com

^{b)}Author to whom correspondence should be addressed. Electronic mail: luca.molinari@mi.infn.it

Lorentzian manifolds with a Ricci tensor of the form

$$R_{ij} = Ag_{ij} + Bu_iu_j, \quad (3)$$

where A and B are scalar fields and $u_iu^i = -1$, are often named *perfect fluid space-times*. It is well known that any Robertson-Walker space-time is a perfect fluid space-time,²⁷ and for $n = 4$, a GRW space-time is a perfect fluid if and only if it is a Robertson-Walker space-time.

Form (3) of the Ricci tensor is implied by Einstein's equation if the energy-matter content of space-time is a perfect fluid with velocity vector field u . The scalars A and B are linearly related to the pressure p and the energy density μ measured in the locally comoving inertial frame. They are not independent because of the Bianchi identity $\nabla^m R_{im} = \frac{1}{2}\nabla_i R$, which translates into

$$\nabla^m(Bu_ju_m) = \frac{1}{2}\nabla_j[(n-2)A - B]. \quad (4)$$

Geometers identify special form (3) of the Ricci tensor as the defining property of quasi-Einstein manifolds (with any metric signature). The Riemannian ones were introduced by Defever and Deszcz in 1991¹³ (see also Refs. 15 and 16, and Chaki *et al.*⁷). In Ref. 14, Deszcz proved that a quasi-Einstein Riemannian manifold, with null Weyl tensor and few other conditions, is a warped product $(+1) \times q^2 \mathcal{M}^*$, where \mathcal{M}^* is an $(n-1)$ -dimensional Riemannian manifold of constant curvature.

Pseudo-Riemannian quasi-Einstein spaces arose in the study of exact solutions of Einstein's equations. Robertson-Walker space-times are quasi-Einstein (see Refs. 4 and 34 and references therein).

Shepley and Taub studied a perfect-fluid space-time in dimension $n = 4$, with equation of state $p = p(\mu)$ and the additional condition that the Weyl tensor has null divergence, $\nabla_m C_{jkl}{}^m = 0$. They proved the following: the space-time is conformally flat $C_{jklm} = 0$; the metric is Robertson-Walker; the flow is irrotational, shear-free, and geodesic.³³

A related result was obtained by Sharma³² (corollary p. 3584): if a perfect-fluid space-time in $n = 4$ with $\nabla_m C_{jkl}{}^m = 0$ admits a proper conformal Killing vector, i.e., $\nabla_i X_j + \nabla_j X_i = 2\rho g_{ij}$, then it is conformally flat ($C_{ijkl} = 0$). In the framework of Yang's gravitational theory, Guilfoyle and Nolan proved that a $n = 4$ perfect fluid space-time with $p + \mu \neq 0$ is a Yang pure space (i.e., $\nabla_m C_{jkl}{}^m = 0$ and $\nabla_k R = 0$) if and only if it is a Robertson-Walker space-time.¹⁹

Coley proved that any perfect fluid solution of Einstein's equations satisfying a barotropic equation of state $p = p(\mu)$ and $p + \mu \neq 0$, which admits a proper conformal Killing vector parallel to the fluid 4-velocity, is locally a Friedmann-Robertson-Walker model.¹⁰

De *et al.*¹² showed that $n = 4$ conformally flat almost pseudo-Ricci-symmetric space-times, i.e., $\nabla_k R_{ij} = (a_k + b_k)R_{ij} + a_j R_{ik} + a_i R_{jk}$, are Robertson-Walker space-times.

Riemannian spaces equipped with a torse-forming vector field were studied by Yano as early as 1944;³⁷ his results were extended to pseudo-Riemannian spaces by Sinyukov.³⁵ They showed that the existence of such a vector implies the following local shape of the metric: $ds^2 = \pm(dx^1)^2 + F(x_1, \dots, x_n)d\tilde{s}^2$, where $d\tilde{s}^2$ is the metric of the submanifold parametrized by x_2, \dots, x_n . If the vector field is concircular (then it is rescalable to $\nabla_k X_j = \rho g_{kj}$) then F is a function of x_1 only.

De and Ghosh¹¹ showed that if $R_{ij} = Ag_{ij} + Bu_iu_j$ with u_i closed and $C_{ijkl} = 0$, then u is a concircular vector. The results were extended by Mantica and Suh to pseudo-Z-symmetric spaces²⁴ and to weakly Z-symmetric spaces.²³

In this paper, the theorem by Shepley and Taub is generalised to perfect-fluid space-times of dimension $n \geq 4$. The converse is also proven: a GRW space-time with $\nabla_m C_{jkl}{}^m = 0$ is a perfect-fluid space-time. In the conclusion, some consequences for physics are presented.

II. THE THEOREM

Theorem 2.1. *Let \mathcal{M} be perfect fluid-space-time, i.e., a Lorentzian manifold (of dimension $n > 3$) with Ricci tensor $R_{kl} = Ag_{kl} + Bu_ku_l$, where A and B are scalar fields, u is a time-like unit vector field $u^i u_j = -1$.*

If $\nabla_k u_j - \nabla_j u_k = 0$ (u is closed) and if $\nabla_m C_{jkl}{}^m = 0$, then

- (i) *u is a concircular vector and it is rescalable to a time-like conformal Killing vector X such that*

$$\nabla_k X_j = \rho g_{kj} \quad \text{and} \quad \nabla_k \rho = \frac{A-B}{1-n} X_k; \quad (5)$$

- (ii) \mathcal{M} is a generalised Robertson-Walker space-time whose sub-manifold (\mathcal{M}^*, g^*) is a Riemannian Einstein space.
- (iii) $C_{jklm} u^m = 0$.

Proof. The condition $\nabla_m C_{jkl}{}^m = 0$ implies: $\nabla_k R_{jl} - \nabla_l R_{jk} = \frac{1}{2(n-1)}(g_{jl} \nabla_k R - g_{jk} \nabla_l R)$. With the explicit form of the Ricci tensor, it becomes

$$\nabla_k (Bu_j u_l) - \nabla_l (Bu_j u_k) = -\frac{g_{jl} \nabla_k \gamma - g_{jk} \nabla_l \gamma}{2(n-1)} \quad (6)$$

being $\gamma = (n-2)A + B$. By transvecting with $u^j u^l$ and using $u^l \nabla_k u_l = 0$, we obtain

$$(\nabla_k + u_k u^l \nabla_l) B + Bu^l \nabla_l u_k = \frac{1}{2(n-1)} (\nabla_k + u_k u^l \nabla_l) \gamma. \quad (7)$$

Contraction of the identity (4) with u^j gives $-B \nabla_m u^m = \frac{1}{2} u^m \nabla_m \gamma$, which rewrites identity (4) as

$$(\nabla_k + u_k u^i \nabla_i) B + Bu^m \nabla_m u_k = \frac{1}{2} (\nabla_k + u_k u^i \nabla_i) \gamma. \quad (8)$$

Equations (7) and (8) imply

$$(\nabla_j + u_j u^k \nabla_k) \gamma = 0, \quad (9)$$

$$(\nabla_j + u_j u^k \nabla_k) B + Bu^m \nabla_m u_j = 0. \quad (10)$$

Contraction of (6) with u^l gives

$$\begin{aligned} & -u_j (\nabla_k + u_k u^l \nabla_l) B - B \nabla_k u_j - u_j Bu^l \nabla_l u_k - u_k Bu^l \nabla_l u_j \\ & = -\frac{1}{2(n-1)} (u_j \nabla_k - g_{jk} u^l \nabla_l) \gamma. \end{aligned}$$

By use of Eq. (10), it simplifies to

$$B (\nabla_k + u_k u^m \nabla_m) u_j = \frac{1}{2(n-1)} (u_j \nabla_k - g_{jk} u^l \nabla_l) \gamma. \quad (11)$$

If u is closed, it is $u^m \nabla_m u_j = u^m \nabla_j u_m = 0$. Eq. (11) simplifies and shows that u is a torse-forming vector,

$$\nabla_k u_j = \frac{\nabla_k \gamma}{2B(n-1)} u_j - \frac{u^m \nabla_m \gamma}{2B(n-1)} g_{kj} \equiv \omega_k u_j + f g_{kj}. \quad (12)$$

Let us show that u is a concircular vector, i.e., that ω_k is closed,

$\nabla_j \omega_k - \nabla_k \omega_j = -\frac{1}{B} (\omega_k \nabla_j - \omega_j \nabla_k) B = -(\omega_k u_j - \omega_j u_k) u^m \nabla_m B$ by (10). Eq. (9) gives the relation $\omega_k = -u_k u^m \omega_m$, then $\omega_k u_j - \omega_j u_k = 0$.

Being closed, ω_k is locally the gradient of a scalar function: $\omega_k = \nabla_k \sigma$. Let $X_l = u_l e^{-\sigma}$; we have $\nabla_k X_l = e^{-\sigma} (-u_l \nabla_k \sigma + \omega_k u_l + f g_{kl}) = e^{-\sigma} f g_{kl}$ and, consequently,

$$\nabla_k X_l = \rho g_{kl} \quad (13)$$

being $\rho = e^{-\sigma} f$ and $X_j X^j = -e^{-2\sigma} < 0$ (time-like vector). The symmetrized equation $\nabla_k X_j + \nabla_j X_k = 2\rho g_{kj}$ shows that X_j is a conformal Killing vector.³⁴

According to Chen's theorem, (13) is a sufficient condition for the space-time to be a GRW. In appropriate coordinates, $\mathcal{M} = (-1) \times q^2 \mathcal{M}^*$. The additional condition $\nabla_m C_{jkl}{}^m = 0$ assures that the $(n-1)$ -dimensional Riemannian space \mathcal{M}^* is an Einstein space, by Gębarowski's lemma.¹⁸

Another derivative and the Ricci identity give: $(\nabla_j \nabla_k - \nabla_k \nabla_j) X_l = R_{jkl}{}^m X_m = g_{kl} \nabla_j \rho - g_{jl} \nabla_k \rho$. Contraction with g^{kl} : $R_{jm} X^m = (1-n) \nabla_j \rho$. However, for the perfect fluid (3), $R_{jm} X^m = (A-B) X_j$, then

$$\nabla_j \rho = \frac{A-B}{1-n} X_j, \quad (14)$$

(this is an explicit expression for a relation obtained by Chen). Therefore, if $A \neq B$, the conformal killing vector X is proper; if $A = B$, it is homothetic. Moreover,

$$R_{jklm}X^m = \frac{A - B}{1 - n}(X_jg_{kl} - X_kg_{jl}). \quad (15)$$

The Weyl tensor is

$$C_{jklm} = R_{jklm} + \frac{1}{n-2}(g_{jm}R_{kl} - g_{km}R_{jl} + R_{jm}g_{kl} - R_{km}g_{jl}) - \frac{(g_{jm}g_{kl} - g_{mk}g_{jl})R}{(n-1)(n-2)}.$$

The previous equations and a little algebra imply that $C_{jklm}X^m = 0$, so that $C_{jkl}{}^m u_m = 0$. It follows that the Weyl tensor is purely electric.²¹

In $n = 4$ the condition is equivalent to $u_i C_{jklm} + u_j C_{kilm} + u_k C_{ijlm} = 0$ (see Lovelock and Rund²² page 128). Multiplication by u^i gives $C_{ijkl} = 0$. \square

Remark 2.2. Eq. (9) gives $u_j \nabla_k \gamma = u_k \nabla_j \gamma$. In the antisymmetric part of Eq. (11), $B(\nabla_k u_j - \nabla_j u_k) + B(u_k u^m \nabla_m u_j - u_j u^m \nabla_m u_k) = 0$, the last terms are replaced with the help of (10) to give:

$$\nabla_k (B u_j) = \nabla_j (B u_k). \quad (16)$$

Remark 2.3. The case $A = 0$, i.e., $R_{ij} = B u_i u_j$, was studied in Ref. 25. Since $\gamma = -B$, the property $u_j \nabla_k \gamma = u_k \nabla_j \gamma$ and (16) imply that u is closed.

If $A \neq 0$, the condition that u is closed is necessary for proving the theorem. However, if a one-to-one differentiable relation $A(x) = F(B(x))$ exists, one proves that u is closed.

Remark 2.4. In Refs. 5, 6, and 17, a notion of quasi-Einstein manifold different from (3) was introduced. It emerges from generalizations of Ricci solitons. More generally, they defined a generalized quasi-Einstein manifold by the condition

$$R_{ij} + \nabla_i \nabla_j \theta - \eta (\nabla_i \theta) (\nabla_j \theta) = \lambda g_{ij}, \quad (17)$$

where θ, η, λ are smooth functions. If $\lambda = \text{const}$ and $\eta = 0$, it is named gradient Ricci soliton, if $\lambda = \text{const}$. and $\eta = \text{const}$., it is named quasi-Einstein.

In the present case, the condition that u is closed means that locally $u_k = \nabla_k \theta$, for some function θ . Then (3) takes the form $R_{ij} = A g_{ij} + B (\nabla_i \theta) (\nabla_j \theta)$. At the same time, Eq. (12) can be written $\nabla_i \nabla_j \theta = f (\nabla_i \theta) (\nabla_j \theta) + f g_{ij}$ (since $f = -u^k \omega_k$ and $\omega_i = -u_i u^k \omega_k$ by Eq. (9), it is $\omega_i = f u_i$). The sum of the equations yields a Ricci tensor of the form (17) with $\lambda = A + f$ and $\eta = B + f$, i.e., the manifold is generalized quasi-Einstein in the sense of Refs. 5, 6, and 17. A gradient Ricci soliton is recovered if $A + f = \text{const}$. and $B + f = 0$.

In Ref. 5, it was proven that locally conformally flat Lorentzian quasi-Einstein manifolds are globally conformally equivalent to a space form, or locally isometric to a warped product of Robertson-Walker type, or a pp-wave. Catino⁶ proved that a complete (i.e., $A + f$ is a smooth function) generalized quasi-Einstein Riemannian manifold with harmonic Weyl tensor and zero radial curvature, is locally a warped product with $(n - 1)$ dimensional Einstein fibers.

An inverse statement of the theorem is proven.

Theorem 2.5. A generalized Robertson-Walker space-time with $\nabla_m C_{jkl}{}^m = 0$ is a quasi-Einstein space-time.

Proof. A GRW is characterized by the metric (1). The explicit form of the Ricci tensor R_{ij} is reported, for example, in Arslan *et al.*³: $R_{1\alpha} = R_{\alpha 1} = 0$,

$$R_{11} = -(n-1) \frac{q'}{q}, \quad R_{\alpha\beta} = R_{\alpha\beta}^* + g_{\alpha\beta}^* [q'^2(n-2) + qq''], \quad \alpha, \beta = 2 \dots n.$$

Gębarowski proved that $\nabla_m C_{jkl}{}^m = 0$ if and only if $R_{\alpha\beta}^* = g_{\alpha\beta}^* \frac{R^*}{n-1}$, then

$$R_{\alpha\beta} = g_{\alpha\beta}^* \left[\frac{R^*}{n-1} + q'^2(n-2) + qq'' \right].$$

Following the trick in Ref. 9, in the local frame where (1) holds, define the vector $u^1 = 1$ and $u^\alpha = 0$ ($u_1 = -1$). It is $u_j u^j = -1$ in any frame. The components of the Ricci tensor gain the covariant expression $R_{ij} = Ag_{ij} + Bu_i u_j$, where

$$A = \frac{1}{q^2} \left[\frac{R^*}{n-1} + q'^2(n-2) + qq'' \right], \quad B = -(n-1) \frac{q'}{q} + A. \quad (18)$$

The expression is such in all coordinate frames and characterizes a quasi-Einstein Lorentzian manifold. \square

III. SOME NOTES ON PHYSICS

We transpose some of the results to physics (we use units $c = 1$). Consider a perfect fluid with energy momentum tensor $T_{ij} = (p + \mu)g_{ij} + \mu u_i u_j$, where u_j is the velocity vector field, p is the isotropic pressure field and μ is the energy density. By Einstein's equations $R_{ij} - \frac{1}{2}Rg_{ij} = \kappa T_{ij}$ ($\kappa = 8\pi G$ is the gravitational constant) the Ricci tensor is

$$R_{ij} = \kappa(p + \mu)u_i u_j + \kappa \frac{p - \mu}{2 - n} g_{ij}.$$

Comparison with the form (3) identifies $A = \kappa(p - \mu)/(2 - n)$, $B = \kappa(p + \mu)$. Then $\gamma = (n - 2)A + B = 2\kappa\mu$.

As is well known (see Wald³⁶) in general relativity, the equations of motion $\nabla_k T^{kj} = 0$ result from the Bianchi identity in Einstein's equations. For a perfect fluid, the projection along u and its complementary part are

$$u^k \nabla_k \mu + (p + \mu) \nabla_k u^k = 0, \quad (19)$$

$$(\nabla_j + u_j u^k \nabla_k) p + (p + \mu) u^k \nabla_k u_j = 0. \quad (20)$$

By taking into account the results of Sec. II, we prove the following:

Proposition 3.1. *A perfect fluid space-time in dimension $n \geq 4$, with differentiable equation of state $p = p(\mu)$, $p + \mu \neq 0$, and with null divergence of the Weyl tensor, $\nabla_m C_{jkl}{}^m = 0$, is a generalized Robertson-Walker space-time.*

The velocity vector field is irrotational ($\nabla_k u_l - \nabla_l u_k = 0$), geodesic ($u^k \nabla_k u_j = 0$), and it annihilates the Weyl tensor ($C_{jkl}{}^m u_m = 0$).

Proof. We prove that u is irrotational and geodesic. Then, by the main theorem 2.1, it follows that the manifold is a generalized Robertson-Walker space-time and that u annihilates the Weyl tensor.

If $p'(\mu) \neq 0$, then $\nabla_k p = p'(\mu) \nabla_k \mu$. The eqs. $u_j \nabla_k \gamma = u_k \nabla_j \gamma$ (see remark 2.2) and (16) become: $u_j \nabla_k \mu = u_k \nabla_j \mu$ and $\nabla_j [(p + \mu)u_k] = \nabla_k [(p + \mu)u_j]$. Being $\nabla_k p = p'(\mu) \nabla_k \mu$, it follows that $\nabla_k u_j = \nabla_j u_k$.

Eq. (9) is $\nabla_j \mu + u_j u^m \nabla_m \mu = 0$, and translates to $\nabla_j p + u_j u^m \nabla_m p = 0$. This is used in (20) to annihilate the first term. The equation of a geodesic is obtained: $(p + \mu) u^k \nabla_k u_j = 0$. If $\nabla_k p = 0$, Eq. (19) again gives $(p + \mu) u^k \nabla_k u_j = 0$. \square

The special case $A = B$ in (14) characterizes a homothetic conformal Killing field ($\nabla_j X_k = \rho g_{jk}$ with $\nabla_j \rho = 0$). In terms of pressure and density, this means

$$p = \frac{3 - n}{n - 1} \mu,$$

which, when $n = 4$, is $p = -\mu/3$.

¹ Alías, L. J., Romero, A., and Sánchez, M., "Uniqueness of complete spacelike hypersurfaces of constant mean curvature in generalized Robertson-Walker space-times," *Gen. Relativ. Gravitation* **27**(1), 71–84 (1995).

² Alías, L., Romero, A., and Sánchez, M., "Compact spacelike hypersurfaces of constant mean curvature in generalized Robertson-Walker spacetimes," in *Geometry and Topology of Submanifolds VII*, edited by Dillen, F. (World Scientific, River Edge, NJ, USA, 1995), pp. 67–70.

- ³ Arslan, K., Deszcz, R., Ezentaş, R., Hotłoś, M., and Murathan, C., "On generalized Robertson-Walker spacetimes satisfying some curvature condition," *Turk. J. Math.* **38**, 353–373 (2014).
- ⁴ Beem, J. K., Ehrlich, P. E., and Easley, K. L., *Global Lorentzian Geometry*, 2nd ed. Pure and Applied Mathematics (Marcel Dekker, New York, 1996), Vol. 202.
- ⁵ Brozos-Vázquez, M., García-Río, E., and Gavino Fernández, S., "Locally conformally flat Lorentzian quasi-Einstein manifolds," *Monatsh. Math.* **173**, 175–186 (2014).
- ⁶ Catino, G., "Generalized quasi-Einstein manifolds with harmonic Weyl tensor," *Math. Z.* **271**(3-4), 751–756 (2012).
- ⁷ Chaki, M. C. and Maity, R. K., "On quasi Einstein manifolds," *Publ. Math. Debrecen* **57**, 257–306 (2000).
- ⁸ Chen, B.-Y., "A simple characterization of generalized Robertson-Walker manifolds," *Gen. Relativ. Gravitation* **46**, 1833 (2014).
- ⁹ Chojnacka-Dulas, J., Deszcz, R., Głogowska, M., and Prvanovic, M., "On warped product manifolds satisfying some curvature conditions," *J. Geom. Phys.* **74**, 328–341 (2013).
- ¹⁰ Coley, A. A., "Fluid spacetimes admitting a conformal killing vector parallel to the velocity vector," *Classical Quantum Gravity* **8**, 955–968 (1991).
- ¹¹ De, U. C. and Ghosh, S. K., "On conformally flat pseudo-symmetric spaces," *Balk. J. Geom. Appl.* **5**(2), 61–64 (2000), <http://www.emis.de/journals/BJGA/v05n2/B05-2-DE.pdf>.
- ¹² De, A., Özgür, C., and De, U. C., "On conformally flat almost pseudo-Ricci symmetric spacetimes," *Int. J. Theor. Phys.* **51**(9), 2878–2887 (2012).
- ¹³ Defever, F. and Deszcz, R., "On semi-Riemannian manifolds satisfying a condition $R \cdot R = Q(S, R)$," in *Geometry and Topology of Submanifolds III* (World Scientific Publishing, Singapore, 1991), pp. 108–130.
- ¹⁴ Deszcz, R., "On conformally flat Riemannian manifolds satisfying certain curvature conditions," *Tensor (N.S.)* **49**, 134–145 (1990).
- ¹⁵ Deszcz, R., Dillen, F., Verstraelen, L., and Vrancken, L., "Quasi-Einstein totally real submanifolds of the nearly Kähler 6-sphere," *Tohoku Math. J.* **51**(4), 461–478 (1999).
- ¹⁶ Deszcz, R., Głogowska, M., Hotłoś, M., and Sentürk, Z., "On certain quasi-Einstein semisymmetric hypersurfaces," *Ann. Univ. Sci. Budapest Eötvös Sect. Math.* **41**, 151–164 (1998).
- ¹⁷ Gavino-Fernández, S., "The geometry of Lorentzian Ricci solitons," Ph.D. thesis, Publicaciones del Departamento de Geometría y Topología, Universidad de Santiago de Compostela, 2012.
- ¹⁸ Gębarowski, A., "On nearly conformally symmetric warped product spacetimes," *Soochow J. Math.* **20**(1), 61–75 (1994).
- ¹⁹ Guilfoyle, B. S. and Nolan, B. C., "Yang's gravitational theory," *Gen. Relativ. Gravitation* **30**(3), 473–495 (1998).
- ²⁰ Gutiérrez, M. and Olea, B., "Global decomposition of a Lorentzian manifold as a generalized Robertson-Walker space," *Differ. Geom. Appl.* **27**, 146–156 (2009).
- ²¹ Hervik, S., Ortaggio, M., and Wylleman, L., "Minimal tensors and purely electric and magnetic spacetimes of arbitrary dimensions," *Classical Quantum Gravity* **30**, 165014 (2013).
- ²² Lovelock, D. and Rund, H., *Tensors, Differential Forms and Variational Principles*, Reprinted Edition (Dover, 1988).
- ²³ Mantica, C. A. and Molinari, L. G., "Weakly Z symmetric manifolds," *Acta Math. Hung.* **135**(1-2), 80–96 (2012).
- ²⁴ Mantica, C. A. and Suh, Y. J., "Pseudo Z symmetric Riemannian manifolds with harmonic curvature tensors," *Int. J. Geom. Methods Mod. Phys.* **9**(1), 1250004 (21pp) (2012).
- ²⁵ Mantica, C. A., Suh, Y. J., and De, U. C., "A note on generalized Robertson-Walker spacetimes," *Int. J. Geom. Methods Mod. Phys.* (submitted), private communication (2015).
- ²⁶ Mikeš, J. and Rachůnek, L., "Torse-forming vector fields in T-symmetric Riemannian spaces," in *Steps in Differential Geometry, Proceedings of the Colloquium on Differential Geometry, Debrecen, July 25-30, 2000* (Debrecen, Hungary), edited by L. Kozma, P. T. Nagy, and L. Tamássy, pp. 219–229, http://www.emis.de/proceedings/CDGD2000/pdf/K_MikesRach.pdf.
- ²⁷ O'Neill, B., *Semi-Riemannian Geometry with Applications to Relativity* (Academic Press, New York, 1983).
- ²⁸ Rachůnek, L. and Mikeš, J., "On tensor fields semi-conjugated with torse-forming vector fields," *Acta Univ. Palacki. Olomuc., Fac. rer. nat. Mathematica* **44**, 151–160 (2005), <http://kma.upol.cz/data/xinha/ULOZISTE/ActaMath/1305.pdf>.
- ²⁹ Romero, A., Rubio, R. N., and Salamanca, J. J., "Uniqueness of complete maximal hypersurfaces in spatially parabolic generalised Robertson-Walker space times," *Classical Quantum Gravity* **30**(11), 115007 (2013).
- ³⁰ Sánchez, M., "On the geometry of generalized Robertson-Walker spacetimes: Geodesics," *Gen. Relativ. Gravitation* **30**, 915–932 (1998).
- ³¹ Sánchez, M., "On the geometry of generalized Robertson-Walker spacetimes: Curvature and killing fields," *Gen. Relativ. Gravitation* **31**, 1–15 (1999).
- ³² Sharma, R., "Proper conformal symmetries of space-times with divergence-free Weyl tensor," *J. Math. Phys.* **34**, 3582–3587 (1993).
- ³³ Shepley, L. C. and Taub, A. H., "Space-times containing perfect fluids and having a vanishing conformal divergence," *Commun. Math. Phys.* **5**, 237–256 (1967).
- ³⁴ Stephani, H., Kramer, D., MacCallum, M., Hoenselaers, C., and Hertl, E., *Exact Solutions of Einstein's Field Equations*, 2nd ed. Cambridge Monographs on Mathematical Physics (Cambridge University Press, 2003).
- ³⁵ Sinyukov, N. S., *Geodesic Mappings of Riemannian Spaces* (Nauka, Moscow, 1979).
- ³⁶ Wald, R. M., *General Relativity* (The University of Chicago Press, 1984).
- ³⁷ Yano, K., "On the torse-forming directions in Riemannian spaces," *Proc. Imp. Acad. (Tokyo)* **20**, 340–345 (1944).