

Table 1. Characteristics of non-pregnant women with endometriosis-associated acute hemoperitoneum. Literature data 1970-2016.

Source	Year	No. of patients	Previous diagnosis of endometriosis	Medical treatment	Surgical Treatment	Surgical findings
Ranney B. (6)	1970	4	No	No	Laparotomy	Case 1. Bleeding endometriotic peritoneal deposits Case 2. Ruptured right endometrioma of 20-25 cm Case 3. Ruptured left endometrioma Case 4. Bleeding endometriotic peritoneal deposits
Carmichael <i>et al.</i> (7)	1972	1	No	No	Laparotomy	Active bleeding from endometriosis deposits of the posterior surface of the left broad ligament
Kumar S. (8)	1996	1	No	Oral contraceptives	Laparotomy	Active bleeding from endometriotic deposits in the pouch of Douglas and along both uterosacral ligaments
Harmanli <i>et al.</i> (9)	1998	1	No	No	Laparotomy	Tubal endometriosis implants and a large, ruptured, hemorrhagic endometriotic cyst
Janicki <i>et al.</i> (10)	2002	1	Yes	No	Laparotomy (stage III endometriosis diagnosed at prior laparoscopy)	Left endometrioma (5x6 cm) Erosion of the left uterine artery
Fiadjoe <i>et al.</i> (11)	2008	1	Not specified	No	Laparotomy	Ruptured right endometrioma eroding into the broad ligament, pelvic side wall and uterine artery

Table 1. Continued

Source	Year	No. of patients	Previous diagnosis of endometriosis	Medical treatment	Surgical treatment	Surgical findings
Evangelinakis <i>et al.</i> (12)	2009	16	Not specified	Not specified	Laparoscopy	Ruptured left ($n = 11$) and right ($n = 5$) endometrioma
Mutihir <i>et al.</i> (13)	2010	1	Yes (umbilical endometriosis)	Discontinued Norethisterone 6 days before the event	Laparotomy	Active bleeding from endometriotic deposits in the pouch of Douglas and of the sigmoid colon
Togami <i>et al.</i> (14)	2015	1	Yes (site and prior procedures not specified)	No	Laparoscopy	Active bleeding from endometriotic peritoneal deposits
Present report	2016	1	No	Oral contraceptives	Laparoscopy	Fissured left uterine artery due to the presence of deep infiltrating endometriosis