

# Initial outcomes of a harmonized approach to collect welfare data in sport and leisure horses

E. Dalla Costa<sup>1†</sup>, F. Dai<sup>1</sup>, D. Lebelt<sup>2</sup>, P. Scholz<sup>2</sup>, S. Barbieri<sup>1</sup>, E. Canali<sup>1</sup> and M. Minero<sup>1</sup>

<sup>1</sup>Dipartimento di Medicina Veterinaria, Università degli Studi di Milano, via Celoria 10, 20133 Milan, Italy; <sup>2</sup>Pferdeklinik Havelland/Havelland Equine Hospital, Hohenferchesarner Straße 49, D-14778 Beetzsee OT Brielow, Germany

(Received 24 November 2015; Accepted 25 May 2016)

---

*A truthful snapshot of horse welfare conditions is a prerequisite for predicting the impact of any actions intended to improve the quality of life of horses. This can be achieved when welfare information, gathered by different assessors in diverse geographical areas, is valid, comparable and collected in a harmonized way. This paper aims to present the first outcomes of the Animal Welfare Indicators (AWIN) approach: the results of on-farm assessment and a reliable and harmonized data collection system. A total of 355 sport and leisure horses, stabled in 40 facilities in Italy and in Germany, were evaluated by three trained assessors using the AWIN welfare assessment protocol for horses. The AWINHorse app was used to collect, store and send data to a common server. Identified welfare issues were obesity, unsatisfactory box dimensions, long periods of confinement and lack of social interaction. The digitalized data collection was feasible in an on-farm environment, and our results suggest that this approach could prove useful in identifying the most relevant welfare issues of horses in Europe or worldwide.*

---

**Keywords:** AWIN, animal-based indicator, data collection, horse, welfare assessment

## Implications

This study reports the initial outcomes deriving from the application of the approach developed in the Animal Welfare Indicators (AWIN) project to assess the welfare status of stabled sport and leisure horses. For the first time, the authors present an innovative approach to collecting welfare data in a harmonized way that could prove useful for creating a common database of the general welfare status in different horse categories.

## Introduction

Horses are a peculiar species: they can be classified as farm or companion animals; they can be owned for several reasons ranging, for example, from use in animal-assisted therapies to food production. It follows that horses are managed heterogeneously and they are exposed to diverse welfare issues. For the same reasons, the assessment of horse welfare poses several challenges to researchers, for example, collecting data in different housing systems, or linking equids to their responsible person (World Horse Welfare and Eurogroup for Animals, 2015). A scientifically sound method to assess horse welfare on-farm represents

the foundation for a strategic plan aimed at improving the welfare of horses. A subsequent step includes the collection of reliable and uniform data in different countries to identify the most relevant welfare issues. By definition, harmonization is to create the possibility to combine data from heterogeneous sources into integrated, consistent and unambiguous information products. Harmonized data collection is essential in order to obtain a reliable picture of horse welfare conditions in different countries on which to base a roadmap to its improvement. Although significant progress has been made in this field over the past 15 years (Pritchard *et al.*, 2005; Burn *et al.*, 2010; Neijenhuis *et al.*, 2011; Vervaecke *et al.*, 2011; Popescu and Diugan, 2013; Visser *et al.*, 2014), limited data on welfare of European horses is yet available.

The AWIN project, funded by the European Commission in the Seventh Framework Programme, aimed to improve the welfare of several species, including horses, by developing scientifically sound and practical on-farm welfare assessment protocols (Battini *et al.*, 2015; Dalla Costa *et al.*, 2016). AWIN research was grounded on and progressed from the approach defined in the Welfare Quality<sup>®</sup> research project (Botreau *et al.*, 2007; Blokhuis *et al.*, 2010; Rushen *et al.*, 2011) and by Visser *et al.* (2014). As for horses, AWIN research was not only focussed on developing a welfare assessment protocol grounded on valid animal-based

---

<sup>†</sup> E-mail: emanuela.dallacosta@unimi.it

indicators (Dalla Costa *et al.*, 2014), but also on finding innovative methods to standardize on-farm data collection (Dai *et al.*, 2014). This study aims to report the first outcomes of the AWIN approach on the development of a strategy to improve horse welfare: it presents the results of on-farm assessment and the use of a harmonized data collection system.

## Material and methods

### *Facilities and horses*

The welfare of horses stabled in 40 facilities in Italy ( $n = 20$ ) and in Germany ( $n = 20$ ) was assessed between March and July 2014. Considering that the number of horses stabled in each facility can be very different and that this can be associated with specific welfare issues, a stratified random sample of very small ( $\leq 4$  horses), small (5 to 10 horses), medium (11 to 30 horses) and large ( $> 31$  horses) horse facilities was selected. All the selected facilities were contacted over the phone and participated in the study on a voluntary basis. In each facility, all the single-stabled horses (for at least half of the day)  $\geq 5$  years old were included in the study. A total of 355 sport and leisure horses (females = 146; geldings = 190; stallions = 19), of different breed and riding discipline (show jumping = 118; dressage = 22; western = 18; endurance = 8; eventing = 8; school = 28; leisure = 73; pet = 12; other = 40; NA = 28), aged between 5 and 33 years old (mean = 12.7) were assessed.

### *Assessors*

Three veterinarians (two females and one male), aged between 30 and 37 years, experienced in horse behaviour and welfare, were recruited to perform the assessments. Before carrying out the on-farm evaluation, they underwent a joint training period to learn how to perform and score all the indicators included in the AWIN welfare assessment protocol for horses (AWIN, 2015a). The training of assessors consisted of two phases: first e-learning and then face to face. The e-learning phase was developed in order to reduce time and costs related to the face-to-face training, without losing accuracy in the assessment. Each welfare indicator was transferred into a learning object organized in different sections: description, how to assess, how to score, examples and self-assessment exercises. The online material was available to the trainees for 1 month (December 2013). Then, a face-to-face training phase lasting 2 days and consisting of theoretical and practical on-farm training was performed in order to acquire the practical skills necessary to perform and score all the indicators accurately and reliably. During the training on-farm, assessors worked in pairs with silver standard, that is, two AWIN researchers with experience in assessing horse welfare. Both phases ended with an assessment of learning: as for the e-learning phase, assessors had to answer 58 questions (including videos and/or pictures); whilst at the end of the face-to-face phase, assessors conducted live assessments of horses until they performed a

minimum of five consecutively accurate assessments. The training was considered complete when the assessors achieved  $\geq 80\%$  agreement with the silver standard, on both e-learning and live scoring.

### *Welfare assessment*

The assessment was conducted using the AWIN welfare assessment protocol for horses. This document (AWIN, 2015a) reports the description, the assessment and scoring methods of the welfare indicators used in the current work. The AWIN welfare assessment protocol for horses is based on the Welfare Quality<sup>®</sup> principles and criteria, developed following the methods reported by Dalla Costa *et al.* (2016) and includes animal, resource and management-based indicators (Table 1) evaluated in terms of their validity, reliability and feasibility. In the current study, the welfare assessment was conducted at least 1 h after feed distribution, as recommended in the protocol.

### *Data collection*

A digital system to collect, store and download the indicators included in the AWIN welfare assessment protocol for horses was created (Dai *et al.*, 2015b); the AWINHorse app was developed for Android devices and it is now freely available on Google Play Store (AWIN, 2015b). A preliminary version of the AWINHorse app was installed on tablets or smartphones and tested on-farm in order to evaluate its feasibility during the assessments. The welfare assessors learned how to use the app to collect data on-farm during their training.

### *Statistical analysis*

Data collected on-farm by assessors were downloaded from the app to a comma-separated value file before performing descriptive statistics using SPSS statistical package (IBM Corp., 2012). The proportion of horses with different scores for each welfare indicator was calculated. For the fear test (Dai *et al.*, 2015a; Gorecka-Bruzda *et al.*, 2011), minimum, maximum values and means and standard deviations were calculated; an ANOVA was used to identify possible links between fear behaviour shown by horses (e.g. freezing, prancing, vigilance, defecation/urination) (Le Sclan *et al.*, 1997; Wolff *et al.*, 1997; Christensen *et al.*, 2005; Forkman *et al.*, 2007) and the latency to approach the novel object in the fear test. Latency was considered the dependent variable, and fear behaviour was introduced in the model as fixed effect. Data were tested for normality using a Kolmogorov–Smirnov test. As variables were not normally distributed, a  $\chi^2$  test was used to identify a possible relationship between variables.

## Results

This section initially reports, for each welfare principle presented in Table 1, the results of the welfare assessment; then explores feasibility aspects related to the data collection system.

**Table 1** Names of the Animal Welfare Indicators (AWIN) for horse welfare assessment presented by principles and criteria

Welfare principles	Welfare criteria	Welfare indicators
Good feeding	Appropriate nutrition	Body condition score
	Absence of prolonged thirst	Water availability
Good housing	Comfort around resting	Bucket test
		Bedding
		Box dimensions
Good health	Ease of movement	Exercise
	Absence of injuries	Integument alterations
		Swollen joints
		Lameness
		Prolapse
		Hair coat condition
		Discharges
		Consistency of manure
		Abnormal breathing
		Coughing
		Absence of pain and pain induced by management procedures
	Signs of hoof neglect	
Appropriate behaviour	Expression of social behaviour	Lesions at mouth corners
	Expression of other behaviours	Social interaction
		Stereotypies
		Fear test
	Good human–animal relationship	Human–animal relationship tests

Detailed descriptions of each welfare indicator are reported in the AWIN welfare assessment protocol for horses (AWIN, 2015a).

*On-farm welfare assessment*

No safety issues were encountered and time required to perform the assessment varied from 5 to 25 min/horse. For each welfare indicator, the proportions of horses with different scores are reported in Table 2 (for the principle ‘good feeding’), Table 3 (‘good housing’), Table 4 (‘good health’) and Table 5 (‘appropriate behaviour’).

As regards the principle ‘good feeding’ (Table 2), most of the assessed animals enjoyed appropriate nutrition (body condition score (BCS) = 3). Extreme scores, BCS = 1 and BCS = 5, were observed in a few cases, whereas dressage horses presented a significantly higher prevalence (54.5%;  $\chi^2 P = 0.002$ ) of overweight subjects (BCS > 3) compared with different riding disciplines. Our results show that the vast majority of horses had free access to a water point. However, the water points were often dirty or partially dirty, meaning that they were not checked or cleaned regularly by the stable staff.

As regards the principle ‘good housing’ (Table 3), bedding provided to the horses was in the majority of cases sufficient and clean; however, box dimensions were scored as satisfactory only in 68.6% of cases.

About half the horses had the possibility to exercise (free or ridden) on a daily basis. Remarkably, this study uncovered that 9.3% of subjects did not have the chance to get out of their box. In these cases, reported justifications were as follows: ‘the owner does not have time to ride the horse’, ‘the horse is old and no paddock for free exercise is available’, ‘giving the horse the possibility to spend free time in the paddock increases the risk of injuries’. Sometimes, it

**Table 2** Results of the Animal Welfare Indicators (AWIN) welfare assessment protocol for horses related to the principle ‘good feeding’

Welfare indicators	Score	Prevalence of each score (%)
Body condition score	1	0.6
	2	8.5
	3	58.8
	4	28.0
	5	4.2
Water availability: type of water point	Not present	0.8
	Trough	7.3
	Automatic drinker	91.8
Water availability: cleanliness of water point	Dirty	17.5
	Partially dirty	24.5
	Clean	53.0
	NA	5.1
Water availability: functioning of automatic drinkers	Not functioning	0.6
	Functioning	91.0
	NA	8.5

NA = not applicable.

Detailed descriptions of different scores for each welfare indicator are reported in the AWIN welfare assessment protocol for horses (AWIN, 2015a).

was not possible to gather information about exercise (proportion of NA in Table 3) because the stable manager was not always available to answer the questions for every horse present in the stable. In the considered sample, the majority of show jumping horses (60.5%) spent <2 h/day outside their box, the situation was different for dressage and leisure horses with a proportion of 22.7% and 16.4%, respectively.

**Table 3** Results of the Animal Welfare Indicators (AWIN) welfare assessment protocol for horses related to the principle 'good housing'

Welfare indicators	Score	Prevalence of each score (%)
Bedding: quantity	No bedding	0.3
	Insufficient <sup>1</sup>	19.2
	Sufficient/rubber mat	80.6
Bedding: cleanliness	Dirty	11.0
	Clean	88.5
	NA	0.6
Box dimensions	Not satisfactory <sup>2</sup>	30.4
	Satisfactory	68.2
	NA	1.4
Exercise	Never	9.3
	Sometimes (<1/week)	1.7
	Weekly (1 to 4 times/week)	28.2
	Daily	51.5
	NA	9.3

NA = not applicable.

Detailed descriptions of different scores for each welfare indicator are reported in the AWIN welfare assessment protocol for horses (AWIN, 2015a).

<sup>1</sup>Insufficient bedding = floor areas not covered by bedding are clearly visible.

<sup>2</sup>Not satisfactory = the area of the box is less than the satisfactory dimensions reported in the Swiss Animal Welfare Ordinance (2008).

As regards the principle 'good health' (Table 4), the majority of the horses did not present swollen joints, lameness, prolapse, unhealthy coat, discharges, abnormal breathing, coughing, signs of pain (Horse Grimace Scale (HGS) score < 2), hoof neglect and lesions at mouth corners. The most frequent integument alteration was alopecia, followed by superficial skin lesions and swellings. The HGS score was  $\geq 2$  in 2% of cases, always linked with other signs of pain (e.g. lameness). Lesions at mouth corners sometimes were not assessed (NA) because handlers were not available to hold the horse's head. In some cases, the horse was head shy and it was not safe to touch the corners of the mouth to assess the possible presence of lesions. It was not possible to evaluate lameness (NA) for 14.4% of horses. Another indicator that could not be assessed in quite a high number of cases was the faeces evaluation (manure), as most of the time boxes were clean at the moment of the inspection and no faeces were present.

Table 5 reports results regarding the principle 'appropriate behaviour'. In 22.3% of cases, horses had no possibility to interact with conspecifics, not even visually.

In our sample, evidence of stereotypic behaviour (e.g. cribbing, weaving) was significantly related to the reduced possibility of social contact ( $\chi^2 P = 0.001$ ). Most particularly, western riding horses presented the highest prevalence of these behaviours (27.8%) and they also had the highest prevalence of lack of social contact (27.8%).

Most of the assessed horses showed a positive reaction to an unknown human interacting with them during three behavioural tests, with a small prevalence of horses showing avoidance or negative reaction. Testing the avoidance distance to a human approaching the box door was not possible in 23.3% of the cases, mostly when horses were inattentive to the human presence.

**Table 4** Results of the Animal Welfare Indicators (AWIN) welfare assessment protocol for horses related to the principle 'good health'

Welfare indicators	Score	Prevalence of each score (%)
Integument alterations	Present	34.6
	Absent	65.4
Swollen joints	Present	2.3
	Absent	97.7
Lameness	Lame	1.7
	Not lame	83.9
	NA	14.4
Prolapse	Absent	100
Hair coat condition	Unhealthy <sup>1</sup>	2.8
	Healthy	96.9
	NA	0.3
Ocular discharge	Present	1.1
	Absent	98.9
Nasal discharges	Present	0.3
	Absent	99.7
Discharge from vulva or penis	Absent	100
Consistency of manure	Abnormal	2.0
	Normal	75.2
	NA	22.8
Abnormal breathing <sup>2</sup>	Present	0.6
	Absent	99.4
Coughing	Coughing	0.6
	No coughing	99.4
HGS <sup>3</sup>	Signs of pain (HGS $\geq 2$ )	2.0
	No signs of pain (HGS < 2)	97.7
	NA	0.3
Signs of hoof neglect <sup>4</sup>	Present	3.1
	Absent	96.9
Lesions at mouth corners	Open wounds	0.8
	Redness	0.8
	Hardened spots	7.6
	No lesions	82.8
	NA	7.9

NA = not applicable; HGS = Horse Grimace Scale.

Detailed descriptions of different scores for each welfare indicator are reported in the AWIN welfare assessment protocol for horses (AWIN, 2015a).

<sup>1</sup>Unhealthy coat = dull, dry coat with or without rough coat.

<sup>2</sup>Abnormal breathing = it is characterized by an exaggerated effort to breathe under standard climate conditions and at rest.

<sup>3</sup>HGS = it is a standardized method to evaluate changes in a horse facial expression due to pain (Dalla Costa *et al.*, 2014).

<sup>4</sup>Signs of hoof neglect = hooves are overgrown, rarely trimmed or trimmed incorrectly.

In the fear test, after the novel object was dropped in the box, the horses needed  $34 \pm 52$  s (MIN = 0 s, MAX = 285 s) to approach it. Latencies were significantly related to the presence of behavioural fear reactions (e.g. freezing, prancing, vigilance, defecation/urination) (ANOVA,  $P < 0.01$ ).

#### Data collection system

All three assessors successfully completed both phases of training, reaching a good level of agreement with the silver standard ( $\geq 80\%$ ).

**Table 5** Results of the Animal Welfare Indicators (AWIN) welfare assessment protocol for horses related to the principle 'appropriate behaviour'

Welfare indicators	Score	Prevalence of each score (%)
Social interaction	No possibilities for visual or physical contact	22.3
	Possibility to have visual contact	38.6
	Possibility to sniff other horses	29.0
	Possibility to nibble and partly groom	9.8
	NA	0.3
Stereotypies	Evidence of stereotypies	19.4
	No evidence of stereotypies	80.6
Avoidance distance <sup>1</sup>	Avoidance behaviour	6.2
	No avoidance	70.1
	NA	23.7
Voluntary animal approach	Negative signs <sup>2</sup>	2.3
	No interest	18.6
	Positive signs	72.4
	NA	6.8
Forced human approach	Negative signs <sup>2</sup>	3.4
	Avoidance	16.1
	Positive signs	78.9
	NA	1.7

NA = not applicable.

Detailed descriptions of different scores for each welfare indicator are reported in the AWIN welfare assessment protocol for horses (AWIN, 2015a).

<sup>1</sup>Avoidance distance = presence of any avoidance behaviour.

<sup>2</sup>Negative signs = any signs of aggressive behaviours such as trying to bite and/or kick.

All our assessors quickly learned how to use the digitalized system to collect data and upload it to a server. Thanks to the use of tablets or smartphones, in 1 or 2 min the assessors were able to insert the data for a particular horse and upload it to a central server. When an internet connection was not available on-farm, data could be stored on the device and sent to the server later on. The use of tablets on-farm caused some minor difficulties: as these devices could be cumbersome, horses sometimes approached and sniffed them, thus interfering with some of the behaviour tests; the use of smartphones that can be safely stored in a pocket could reduce these problems.

## Discussion

The results highlight that the approach described in this paper was useful to assess the welfare status of stabled sport and leisure horses. Even though no cases of cruelty and no major welfare problems were encountered, the authors consistently uncovered issues such as being overweight, unsatisfactory box dimensions, long periods of confinement and lack of social interaction. These issues are unsurprisingly similar to those reported as commonly perceived by respondents to a recent European survey (World Horse Welfare and Eurogroup for Animals, 2015). The number of facilities assessed was relatively limited; for this reason, the sample of horses assessed does not necessarily represent the welfare status of all horses in Europe, because conditions vary within and between countries. Obesity is a serious and largely under-reported equine welfare and health problem (Wyse *et al.*, 2008); overweight horses are predisposed to the

development of several pathological conditions such as hyperlipaemia, laminitis and osteoarthritis (Watson *et al.*, 1992; Geor, 2008). The prevalence of overweight horses found in the current study is in line with the findings of Visser *et al.* (2014), confirming that, fat subjects are becoming more common than thin ones. Unlike other issues, this one can be addressed by changes to husbandry practices such as nutrition management and exercise routine.

Box dimension can affect the lying behaviour of horses: Raabymagle and Ladewig (2006) observed that when insufficient lying space is provided, horses do not lie in lateral recumbency and they are unlikely to achieve paradoxical sleep. As for the AWIN protocol, box dimensions were considered satisfactory when responding to the requirements of the Swiss Animal Welfare Ordinance (2008). In order to prefigure the impact and consequences deriving from any possible large-scale prescriptions about horse box dimensions, it is essential to obtain a representative picture of their actual characteristics in the relevant geographical areas.

Our results show that horses can be confined for long period of time. Spending some time outside the box is important to prevent health and behavioural problems (Cooper and Albentosa, 2005; Visser *et al.*, 2014), as for example, it has long been known that respiratory problems are closely associated with improper housing microclimate (Halliwell *et al.*, 1993) and that activity level affects the presence of abnormal locomotory behaviours (McGreevy *et al.*, 1995a). An increasing number of organizations of the equine sector nowadays agree that many common horse welfare problems, including the lack of adequate free exercise, would be effectively improved by the promotion of

responsible ownership (World Horse Welfare and Eurogroup for Animals, 2015). Horses are social animals, in nature they rely on survival strategies centred on the formation of cohesive social bonds within their family or bachelor bands (Mills and Nankervis, 1999), and interacting with conspecifics is a basic behavioural need. Frustration of this need represents a serious welfare issue that can lead to the development of undesired behaviours (McGreevy *et al.*, 1995a; Cooper and Alentosa, 2005). In our sample, lack of social interaction is common. The prevalence of stereotypic behaviour, we found, is in line with findings of previous studies on sport and leisure horses (Kennedy *et al.*, 1993; McGreevy *et al.*, 1995b; Muñoz-Alonzo *et al.*, 2015). For example, McGreevy *et al.* (1995b) reported that the prevalence of stereotypies for dressage, eventing and endurance horses ranged from 19.5% to 32.5% and found that the time a horse spends out of the stable is negatively correlated with an increased risk of abnormal behaviour. Therefore, if the stable is constructed so that horses cannot have any social contact, it should be remodelled as soon as possible and, in the meanwhile, different solutions should be implemented, for example, offering daily access to paddocks where horses can interact with conspecifics.

Other minor welfare issues encountered were that water points were not regularly cleaned by the stable staff.

In this study, as in the case of overweight horses, a key role is played by education on the importance of evaluation of body condition and regular checks to ensure that water points are clean and no blockages are present.

As regards the principle good health, that is patently of primary importance for horse welfare, some of the present findings seem to differ from what is described elsewhere in the literature (Neijenhuis *et al.*, 2011; Asknes and Mejdell, 2012; Visser *et al.*, 2014). For example, Visser *et al.* (2014) in their inspiring paper reported a prevalence of lameness (4.8%), higher than that registered by the authors (1.7%). The discrepancies with these studies are probably due to the different degrees of sensitivity of their indicators, which required more extensive training (i.e. 1 week on-farm) and a deeper understanding of animal handling (i.e. evaluation of horses trotting in a straight line on a firm underground for 40 m). In fact, grade 3 lameness on the American Association of Equine Practitioners (AAEP) Lameness Grading Scale is difficult to observe at a walk (Dyson, 2011). It is known that back pain is a potentially highly relevant welfare problem in sport and leisure horses (Visser *et al.*, 2014); however, its accurate assessment requires extensive training and considerable animal handling. For this reason, a specific indicator for back pain was not included in the AWIN welfare assessment protocol. Despite the commitment to select and develop highly feasible indicators, some practical issues posed difficulties during this research. Lameness assessment consisted of walking the horse outside its box on a hard and even surface. The main feasibility issue in this case was that the stable manager would not always take the responsibility for moving the horse out of its box without the owner's formal consent. Whereas on the one hand this result

highlights the possible constraints that can be faced assessing animal-based measures on-farm, on the other it suggests that handling problems that could lead to safety issues are relatively frequent. This should be addressed by teaching the owners how to prevent them through the adoption of appropriate handling techniques, taking into account the behaviour and learning characteristics of the horses.

As the relationship between horses and humans relies on repeated interactions, our results indicate that most of our horses maintained a positive relationship with humans (showing positive reaction in all the tests), this being of paramount importance to prevent negative emotional states and reactions potentially leading to accidents (Hausberger *et al.*, 2008). Sometimes, we found that horses were inattentive towards humans, even if the assessors signalled their presence (clicking three times with the tongue); this result may be because, in some cases, the human–horse interactions are neutral, even if neither negative nor positive.

The training formula, online and face to face, proved to be valuable, as it limited the time spent with the trainer to 2 days. In previous studies, the time needed for reliable training of assessors ranged from 8 days to 6 months (Burn *et al.*, 2009; Visser *et al.*, 2014). Our experience demonstrated that the AWINHorse app is friendly and practical to use and it does not require a long training period. The app permits insertion of data reducing possible errors of transcription and immediately provides an output of welfare status that can be used to open a dialogue with the stable manager. Furthermore, once uploaded, data are immediately available for further analysis on the server. Therefore, the app would be a useful tool for the development of an accessible data repository on animal welfare and for increasing data and knowledge accessibility to all European countries. The digitalized data collection system proved to be feasible on-farm, it should however be applied in other contexts (e.g. working horses, competitions) to further evaluate its potential in improving the efficiency of welfare data collection.

Our findings suggest that education of horse owners and stable managers is needed to make them more aware of some aspects of horse welfare and incentivize the uptake of the information. This study presents, for the first time, an innovative data collection system that could prove useful in creating a larger and more geographically distributed database of welfare issues in horses. As regards the welfare assessment, there are still some important challenges to address. Although the AWIN protocol reports suggestions for adaptation to some specific management situations (group housed horses), further scientific research is needed and it is forecast that the protocol will be updated for use in different conditions in the light of new scientific knowledge.

### Acknowledgements

The authors thank Alberto Carrascal at Daia Solutions for his programming expertise and collaboration in developing the AWINHorse app. The authors are grateful to all the horse owners who allowed them to enter their facilities. Finally, the authors

thank Kirk Ford for his extensive and professional revisions of language. The AWIN project (FP7-KBBE-2010-4) has received funding from the European Union Seventh Framework Programme for research, technological development and demonstration.

## References

- Animal Welfare Indicators (AWIN) 2015a. AWIN welfare assessment protocol for horses. Università degli Studi di Milano, Milan, Italy.
- Animal Welfare Indicators (AWIN) 2015b. AWINHorse app. Retrieved on 23 November 2015 from <https://play.google.com/store/apps/details?id=com.daia.awinhorse>.
- Asknes F and Mejdell C 2012. Back soreness is common among 'healthy' riding horses. Proceedings of the 46th Congress of the International Society for Applied Ethology, 31 July–4 August 2012, Vienna, Austria, 236pp.
- Battini M, Stilwell G, Vieira A, Barbieri S, Canali E and Mattiello S 2015. On-farm welfare assessment protocol for adult dairy goats in intensive production systems. *Animals (Basel)* 5, 934–950.
- Blokhuys HJ, Veissier I, Miele M and Jones B 2010. The Welfare Quality® project and beyond: safeguarding farm animal well-being. *Acta Agriculturae Scandinavica, Section A – Animal Science* 60, 129–140.
- Botreau R, Veissier I, Butterworth A, Bracke MBM and Keeling LJ 2007. Definition of criteria for overall assessment of animal welfare. *Animal Welfare* 16, 225–228.
- Burn CC, Dennison TL and Whay HR 2010. Relationships between behaviour and health in working horses, donkeys, and mules in developing countries. *Applied Animal Behaviour Science* 126, 109–118.
- Burn CC, Pritchard J and Whay HR 2009. Observer reliability for working equine welfare assessment: problems with high prevalences of certain results. *Animal Welfare* 18, 177–187.
- Christensen JW, Keeling LJ and Nielsen BL 2005. Responses of horses to novel visual, olfactory and auditory stimuli. *Applied Animal Behaviour Science* 93, 53–65.
- Cooper J and Albentosa M 2005. Behavioural adaptation in the domestic horse: potential role of apparently abnormal responses including stereotypic behaviour. *Livestock Production Science* 92, 177–182.
- Dai F, Cogi NH, Heinzl EUL, Dalla Costa E, Canali E and Minero M 2015a. Validation of a fear test in sport horses using infrared thermography. *Journal of Veterinary Behavior: Clinical Applications and Research* 10, 128–136.
- Dai F, Dalla Costa E, Battini M, Barbieri S, Ferrari L, Minero M, Mattiello S and Canali E 2015b. New frontiers in welfare data collection: AWINGoat and AWINHorse app. Proceedings of AWSELVA-ECAWBM-ESVCE Congress, 30 September–3 October 2015, Bristol, United Kingdom, 42pp.
- Dai F, Dalla Costa E, Battini M, Barbieri S, Minero M, Mattiello S and Canali E 2014. An innovative tool for on-farm data collection and information sharing. Proceedings of the 6th International Conference on the Assessment of Animal Welfare at Farm and Group Level, 3–5 September 2014, Clermont-Ferrand, France, 153pp.
- Dalla Costa E, Dai F, Lebelt D, Scholz P, Barbieri S, Canali E, Zanella AJ and Minero M 2016. Welfare assessment of horses: the AWIN approach. *Animal Welfare* (in press).
- Dalla Costa E, Minero M, Lebelt D, Stucke D, Canali E and Leach MC 2014. Development of the Horse Grimace Scale (HGS) as a pain assessment tool in horses undergoing routine castration. *PLoS ONE* 9, e92281.
- Dalla Costa E, Murray LAM, Dai F, Canali E and Minero M 2014. Equine on-farm welfare assessment: a review of animal-based indicators. *Animal Welfare* 23, 323–341.
- Dyson S 2011. Can lameness be graded reliably? *Equine Veterinary Journal* 43, 379–382.
- Forkman B, Boissy A, Meunier-Salaün M-C, Canali E and Jones RB 2007. A critical review of fear tests used on cattle, pigs, sheep, poultry and horses. *Physiology & Behavior* 92, 340–374.
- Geor RJ 2008. Metabolic predispositions to laminitis in horses and ponies: obesity, insulin resistance and metabolic syndromes. *Journal of Equine Veterinary Science* 28, 753–759.
- Górecka-Bruzda A, Jastrzębska E, Sosnowska Z, Jaworski Z, Jezierski T and Chruszczewski MH 2011. Reactivity to humans and fearfulness tests: field validation in Polish Cold Blood Horses. *Applied Animal Behaviour Science* 133, 207–215.
- Halliwel REW, McGorum BC, Irving P and Dixon P 1993. Local and systemic antibody production in horses affected with chronic obstructive pulmonary disease. *Veterinary Immunology and Immunopathology* 38, 201–215.
- Hausberger M, Roche H, Henry S and Visser EK 2008. A review of the human–horse relationship. *Applied Animal Behaviour Science* 109, 1–24.
- IBM Corp 2012. IBM SPSS statistics for Windows. IBM Corp., Armonk, New York, USA.
- Kennedy MJ, Schwabe AE and Broom DM 1993. Crib-biting and wind-sucking stereotypies in the horse. *Equine Veterinary Education* 5, 142–147.
- Le Scolan N, Hausberger M and Wolff A 1997. Stability over situations in temperamental traits of horses as revealed by experimental and scoring approaches. *Behavioural Processes* 41, 257–266.
- McGreevy PD, Cripps PJ, French NP, Green LE and Nicol C 1995a. Management factors associated with stereotypic and redirected behaviour in the Thoroughbred horse. *Equine Veterinary Journal* 27, 86–91.
- McGreevy PD, French NP and Nicol CJ 1995b. The prevalence of abnormal behaviours in dressage, eventing and endurance horses in relation to stabling. *The Veterinary Record* 137, 36–37.
- Mills DS and Nankervis KJ 1999. *Equine behaviour: principles and practice*. Blackwell Science, Oxford, UK.
- Muñoz-Alonzo LE, Medina Vera MP, Cruces Leal J and Briones Luengo M 2015. Frequency of classic stereotypies in endurance horses. *Scientia Agropecuaria* 6, 119–124.
- Neijenhuis F, De Graaf-Roelfsema E, Wesselink H, Van Reenen C and Visser EK 2011. Towards a welfare monitoring system for horses in the Netherlands: prevalence of several health matters. Proceedings of 7th International Equitation Science Conference – Equitation Science: Principles and Practice – Science at Work, 27–29 October 2011, Hooge Mierde, The Netherlands, 80pp.
- Popescu S and Diugan EAA 2013. The relationship between behavioral and other welfare indicators of working horses. *Journal of Equine Veterinary Science* 33, 1–12.
- Pritchard JC, Lindberg AC, Main DCJ and Whay HR 2005. Assessment of the welfare of working horses, mules and donkeys, using health and behaviour parameters. *Preventive Veterinary Medicine* 69, 265–283.
- Raaby Magle P and Ladewig J 2006. Lying behavior in horses in relation to box size. *Journal of Equine Veterinary Science* 25, 502–504.
- Rushen J, Butterworth A and Swanson JC 2011. Animal behavior and well-being symposium. Farm animal welfare assurance: science and application. *Journal of Animal Science* 89, 1219–1228.
- Swiss Animal Welfare Ordinance 2008. Swiss Animal Welfare Ordinance (TSchV) of 23 April 2008 (position as at 1 April 2011). Retrieved on 11 November 2015 from [http://www.blv.admin.ch/themen/tierschutz/index.html?lang=en&download=NHZLpZeg7t,Inp6I0NTU042I2Z6In1ad1Izn4Z2qZpnO2Yuq2Z6gpjCFd3t2fmy m162epYbg2c\\_JjKbNoKSn6A-](http://www.blv.admin.ch/themen/tierschutz/index.html?lang=en&download=NHZLpZeg7t,Inp6I0NTU042I2Z6In1ad1Izn4Z2qZpnO2Yuq2Z6gpjCFd3t2fmy m162epYbg2c_JjKbNoKSn6A-).
- Vervaecke H, Boydens M, De Nil M and Laevens H 2011. Pilot study on the occurrence of pressure marks on the body and mouth lesions in riding horses in Flanders. In 7th International Equitation Science Conference – equitation science: principles and practice – science at work (ed. M van Dierendonck, P de Cocq and E Visser), p. 37. Academic Publishers, Wageningen, Hooge Mierde, The Netherlands.
- Visser EK, Neijenhuis F, de Graaf-Roelfsema E, Wesselink HGMH, de Boer J, van Wijhe-Kiezebrink MCM, Engel B and van Reenen CG 2014. Risk factors associated with health disorders in sport and leisure horses in the Netherlands. *Journal of Animal Science* 92, 844–855.
- Watson T, Murphy D and Love S 1992. Equine hyperlipaemia in the United Kingdom: clinical features and blood biochemistry of 18 cases. *The Veterinary Record* 131, 48–51.
- Wolff A, Hausberger M and Le Scolan N 1997. Experimental tests to assess emotionality in horses. *Behavioural Processes* 40, 209–221.
- World Horse Welfare and Eurogroup for Animals 2015. Removing the blinkers: the health and welfare of European Equidae in 2015. Retrieved on 11 November 2015 from <http://www.worldhorsewelfare.org/Removing-the-Blinkers>.
- Wyse CA, McNie KA, Tannahill VJ, Murray JK and Love S 2008. Prevalence of obesity in riding horses in Scotland. *The Veterinary Record* 162, 590–591.