

TITLE PAGE

Evidence of a bacterial core in the stored products pest *Plodia interpunctella*: the influence of different diets

Matteo Montagna^{1*}, Valeria Mereghetti¹⁺, Giorgio Gargari²⁺, Simone Guglielmetti², Franco Faoro¹, Giuseppe Lozzia¹, Daria Locatelli², Lidia Limonta².

¹Dipartimento di Scienze Agrarie e Ambientali - Università degli Studi di Milano, Milan, Italy.

²Dipartimento di Scienze per gli Alimenti la Nutrizione, l'Ambiente - Università degli Studi di Milano, Milan, Italy.

⁺these authors contributed equally to this work.

Corresponding author: Matteo Montagna

Dipartimento di Scienze Agrarie e Ambientali

Via Celoria 2, I-20133 Milano, Italy

Telephone: 0039 02 50316782; Fax: 0039 02 50316781; Email: matteo.montagna@unimi.it

Running title: Core microbiota in the Indian Meal Moth

Summary

The potential influence of insects' feeding behaviour on their associated bacterial communities is currently a matter of debate. Using the major pest of commodities,

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the Version of Record. Please cite this article as an 'Accepted Article', doi: 10.1111/1462-2920.13450

Plodia interpunctella, as a model and adopting a culture-independent approach, the impact of different diet on the host-associated microbiota was evaluated. An analysis of similarity showed differences among the microbiotas of moths fed with five substrates and provided evidence that diet represents the only tested factor that explains this dissimilarity. Bacteria shared between food and insects provide evidence for a limited conveyance to the host of the bacteria derived by the diet; more likely, the content of carbohydrates and proteins in the diets promotes changes in the insect's microbiota. Moth microbiotas were characterised by two robust entomotypes, respectively associated with a carbohydrate-rich diet and a protein-rich diet. These results were also confirmed by the predicted metagenome functional potential. A core microbiota, composed of six taxa, was shared between eggs and adults, regardless of the origin of the population. Finally, the identification of possible human and animal pathogens on chili and associated with the moths that feed on it highlights the possibility that these bacteria may be conveyed by moth frass.

Keywords: Indian meal moth; Lepidoptera; Pyralidae; microbiota; symbiosis; bacterial 16S rRNA pyrotag

Introduction

The Indian meal moth (IMM), *Plodia interpunctella* (Hübner 1813) (Lepidoptera: Pyralidae), is a cosmopolitan moth (Rees, 2004) that infests a multitude of stored-products, such as different types of cereals, nuts and dried fruits but also spices and dried meat (Hamlin et al., 1931; Sedlacek et al., 1996; Nansen et al., 2004; Mohandass et al., 2007; Fontenot et al., 2012). Due to the wide spectrum of infested products, *P. interpunctella* is considered a major pest of commodities, causing

extensive economic damages by reducing the quality grade of harvested products. Direct damage is represented by the feeding activity through all the larval instars, whereas indirect damage is due to the presence of larval frasses, exuviae and the silk-web (Allotey and Goswami, 1990). The capability of the IMM to efficiently colonise a wide spectrum of commodities poses the question of whether these metabolic capabilities are host constitutive or if they are provided or improved by the host-associated bacterial communities.

The establishment of mutualistic associations between the insect and microorganisms (bacteria, protozoa and fungi) plays an important role in the evolution of many insect lineages, such as termites and cockroaches (e.g., Bandi et al., 1995; Brune, 1998; López-Sánchez, 2009; Köhler et al., 2012). Consortia of mutualistic bacteria provide the host with essential compounds, such as vitamins, amino acids and sterols, but they also contribute to the digestion of ingested materials (e.g., Moran et al., 2003; Moran, 2006; Douglas, 2009; López-Sánchez et al., 2009; McCutcheon et al., 2009). In addition, it was recently postulated that a diversified host-associated microbiota may confer selective advantages to their host in changing environments, providing the capability to exploit different food sources, and thus, adapt to new ecological niches (Montagna et al., 2015a; Sudakaran et al., 2015).

Although the importance of bacterial communities on insect physiology is well recognised, the evolutionary and ecological determinants (e.g., diet, life stage, gender and host environment) that shape these communities are not well understood. The intuitive role played by the host's food substrate in shaping the associated bacterial communities was previously established in several insects, such as the cockroach *Periplaneta americana*, the cotton bollworm *Helicoverpa armigera*, the mealybug *Planococcus ficus*, and the red palm weevil *Rhynchophorus ferrugineus* (Kane and

Breznak, 1991; Priya et al., 2012; Iasur-Kruh et al., 2015; Montagna et al., 2015a). Nevertheless, a limited or even null impact on the host-associated bacterial community was recovered in the case of *Spodoptera littoralis* (Tang et al., 2012) and in the closely-related leaf beetles species (Montagna et al., 2015b); in the latter case, the altitude of the sites where the specimens were collected generated the ecological trait that affected the insect's microbiota. These results suggest that the host's microbiota is influenced by a multitude of biological factors, such as the host trophic guilds, and environmental factors, such as the altitude of the collecting site.

In this study, using a culture-independent approach based on 454 pyrosequencing targeting the bacterial 16S rRNA gene, we used the IMM *P. interpunctella* as a study model to address the following biological questions: *i*) Are the microbiotas associated with insects reared on five types of diet different? *ii*) Is there a core microbiota associated with eggs that is retained until adulthood, regardless of diet and population origin? *iii*) Are the microbiotas associated with adults vehiculated through the ingestion of diet? In addition, the presence of bacteria that is potentially hazardous for human and/or animal health was also evaluated.

Materials and Methods

Experimental design

All of the *Plodia interpunctella* specimens used in the assays were laboratory reared on different substrates under controlled conditions of light (light-dark cycle 16:8), humidity (RH = 70 ± 5%) and temperature (27° ± 1°C). In order to evaluate the impact of the diet in shaping the microbiotas associated with the IMM, 90 out of 120 eggs collected randomly from the same cohort of the laboratory-reared population, from a line maintained since 1980 in the rearing facilities established at the DEFENS,

Accepted Article

were reared until adulthood on the following different substrates: *a*) 30 eggs on an artificial diet prepared according to Stampini and Locatelli (2007); *b*) 30 eggs on dried *Moringa oleifera* leaves (hereafter named moringa); and *c*) 30 eggs on *Vicia faba* beans (hereafter fava bean). The remaining 30 eggs were directly analysed in order to provide information on the native microbiotas associated with the laboratory-reared population. Newborn adults of two distinct wild populations collected in Italy and living respectively on dried *Capsicum annuum* (chili pepper; hereafter named chili) and on noodles of white-black buckwheat flour (named pizzoccheri) were processed for DNA extraction to better investigate the variability of the bacterial communities associated with the IMM. In order to avoid possible shifts in the composition and structure of the insect-associated microbiotas due to a seasonal variability (Jia et al., 2013), all of the analysed moth specimens were collected during the first twenty days of May 2014.

DNA extraction

A total of five new-born adults (three females and two males) from each population were surface sterilised and DNA extraction was performed on the whole body (Montagna et al., 2015b). In order to characterise their internal and external microbiotas, the eggs were not surface sterilised because the outer shell of an insect's egg, called the chorion, is the first meal of the newborn larvae. The DNA was extracted from each sample using the DNeasy Blood and Tissue Kit (Qiagen) following the manufacturer's instructions. The concentration and purity of the extracted DNA were determined by a NanoDrop ND-1000 spectrophotometer (NanoDrop Technologies Inc., Wilmington, Delaware). In order to characterise the bacteria associated with the five substrates used as food sources, DNA was extracted

using the DNeasy Plant Kit (Qiagen) starting from 0.5 g of food after homogenisation with liquid nitrogen of the whole amount of substrate on which the IMM were developed.

Pyrosequencing

The bacterial communities associated with the insects and food samples were characterised by targeting the V1-V3 variable regions of the 16S rRNA gene using universal primers for bacteria as reported in a previous studies (Mazza et al., 2014; Montagna et al., 2015a,b). A commercial service performed the PCR reactions and the 454 pyrosequencing by Roche 454 GS FLX Titanium (MR DNA, Shallowater, TX—U.S.). The 16S rRNA gene sequences obtained by the 454 pyrosequencing assays were deposited in the European Nucleotide Archive (accession number PRJEB14361), mapping file is provided in Table S1.

Using a dedicated filtering script working under the QIIME platform (Camporaso et al., 2010), the obtained raw 16S rRNA V1-V3 gene sequences were trimmed to remove adaptors, low quality base calls (<30 Phred score) and short-sized sequences (retained sequences between 350 and 500 bp). Chimeras were removed with *Chimeraslayer*, and the high quality non-chimeric sequences were clustered into operational taxonomic units (OTUs), based on a sequence identity threshold of 97%, using Uclust (Edgar 2010). PyNast (Camporaso et al., 2010) was used to align a representative sequence from each OTU with Greengenes (<http://greengenes.lbl.gov/>); these representative sequences were then taxonomically classified by BLASTn-based comparisons to the Greengenes and Silva databases. After a first analysis, the OTUs identified as plant chloroplast and mitochondria were removed using an *ad hoc* perl script. This output was used for a second run of QIIME analyses.

Diversity and statistical analyses

The resulting set of OTUs was used as input for the diversity analyses carried out with different R packages (R Project 3.0.2; <http://cran.r-project.org/>).

The diversity indices and the following analysis (exceptions are specified) were estimated using the R package *vegan* (Dixon, 2003). The Shannon H index (Shannon, 1948), Pielou's evenness (Pielou, 1975) and the total species richness index Chao 1 (Chao, 1984; Chao et al., 1992) were estimated. The bacterial communities associated with the moth samples were ordinated, according to OTU composition similarity, using the distance-based Non-Metric Multi-Dimensional Scaling (NMDS; Kruskal, 1964) with the Chao probabilistic distance (Chao et al., 2005) as described in a previous work (Montagna et al., 2015a,b). Correlations between the diet typologies (i.e., artificial diet, moringa, fava bean, chili and pizzoccheri), the origin of the samples (i.e. lab reared specimens vs wild populations) and the specimen's gender with the insect-associated bacterial communities were tested by fitting the NMDS scores using the *envfit* function. The significance of the fitted factors was assessed by permutations (9999) of the dissimilarity OTUs matrix. The same analyses were also performed on the pairwise UniFrac distance matrix (rarefaction on the minimum samples size) calculated using the taxonomy-independent weighted UniFrac metric (Lozupone et al., 2007), which accounts for similarities in the phylogenetic structure of the communities associated to each moth samples. The best number of clusters identified by the NMDS analyses was evaluated using the Calinski-Harabasz criterion (Calinski and Harabasz, 1974), and their consistency was evaluated using silhouette scores (Rousseeuw, 1987).

The differences among the bacterial communities associated with the IMM fed with

different substrates were estimated by a nonparametric one-way analysis of similarity (ANOSIM; Clarke, 1993). Shifts in the bacterial community structure associated with the three groups of moths reared in laboratory on different substrates (i.e., artificial diet, moringa leaves and fava beans) were estimated with the R package *betapart* (Baselga and Orme, 2012) using Simpson's dissimilarity index as described in Montagna et al., (2015b).

In order to evaluate the bacterial OTUs associated with the eggs and those retained until adulthood, as well as those shared between the insects and their food, a co-occurrence (i.e., eggs-IMM and IMM-food microbiotas) table was obtained. The OTU table was filtered using the following criterion: the co-occurrence was assigned when a specific bacterial OTU of the eggs or food microbiotas was present in the microbiota of at least three insects feeding on the same resources. Venn diagrams, acquired with the R package *gplots*, provide a visualisation of the bacteria OTUs numbers, establishing the IMM bacterial core and the bacterial community shared among the adult insects, its food substrates and the eggs. In the Venn diagrams, the presence of a bacterial OTU is assigned to the group of insects feeding on the same resources when: *i*) the OTU is reported for at least one specimen of the group; *ii*) the OTU is reported for at least three specimens of the group, as for the co-occurrence table.

Predictive functional profiling

The functional profiles of the bacterial communities were investigated using PICRUST (Langille et al., 2013). The OTUs were closed-reference picked against Greengenes (GG version 13.5) using QIIME v 1.9 according to the online protocol. The bacterial metagenome was predicted for the bacterial communities associated with the IMM adults.

The table with the predicted L3-functions counts per-samples, according to the Cluster of Orthologous Groups (Tatusov et al., 1997) and identifiers adopted by KEGG Orthology (Kanehisa et al., 2012), was cleaned up by removing the categories not related to the bacterial physiology/metabolism and the null categories. The Nearest Sequenced Taxon Index median value for the PICRUSt predictions was 0.08 (sd = 0.04), reflecting good availability for the reference genomes on which the metagenome function predictions were based. The overall differences in the amount of counts were estimated by the non-parametric Kruskal-Wallis test (Kruskal and Wallis, 1952) after assessing the homogeneity of the variance among the groups through the Levene test (Levene, 1960). If the *p-value* of Kruskal-Wallis test resulted in a significance level that was less than a 0.05, then a Mann-Whitney-Wilcoxon test with a Hochberg correction (Mann and Whitney, 1947) was applied to compare the pairs of groups. These tests were performed using the lawstat-package (Hui et al., 2008) in the R software, and the results are visualised by a table.

Results and Discussion

α , β -diversity and community structure

In total, the analysed samples yielded approximately 200,000 16S rRNA reads. After the removal of the low-quality bases, chimeras, mitochondrial and chloroplast sequences, a total of 79,933 16S rRNA bacterial sequences were retained, of which 78,278 of the bacterial reads were associated with *P. interpunctella* specimens (average 3,011) and 1,655 were associated with their food (average 331). The 16S rRNA bacterial gene sequences were clustered into 1,611 bacterial operational taxonomic units (OTUs; Table 1). The bacterial communities associated with *P. interpunctella* reared on an artificial diet and pizzoccheri and characterised by the

highest carbohydrate content had the highest species diversity (Table 1), with an average number (\pm SD) of 195 ± 49 and 192 ± 89.7 detected OTUs. Only 62 OTUs were associated with the insects reared on fava beans (high protein content). A relationship between the protein content of the insect's diet and the bacterial diversity associated with the host was previously observed in different species, such as in *Drosophila* spp., *Lymantria dispar* and *Blattella germanica* (Chandler et al., 2011; Mason and Raffa, 2014; Pérez-Cobas et al., 2015). The bacterial communities associated with the IMMs analysed in this study showed, on average, a significantly higher number of bacterial OTUs compared with OTU abundance reported in other studies for different caterpillars (Broderick et al., 2004; Yu et al., 2008; Pinto-Tomás et al., 2011). The recovered differences in the OTU richness may be explained by the fact that the IMMs, feeding on a variety of stored products, do not require a selected microbiota but will benefit from the diversified metabolic capabilities provided by a differentiated microbiota, whereas the latter feed on plant tissues, which are potentially rich in secondary compounds or allelochemicals that benefit a selected microbiota to overcome plant constitutive biochemical barriers.

The highest values of the Shannon and Pielou's evenness indices (Table 1) were recovered in the microbiotas associated with specimens feeding on moringa leaves and chili (3.70 and 3.36, 0.77 and 0.75, respectively), whereas the bacterial communities of the *P. interpunctella* feeding on an artificial diet and pizzoccheri had the lowest values (Table 1). The fact that the bacterial communities associated with these latter samples are unbalanced was also confirmed by the abundance of the most dominant OTU, which respectively establishes 45.4% and 57.7% of the total retrieved bacterial 16S rRNA sequences (Fig. 1c). With respect to the microbiotas of the food

sources, 128 bacterial OTUs were found in the artificial diet, whereas only one and two OTUs were detected in moringa and chili, respectively.

The β -diversity, measured as Sørensen's dissimilarity, was estimated over the three groups of IMM from the laboratory population sharing a common origin and genotype. The values obtained for the two components of the β -diversity, the turnover and the nestedness (respectively $\beta_{SIM} = 0.56$ and $\beta_{NES} = 0.2$), indicate that only approximately 20% of the bacterial OTUs is shared among the three analysed groups. The differences in the bacterial communities associated with all five groups of *P. interpunctella* were confirmed by an ANOSIM analysis ($P < 0.001$).

Factors affecting the bacterial community structure

The Non-Metric Multi-Dimensional scaling analyses, performed on the moth-associated bacterial OTUs (Chao and Weighted UniFrac distance matrices), were fitted with factors potentially affecting the microbiota composition and structure as follows: *i*) the diet; *ii*) the IMM population of origin and *iii*) the IMM gender (Figure 2). The diet was the only tested trait able to explain the dissimilarity among the moth-associated bacterial communities (NMDS_{CHAO} $R^2 = 0.49$, $P = 0.0002$, stress = 0.08; NMDS_{UNIFRAC} $R^2 = 0.63$, $P = 0.0001$, stress = 0.09), whereas the gender and population origin did not significantly affect the insect microbiotas. This result is in agreement with those recently achieved in the case of higher termites, *Blattella germanica* and in the larvae of *Melitaea cinxia* (Mikaelyan et al., 2015; Pérez-Cobas et al., 2015; Ruokolainen et al., 2016) and provides further evidence for the relevant role of the host's diet in moulding the associated bacterial community. A possible hypothesis, supported by published studies and that should be tested in a rigorous framework, is that the microbiota associated with a generalist insect, in terms of exploitation of food

resources, seems to be more influenced by diet typologies compared to those that are more specialised. An exemplary case of the latter is represented by the pine weevil (*Hylobius abietis*), where its microbiota remains almost unchanged across Europe, and interestingly, it differs from closely related weevil not associated with conifers but is highly similar to that of bark beetles that feed on conifers (Berasategui et al., 2016).

On the basis of the Calinski-Harabasz index and a silhouette score of 0.38, two consistent groups of bacteria (hereafter, entomotypes from the Greek *entomon* meaning insect and referred to as consistent groups of bacteria present within the volume delimited by the exoskeleton), were identified within the IMM microbiotas.

The two groups were clearly segregated by the first NMDS axis and were independent of the moth population of origin (Figure 2). The first group is associated with the specimens that fed on an artificial diet and pizzoccheri, whereas, in the second group, the specimens fed on moringa, fava beans and chili. The taxonomic composition and the relative taxa abundance of the two groups are reported in the next section.

Bacterial taxonomic composition of the IMM and food microbiotas

The microbiotas associated with the IMM fed an artificial diet and pizzoccheri as well as the bacterial community associated with the eggs were dominated by taxa belonging to Firmicutes (61.6%, 77% and 51%, respectively; Fig. 1b, Table S2), whereas the microbiotas associated with the specimens fed chili, fava beans and moringa did not show a dominant taxonomic phylum because the relative abundances of Proteobacteria, Firmicutes and Actinobacteria were roughly equivalent (Fig. 1b, Table S2). Firmicutes was the dominant taxon and was also in the microbiota associated with *Spodoptera littoralis* (Tang et al., 2012), whereas their abundance is

limited in the bacterial communities associated with tropical saturniid caterpillars and in the cabbage root fly larvae (Pinto-Tomás et al., 2011; Welte et al., 2016).

Members of the family Aerococcaceae were the dominant component of the microbiotas associated with the moths reared on an artificial diet and the wild population that fed on pizzoccheri ($52.5\% \pm 1.77$ and $65.4\% \pm 5.76$, respectively; Fig. 3, Table S3). On the contrary, the bacterial communities associated with the *P. interpunctella* feeding on chili, fava bean and moringa were dominated by members of Propionibacteriaceae (20.7, 17.6, and 15 %, respectively; Fig. 3). A second major component in the microbiota of the moths feeding on chili was represented by Comamonadaceae (14.6%; Fig. 3). Streptococcaceae (19.9%) and Propionibacteriaceae (8.34%), which were the dominant families in the egg microbiotas (Fig. 3).

The genus *Atopococcus* (family Carnobacteriaceae) characterised the entomotype of the *P. interpunctella* reared on an artificial diet and pizzoccheri, representing $45.4\% \pm 1.42$ and $56.2\% \pm 4.67$, respectively, of the 16S rRNA gene reads (Fig. 4A; Table 2 and S4). The relative abundance of *Atopococcus* in the moths fed chili, moringa, fava beans and eggs was lower than that in the insects reared on an artificial diet and pizzoccheri (Table 2 and S4); nevertheless, the presence of this bacterial genus in the majority of the analysed samples allows us to speculate that *Atopococcus* might represent a potential symbiont of *P. interpunctella*.

Lactic acid bacteria (LAB) is found as a typical inhabitant in the insect gut (especially the foregut), such as in the wood- and soil-feeding termites (Bauer et al., 2000), cockroaches (Kane and Breznak, 1991) and honeybees (*Apis* spp.; Vásquez et al., 2009; Olofsson et al., 2011). The biological role of LAB with respect to insect physiology remains, at present, poorly investigated and understood. In cockroaches,

lactate produced by LAB is used to support the animal's respiratory requirements (Kane and Breznak, 1991) or plays a fundamental role in the insect's aggregative behaviour (McFarlane and Alli, 1986); in honeybees, LAB is involved in the defense against the insect's pathogens (Vásquez et al., 2012). Based on previous evidence, we hypothesise that LAB-mediated pathogen protection represents a useful adaptation for the generalist trophic behaviour of IMM; the involvement of LAB lactate in the IMM aggregating semiochemicals cannot be excluded.

Metabolic potential

The metagenomic functional potential of the bacterial communities associated with the IMMs fed from different resources was inferred from the bacterial 16S rRNA sequences using PICRUSt (Langille et al., 2013; Table S5). Among the five groups of IMMs significant differences were recovered by the Kruskal-Wallis test ($P < 0.05$) in a total of 38 functional categories (Fig. 5; Table S6). The majority of these categories are related to metabolism (31), while only few cases were observed for genetic (3) and environmental (1) information processing (Fig. 5). Within the functional categories related to metabolism, the majority related to the metabolism of amino acids, cofactors and vitamins, including both biosynthesis and catabolism (Fig. 5).

Interestingly, no differences in functional potential were assessed by the Mann-Whitney-Wilcoxon test with Hochberg correction for pairwise comparisons between the IMMs reared on moringa leaves, fava beans and chili pepper powder (Fig. 5, Table S7). This result represents a metabolic confirmation of the pattern achieved by the NMDS analyses (Figs. 2, 4), where these three groups of insects belong to the same cluster and are characterised by the *Propionibacterium* entomotype. Similar results, with no differences in a total of 20 of 38 functional categories, were obtained

for comparisons between the IMM reared on an artificial diet and pizzoccheri and the two groups characterised by the *Atopococcus* entomotype and recovered in the same cluster by NMDS analyses (Figs. 2, 4). All of the remaining pairwise comparisons possessed the highest number of differences in the functional categories.

The achieved pattern is congruent with the detected bacterial entomotypes (Fig. 4) as well as with the differences in the nutrient composition of the food resources. An artificial diet and pizzoccheri possessed the highest amount of carbohydrates (more than 50%; Table S8) with a low protein content (high C:N ratio; Table S8), whereas, on the contrary, the moringa leaves, chili powder and fava beans are characterised by low carbohydrates and a high protein content (low C:N ratio; Table S8).

Impact of the eggs and diet microbiotas on the adult bacterial communities

The bacterial OTUs associated with the eggs/food resources and retained by a majority of the adult specimens (i.e., in at least three out of five specimens) were inferred from the OTU table and were then visualised through a co-occurrence table (Fig. 6; Table S9, S10). The moths reared on an artificial diet, moringa leaves and fava beans, which came from the same population, retained, until adulthood, 35, 37 and 10 bacterial OTUs of the eggs microbiota, respectively (average number of OTU reads per sample of 109, 26 and 33; Fig. 6, Supplementary Figure S1, S2); in contrast, only a few (11 in the case of artificial diet) or none (in moringa and fava beans) of the bacterial OTUs associated with food resources were recovered in the insect microbiotas. Similar results were also achieved in the case of the IMM from the population fed with pizzoccheri (41 OTUs were retained from the eggs and five from the food, with an average number of OTU reads per sample of 102; Fig. 6, Supplementary Figure S1, S2). A slightly different result was achieved in the case of

the moths reared on chili powder, where the same number of OTUs (15), even if with taxonomic differences, was shared between the eggs (average number of OTU reads per sample of 42) and the diet microbiotas. The co-occurrence of diet-associated bacteria in the host could be the result of two effects that are not easily distinguishable: *i*) the inoculation of the insect with food-associated bacteria through ingestion, or alternatively, *ii*) the contamination of food resources by insect frass conveying host bacteria and/or their DNA. Even if conclusive results could not be reached, it is reasonable to interpret our findings as a result of the second effect because digestive enzymes, pH and reactive oxygen species present in the insect gut usually kill most of the bacteria ingested with food (Vallet-Gely et al., 2008; Garcia et al., 2010). For this reason, the IMMs may convey bacteria to humans or animals on the infested harvested products, especially in the case of food consumed raw, such as spices, nuts and dry meat. In the present study, the OTUs identified as *Staphylococcus* (otu 1242 and 3023; Fig. 6) and *Streptococcus* (otu 4214; Fig. 6), which are two genera that include species pathogenic to humans and animals (Schleifer and Bell, 2009; Whiley and Hardie, 2009), were found in association with the infested chili pepper. Because our results are based only on a fragment of approximately 400 bp of the 16S rRNA and this topic is outside of the scope of our research, we believe that further investigations are required to isolate and identify these bacteria.

All of the groups of insects, at least in three specimens out of the analysed five, retained, until adulthood, six bacterial OTUs of the eggs microbiota (average number of OTU reads per sample of 53 and 49, in the adults and eggs, respectively). These taxa, represented by *Propionibacterium* (three OTUs), *Staphylococcus* (two OTUs) and one OTU assigned to the family Xanthomonadaceae, are regarded as the core microbiota, i.e., the number and identity of the bacteria associated with the eggs and

retained until insect adulthood independently from the population of origin (Fig. 6).

The co-occurrence of bacterial phlotypes between sterilised eggs and the midgut of caterpillars was also observed in the case of *Rothschildia lebeau* (Pinto-Tomás et al., 2011). The mechanisms behind the transmission of the core microbiota from egg to adult were not elucidated in this study; however, we can hypothesise that the newborn larvae, using the egg corion as the first meal after hatching, acquires the associated bacterial community from which partially derives the adult microbiota.

Venn diagrams, reporting the bacterial OTUs shared among the five groups of IMMs, highlighted the presence of 42 OTUs shared among the five groups of samples (Supplementary Figure S3). This number includes the core microbiota shared with the eggs plus 36 further bacterial OTUs.

Conclusion

The bacterial communities associated with the analysed Indian meal moths were relatively complex and diversified (see Table 1), with the maximum number of bacterial OTUs recovered in the moths reared on an artificial diet ($OTU_{AVG} = 195$) and the minimum in the case of the moths fed with fava beans ($OTU_{AVG} = 62$).

Noteworthy, the microbiotas associated with *P. interpunctella* were characterised by a relatively high species richness if compared with that of phytophagous caterpillars (e.g., Broderick et al., 2004; Yu et al., 2008; Pinto-Tomás et al., 2011; Tang et al., 2012). These results are in agreement with the hypothesis that diversified host-associated microbiotas may confer essential metabolic capabilities in exploiting highly different food sources, as in the case of *P. interpunctella*.

As reported for other insects (e.g., cockroaches, the cotton bollworm and the red palm weevil), and also in the case of *P. interpunctella*, the diet on which larvae have

developed significantly contributes to the shaping of the structure and taxonomy of the associated bacterial communities. Nonetheless, considering the bacteria associated with food and shared with the insect, our findings provide evidences for a limited role of the diet in the active conveying of bacteria to the host. Conversely, the data from this study support the alternative hypothesis that diets with different compositions indirectly promote changes in the host-associated bacterial communities. The comparison of the microbiotas associated with eggs from the laboratory population with those associated with adult from different populations, including those from the laboratory, highlights the presence of a six-OTU bacterial core. Future studies, with the inclusion of specimens from different geographic areas and developed on different substrates, are required to confirm the presence of the identified bacterial core and, in the case of confirmation, to elucidate the possible mechanisms of its transmission through generations (i.e., by feeding the egg corion or by a transovarian transmission). Interestingly, the five groups of IMMs were characterised by two different entomotypes (Figs. 2, 4). The first was dominated by the bacteria of the genus *Atopococcus*, and the second was dominated by *Propionibacterium* and *Streptococcus*. The identification of LAB in the microbiota of *P. interpunctella* represent an interesting discovery because it was demonstrated that these bacteria could play an important role in insect biology. Apart from the possible biological role played by the LAB, these bacteria represent a possible target to control IMMs by developing symbiotic control strategies.

The functional potential of the IMMs microbiotas, predicted on the basis of 16S rRNA gene sequences, confirmed, from a metabolic point of view, the separation of bacterial communities associated with the IMM in two distinct groups (Figs 2, 4, 5) and characterised by two different entomotypes. The majority of the between-group

differences are related to metabolism (Fig. 5), and in particular, with the metabolism of amino acids, cofactors and vitamins. The achieved pattern is also congruent with the differences of the food resources in nutrient composition, with the artificial diet and pizzoccheri (high carbohydrates and low proteins) on one side and moringa leaves, chili and fava beans (low carbohydrates and high protein) on the other.

Further investigations are required to understand the possible involvement of IMM in conveying bacterial pathogens to the food substrates used for development.

Acknowledgments

We acknowledge Massimino Pajoro to provide essential samples. The work was partially supported by Linea 2-A Piano di Sostegno alla Ricerca 2015-2017 (grant 15-6-3014000-12).

References

- Allotey, J., and Goswami, L. (1990) Comparative biology of two phycitid moths, *Plodia interpunctella* (Hübner) and *Ephestia cautella* (Wlk.) on some selected food media. *Insect Sci App* 11: 209-215.
- Bandi, C., Sironi, M., Damiani, G., Magrassi, L., Nalepa, C.A., Laudani, U. and Sacchi, L. (1995) The establishment of intracellular symbiosis in an ancestor of cockroaches and termites. *Proc R Soc B* 259: 293-299.
- Berasategui, A., Axelsson, K., Nordlander, G., Schmidt, A., Borg-Karlson, A.K., Gershenzon, J., Terenius, O., and Kaltenpoth, M. (2016) The gut microbiota of the pine weevil is similar across Europe and resembles that of other conifer-feeding beetles. *Mol Ecol* doi: 10.1111/mec.13702. [Epub ahead of print].

- Baselga, A., and Orme, C.D.L. (2012) Betapart: an R package for the study of beta diversity. *Methods Ecol Evol* 3: 808-812.
- Bauer, S., Tholen, A., Overmann, J., and Brune, A. (2000) Characterization of abundance and diversity of lactic acid bacteria in the hindgut of wood- and soil-feeding termites by molecular and culture-dependent techniques. *Arch Microbiol* 173:126-37.
- Broderick, N.A., Raffa, K.F., Goodman, R.M., and Handelsman, J. (2004) Census of the bacterial community of the Gypsy Moth larval midgut by using culturing and culture-independent methods. *App Environ Microbiol* 70: 293-300
- Brune, A. (1998) Termite guts: the world's smallest bioreactors. *Trends Biotechnol* 16: 16-21.
- Calinski, T., and Harabasz, J. (1974) A dendrite method for cluster analysis. *Communications in Statistics*. 3: 1-27.
- Caporaso, J.G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F.D., et al. (2010) QIIME allows analysis of high-throughput community sequencing data. *Nat Methods* 7: 335-336.
- Chandler, J.A., Lang, J.M., Bhatnagar, S., Eisen, J.A. and Kopp, A. (2011) Bacterial communities of diverse *Drosophila* species: ecological context of a host-microbe model system. *PLoS Genet* 7: e1002272.
- Chao, A. (1984) Nonparametric estimation of the number of classes in a population. *Scand J Stat* 11: 265-270.
- Chao, A., and Lee, S.M. (1992) Estimating the number of classes via sample coverage. *J Am Stat Assoc* 87: 210-217.

- Chao, A., Chazdon, R.L., Colwell, R.K., and Shen, T. (2005) A new statistical approach for assessing similarity of species composition with incidence and abundance data. *Ecology Letters* 8: 148-159.
- Clarke, K. (1993) Non-parametric multivariate analysis of changes in community structure. *Aust J Ecol* 18: 117-143.
- Dixon, P. (2003) VEGAN, a package of R functions for community ecology. *J Veg Sci* 6: 927-930.
- Douglas, A.E. (2009) The microbial dimension in insect nutritional ecology. *Funct Ecol* 23: 38-47.
- Edgar, R.C. (2010) Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* 26: 2460-2461.
- Fontenot, E.A., Arthur, F.H., Nechols, J.R., and Throne, J.E. (2012) Using a population growth model to simulate response of *Plodia interpunctella* Hübner to temperature and diet. *J. Pest Sci.* 85: 163-167.
- Garcia, E.S., Castro, D.P., Figueiredo, M.B., and Azambuja, P. (2010) Immune homeostasis to microorganisms in the guts of triatomines (Reduviidae): a review. *Mem Inst Oswaldo Cruz* 105: 605-610.
- Hamlin, J.C., Reed, W.D., and Phillips, M.E. (1931) Biology of the Indian meal moth on dried fruits in California. USDA Technical Bulletin No. 242, pp. 27.
- Hui, W., Gel, Y.R., and Gastwirth, J.L. (2008) Lawstat: an R package for law, public policy and biostatistics. *J Stat Softw* 28: 1-26.
- Iasur-Kruh, L., Taha-Salaime, L., Robinson, W.E., Sharon, R., Droby, S., Perlman, S.J., and Zchori-Fein, E. (2015) Microbial associates of the vine mealybug *Planococcus ficus* (Hemiptera: Pseudococcidae) under different rearing conditions. *Microb Ecol.* 69: 204-214.

Jia, S., Zhang, X., Zhang, G., Yin, A., Zhang, S., et al. (2013) Seasonally variable intestinal metagenomes of the red palm weevil (*Rhynchophorus ferrugineus*). *Environ Microb* 15: 3020-3029.

Johnson, J.A., Wofford, P.L., and Whitehand, L.C. (1992) Effect of diet and temperature on development rates, survival and reproduction of the Indian meal moth (Lepidoptera: Pyralidae). *J Econ Entomol* 85: 561-566.

Kane, M.D., and Breznak, J.A. (1991) Effect of host diet on production of organic acids and methane by cockroach gut bacteria. *Appl Environ Microbiol* 57: 2628-2634.

Kanehisa, M., Goto, S., Sato, Y., Furumichi, M., and Tanabe, M. (2012) KEGG for integration an interpretation of large-scale molecular data sets. *Nucl Acids Res* 40: D109-D114.

Köhler, T., Dietrich, C., Scheffrahn, R.H., and Brune, A. (2012) High-resolution analysis of gut environment and bacterial microbiota reveals functional compartmentation of the gut in wood-feeding higher termites (*Nasutitermes* spp.). *Appl Environ Microb* 7: 4691-4701.

Kruskal, J.B. (1964) Nonmetric multidimensional scaling: a numerical method. *Psychometrika* 29:115-129.

Kruskal, W.H., and Wallis, W.A. (1952) Use of Ranks in One-Criterion Variance Analysis. *J Am Stat Assoc* 47: 583-621.

Langille, M.G.I, Zaneveld, J., Caporaso, J.G., McDonald, D., Knights, D., et al. (2013) Predictive functional profiling of microbial communities using 16s rRNA marker gene sequences. *Nat Biotechnol* 31: 814-821.

Levene, H. (1960) Robust tests for equality of variances. In: *Contributions to Probability and Statistics: Essays in Honor of Harold Hotelling*. Olkin, I.,

- Ghurye, S.G., Hoeffding, W., Madow, W.G. & Mann, H.B. (eds). Stanford University Press, Stanford, CA. pp: 278–292.
- López-Sánchez, M.J., Neef, A., Peretó, J., Patiño-Navarrete, R., Pignatelli, M., Latorre A., and Moya, A. (2009) Evolutionary convergence and nitrogen metabolism in *Blattabacterium* strain Bge, primary endosymbiont of the cockroach *Blattella germanica*. PLoS Genet. 5: e1000721.
- Lozupone, C.A., Hamady, M., Kelley, S.T., and Knight, R. (2007) Quantitative and qualitative β diversity measures lead to different insights into factors that structure microbial communities. Appl Environ Microbiol. 73: 1576-1585.
- Mann, H.B., and Whitney, D.R. (1947) On a test of whether one of two random variables is stochastically larger than the other. Ann Math Stat 18: 50-60.
- Mason, C.J. and Raffa, K.F. (2014) Acquisition and structuring of midgut bacterial communities in gypsy moth (Lepidoptera: Erebidae) larvae. Environ Entomol 43: 595-604.
- Mazza, G., Chouaia, B., Lozzia, G.C., Montagna, M. (2014). The bacterial community associated to an Italian population of *Psacotha hilaris*: a preliminary study. Bull Insectology 67: 281-285.
- Mbata, G.N. (1985) Some physical and biological factors affecting oviposition by *Plodia interpunctella* (Hübner) (Lepidoptera: Phycitidae). Insect Sci App 6: 597-604.
- McCutcheon, J.P., McDonald, B.R., and Moran, N.A. (2009) Convergent evolution of metabolic roles in bacterial co-symbionts of insects. Proc Natl Acad Sci U.S.A. 106: 15394-15399.
- McFarlane, J.E., and Alli, I. (1986) Aggregation of larvae of *Blattella germanica* (L.) by lactic acid present in excreta. J Chem Ecol 12: 1369-1375.

Mikaelyan, A., Dietrich, C., Köhler, T., Poulsen, M., Sillam-Dussès, D., and Brune, A. (2015) Diet is the primary determinant of bacterial community structure in the guts of higher termites. *Mol Ecol.* 24: 5284-5295.

Mohandass, S., Arthur, F.H., Zhu, K.Y., and Throne, J.E. (2007) Biology and management of *Plodia interpunctella* (Lepidoptera: Pyralidae) in stored products. *J Stored Prod Res.* 43: 302-311.

Montagna, M., Chouaia, B., Mazza, G., Prosdocimi, E.M., Crotti, E., Mereghetti, V., et al. (2015a) Effects of the diet on the microbiota of the red palm weevil (Coleoptera: Dryophthoridae). *PLoS One* 10: e0117439.

Montagna, M., Gómez-Zurita, J., Giorgi, A., Epis, S., Lozzia, G., and Bandi, C. (2015b) Metamicrobiomics in herbivore beetles of the genus *Cryptocephalus* (Chrysomelidae): toward the understanding of ecological determinants in insect symbiosis. *Insect Sci* 22: 340-352.

Moran, N.A., Plague, G.R., Sandström, J.P., and Wilcox, J.L. (2003) A genomic perspective on nutrient provisioning by bacterial symbionts of insects. *Proc Natl Acad Sci U.S.A.* 100: 14543-14548.

Moran, N.A. (2006) Symbiosis. *Curr Biol* 16: 866-871.

Nansen, C., Phillips, T.W., and Palmer, M.W., (2004) Analysis of the insect community in a stored-maize facility. *Ecol Res* 19: 197-207.

Olofsson, T.C., Vásquez, A., Sammataro, D., and Macharia, J. (2011) A scientific note on the lactic acid bacterial flora within the honeybee subspecies; *Apis mellifera* (Buckfast), *A. m. scutellata*, *A. m. mellifera*, and *A. m. monticola*. *Apidologie*: 1-4.

Pérez-Cobas, A.E., Maiques, E., Angelova, A., Carrasco, P., Moya, A., and Latorre, A. (2015) Diet shapes the gut microbiota of the omnivorous cockroach *Blattella germanica*. FEMS Microbiol Ecol 91: fiv022.

Pielou, E.C. (1975) Ecological Diversity. Wiley & Sons, New York.

Pinto-Tomás, A.A., Sittenfeld, A., Uribe-Lorío, L., Chavarría, F., Mora, M., Janzen, D.H., Goodman, R.M., and Simon, H.M. (2011) Comparison of midgut bacterial diversity in tropical caterpillars (Lepidoptera: Saturniidae) fed on different diets. Environ Entomol. 40: 1111-1122.

Priya, N.G., Ojha, A., Kajla, M.K., Raj, A., and Rajagopal, R. (2012) Host plant induced variation in gut bacteria of *Helicoverpa armigera*. PLoS One. 7: e30768.

Rees, D., (2004) Insects of Stored Products. CSIRO Publishing, Collingwood, Victoria, Australia.

Rousseeuw, P.J. (1987) Silhouettes: A graphical aid to the interpretation and validation of cluster analysis. J Comp App Math 20: 53-65.

Ruokolainen, L., Ikonen, S., Makkonen, H., and Hanski, I. (2016) Larval growth rate is associated with the composition of the gut microbiota in the Glanville fritillary butterfly. Oecologia doi:10.1007/s00442-016-3603-8 [Epub ahead of print].

Schleifer, K.A., and Bell J.A. (2009) Family VIII. Staphylococcaceae fam. nov. Karl-Heinz Schleifer and Julia A. Bell. In: De Vos, P., Garrity, G.M., Jones, D., Krieg, N.R., Ludwig, W., Rainey, F.A., Schleifer K.H., Whitman, W.B. (Eds), Bergey's manual of systematic bacteriology 2nd edition volume 3 – The Firmicutes. Springer-Verlag, New York, pp. 392-433.

- Sedlacek, J.D., Weston P.A., and Barney, J. (1996) Lepidoptera and Psocoptera. In: Subramanyam, Bh., Hagstrum, D.W. (Eds.), Integrated Management of Insects in Stored Products. Marcel Dekker, Inc., New York, pp. 41-70.
- Shannon, C.E. (1948) A mathematical theory of communication. Bell Sys Tech J 27: 379-423, 623-656.
- Stampini M., and Locatelli D.P. (2007) Considerazioni sulle preferenze alimentari di *Idaea inquinata* (Lepidoptera Geometridae) per la messa a punto di una dieta artificiale. In: XXI Congresso Nazionale di Entomologia, Campobasso, Italy, 11-16 June 2007, p. 322.
- Sudakaran, S., Retz, F., Kikuchi, Y., Kost, C., and Kaltenpoth, M. (2015) Evolutionary transition in symbiotic syndromes enabled diversification of phytophagous insects on an imbalanced diet. ISME Journal 9: 2587-2604.
- Tang, X., Freitak, D., Vogel, H., Ping, L., Shao, Y., Cordero, E.A., Andersen, G., Westermann, M., Heckel, D.G., and Boland, W. (2012) Complexity and variability of gut commensal microbiota in polyphagous lepidopteran larvae. PLoS One. 7: e36978.
- Tatusov, R.L., Koonin, E.V., and Lipman, D.J. (1997) A genomic perspective on protein families. Science 278: 631-637.
- Vallet-Gely, I., Lemaitre, B., and Boccard, F. (2008) Bacterial strategies to overcome insect defences. Nat Rev Micro 6: 302-313.
- Vásquez, A., Olofsson, T.C., and Sammataro, D. (2009) A scientific note on the lactic acid bacterial flora in honeybees in the USA – a comparison with bees from Sweden. Apidologie 40: 26-8.

Vásquez, A., Forsgren, E., Fries, I., Paxton, R.J., Flaberg, E., Szekely, L., and Olofsson, T.C. (2012) Symbionts as major modulators of insect health: lactic acid bacteria and honeybees. PLoS One 7: e33188.

Welte, C.U., de Graaf, R.M., van den Bosch, T.J., Op den Camp, H.J., van Dam, N.M. and Jetten M.S. (2016) Plasmids from the gut microbiome of cabbage root fly larvae encode SaxA that catalyses the conversion of the plant toxin 2-phenylethyl isothiocyanate. Environ Microbiol 18: 1379-1390.

Whiley, R.A., and Hardie, J.M. (2009) Family VI. Streptococcaceae Deibel and Seeley 1974, 490^{AL}. In: De Vos, P., Garrity, G.M., Jones, D., Krieg, N.R., Ludwig, W., Rainey, F.A., Schleifer K.H., and Whitman, W.B. Bergey's manual of systematic bacteriology 2nd edition volume 3 – The Firmicutes. (eds) Springer-Verlag, New York, pp. 655-735.

Yu, H., Wang, Z., Liu, L., Xia, Y., Cao, Y., and Yin, Y. (2008) Analysis of the intestinal microflora in *Hepialus gonggaensis* larvae using 16S rRNA sequences. Curr Microbiol 56: 391-396.

Table 1. Diversity indices estimated for the bacterial communities associated with the analysed samples: eggs, adults and food substrates.

Identifier	Gender	Food	S _{OBSERVED} ^a	S _{CHAO} ^b	Shannon <i>H'</i>	Pielou <i>J'</i>
Eggs	-	-	121	215.1	3.79	0.79
Pad.1F	+	artificial diet	216	368.8	2.63	0.49
Pad.2F	+	artificial diet	158	438.1	2.70	0.53
Pad.3F	+	artificial diet	264	535.4	2.75	0.49
Pad.1M	♂	artificial diet	140	217	2.37	0.48
Pad.2M	♂	artificial diet	196	359.8	2.65	0.50
Ppiz.1F	+	pizzoccheri	177	331.1	2.26	0.44
Ppiz.2F	+	pizzoccheri	105	181.2	2.34	0.50
Ppiz.3F	+	pizzoccheri	162	300.8	2.31	0.45
Ppiz.1M	♂	pizzoccheri	173	278.6	2.16	0.42
Ppiz.2M	♂	pizzoccheri	344	552	2.18	0.37
Pchil.1F	+	chili	107	197	3.79	0.81
Pchil.2F	+	chili	138	207.5	3.74	0.76
Pchil.3F	+	chili	87	153	3.44	0.77
Pchil.1M	♂	chili	44	70.3	2.91	0.77
Pchil.2M	♂	chili	89	114	2.92	0.65
Pfav.1F	+	fava bean	60	90	3.13	0.76
Pfav.2F	+	fava bean	68	102.5	3.06	0.73
Pfav.3F	+	fava bean	19	21.5	1.98	0.67
Pfav.1M	♂	fava bean	77	94.3	3.49	0.80
Pfav.2M	♂	fava bean	86	117	2.94	0.66
Pmor.1F	+	moringa	99	162	3.50	0.76
Pmor.2F	+	moringa	130	161.7	3.81	0.78
Pmor.3F	+	moringa	104	128.5	3.89	0.84
Pmor.1M	♂	moringa	111	166.5	3.46	0.74
Pmor.2M	♂	moringa	160	222.7	3.83	0.76
Food_ad		artificial diet	128	197.8	3.51	0.72
Food_piz		pizzoccheri	52	98.8	3.35	0.85
Food_chil		chili	80	111.1	2.32	0.53
Food_fav		fava bean	1	1	0	-
Food_mor		moring	2	2	0.64	0.92

^a Number of OTUs observed in the microbiota of each sample; ^b Number of OTUs estimated to be present in the microbiota of each sample. P: *Plodia interpunctella*; ad: artificial diet; piz: pizzocchero; chil: chili; fav: fava beans; mor: moringa leaves; M: male; F: female.

<i>Comomonas</i>	-	-	0.2	-	-	1	2.1	0.3	1.8	1.2	1.8	0.4	0.2	0.6	1	-	0.9	3.3	-	-	-	-	0.8	5.7	0.6	0.1	0.5	-	0.9	
<i>Tepidimonas</i>	-	-	0.3	-	-	-	0.2	-	-	0.1	-	-	0.5	-	0.1	2.9	-	2.1	6.3	2.8	1.4	-	-	12	0.1	-	0.6	-	-	0.2
<i>Haemophilus</i>	-	-	-	-	-	0.1	-	-	-	-	-	-	0.1	0.1	0.2	0.1	0.2	-	-	-	-	-	-	-	1	-	0.3	24.3	0.4	
<i>Acinetobacter</i>	0.3	-	0.6	-	-	3.7	8.2	11.7	7	8.4	2.5	2.5	2.7	2.4	0.5	-	-	0.4	-	0.4	1.1	-	-	-	2.5	-	0.4	1.9	-	0.1
<i>Pseudomonas</i>	11.9	20.6	32.4	-	10	10.6	8.4	7.7	17.5	14.3	5.4	11.6	5.6	5	2.7	3.3	5.1	8.8	0.2	1.6	-	-	-	0.8	0.1	2.5	2.5	1.3	2.1	2

^a A: artificial diet, p: pizzochero, c: chili powder, f: fava beans, m: moringa leaves.

Figure 1. Experimental design, taxonomic composition of the microbiotas and cumulative abundance of the dominant bacterial OTUs.

A. *Plodia interpunctella* from different populations (laboratory-reared and wild) were reared until adulthood on different substrates (artificial diet, *Moringa oleifera* leaves, *Vicia faba* beans, *Capsicum annum* powder, and white-black buckwheat noodles).

DNA was extracted from the eggs of the laboratory-reared population, from five adults (3 females and 2 males) of each group and from the food resources.

B. The taxonomical composition of the microbiotas associated with the analysed samples (eggs, adults and food substrate) reported at the phylum and family levels (upper and lower bars, respectively). The average abundance across the five adult specimens is reported; only the bacterial families with an average abundance $\geq 5\%$ in the IMM group are reported, and bacterial families with an abundance lower than 5% are grouped in the category *other families*.

C. Box-plot of the cumulative abundance of the dominant bacterial OTUs (first five) associated with the adult moths reared on different substrates; yellow: moringa leaves; lilac: fava beans; green: artificial diet; red: chili powder; and cyan: white-black buckwheat.

Figure 2. Non-metric multidimensional scaling analysis of the bacterial community structure using the Chao (A) and Weighted Unifrac (B) metrics. The colour of rectangles indicates the different food resources; yellow: moringa leaves; lilac: fava beans; green: artificial diet; red: chili powder; and cyan: white-black buckwheat. The circular open dots indicate the single organism while the crosses, where present, represent the identified OTUs. The blue lines connect the individual microbiotas to the centroid values of each group. The ellipses represent a 95% confidence area

around the mean of the group.

Figure 3. Histogram representing the taxonomic assignment of the 16S rRNA gene sequences to the bacterial family for the IMM eggs and adults. Only the bacterial families with an abundance $\geq 5\%$ in at least one moth specimen are reported, and bacterial families with an abundance lower than 5% are grouped in the category *other families*. Each bar corresponds to a single individual.

Figure 4. Boxplots of the bacterial entomotypes characterising the two groups of IMMs identified by the NMDS and clustering analyses. A. *Atopococcus* entomotype characterising the *P. interpunctella* specimens reared on an artificial diet and pizzoccheri. B. *Propionibacterium* entomotype associated with the moths fed chili, fava beans and moringa.

Figure 5. Mann-Whitney-Wilcoxon table reporting the categories of functional potential, inferred from the bacterial 16S rRNA gene sequences for which differences among the five-group were recovered by the Kruskal-Wallis test. Mann-Whitney-Wilcoxon test were performed to assess the differences in the pairwise comparisons of IMM groups: the white squares indicate a $P \geq 0.05$, while the grey squares indicate a $P < 0.05$. EIP: environmental information processing; GIP: genetic information processing; M: metabolism; U: unclassified; MT: membrane transport; FSD: folding sorting and degradation; RR: replication and repair; T: translation; AAM: amino acid metabolism; BOSM: biosynthesis of other secondary metabolites; CM: carbohydrate metabolism; EM: energy metabolism; GBM: glycan biosynthesis and metabolism; LM: lipid metabolism; MCV: metabolism of cofactors and vitamins; MOAA:

metabolism of other amino acids; MTP: metabolism of terpenoids and polyketides; XBM: xenobiotics biodegradation and metabolism; CPS: cellular processes and signaling.

Figure 6. Co-occurrence OTUs table. For each IMM group, the co-occurrence table reports the IMM bacterial OTUs shared between the eggs from the laboratory-established population (left) and the food resource used for larval development (right). The colour of the rectangles indicates the different IMM groups: yellow: moringa leaves; lilac: fava beans; green: artificial diet; red: chili powder; and cyan: white-black buckwheat. The red squares of the table indicate the OTU co-occurrence, while the black squares indicate no co-occurrence of the bacterial OTU.

Supplementary figure S1. Venn diagram pairs showing the bacterial OTUs (at 97% similarity) shared by eggs and moth groups. The presence of a bacterial OTU is assigned to the group of insects feeding on the same resources when it is reported for at least three specimens of the group.

Supplementary figure S2. Venn diagrams showing the bacterial OTUs (at 97% similarity) shared by eggs and the three groups of adult moths from the laboratory population. The presence of a bacterial OTU is assigned to the group of insects feeding on the same resources when it is reported for at least one specimen of the group (A) and at least in three specimens of the group (B).

Supplementary figure S3. Venn diagrams showing the bacterial OTUs (at 97% similarity) shared by the five groups of adult IMM. The presence of a bacterial OTU

is assigned to the group of insects feeding on the same resources when it is reported for at least three specimens of the group (A) and one specimen of the group (B).




Figure 2. Non-metric multidimensional scaling analysis of the bacterial community structure using the Chao (A) and Weighted UniFrac (B) metrics. The colour of rectangles indicates the different food resources; yellow: moringa leaves; lilac: fava beans; green: artificial diet; red: chili powder; and cyan: white-black buckwheat. The circular open dots indicate the single organism while the crosses, where present, represent the identified OTUs. The blue lines connect the individual microbiotas to the centroid values of each group. The ellipses represent a 95% confidence area around the mean of the group.

78x36mm (300 x 300 DPI)

Accepted




Figure. 3. Histogram representing the taxonomic assignment of the 16S rRNA gene sequences to the bacterial family for the IMM eggs and adults. Only the bacterial families with an abundance $\geq 5\%$ in at least one moth specimen are reported, and bacterial families with an abundance lower than 5% are grouped in the category other families. Each bar corresponds to a single individual.
149x112mm (300 x 300 DPI)

Accep




Figure 4. Boxplots of the bacterial entomotypes characterising the two groups of IMM identified by the NMDS and clustering analyses. A. Atopococcus entomotype characterising the *P. interpunctella* specimens reared on an artificial diet and pizzoccheri. B. Propionibacterium entomotype associated with the moths fed chili, fava beans and moringa.
44x11mm (300 x 300 DPI)

Accepted A




Figure 5. Mann-Whitney-Wilcoxon table reporting the categories of functional potential, inferred from the bacterial 16S rRNA gene sequences for which differences among the five-group were recovered by the Kruskal-Wallis test. Mann-Whitney-Wilcoxon test were performed to assess the differences in the pairwise comparisons of IMM groups: the white squares indicate a $P \geq 0.05$, while the grey squares indicate a $P < 0.05$. EIP: environmental information processing; GIP: genetic information processing; M: metabolism; U: unclassified; MT: membrane transport; FSD: folding sorting and degradation; RR: replication and repair; T: translation; AAM: amino acid metabolism; BOSM: biosynthesis of other secondary metabolites; CM: carbohydrate metabolism; EM: energy metabolism; GBM: glycan biosynthesis and metabolism; LM: lipid metabolism; MCV: metabolism of cofactors and vitamins; MOAA: metabolism of other amino acids; MTP: metabolism of terpenoids and polyketides; XBM: xenobiotics biodegradation and metabolism; CPS: cellular processes and signaling.


208x261mm (300 x 300 DPI)



Figure 6. Co-occurrence OTUs table. For each IMM group, the co-occurrence table reports the IMM bacterial OTUs shared between the eggs from the laboratory-established population (left) and the food resource used for larval development (right). The colour of the rectangles indicates the different IMM groups: yellow: moringa leaves; lilac: fava beans; green: artificial diet; red: chili powder; and cyan: white-black buckwheat. The red squares of the table indicate the OTU co-occurrence, while the black squares indicate no co-occurrence of the bacterial OTU.


220x501mm (300 x 300 DPI)

A



Supplementary figure S1. Venn diagram pairs showing the bacterial OTUs (at 97% similarity) shared by eggs and moth groups. The presence of a bacterial OTU is assigned to the group of insects feeding on the same resources when it is reported for at least three specimens of the group.
51x28mm (600 x 600 DPI)


Accepted



Supplementary figure S2. Venn diagrams showing the bacterial OTUs (at 97% similarity) shared by eggs and the three groups of adult moths from the laboratory population. The presence of a bacterial OTU is assigned to the group of insects feeding on the same resources when it is reported for at least one specimen of the group (A) and at least in three specimens of the group (B).

102x111mm (300 x 300 DPI)

AC



Supplementary figure S3. Venn diagrams showing the bacterial OTUs (at 97% similarity) shared by the five groups of adult IMM. The presence of a bacterial OTU is assigned to the group of insects feeding on the same resources when it is reported for at least three specimens of the group (A) and one specimen of the group (B).

145x222mm (300 x 300 DPI)

A