

1 <https://doi.org/10.1017/S1751731118001234>

2  
3 **The effects of superoxide dismutase-rich melon pulp concentrate on**  
4 **inflammation, antioxidant status and growth performance of challenged post-**  
5 **weaning piglets**

6  
7 A.S.M.L. Ahasan<sup>1a</sup>, G. Invernizzi<sup>1</sup>, G. Farina<sup>1</sup>, A. Pilotto<sup>1</sup>, F. Barbé<sup>2</sup>, V. Bontempo<sup>1</sup>, R.  
8 Rossi<sup>1</sup>, F. Bellagamba<sup>1</sup>, C. Lecchi<sup>3</sup>, G. Savoini<sup>1</sup> and A. Agazzi<sup>1</sup>

9  
10 <sup>1</sup>*Dipartimento di Scienze Veterinarie per la Salute, la Produzione Animale e la*  
11 *Sicurezza Alimentare, Università degli Studi di Milano, 20133 Milano, Italy.*

12 <sup>2</sup>*Lallemand SAS, 31700 Blagnac, France.*

13 <sup>3</sup>*Dipartimento di Medicina Veterinaria, Università degli Studi di Milano, 20133 Milano,*  
14 *Italy.*

15  
16 <sup>a</sup>*Present address: Department of Anatomy and Histology. Faculty of Veterinary*  
17 *Medicine, Veterinary and Animal Sciences University. 4225, Chittagong. Bangladesh.*

18  
19 Corresponding author: Alessandro Agazzi. E-mail: [alessandro.agazzi@unimi.it](mailto:alessandro.agazzi@unimi.it)

20  
21 Short title: Melon pulp concentrate fed to challenged piglets

## 22 **Abstract**

23 Piglets can often suffer impaired antioxidant status and poor immune response during  
24 post-weaning, especially when chronic inflammation takes place, leading to lower  
25 growth rates than expected. Oral administration of dietary antioxidant compounds  
26 during this period could be a feasible way to balance oxidation processes and increase  
27 health and growth performance. The aim of the trial was to study the effects of an  
28 antioxidant feed supplement (melon pulp concentrate) that contains high concentration  
29 of the antioxidant superoxide dismutase (**SOD**) on inflammation, antioxidant status and  
30 growth performance of lipopolysaccharide (**LPS**) challenged weaned piglets. Forty-  
31 eight weaned piglets were individually allocated to four experimental groups in a 2 x 2  
32 factorial design for 29 days. Two different dietary treatments were adopted: a) Control  
33 (**CTR**), fed a basal diet, b) Treatment (**MPC**), fed the basal diet plus 30g/ton of melon  
34 pulp concentrate. On days 19, 21, 23 and 25 half of the animals within CTR and MPC  
35 groups were subjected to a challenge with intramuscular injections of an increasing  
36 dosage of LPS from *E. coli* (serotype 0.55:B5) (+) or were injected with an equal  
37 amount of PBS solution (-). Blood samples were collected at the beginning of the trial  
38 and under the challenge period for interleukin 1 $\beta$ , interleukin 6, tumour necrosis factor  
39  $\alpha$ , haptoglobin, plasma SOD activity, total antioxidant capacity, reactive oxygen  
40 species, red blood cells and plasma resistance to haemolysis, and 8-oxo-7, 8-dihydro-  
41 2'-deoxyguanosine. Growth performance was evaluated weekly. A positive effect of  
42 melon pulp concentrate was evidenced on total antioxidant capacity, half-haemolysis  
43 time of red blood cells, average daily gain and feed intake, while LPS challenge  
44 increased proinflammatory cytokines and haptoglobin serum concentrations, with a  
45 reduced feed intake and gain:feed. The obtained results show that oral SOD  
46 supplementation with melon pulp concentrate ameliorates the total antioxidant capacity

47 and the half-haemolysis time in red blood cell of post-weaning piglets, with positive  
48 results on growing performance.

49

50 **Keywords:** Cucumis melo, total antioxidant capacity, growth, lipopolysaccharide, pig.

51

## 52 **Implications**

53 Piglets can show impaired antioxidant status and inflammation processes during post-  
54 weaning, leading to lower growth rates than expected. This is especially true when  
55 occasional presence of pathogens in the farm causes a chronic inflammation status.  
56 In this context superoxide dismutase administration could be a feasible way to  
57 overcome inflammation, impaired antioxidant status, and low performance. The  
58 present study demonstrates that oral supplementation with melon pulp concentrate,  
59 rich in superoxide dismutase, is able to increase the total antioxidant capacity of  
60 weaned pigs, contributing to sustain the health status and the growth rate of these  
61 animals.

62

## 63 **Introduction**

64 Inflammation and oxidative stress are closely linked together (Kick *et al.*, 2012;  
65 Carillon *et al.*, 2013a) and lead to decreased growth rates and feed efficiency in  
66 weaned piglets. This is mainly due to the imbalance in oxidant/antioxidant equilibrium,  
67 the decreased activity of major antioxidant enzymes, and the increase in radical-  
68 mediated lipid peroxidation and protein and DNA oxidation (Campbell *et al.*, 2013;  
69 Gessner *et al.*, 2017). In this context, the use of dietary antioxidant compounds after  
70 weaning seems to be a feasible way to overcome impaired antioxidant status and poor  
71 immune response in piglets (Bontempo *et al.*, 2014; Jiang *et al.*, 2015a,b). Nowadays

72 there is great interest in the therapeutic application of dietary superoxide dismutase  
73 (**SOD**) (Carillon *et al.*, 2013a,b), as a primary antioxidant molecule. A specific  
74 cantaloupe melon (*Cucumis melo L.*) from the *Cucurbitaceae* family, known to be  
75 characterized by high SOD activity, showed antioxidant and anti-inflammatory  
76 properties in *in vitro* and *in vivo* animal models such as rodents and horses (Vouldoukis  
77 *et al.*, 2004a,b; Notin *et al.*, 2010).

78 Although some trials have been conducted on orally melon concentrate  
79 administration in different species, at the present moment the effects of feeding SOD-  
80 rich melon pulp concentrate on piglets subjected to stress are only reported in two  
81 studies by Lallès *et al.* (2011) and Royer *et al.* (2016) on stress proteins along the  
82 gastrointestinal tract and changes in blood oxidative stress biomarkers, respectively.  
83 At experimental level, oxidative and inflammatory stress conditions can be mimed by  
84 chronic immune system stimulation with an increasing dose of lipopolysaccharide  
85 (**LPS**) challenge (Rakhshandeh and de Lange, 2012). The aim of the present study is  
86 to evaluate the effects of oral supplementation of a SOD-rich melon pulp concentrate  
87 in increasing antioxidant status, inflammation response, and growth performance in  
88 LPS-chronically challenged weaning piglets.

89

## 90 **Material and methods**

91

### 92 *Animals housing and experimental design*

93 The present trial was performed at the Centro Clinico-Veterinario e Zootecnico-  
94 Sperimentale d'Ateneo di Lodi, Università degli Studi di Milano. In total, 48 crossbred  
95 female piglets (Topigs 40 x Topdelta) from the same herd were weaned at  $24 \pm 1$  days  
96 of age (BW  $7.79 \pm 0.17$  kg) and divided in four homogeneous experimental groups of

97 twelve animals each in a 2 x 2 factorial arrangement. The piglets were placed in  
98 individual pens (0.47 m<sup>2</sup>) and allocated in the same environmentally-controlled post-  
99 weaning room on slatted floor. Each pen was equipped with one standard nursery pig  
100 bite-style nipple drinker and a self-feeder to allow for *ad libitum* access to water and  
101 feed. Room temperature and ventilation were electronically controlled over a 24 hours  
102 period. Starting room temperature was 28°C with a ventilation of 10 m<sup>3</sup>/hour/piglet and  
103 was decreased by 1°C/week until 25°C at the end of the trial.

104         The first factorial arrangement consisted of the administration of a basal diet or  
105 the same basal diet plus melon pulp concentrate (Melofeed, Lallemand SAS, Blagnac,  
106 France). The second factorial arrangement consisted on a LPS challenge, with  
107 repeated increasing intramuscular injections of LPS from *E. coli* (serotype 055:B5,  
108 Sigma-Aldrich Canada Ltd, Oakville, ON, Canada; cat. no. L2880) (+) to mimic chronic  
109 inflammation, or the injection of an equivalent amount of PBS solution (-). The  
110 challenge was performed starting on day 19 of the trial and subsequent injections were  
111 performed on days 21, 23 and 25. The initial LPS dosage of 60 µg/kg of BW was  
112 increased by 12% at each subsequent injection to reduce endotoxin tolerance  
113 (Rakhshandeh and de Lange, 2012). Specifically, the applied concentrations of LPS  
114 from the second to the fourth injection day were 67.2, 75.26, and 84.30 µg/kg of BW.  
115 Individual body weight was determined prior to each LPS injection to calculate  
116 individual total LPS amount to be injected.

117

118 *Experimental diets*

119 To avoid any other influencing stress factor besides the weaning and the  
120 challenge, in the present trial pigs were fed with a mash wheat-based basal diet (Table  
121 1) for all the trial period with (**MPC**) or without (**CTR**) the inclusion of 30 g/ton of melon  
122 pulp concentrate. Melon pulp concentrate was provided in powder form and  
123 incorporated through the premix at 1.5 kg/ton of feed to get the final content of 30g of  
124 Melofeed/ton of feed (Lalleman SAS, Blagnac, France). The applied dosage of melon  
125 pulp concentrate was based on the previous publication of Lallès *et al.* (2001) where  
126 Promutase (former name of Melofeed) was supplemented at 5 and 20 g/ton feed.  
127 Because a decrease of feed intake was expected due to the LPS challenge, in the  
128 present trial the dosage was increased to 30 g/ton feed during the whole experimental  
129 period.

130 The experimental diets were formulated to be isocaloric and isoproteic on net  
131 energy and CP basis, and were produced with the same batches of feeds by  
132 Tracciaverde S.R.L. (Bonemerse, Italy). Both CTR and MPC diets did not contain any  
133 antimicrobials or growth promoters, being designed to meet or exceed the nutrient  
134 requirements of weaned piglets recommended by National Research Council (NRC,  
135 2012), with the exception of suboptimal concentration of tryptophan. In the present trial  
136 no inclusion of tryptophan was performed besides its content in the feeds in order to  
137 amplify the inflammation effect of the LPS challenge performed in the experimental  
138 animals (Le Floc'h and Seve, 2007).

139

140 *Samples and data collection*

141 Blood samples for pro-inflammatory interleukin 1 $\beta$  (**IL-1 $\beta$** ), interleukin 6 (**IL-6**)  
142 and tumour necrosis factor  $\alpha$  (**TNF- $\alpha$** ) and haptoglobin were collected on days 0, 19,

143 21, 23, 25, and 29, while blood samples for plasma SOD activity, total antioxidant  
144 capacity (**TAOC**), and reactive oxygen species (**ROS**) serum content were obtained  
145 on days 0, 19, 21, 23, 25, 27, and 29. Each blood sample under the challenge period  
146 was collected prior the LPS injection. For sampling procedures a 10 mL clot activator  
147 vacutainer tube (VF-109SP, Venoject®, Terumo Europe N.V., Leuven, Belgium) was  
148 used to yield serum from the cranial vena cava. Blood sample for plasma SOD activity  
149 was collected separately using a 4 mL vacutainer tube containing EDTA (VF054STK,  
150 Venoject®, Terumo Europe N.V., Leuven, Belgium). The serum was obtained by  
151 allowing the whole blood to clot at room temperature for 30 min. The tubes were then  
152 centrifuged at 1500 x g, 10 min at 4°C, the supernatant was placed in 2 mL Eppendorf  
153 tubes (Eppendorf AG, Amburg, Germany) and subsequently stored at -80°C to prevent  
154 changes in antioxidant and inflammation biomarkers. Blood samples for the whole  
155 blood and red blood cells (**RBC**) resistance to haemolysis and plasma contribution by  
156 Kit Radicaux Libres (**KRL**), and 8-oxo-7, 8-dihydro-2'-deoxyguanosine (**8-oxodGuo**)  
157 were collected on days 19, 25, and 29 of the trial. Blood samples for KRL test were  
158 collected by 4 mL vacutainer tubes containing EDTA (VF054STK, Venoject®, Terumo  
159 Europe N.V., Leuven, Belgium) (Rossi *et al.*, 2013), immediately stored at 4°C,  
160 processed within three hours from sampling, and analysed in the next 24 hours after  
161 collection. Blood samples for 8-oxodGuo were collected with a 10 mL clot activator  
162 vacutainer tube (VF-109SP, Venoject®, Terumo Europe N.V., Leuven, Belgium) and  
163 serum was obtained by centrifugation at 1500 x g, 10 min at 4°C. Serum was then  
164 placed in a 2 mL Eppendorf tube (Eppendorf AG, Amburg, Germany), and stored at -  
165 80°C for pending analysis.

166 A sample of the basal diet was collected at the beginning of the trial and stored  
167 at -20°C for pending analyses. Individual piglet BW was recorded on days 0, 8, 15, 19,

168 21, 23, 25, 26 and 29 with an electronic scale (ES100L, Ohaus, Switzerland), and  
169 individual feed intake (**FI**) for 0-19 and 19-29 days periods was calculated by  
170 subtracting the relative excreta to the total daily-administered amount of feed.  
171 Subsequently average daily gain (**ADG**) and gain:feed (**G:F**) were calculated. The  
172 health status of the piglets was daily checked and any sanitary treatment was recorded.  
173 Morbidity and mortality were recorded. Rectal temperature was manually measured  
174 before each LPS injection as a baseline measurement, after two hours from each LPS  
175 injection, and on the last day of trial.

176

### 177 *Chemical analysis*

178 Serum pro-inflammatory cytokines IL-1 $\beta$ , IL-6 and TNF- $\alpha$  were measured by  
179 porcine-specific ELISA kit according to the recommendations of the manufacturer (R  
180 and D Systems Inc., Abingdon Science Park, UK). The serum concentrations of  
181 haptoglobin were determined by colorimetric assay (Tridelta Phasorange serum  
182 haptoglobin assay, Cat. No. TP-801) and expressed on the basis of a standard curve  
183 (Cooke and Arthington, 2013). All samples were assayed in duplicate. Intra-assay  
184 coefficient of variation were 5.71%, 6.33%, 5.83%, and 7.41%, while inter-assay  
185 coefficient of variation were 5.59%, 6.38%, 5.34%, and 6.18% respectively for IL-1 $\beta$ ,  
186 IL-6, TNF- $\alpha$ , and haptoglobin. All the intra- and inter-assay coefficients of variations  
187 were within the range of values declared by the commercial product datasheet. Plasma  
188 SOD activity and serum TAOC were measured using commercial available kits  
189 according to the manufacturer's instructions (Sigma-Aldrich, Cat. No. 19160 and Cat.  
190 No. CS0790, respectively), while ROS were evaluated by Cyt C reduction assay  
191 (Sigma Aldrich, cat. No. C2506), as reported by Sartorelli *et al.* (2000). The total  
192 antiradical activity of whole blood, RBC and the plasma contribution was determined

193 using the KRL biological test (Rossi *et al.*, 2013). Results were expressed as the time  
194 (min) that is required to reach 50% of maximal haemolysis. Half haemolysis time for  
195 total blood cells (**HT<sub>50</sub>WB**) and for red blood cells (**HT<sub>50</sub>RBC**) refers to the whole blood  
196 and the red cell resistance to free radical attack, respectively. Lastly, the plasma  
197 resistance to haemolysis (**HT<sub>50</sub>PC**) was calculated by subtracting HT<sub>50</sub>RBC from  
198 HT<sub>50</sub>WB. Serum concentration of 8-oxodGuo were determined using a competitive  
199 ELISA method based on monoclonal antibody highly specific to 8-OHdG (Japan  
200 Institute for Control Aging Fukuroi, Japan) following the manufacturer's instructions.  
201 For 8-oxodGuo assays six different standard dilutions were used in triplicate for each  
202 ELISA plate. Intra- and inter -assay coefficients of variation were respectively 5.56%  
203 and 8.86%.

204 The chemical composition of the basal diet was analysed at the beginning of the  
205 trial to determine DM (method 930.15), CP (method 984.13), ether extract (method  
206 920.39A), ash (method 942.05), Ca (method 968.08) and P (method 946.06) content  
207 following the relative Association of Official Analytical Chemists methods of analysis  
208 (AOAC, 2005) (Table 1). Neutral detergent fibre content in the diet was determined as  
209 reported by Van Soest *et al.* (1991).

210

### 211 *Statistical analyses*

212 Data relative to inflammatory and oxidative biomarkers, BW, and rectal  
213 temperature were performed by ANOVA using the Proc MIXED for repeated  
214 measurements in SAS (SAS Inst. Inc., Cary, NC), with the piglet as the experimental  
215 unit. The statistical model included the effects of LPS challenge, dietary treatment  
216 (CTR, MPC), time and their interaction as fixed effects. Average daily gain, FI and G:F  
217 considered two different trial periods corresponding to days 0-19 (pre-challenge) and

218 19-29 (challenge and post-challenge) of the trial. Statistical analyses for these  
219 parameters were performed by a GLM procedure in SAS with the piglet as the  
220 experimental unit and the corresponding data relative to the pre-challenge period (0-  
221 19 days) as covariate for 19-29 days period. Differences were considered significant  
222 for  $P < 0.05$ . A tendency toward a significant difference between treatment means was  
223 also considered at  $P < 0.10$ .

224

## 225 **Results**

226

### 227 *Inflammatory and oxidative biomarkers status*

228 In the present trial LPS challenge induced a strong effect on inflammatory status  
229 with increased serum concentration of IL-1 $\beta$  (0.53 ng/mL vs. 0.21 ng/mL,  $P < 0.01$ ), IL-  
230 6 ( $6.99 \times 10^3$  ng/mL vs.  $3.10 \times 10^3$  ng/mL,  $P < 0.01$ ), TNF- $\alpha$  ( $2.56 \times 10^{-2}$  ng/mL vs.  $1.44$   
231  $\times 10^{-2}$  ng/mL,  $P < 0.01$ ), and haptoglobin ( $8.96 \times 10^5$  ng/mL vs.  $7.82 \times 10^5$  ng/mL,  
232  $P < 0.05$ ) (Figure 1), while melon pulp concentrate supplementation did not induce any  
233 significant change in these parameters. Time effect and the interaction between LPS  
234 challenge and time were always significant ( $P < 0.01$ ) (Figure 1). No significant  
235 differences between challenged or not challenged piglets were found at the beginning  
236 of the trial for IL-1 $\beta$ , IL-6, and TNF- $\alpha$ , while a higher haptoglobin content was found in  
237 piglets not subjected to LPS on day 0 ( $9.50 \times 10^5$  ng/mL vs.  $5.62 \times 10^5$  ng/mL;  $P < 0.01$ ).  
238 Starting from day 21 challenged groups showed increased concentration of all  
239 inflammation biomarkers until day 23 for haptoglobin or day 25 for interleukins and  
240 TNF- $\alpha$  ( $P < 0.01$ ).

241 Oxidative biomarkers were not affected by LPS challenge (Table 2), although  
242 some trends were found for decreased TAOC (5.30 mM Trolox equivalent vs. 6.46 mM

243 Trolox,  $P=0.08$ ), and increased HT<sub>50</sub>WB (113.78 min vs. 109.14 min,  $P=0.09$ ) and  
244 HT<sub>50</sub>PC (44.90 min vs. 40.89 min,  $P=0.08$ ). Melon pulp concentrate supplementation  
245 increased serum TAOC (7.22 mM Trolox equivalent vs. 4.54 mM Trolox equivalent,  
246  $P<0.01$ ) and HT<sub>50</sub>RBC (70.71 min vs. 66.41 min,  $P<0.01$ ), but did not induce significant  
247 effects on HT<sub>50</sub>WB, HT<sub>50</sub>PC, 8-oxo-dGuo concentration, SOD activity, and ROS.  
248 Sampling time was always significant ( $P<0.01$ ), with the exception of SOD activity and  
249 HT<sub>50</sub>PC, but no difference was detected considering the interaction between dietary  
250 treatment, challenge and time.

251

#### 252 *Growth performance and rectal temperature*

253 No significant effects of LPS challenge ( $P=0.47$ ) or melon pulp concentrate  
254 supplementation ( $P=0.70$ ) and their interactions ( $P=0.73$ ) were found on BW. The LPS  
255 challenge impaired ADG (291 g/d vs. 490 g/d,  $P<0.01$ ), FI (5.57 kg vs. 7.76 kg,  $P<0.01$ )  
256 and G:F (0.52 vs. 0.62,  $P<0.01$ ) from days 19 to 29. Average daily gain and FI were  
257 increased in pigs fed melon pulp concentrate under the challenge and post-challenge  
258 period (19-29 days on trial) (422 g/d vs. 359 g/d for ADG,  $P<0.05$ ; 7.18 kg vs. 6.15 kg  
259 for FI,  $P<0.01$ ) (Table 3), but no differences were found during the first 19 days of the  
260 trial and overall the experimental period. No significant differences were outlined for  
261 G:F among the dietary treatments, and the diet x challenge interaction was found not  
262 to affect growing performance.

263 Rectal temperature was always increased by LPS challenge at two hours after  
264 injection ( $P<0.01$ , data not shown), while no effect of dietary melon pulp concentrate  
265 supplementation was evidenced, with the exception of a significant reduction at two  
266 hours after the fourth LPS injection (day 25 of the trial) (CTR= 40.25 ± 0.09 °C vs.  
267 MPC= 40.12 ± 0.08 °C,  $P=0.02$ ). No differences in piglet's rectal temperature at the

268 end of the trial (day 29) were found between CTR and MPC groups, also considering  
269 the challenge effect or the diet x challenge interaction. During the trial no antibiotic  
270 treatments were performed on experimental animals and no deaths occurred within  
271 CTR and MPC groups.

272

## 273 **Discussion**

274 Weaning is a critical stress period for piglets and it is characterized by  
275 decreased immune response, FI, and nutrient absorption (Niekamp *et al.*, 2007).  
276 During this period, pigs are subjected to a number of stressors such as a sudden  
277 separation from the sow, transportation and handling, different environment, liquid to  
278 solid feed shift, establishment of a new social hierarchy, and increased exposure to  
279 pathogens. All these factors lead to metabolism, immune system, and intestinal  
280 functions alterations. Several studies report how serum pro-inflammatory cytokines  
281 and their gene expression are increased at weaning (Pié *et al.*, 2004; Lallès *et al.*,  
282 2011). In accordance with this, in the present trial initial high concentrations of IL-6 and  
283 TNF- $\alpha$  were found in all the experimental subjects, although they were decreased just  
284 before the LPS challenge (day 19). At the starting of the trial IL-6 showed half of the  
285 concentration further evidenced during the challenge, while TNF- $\alpha$  concentration  
286 detected on day 0 and during the LPS injection were comparable, outlining how  
287 weaning is able to strongly affect the immune response of piglets.

288 Pro-inflammatory cytokines IL-1 $\beta$ , IL-6, and TNF- $\alpha$  are reported to be important  
289 inducers of the synthesis of acute phase proteins, such as haptoglobin, among others  
290 (Carroll *et al.*, 2004). In the present trial a higher concentration of haptoglobin in both  
291 no-challenge groups compared to challenge groups was found at the beginning of  
292 melon pulp concentrate supplementation. Although all the experimental animals were

293 selected, grouped and placed in the experimental facilities at the same time, this initial  
294 difference seems to be unfortunately related to some un-accounted (not estimable)  
295 stress factors (Salamano *et al.*, 2008; Cray *et al.*, 2009) rather than a direct effect of  
296 high concentration of interleukins that could lead to increased levels of haptoglobin.

297 Besides the strong effect of weaning, occasional presence of pathogens in the  
298 farm can causes a chronic inflammation status that can further impairs the immune  
299 response and the antioxidant status of piglets. This last scenario can be efficiently  
300 obtained from an experimental point of view applying a chronic challenge in the post-  
301 weaning period. With this purpose, in the present trial a chronic challenge procedure  
302 with LPS was adopted to mimic subclinical or mild clinical disease conditions that  
303 frequently occur in the field due to the presence of pathogens that lead to immune  
304 system stimulation (Rakhshandeh and de Lange 2012; de Ridder *et al.*, 2012). As a  
305 result, in the present trial LPS was found to act on the immune system through  
306 increased concentration of serum pro-inflammatory cytokines IL-1 $\beta$ , IL-6, and TNF- $\alpha$   
307 and haptoglobin according to Carroll *et al.* (2001) and Frank *et al.* (2005), while the  
308 administration of melon pulp concentrate did not lead to any significant variation on  
309 immune response.

310 Inflammation and oxidative stress have been reported to be closely linked to  
311 each other, since the pathways generating the mediators of inflammation are all  
312 induced by oxidative stress. (Carillon *et al.*, 2013a). Oxidative stress decreases  
313 antioxidant defences and induces elevated radical-mediated lipid peroxidation driven  
314 by reactive oxygen and nitrogen species (Campbell *et al.*, 2013). At weaning, piglets  
315 can be subjected to an accumulation of ROS that could exceed the antioxidant  
316 capacities of the animal, leading to a high susceptibility to oxidative stress (Zhu *et al.*,  
317 2012).

318 As a strong effect of the LPS challenge on immune response can be highlighted  
319 in this trial, the general lack of significant changes in oxidative biomarkers, with the  
320 exception of a trend for decreased serum content of TAOC, can be attributed to other  
321 numerous factors. Moreover, independently from the applied dosage to mimic a  
322 chronic inflammation in the present trial, it must be noted that the effects of an immune  
323 challenge did not always lead to univocal results on oxidative stress biomarkers,  
324 depending on the quantification of different antioxidant or pro-oxidant components of  
325 oxidative stress (van de Crommenacker *et al.*, 2010). In the present trial oxidative  
326 markers such as plasma SOD activity, ROS and 8-oxo-dGuo were not influenced by  
327 melon pulp administration, while TAOC and HT<sub>50</sub>RBC were improved in MPC groups.  
328 If the obtained results on ROS can be attributed to their reported very short half-life  
329 and high instability, the SOD activity could have been influenced by several factors and  
330 can be interpreted in different ways, as the activity of major antioxidant enzymes (**AOX**)  
331 is quite variable depending on the experimental design adopted, the challenge  
332 performed or the animal species. Increased activity of AOX (SOD, CAT, GPx) can be  
333 related both to the reaction of the organism to oxidative stress, or to the stimulation of  
334 antioxidant defences when supplying antioxidant compounds in the diet. To  
335 discriminate between these two opposite interpretations, additional biomarkers of  
336 oxidative stress, such as oxidized proteins (e.g. protein carbonyls) or oxidized lipids  
337 (e.g. TBARS, MDA, isoprostanes, lipid peroxides) could have been measured. In fact  
338 a study by Royer *et al.* (2016) found that a decreased concentration of oxidative stress  
339 biomarkers and an increased activity of AOX confirmed the efficacy of melon pulp  
340 concentrate supplementation in pigs. Although dietary SOD mechanism of action is not  
341 fully understood at the present moment, Carillon *et al.* (2013a) suggested that melon  
342 pulp concentrate administration could induce increased antioxidant defence, as

343 outlined by TAOC and RBC blood resistance to haemolysis in the present trial, through  
344 the activation of mRNA transcription or the regulation or induction of complex cellular  
345 pathways, involving different transcription factors. These authors hypothesized that the  
346 induction of antioxidant enzymes could be regulated at the transcriptional level through  
347 the nuclear-factor-E2-related factor (Nrf2)/antioxidant response as demonstrated in  
348 some other studies with different antioxidant supplements in humans and *in vitro*  
349 (Muchová *et al.*, 2001; Zhang *et al.*, 2012).

350 In the present study, decreased FI and impaired growth performance during  
351 LPS challenge period were observed according with Gessner *et al.* (2017). Lower  
352 performance under the challenged corresponded to a higher concentration of  
353 inflammation markers and an increased rectal temperature of the piglets at two hours  
354 after each LPS injection as a sign of the presence of a systemic inflammation (Ceciliani  
355 *et al.*, 2012). When an infection occurs, metabolic shifts are characterized by the  
356 redistribution of nutrients away from the growth processes toward the immune system  
357 function, with a subsequent decrease in feed efficiency for growth (Gessner *et al.*,  
358 2017). Moreover the suboptimal concentration of tryptophan supplied in the present  
359 trial (approximately 78% of the requirements on total basis for piglets ranging from 7  
360 to 25 kg BW) (NRC, 2012) could have represent an amplification factor for the  
361 effectiveness of the applied challenge, since tryptophan metabolic demand is strongly  
362 increased by the synthesis of acute phase proteins during immune stimulation, with a  
363 consequent lower availability for growth (de Ridder *et al.*, 2012).

364 Besides the inflammation effect of the applied challenge, the administration of  
365 melon pulp concentrate led to increased performance, which instead was not observed  
366 by Lallès *et al.* (2011). It must be outlined that the total duration of administration in the  
367 work of Lallès *et al.* (2011) was 5 or 12 days, after an initial two-days period of fasting

368 to induce greater stress protein concentrations. Animals were then slaughtered on  
369 days 7 and 14 after weaning. The lack of improved performance in the work of Lallès  
370 *et al.* (2011) can be probably due to the short duration of melon pulp concentrate  
371 supplementation to translate any positive effect in the gastrointestinal tract or on  
372 immune response into improved growth performance.

373 In conclusion, the present study demonstrates that oral supplementation with  
374 melon pulp concentrate rich in SOD during the post-weaning period can enhances the  
375 total antioxidant capacity, half-haemolysis time of red blood cells and growth rate of  
376 post-weaning piglets when oxidative stress and/or inflammation are increased.

377

### 378 **Acknowledgments**

379 Authors would like to thank Dr. Eric Chevaux and Dr. Mathieu Castex (Lallemand SAS,  
380 Blagnac, France) for cooperation and research funding.

381

### 382 **Declaration of interest**

383 The authors declare no conflicts of interest

384

### 385 **Ethics statement**

386 The protocol for care, handling, and sampling of animals defined in the present  
387 study was reviewed and approved by the Università degli Studi di Milano Animal Care  
388 and Use Committee (Protocol No 82/14).

389

### 390 **Software and data repository resources**

391 Data or models from the present work are not deposited in an official repository

392

393 **References**

- 394 Association of Official Analytical Chemists (AOAC) 2005. Official methods of analysis, 18th  
395 edition. AOAC, Arlington, VA, USA.
- 396 Bontempo V, Jiang XR, Cheli F, Lo Verso L, Mantovani G, Vitari F, Domeneghini C and Agazzi  
397 A 2014. Administration of a novel plant extract product via drinking water to post-  
398 weaning piglets: effects on performance and gut health. *Animal* 8, 721–730.
- 399 Campbell JM, Crenshaw JD and Polo J 2013. The biological stress of early weaned piglets.  
400 *Journal of Animal Science and Biotechnology* 4, 19. doi: 10.1186/2049-1891-4-19.
- 401 Carillon J, Rouanet JM, Cristol JP and Brion R 2013a. Superoxide dismutase administration,  
402 a potential therapy against oxidative stress related diseases: several routes of  
403 supplementation and proposal of an original mechanism of action. *Pharmaceutical*  
404 *Research* 30, 2718-2728.
- 405 Carillon J, Fouret G, Feillet-Coudray C, Lacan D, Cristol JP and Rouanet JM 2013b. Short-  
406 term assessment of toxicological aspects, oxidative and inflammatory response to  
407 dietary melon superoxide dismutase in rats. *Food and Chemical Toxicology* 55, 323-  
408 328.
- 409 Carroll JA, Matteri RL, Dyer CJ, Beausang LA and Zannelli ME 2001. Impact of environmental  
410 temperature on response of neonatal pigs to an endotoxin challenge. *American Journal*  
411 *of Veterinary Research* 62, 561-566.
- 412 Carroll JA, Fangman TJ, Hambach AK and Wiedmeyer CE 2004. The acute phase response  
413 in pigs experimentally infected with *Escherichia coli* and treated with systemic  
414 bactericidal antibiotics. *Livestock Production Science* 85, 35-44.
- 415 Cray C, Zaias J and Altman NH 2009. Acute phase response in animals: a review. *Comparative*  
416 *Medicine* 59, 517-526.

417 Ceciliani F, Ceron JJ, Eckersall PD and Sauerwein H 2012. Acute phase proteins in ruminants.  
418 Journal of Proteomics 75, 4207–4231.

419 Cooke RF and Arthington JD 2013. Concentrations of haptoglobin in bovine plasma  
420 determined by ELISA or a colorimetric method based on peroxidase activity. Journal of  
421 Animal Physiology and Animal Nutrition 97, 531-536.

422 de Ridder K, Levesque CL, Htoo JK and de Lange CFM 2012. Immune system stimulation  
423 reduces the efficiency of tryptophan utilization for body protein deposition in growing  
424 pigs. Journal of Animal Science 90, 3485–3491.

425 Frank JW, Mellencamp MA, Carroll JA, Boyd RD and Allee GL 2005. Acute feed intake and  
426 acute-phase protein responses following a lipopolysaccharide challenge in pigs from  
427 two dam lines. Veterinary Immunology and Immunopathology 107, 179-187.

428 Gessner DK, Ringseis R and Eder K 2017. Potential of plant polyphenols to combat oxidative  
429 stress and inflammatory processes in farm animals. Journal of Animal Physiology and  
430 Animal Nutrition 101, 605-628.

431 Jiang XR, Awati A, Agazzi A, Vitari F, Ferrari A, Bento H, Crestani M, Domeneghini C and  
432 Bontempo V 2015a. Effects of a blend of essential oils and an enzyme combination on  
433 nutrient digestibility, ileum histology and expression of inflammatory mediators in  
434 weaned piglets. Animal 9, 417-426.

435 Jiang XR, Agazzi A, Awati A, Vitari F, Bento H, Ferrari A, Alborali GL, Crestani M, Domeneghini  
436 C and Bontempo V 2015b. Influence of a blend of essential oils and an enzyme  
437 combination on growth performance, microbial counts, ileum microscopic anatomy and  
438 the expression inflammatory mediators in weaned piglets following an Escherichia coli  
439 infection. Animal Feed Science and Technology 209, 219-229.

440 Kick AR, Tompkins MB, Flowers WL, Whisnant CS and Almond GW 2012. Effects of stress  
441 associated with weaning on the adaptive immune system in pigs. *Journal of Animal*  
442 *Science* 90, 649-656.

443 Lallès JP, Lacan D and David JC 2011. A melon pulp concentrate rich in superoxide dismutase  
444 reduces stress proteins along the gastrointestinal tract of pigs. *Nutrition* 27, 358-363.

445 Le Floc'h N and Seve B 2007. Biological roles of tryptophan and its metabolism: Potential  
446 implications for pig feeding. *Livestock Science* 112, 23–32.

447 Muchová J, Šustrová M, Garaiová I, Liptáková A., Blazíček P, Kvasnička P, Pueschel S and  
448 Duračková Z 2001. Influence of age on activities of antioxidant enzymes and lipid  
449 peroxidation products in erythrocytes and neutrophils of Down syndrome patients. *Free*  
450 *Radical Biology and Medicine* 31, 499–508.

451 Niekamp SR, Sutherland MA, Dahl GE, and Salak-Johnson JL 2007. Immune responses of  
452 piglets to weaning stress: impacts of photoperiod. *Journal of Animal Science* 85, 93–  
453 100.

454 Notin C, Vallon L, Desbordes F and Leleu C 2010. Oral supplementation with superoxide  
455 dismutase in Standardbred trotters in training: a double blind placebo controlled study.  
456 *Equine Veterinary Journal* 42 (suppl. 38), 375-381.

457 National Research Council (NRC) 2012. Nutrient requirements of swine, 11th revised edition.  
458 National Academic Press, Washington, DC, USA.

459 Pié S, Lallès JP, Blazy F, Laffitte J, Seve B and Oswald IP 2004. Weaning is associated with  
460 an upregulation of expression of inflammatory cytokines in the intestine of piglets. *The*  
461 *Journal of Nutrition* 134, 641–647.

462 Rakhshandeh A and de Lange CF 2012. Evaluation of chronic immune system stimulation  
463 models in growing pigs. *Animal* 6, 305-310.

464 Royer E, Barbé F, Guillou D, Rousseliere Y and Chevaux E 2016. Development of an oxidative  
465 stress model in weaned pigs highlighting plasma biomarkers' specificity to stress  
466 inducers. *Journal of Animal Science* 94, 48-53.

467 Rossi R, Pastorelli G and Corino C 2013. Application of KRL test to assess total antioxidant  
468 activity in pigs: Sensitivity to dietary antioxidants. *Research in Veterinary Science* 94,  
469 372-377.

470 Salamano G, Mellia E, Candiani D, Ingravalle F, Bruno R, Ru G and Doglione L 2008. Changes  
471 in haptoglobin, C-reactive protein and pig-major acute phase protein (porcine) during a  
472 housing period following long distance transport in swine. *The Veterinary Journal*  
473 177,110–115.

474 Sartorelli P, Paltrinieri S and Comazzi S 2000. Non-specific immunity and ketone bodies. II: *In*  
475 *vitro* studies on adherence and superoxide anion production in ovine neutrophils.  
476 *Journal of Veterinary Medicine. A, Physiology, Pathology, Clinical Medicine* 47, 1-8.

477 van de Crommenacker J, Horrocks NP, Versteegh MA, Komdeur J, Tieleman BI and Matson  
478 KD 2010. Effects of immune supplementation and immune challenge on oxidative  
479 status and physiology in a model bird: implications for ecologists. *The Journal of*  
480 *Experimental Biology* 213, 3527-3535.

481 Van Soest PJ, Robertson JB and Lewis BA 1991. Methods for dietary fiber, neutral detergent  
482 fiber, and nonstarch polysaccharides in relation to animal nutrition. *Journal of Dairy*  
483 *Science* 74, 3583–3597.

484 Vouldoukis I, Conti M, Krauss P, Kamate C, Blazquez S, Tefit M, Mazier D, Calenda A and  
485 Dugas B 2004a. Supplementation with gliadin-combined plant superoxide dismutase  
486 extract promotes antioxidant defences and protects against oxidative stress.  
487 *Phytotherapy Research* 18, 957-962.

- 488 Vouldoukis I, Lacan D, Kamate C, Coste P, Calenda A, Mazier D, Conti M and Dugas B 2004b.  
489 Antioxidant and anti-inflammatory properties of a *Cucumis melo* LC extract rich in  
490 superoxide dismutase activity. *Journal of Ethnopharmacology* 94, 67-75.
- 491 Zhang R, Chae S, Lee JH and Hyun JW. 2012. The cytoprotective effect of butin against  
492 oxidative stress is mediated by the up-regulation of manganese superoxide dismutase  
493 expression through a PI3K/Akt/Nrf2-dependent pathway. *Journal of Cellular*  
494 *Biochemistry* 113, 1987–1997.
- 495 Zhu LH, Zhao KL, Chen XL and Xu JX 2012. Impact of weaning and an antioxidant blend on  
496 intestinal barrier function and antioxidant status in pigs. *Journal of Animal Science* 90,  
497 2581–2589.

498 **Tables**

499 **Table 1** *Ingredients (% w/w) and chemical composition (g/100g dry matter) of the basal diet*

500 *fed to the experimental piglets*

501

Ingredients	Feeds
Wheat meal	29.47
Barley meal	23.12
Wheat flaked	14.00
Soybean meal, 48.0% CP	17.00
Sweet whey powder	6.00
Soybean oil	3.00
Corn gluten meal	2.50
Dextrose monohydrate	1.50
Dicalcium phosphate	1.30
L-lysine, 78% Lys	0.57
Calcium carbonate	0.50
Sodium chloride	0.30
Vitamin-mineral premix <sup>1</sup>	0.25
L-Thr	0.23
DL-Met	0.18
Flavour	0.05
Sweetener <sup>2</sup>	0.01
Zinc oxide	0.01
Cu sulphate	0.01
<b>Chemical composition<sup>3</sup> (g/100g DM)</b>	
DM	88.70
CP	18.01

Ether extract	4.52
Ash	5.08
NDF	14.59
Ca	0.75
Total P	0.57
Digestible energy (Mcal/kg DM)	3.44
Net energy, (Mcal/kg DM)	2.45
Lys	1.25
Met+Cyst	0.80
Thr	0.85
Trp	0.21

---

502 Lys = lysine; Thr = threonine; Met = methionine; Cyst = cysteine; Trp = tryptophan

503 <sup>1</sup>The vitamin-mineral premix provided the following quantities of vitamins and micro minerals per  
504 kilogram of complete diet: vitamin A, 10,500 IU; vitamin D3, 2500 IU; vitamin E, 15 mg; vitamin B1, 1.5  
505 mg; vitamin B2, 3.8 mg; vitamin B12, 0.025 mg; vitamin B6, 1.6 mg; calcium pantothenate, 12 mg; nicotinic  
506 acid, 15 mg; biotin, 0.15 mg; folic acid, 0.5 mg; vitamin K3, 3 mg; Fe, 100 mg; Cu, 6 mg; Co, 0.75 mg;  
507 Zn, 150 mg; Mn, 65 mg; I, 0.75 mg; Se, 0.3 mg; ethoxyquin, 150 mg.

508 <sup>2</sup>The sweetener (Optisweet®) was provided by Nutriad (Dendermonde, Belgium)

509 <sup>3</sup>Feeds were analysed for DM, CP, ether extract, ash, NDF, Ca and total P according to AOAC (2005).

510 All other values were calculated from NRC (2012).

511 **Table 2** Mean concentration of oxidative biomarkers<sup>1,2</sup> of post-weaning piglets supplemented with melon pulp concentrate in the diet and subjected  
 512 to a chronic LPS challenge<sup>3,4</sup>.

	LPS challenge				SEM <sup>5</sup>	P-values		
	-		+			Diet	Challenge	Diet X Challenge
	Dietary treatments <sup>2</sup>							
	CTR	MPC	CTR	MPC				
N. of piglets <sup>6</sup>	12	12	12	12				
SOD (IU/mL)	54.50	54.50	54.28	55.54	3.69	0.66	0.77	0.65
TAOC (mM Trolox equivalent)	4.97	7.94	4.11	6.49	1.71	<0.01	0.08	0.66
ROS (Abs)	0.16	0.15	0.15	0.15	0.03	0.73	0.62	0.74
8-oxo-dGuo (ng/mL)	0.89	0.84	0.87	0.91	0.06	0.99	0.40	0.26
HT <sub>50</sub> WB (min)	106.05	112.23	113.38	114.17	4.60	0.20	0.09	0.32
HT <sub>50</sub> RBC (min)	65.31	71.19	67.51	70.23	2.40	<0.01	0.66	0.26
HT <sub>50</sub> PC (min)	40.74	41.03	45.87	43.94	3.89	0.72	0.08	0.62

513 LPS = lipopolysaccharyde; SEM = standard error of the means; SOD = superoxide dismutase; TAOC = total antioxidant activity, ROS = reactive oxygen species;  
 514 8-oxo-dGuo = 8-oxo-7, 8-dihydro-2'-deoxyguanosine, HT<sub>50</sub>WB = half haemolysis time of whole blood, HT<sub>50</sub>RBC = half haemolysis time of red blood cells; HT<sub>50</sub>PC  
 515 = plasma contribution to half haemolysis.

516

517 <sup>1</sup>SOD and TAOC were performed on plasma samples; ROS and 8-oxo-dGuo were performed in serum; HT<sub>50</sub>WB and HT<sub>50</sub>RBC were performed on whole  
518 blood. HT<sub>50</sub>PC was obtained as the difference between HT<sub>50</sub>WB-HT<sub>50</sub>RBC.

519 <sup>2</sup>Total antioxidant activity was performed by KRL (Kit Radicaux Libres) biological test (Rossi *et al.*, 2013). Results are expressed as the time (min) required to  
520 reach 50% of maximal haemolysis (HT<sub>50</sub>), which refers to the whole blood, red blood cells and plasma resistance to free-radical attack.

521 <sup>3</sup>The challenge was performed from day 19 to 25 of the trial with increasing dosages of lipopolysaccharide (LPS from *E. coli* serotype 055:B5). Subsequent  
522 intramuscular injections of LPS were performed on days 19, 21, 23 and 25. Initial concentration of LPS was 60 µg/kg of BW and the dosage was increased by  
523 12% at each subsequent injection to reduce endotoxin tolerance and mimic a chronic inflammation in piglets. The applied concentrations of LPS from the second  
524 to the fourth injection day were 67.2, 75.26, and 84.30 µg/kg of BW. Individual body weight was determined prior each LPS injection to calculate individual total  
525 LPS amount to be injected.

526 <sup>4</sup>Dietary treatments: CTR- = piglets fed the basal diet and not subjected to the LPS challenge; CTR+ = piglets fed the basal diet and subjected to the LPS  
527 challenge; MPC- = piglets fed the basal diet added with 30g/ton of melon pulp concentrate (Melofeed, Lallemand SAS, Blagnac, France) and not subjected to  
528 LPS challenge; MPC+ = piglets fed the basal diet added with 30g/ton of melon pulp concentrate (Melofeed, Lallemand SAS, Blagnac, France) and subjected to  
529 LPS challenge.

530 <sup>5</sup>SEM = pooled SEM. Means are presented as least square means.

531 <sup>6</sup>Piglets were reared in individual pens (0.47 m<sup>2</sup>) with *ad libitum* access to feed and water.

532 **Table 3** Body weight (BW), average daily gain (ADG), feed intake (FI) and gain:feed (G:F) of post-weaning piglets supplemented with melon  
 533 pulp concentrate in the diet and subjected LPS challenge<sup>1</sup>

	LPS challenge				SEM <sup>4</sup>	P-values		
	-		+			Diet	Challenge	Diet X Challenge
	Dietary treatments <sup>2</sup>							
	CTR	MPC	CTR	MPC				
N. of piglets <sup>3</sup>	12	12	12	12				
BW (kg)								
Day 0	7.79	7.79	7.78	7.79	0.71	0.70	0.47	0.73
Day 29	17.94	18.48	16.98	17.37				
ADG (g/d)								
Days 0 to 19	298	297	343	328	51.98	0.79	0.13	0.78
Days 19 to 29	464	517	255	327	60.69	0.05	<0.01	0.75
Days 0 to 29	350	368	317	331	56.34	0.50	0.14	0.91
FI <sup>4</sup> (kg/period)								
Days 0 to 19	8.56	8.55	9.50	9.31	1.24	0.87	0.18	0.89
Days 19 to 29	7.31	8.22	4.99	6.15	0.67	<0.01	<0.01	0.73

Days 0 to 29	15.74	16.63	14.65	15.56	1.75	0.31	0.22	0.10
G:F								
Days 0 to 19	0.64	0.65	0.69	0.66	0.04	0.71	0.12	0.32
Days 19 to 29	0.62	0.63	0.51	0.53	0.05	0.64	<0.01	0.72
Days 0 to 29	0.64	0.64	0.63	0.61	0.03	0.69	0.18	0.41

---

534 LPS = lipopolysaccharyde; SEM = standard error of the means

535 <sup>1</sup>The challenge was performed from d 19 to 25 of the trial with increasing dosages of lipopolysaccharide (LPS from *E. coli* serotype 055:B5). Subsequent  
536 intramuscular injections of LPS were performed on days 19, 21, 23 and 25. Initial concentration of LPS was 60 µg/kg of BW and the dosage was increased by  
537 12% at each subsequent injection to reduce endotoxin tolerance and mimic a chronic inflammation in piglets. The applied concentrations of LPS from the second  
538 to the fourth injection day were 67.2, 75.26, and 84.30 µg/kg of BW. Individual body weight was determined prior each LPS injection to calculate individual total  
539 LPS amount to be injected.


540 <sup>2</sup>Dietary treatments: CTR-= piglets fed the basal diet and not subjected to the LPS challenge; CTR+= piglets fed the basal diet and subjected to the LPS  
541 challenge; MPC-=piglets fed the basal diet added with 30g/ton of melon pulp concentrate (Melofeed, Lallemand SAS, Blagnac, France) and not subjected to  
542 LPS challenge; MPC+= piglets fed the basal diet added with 30g/ton of melon pulp concentrate (Lallemand SAS, Blagnac, France) and subjected to LPS  
543 challenge.

544 <sup>3</sup>Piglets were reared in individual pens (0.47 m<sup>2</sup>) with *ad libitum* access to feed and water.

545 <sup>4</sup>SEM = pooled SEM. Means are presented as least square means.

546 **List of figure captions**

547 **Figure 1** Challenge effect on Interleukin 1 $\beta$  (IL-1 $\beta$ ), Interleukin 6 (IL-6), Tumour Necrosis Factor  $\alpha$  (TNF- $\alpha$ ) and haptoglobin serum  
548 concentration in post-weaning piglets supplemented with melon pulp concentrate<sup>1</sup> in the diet and subjected to chronic LPS challenge<sup>2,3</sup>



549

550 <sup>1</sup>Dietary treatments: CTR-= piglets fed the basal diet and not subjected to the LPS challenge; CTR+= piglets fed the basal diet and subjected to the LPS  
551 challenge; MPC-=piglets fed the basal diet added with 30g/ton of melon pulp concentrate (Melofeed, Lallemand SAS, Blagnac, France) and not subjected to  
552 LPS challenge; MPC+= piglets fed the basal diet added with 30g/ton of melon pulp concentrate (Melofeed, Lallemand SAS, Blagnac, France) and subjected to  
553 LPS challenge.

554 <sup>2</sup>The challenge was performed from day 19 to 25 of the trial with increasing dosages of lipopolysaccharide (LPS from *E. coli* serotype 055:B5). Subsequent  
555 intramuscular injections of LPS were performed on days 19, 21, 23 and 25. Initial concentration of LPS was 60 µg/kg of BW and the dosage was increased by  
556 12% at each subsequent injection to reduce endotoxin tolerance and mimic a chronic inflammation in piglets. The applied concentrations of LPS from the second  
557 to the fourth injection day were 67.2, 75.26, and 84.30 µg/kg of BW. Individual body weight was determined prior each LPS injection to calculate individual total  
558 LPS amount to be injected.

559 <sup>3A,B</sup>Different letters refer to significant differences between challenged and not challenge piglets for  $P<0.01$ .