

Epidemiology and Psychiatric Sciences

<http://journals.cambridge.org/EPS>


Additional services for *Epidemiology and Psychiatric Sciences*:

Email alerts: [Click here](#)

Subscriptions: [Click here](#)

Commercial reprints: [Click here](#)

Terms of use : [Click here](#)



The dysregulation profile in children and adolescents: a potential index for major psychopathology?

M. Bellani, G. A. L. Negri and P. Brambilla

Epidemiology and Psychiatric Sciences / Volume 21 / Issue 02 / June 2012, pp 155 - 159
DOI: 10.1017/S2045796011000849, Published online: 19 January 2012

Link to this article: http://journals.cambridge.org/abstract_S2045796011000849

How to cite this article:

M. Bellani, G. A. L. Negri and P. Brambilla (2012). The dysregulation profile in children and adolescents: a potential index for major psychopathology?. *Epidemiology and Psychiatric Sciences*, 21, pp 155-159 doi:10.1017/S2045796011000849

Request Permissions : [Click here](#)

This Section of *Epidemiology and Psychiatric Sciences* regularly appears in each issue of the Journal to describe relevant studies investigating the relationship between neurobiology and psychosocial psychiatry in major psychoses. The aim of these Editorials is to provide a better understanding of the neural basis of psychopathology and clinical features of these disorders, in order to raise new perspectives in every-day clinical practice.

Paolo Brambilla, *Section Editor* and Michele Tansella, *Editor EPS*

The dysregulation profile in children and adolescents: a potential index for major psychopathology?

M. Bellani^{1*}, G. A. L. Negri² and P. Brambilla^{2,3}

¹ Department of Public Health and Community Medicine, Section of Psychiatry and Clinical Psychology, Inter-University Center for Behavioural Neurosciences (ICBN), University of Verona, Verona, Italy

² Department of Experimental Clinical Medicine, Inter-University Center for Behavioural Neurosciences (ICBN), University of Udine, Udine, Italy

³ IRCCS 'E. Medea' Scientific Institute, Udine, Italy

We here review the literature on Child Behaviour Checklist-Dysregulation Profile (CBCL-DP) index, which potentially represents a developmental profile of major psychopathology in early adulthood. The understanding of the neural underpinnings of children and adolescents with altered regulation of affect and behaviour may ultimately help in planning strategies to prevent psychiatric syndromes during development.

Received 30 November 2011; Revised 12 December 2011; Accepted 13 December 2011

Key words: CBCL, emotional dysregulation, Juvenile Bipolar Disorder.

The Child Behaviour Checklist (CBCL) is a widely used questionnaire in research on developmental psychopathology (Achenbach & Rescorla, 2001). It is composed of 118 questions filled by the parents and provides T scores for three general domains (Total, Internalizing and Externalizing problems), eight Empirically Based Syndromes Scales (Anxious/Depressed, Withdrawn/Depressed, Somatic Complaints, Social Problems, Thought Problems, Attention Problems, Rule-Breaking Behaviour and Aggressive Behaviour) and six Diagnostic and Statistical Manual of Mental Disorders (DSM)-oriented scales (Affective Problems, Anxiety Problems, Somatic Problems, Attention Deficit/Hyperactivity Problems, Oppositional Defiant Problems and Conduct Problems). In the past few

years, it has been proposed that the sum of T scores in three Syndromes scales (i.e. Aggressive Behaviour, Anxious/Depressed and Attention Problems) could be used to identify CBCL-Juvenile Bipolar Disorder index (CBCL-JBD index; Biederman *et al.* 2009). However, the diagnostic value of this index in detecting JBD has been disconfirmed afterwards because of its low sensitivity and specificity (Volk & Todd, 2007). Recently, the CBCL-JBD index has been renamed as 'CBCL-Dysregulation Profile' (CBCL-DP) index, as it better detects children with 'a persisting deficit of self-regulation of affect and behaviour' (Holtmann *et al.* 2011). Several studies confirmed that CBCL-DP does not represent an early manifestation of JBD and it is not linked to a specific psychiatric diagnosis (Ayer *et al.* 2009), rather it is a developmental risk marker of poor overall functioning in adult life (Holtmann *et al.* 2011, Table 1). In large population-based samples (Ayer *et al.* 2009; Althoff *et al.* 2010, Table 1), this index usually indicates a more severe psychopathology, and studies on clinical populations also confirmed these

*Address for correspondence: Dr M. Bellani, Department of Public Health and Community Medicine, Section of Psychiatry and Clinical Psychology, University of Verona, Piazzale L.A. Scuro 10, 37134 Verona, Italy.

(Email: marcella.bellani@univr.it, paolo.brambilla@uniud.it)

Table 1. Summary of the studies described in this review

Study	Sample	Age range	Type of study	Assessment and diagnostic	Findings
Althoff <i>et al.</i> (2010)	1576 subjects population-based sample	3–17 at baseline	Longitudinal (14 years)	CIDI	CBCL-DP was associated with anxiety and disruptive behaviour disorders after controlling for co-occurring disorders in adulthood.
Ayer <i>et al.</i> (2009)	2029 subjects population-based sample	6–18	Psychometric	CBCL SES	High overlap between CBCL-DP and CBCL-PTSD scale.
Biederman <i>et al.</i> (2009)	204 ADHD	6–18 at baseline	Follow-up (7.4 years)	CBCL KSADS-E	CBCL-DP predicted subsequent diagnoses of BD, major depression and conduct disorder. Associated with impaired psychosocial functioning and psychiatric hospitalization.
Doyle <i>et al.</i> (2010)	765 individuals from 154 families enrolled in a linkage study of ADHD	11.1 mean age	Genetic	GAF SAICA SCID SES CBCL CBCL	Dysregulation profile is highly inheritable. Peaks on 1p21.1, 6p21.3 and 8q21.13 surpassed the threshold for suggestive linkage.
Halperin <i>et al.</i> (2011)	90 ADHD	7–11 at baseline	Follow-up	KSADS-E IOWA Conners	CBCL-DP did not significantly predict Axis I disorders, but it significantly predicted the presence of a cluster C personality disorder.
Holtmann <i>et al.</i> (2011)	325 young adults at risk for psychiatric illnesses	0.3 at baseline	Longitudinal (19 years)	DISC K-SADS-PL SCID-II CBCL YSR MEI	CBCL-DP scores in childhood are at increased risk for ADHD, mood and substance use disorders, suicidality and poorer overall functioning.
				SCID-I GAF SUQ FTND BDI	

Jucksch <i>et al.</i> (2011)	9024 clinical subjects	4–18	High DP <i>v.</i> other clinical samples	MEL CBCL GAPD Psychosocial axis of ICD-10 (consensus among clinicians) CBCL	CBCL-DP subjects were significantly more impaired and had more psychosocial adversities than other subgroups.
Meyer <i>et al.</i> (2008)	74 offspring of parents with mood disorders 27 controls	1.5–7	Longitudinal (23 years)	CAS DICA SCID-I IPDE GAF CBCL	CBCL-DP had significantly lower levels of social and occupational functioning. They were at risk for anxiety disorders, ADHD, cluster B personality disorders and BD, drug abuse and suicidal ideation.
Rich <i>et al.</i> (2011)	20 BD 20 severe mood dysregulation 20 controls	8–17	Neuroimaging (MEG) Affective Posner Task.	KSADS-PL CDRS YMRS CGAS	SMD youth showed greater arousal after negative feedback than both BD and controls, and they responded to negative feedback with significantly greater activation of the ACC and MFG than controls.
Spencer <i>et al.</i> (2011)	197 ADHD 224 controls	6–18	Group comparison	KSADS-E SAICA CBCL	Higher CBCL-DP in ADHD is associated with higher rates of psychiatric comorbidity and with significant impairment in social function.

ACC, anterior cingulate cortex; BDI, Beck Depression Inventory; CAS, Child Assessment Schedule; CDRS, Children’s Depression Rating Scale; CGAS, Children’s Global Assessment Scale; CIDI, Composite International Diagnostic Interview; DICA, Diagnostic Interview for Children and Adolescents; DISC, Diagnostic Interview Schedule for Children; FTND, Fagerström Test for Nicotine Dependence; GAF, Global Assessment of Functioning; GAPD, Global Assessment of Psychosocial Disability; IPDE, International Personality Disorder Examination; KSADS-E, Epidemiological version of Kiddie-Schedule for Affective Disorder and Schizophrenia for Children; MEG, Magneto-Encephalography; MEI, Mannheim Parent Interview; MEL, Munich Events List; SAICA, Social Adjustment Inventory for Children and Adolescents; SES, Socio-Economic Status; SCID, Structured Clinical Interview for DSM-IV; SMD, Severe Mood Dysregulation; SUQ, Substance Use Questionnaire; YMRS, Young Mania Rating Scale; YSR, Youth Self-Report.

results. Most of the studies on clinical subjects involved Attention Deficit Hyperactivity Disorder (ADHD). Spencer *et al.* (2011, Table 1) found that children with ADHD and CBCL-DP were more socially impaired and they had increased probability of comorbid disorders (such as anxiety and disruptive behaviour disorders) compared to non-CBCL-DP ADHD. Consistent findings were reported by follow-up studies on ADHD children (Biederman *et al.* 2009; Halperin *et al.* 2011, Table 1).

Studies on heterogeneous clinical populations found results in the same direction. Meyer *et al.* (2008) analysing data from a 23 years longitudinal study on at-risk youth for psychiatric disorders (i.e. offspring of parents with mood disorders), found that CBCL-DP during childhood is associated with later anxiety disorders, ADHD, cluster-B personality disorders, bipolar disorder (BD), drug abuse and suicidal ideation (Table 1). Similar results were found on 9024 clinical subjects with non-specified psychiatric pathologies (Jucksch *et al.* 2011) and on 325 children that presented risk factors associated with obstetric complications and psychosocial status (Holtmann *et al.* 2011, Table 1).

As to the neurobiological bases of this syndrome, so far few studies have investigated the biological and neural substrates of the CBCL-DP. It has been shown that this profile is highly inheritable (Doyle *et al.* 2010) and that it is associated with genetic regions found in adult BD, schizophrenia, autism and ADHD, like 1p21 (Table 1).

In parallel studies, Rich *et al.* (2011) using the magnetoencephalography (MEG) compared children with Severe Mood Dysregulation (SMD), characterized by abnormal mood, hyper-arousal and increased reactivity to negative emotional stimuli, being therefore very similar to the CBCL-DP (Leibenluft, 2011), to children with BD and controls during a Posner Affective Task. This task consists of a modified Posner Task, in which error feedbacks are modified to elicit frustration. The authors found that SMD youth showed greater arousal following negative feedback than both BD and controls, and they responded to negative feedback with significantly greater activation of the anterior cingulate cortex (ACC) and the medial frontal gyrus (MFG) than controls, while BD youth showed dysfunction in the superior frontal gyrus (SFG) and insula (Table 1). It seems therefore that specific cerebral substrates are involved in childhood SMD, thus supporting the hypothesis of a specific syndrome not related to JBD (Leibenluft, 2011).

In conclusion, there is evidence that the CBCL-DP represents a useful index for identifying children and adolescents at risk for psychiatric problems in early adulthood. Given the ease of administration of the CBCL and its widespread use, this could become a

powerful and low-cost tool to detect children who need early intervention for the prevention of mental illnesses, reducing the subsequent cost of treatment for psychiatric disorders (Kessler *et al.* 2009). However, few studies have been carried out so far to explore the potential neurobiological bases of patients with CBCL-DP. More neuroimaging investigations, coupled with neuropsychology and genetics, are therefore expected to further delineate the neural underpinnings of CBCL-DP, which may ultimately help in planning specific strategies to prevent major psychopathology during development.

References

- Achenbach TM, Rescorla LA (2001). *Manual for the ASEBA School-Age Forms and Profiles*. University of Vermont, Research Center for Children, Youth, and Families: Burlington, VT.
- Althoff RM, Verhulst FC, Rettew DC, Hudziak JJ, van der Ende J (2010). Adult outcomes of childhood dysregulation: a 14-year follow-up study. *Journal of the American Academy of Child and Adolescent Psychiatry* **49**, 1105–1116.
- Ayer L, Althoff R, Ivanova M, Rettew D, Waxler E, Sulman J, Hudziak J (2009). Child behavior checklist juvenile bipolar disorder (CBCL-JBD) and CBCL posttraumatic stress problems (CBCL-PTSP) scales are measures of a single dysregulatory syndrome. *Journal of Child Psychology and Psychiatry* **50**, 1291–1300.
- Biederman J, Petty C, Monuteaux MC, Evans M, Parcell T, Faraone SV, Wozniak J (2009). The CBCL-pediatric bipolar disorder profile predicts a subsequent diagnosis of bipolar disorder and associated impairments in ADHD youth growing up: a longitudinal analysis. *Journal of Clinical Psychiatry* **70**, 732–740.
- Doyle AE, Biederman J, Ferreira MAR, Wong P, Smoller JW, Faraone SV (2010). Suggestive linkage of the CBCL juvenile bipolar disorder phenotype to 1p21, 6p21 and 8q21. *Journal of the American Academy of Child and Adolescent Psychiatry* **49**, 378–387.
- Halperin JM, Rucklidge JJ, Powers RL, Miller CJ, Newcorn JH (2011). Childhood CBCL bipolar profile and adolescent/young adult personality disorders: a 9-year follow-up. *Journal of Affective Disorders* **130**, 155–161.
- Holtmann M, Buchmann AF, Esser G, Schmidt MH, Banaschewski T, Laucht M (2011). The child behaviour checklist-dysregulation profile predicts substance use, suicidality, and functional impairment: a longitudinal analysis. *Journal of Child Psychology and Psychiatry* **52**, 139–147.
- Jucksch V, Salbach-Andrae H, Lenz K, Goth K, Döpfner M, Poustka F, Freitag CM, Lehmkuhl G, Lehmkuhl U, Holtmann M (2011). Severe affective and behavioural dysregulation is associated with significant psychosocial adversity and impairment. *Journal of Child Psychology and Psychiatry* **52**, 686–695.

- Kessler RC, Aguilar-Gaxiola S, Alonso J, Chatterji S, Lee S, Ormel JT, Üstün TB, Wang PS** (2009). The global burden of mental disorders: an update from the WHO World Mental Health (WMH) Surveys. *Epidemiologia e Psichiatria Sociale* **18**, 23–33.
- Leibenluft MD** (2011). Severe mood dysregulation, irritability, and the diagnostic boundaries of bipolar disorder in youths. *American Journal of Psychiatry* **168**, 129–142.
- Meyer SE, Carlson GA, Youngstrom E, Ronsaville DS, Martinez PE, Gold PW, Hakak R, Radke-Yarrow M** (2008). Long-term outcomes of youth who manifested the CBCL-pediatric bipolar disorder phenotype during childhood and/or adolescence. *Journal of Affective Disorders* **113**, 227–235.
- Rich BA, Carver FW, Holroyd T, Rosen HR, Mendoza JK, Cornwell BR, Fox NA, Pine DS, Coppola R, Leibenluft E** (2011). Different neural pathways to negative affect in youth with pediatric bipolar disorder and severe mood dysregulation. *Journal of Psychiatric Research* **45**, 1283–1294.
- Spencer TJ, Faraone SV, Surman CB, Petty C, Clarke A, Batchelder H, Wozniak J, Biederman J** (2011). Toward defining deficient emotional self-regulation in children with attention-deficit/hyperactivity disorder using the child behaviour checklist: a controlled study. *Postgraduate Medicine* **123**, 50–59.
- Volk HE, Todd RD** (2007). Does the child behaviour checklist juvenile bipolar disorder phenotype identify bipolar disorder? *Biological Psychiatry* **62**, 115–120.