

Commun. Korean Math. Soc. **34** (2019), No. 2, pp. 633–635
<https://doi.org/10.4134/CKMS.c180138>
pISSN: 1225-1763 / eISSN: 2234-3024

A NOTE ON CONCIRCULAR STRUCTURE SPACE-TIMES

CARLO ALBERTO MANTICA AND LUCA GUIDO MOLINARI

Reprinted from the
Communications of the Korean Mathematical Society
Vol. 34, No. 2, April 2019

©2019 Korean Mathematical Society

A NOTE ON CONCIRCULAR STRUCTURE SPACE-TIMES

CARLO ALBERTO MANTICA AND LUCA GUIDO MOLINARI

ABSTRACT. In this note we show that Lorentzian Concircular Structure manifolds $(LCS)_n$ coincide with Generalized Robertson-Walker space-times.

Generalized Robertson-Walker (GRW) space-times were introduced in 1995 by Alías, Romero and Sánchez [1] as the warped product $-1 \times_{q^2} M^*$, where (M^*, g^*) is a Riemannian submanifold. In other terms, they are Lorentzian manifolds characterised by a metric

$$(1) \quad g_{ij} dx^i dx^j = -(dt)^2 + q(t)^2 g_{\mu\nu}^*(x^1, \dots, x^{n-1}) dx^\mu dx^\nu.$$

They are interesting not only for geometry [2, 5, 9–11], but also for physics: they include relevant space-times such as Robertson-Walker, Einstein-de Sitter, static Einstein, de-Sitter, the Friedmann cosmological models. They are a wide generalization of space-times for cosmological models.

In 2003 A. A. Shaikh [12] introduced the notion of *Lorentzian Concircular Structure* $(LCS)_n$. It is a Lorentzian manifold endowed with a unit time-like concircular vector field, i.e., $u^i u_i = -1$ and

$$(2) \quad \nabla_k u_j = \varphi(u_k u_j + g_{kj}),$$

where $\varphi \neq 0$ is a scalar function obeying

$$(3) \quad \nabla_j \varphi = \mu u_j$$

being μ a scalar function. Various authors studied the properties of $(LCS)_n$ manifolds [6, 13–15].

We show that GRW and $(LCS)_n$ are the same space-times.

We recall few definitions that will be used in this note. The first ones are the definitions of “torse-forming” and “concircular” vector fields, by Yano:

Definition 1 (Yano, [16, 17]). A vector field X_j is named torse-forming if $\nabla_k X_j = \omega_k X_j + \varphi g_{kj}$, being φ a scalar function and ω_k a non vanishing one-form. It is named concircular if ω_k is a gradient or locally a gradient of a scalar function.

Received April 5, 2018; Accepted May 16, 2018.

2010 *Mathematics Subject Classification*. Primary 53B30; Secondary 53C25.

Key words and phrases. Generalized Robertson-Walker space-time, Lorentzian concircular structure, torse-forming vector, concircular vector.

Fialkow gave a definition different from Yano's:

Definition 2 (Fialkow [4]). A vector field X_j is named concircular if it satisfies $\nabla_k X_j = \rho g_{kj}$, being ρ a scalar function.

The following simple but deep result was recently proven:

Theorem 3 (Bang-Yen Chen, [3]). *A $n > 3$ dimensional Lorentzian manifold is a GRW space-time if and only if it admits a time-like concircular vector field (in the sense of Fialkow).*

It is worth noticing that for a unit time-like vector field, the torse-forming property by Yano becomes precisely Eq. (2), with generic scalar field φ . Based on Chen's theorem we proved:

Proposition 4 (Mantica and Molinari, [7,8]). *A $n > 3$ dimensional Lorentzian manifold is a GRW space-time if and only if it admits a unit time-like torse-forming vector, (2), that is also an eigenvector of the Ricci tensor.*

Now comes the equivalence: from (2) (holding either for GRW and $(LCS)_n$ space-times) we evaluate

$$R_{jkl}{}^m u_m = [\nabla_j, \nabla_k]u_l = (h_{kl}\nabla_j - h_{jl}\nabla_k)\varphi - \varphi^2(u_j g_{kl} - u_k g_{jl}),$$

where $h_{kl} = u_k u_l + g_{kl}$. Contraction with g^{jl} gives

$$(4) \quad R_k{}^m u_m = u_k [u^m \nabla_m \varphi + (n-1)\varphi^2] - (n-2)\nabla_k \varphi.$$

If (3) holds, then u_k is an eigenvector of the Ricci tensor, and we conclude that a $(LCS)_n$ manifold is a GRW space-time.

If $R_{km}u^m = \xi u_k$ it is $(n-2)\nabla_k \varphi = \alpha u_k$ for some scalar field α , i.e., (3) holds. Then we conclude that a GRW space-time is a $(LCS)_n$ manifold.

References

- [1] L. Alías, A. Romero, and M. Sánchez, *Uniqueness of complete spacelike hypersurfaces of constant mean curvature in generalized Robertson-Walker spacetimes*, Gen. Relativity Gravitation **27** (1995), no. 1, 71–84.
- [2] K. Arslan, R. Deszcz, R. Ezentas, M. Hotłoś, and C. Murathan, *On generalized Robertson-Walker spacetimes satisfying some curvature condition*, Turk. J. Math. **38** (2014) 353–373.
- [3] B.-Y. Chen, *A simple characterization of generalized Robertson-Walker spacetimes*, Gen. Relativity Gravitation **46** (2014), no. 12, Art. 1833, 5 pp.
- [4] A. Fialkow, *Conformal geodesics*, Trans. Amer. Math. Soc. **45** (1939), no. 3, 443–473.
- [5] M. Gutiérrez and B. Olea, *Global decomposition of a Lorentzian manifold as a generalized Robertson-Walker space*, Differential Geom. Appl. **27** (2009), no. 1, 146–156.
- [6] S. K. Hui, *On ϕ -pseudo symmetries of $(LCS)_n$ -manifolds*, Kyungpook Math. J. **53** (2013), no. 2, 285–294.
- [7] C. A. Mantica and L. G. Molinari, *On the Weyl and Ricci tensors of generalized Robertson-Walker space-times*, J. Math. Phys. **57** (2016), no. 10, 102502, 6 pp.
- [8] ———, *Generalized Robertson-Walker spacetimes—a survey*, Int. J. Geom. Methods Mod. Phys. **14** (2017), no. 3, 1730001, 27 pp.

- [9] A. Romero, R. M. Rubio, and J. J. Salamanca, *Uniqueness of complete maximal hypersurfaces in spatially parabolic generalized Robertson-Walker spacetimes*, Classical Quantum Gravity **30** (2013), no. 11, 115007, 13 pp.
- [10] M. Sánchez, *On the geometry of generalized Robertson-Walker spacetimes: geodesics*, Gen. Relativity Gravitation **30** (1998), no. 6, 915–932.
- [11] ———, *On the geometry of generalized Robertson-Walker spacetimes: curvature and Killing fields*, J. Geom. Phys. **31** (1999), no. 1, 1–15.
- [12] A. A. Shaikh, *On Lorentzian almost paracontact manifolds with a structure of the concircular type*, Kyungpook Math. J. **43** (2003), no. 2, 305–314.
- [13] ———, *Some results on $(LCS)_n$ -manifolds*, J. Korean Math. Soc. **46** (2009), no. 3, 449–461.
- [14] A. A. Shaikh and K. K. Baishya, *On concircular structure spacetimes*, J. Math. Stat. **1** (2005), no. 2, 129–132.
- [15] ———, *On concircular structure spacetimes II*, American J. Appl. Sci. **3** (2006), no. 4, 1790–1794.
- [16] K. Yano, *Concircular geometry I-IV*, Proc. Imp. Acad. Tokyo **16** (1940) 195–200, 354–360, 442–448, 505–511.
- [17] ———, *On the torse-forming directions in Riemannian spaces*, Proc. Imp. Acad. Tokyo **20** (1944), 340–345.

CARLO ALBERTO MANTICA
I.I.S. LAGRANGE
VIA L. MODIGNANI 65, 20161, MILANO, ITALY
Email address: `carloalberto.mantica@libero.it`

LUCA GUIDO MOLINARI
PHYSICS DEPARTMENT
UNIVERSITÀ DEGLI STUDI DI MILANO AND I.N.F.N. SEZ. MILANO
VIA CELORIA 16, 20133 MILANO, ITALY
Email address: `luca.molinari@unimi.it`