

RESEARCH ARTICLE

Running Author: CHIARELLI PERDOMO et al.

Continuous preparation of flavour-active acetate esters by direct biocatalytic esterification

Igor Chiarelli Perdomo¹

Martina Letizia Contente^{1,2}

Andrea Pinto¹

Diego Romano¹

Pedro Fernandes^{3,4}

Francesco Molinari^{1,✉}

Email francesco.molinari@unimi.it

AQ1

AQ2

¹ Department of Food, Environmental and Nutritional Sciences (DeFENS), University of Milan, Milano, Italy

² School of Chemistry, University of Nottingham, University Park, Nottingham, UK

³ Department of Bioengineering and IBB Institute for Bioengineering and Biosciences, Instituto Superior Técnico, Universidade de Lisboa, Lisboa, Portugal

⁴ Faculty of Engineering and DREAMS, Universidade Lusófona de Humanidades e Tecnologias, Lisboa, Portugal

Correspondence

Francesco Molinari, Department of Food, Environmental and Nutritional Sciences (DeFENS), University of Milan, via Mangiagalli 25, 20133 Milano, Italy.


Email: francesco.molinari@unimi.it

Abstract

Mycelium-bound lipase(s) from *Aspergillus oryzae* catalysed direct esterification of isoamyl alcohol and cinnamyl alcohol with acetic acid in organic solvent, showing high stability towards substrates and products. Water produced during the esterification did not significantly affect the equilibrium of the reaction, allowing for high conversions. These features were exploited for preparing flavour-active acetate esters (isoamyl and cinnamyl acetate) in batch and continuous systems. A continuous stirred tank membrane reactor (CST-MR) was developed securing good reactor productivity and high biocatalyst stability.

Graphical Abstract

xxxxxxxxxxxx.



AQ3

Received: 13 September 2019 | Revised: 12 October 2019 | Accepted: 19 October 2019

Keywords

Aspergillus oryzae; biocatalysis; continuous reactor; enzymatic acetylation; flavour esters

1. INTRODUCTION

Many acetate esters of primary alcohols are components of natural flavours and are used as flavour and fragrance ingredients.[1] They can be obtained by extraction from natural sources or by enzymatic preparation starting from naturally occurring substrates (eg alcohols and carboxylic acids), thus ensuring the definition of "natural". Lipase-catalysed transesterification (or interesterification) in organic solvents or in solvent-free environments is one of the most common enzymatic methods for the obtainment of acetate esters.[2, 3, 4] Alternatively, acylation of alcohol and amines in water can be efficiently catalysed by specific acyltransferase (eg the one from *Mycobacterium smegmatis*, MsAct), furnishing excellent yields of

different flavour esters.[5, 6, 7, 8, 9, 10]

Direct esterification with free carboxylic acids is a preferred option since many acids are available as natural molecules, but the biotransformation is often hampered by formation of water, which may increase water activity of the medium thus negatively affecting the equilibrium of the reaction.[11, 12, 13, 14] Different strategies have been developed for controlling water activity of the medium and improving the final conversion in the desired esters. A large-scale enzymatic production of low molecular weight flavour esters in organic solvent was developed, where solutions for the elimination of substrate and product inhibitions are presented; water produced during the process was continuously removed by azeotropic distillation or by using molecular sieves, hence enabling high yields of the produced esters.[15] A major drawback of direct acetylation with acetic acid is the deactivation of lipases in the presence of the free acid.[16] Lipase stabilisation in the presence of acetic acid has often been achieved through immobilisation.[17, 18, 19, 20, 21, 22, 23, 24] Novozym 435 (the commercial preparation of lipase B from *Candida antarctica*, CALB, immobilised on a macroporous acrylic resin) has been often used as stable and efficient biocatalyst for acetate esters preparation.[19, 22]

Towards the development of a continuous methodology, different type of reactors can be set up for the production of esters by direct esterification catalysed by lipases in organic solvents.[25, 26] The final process productivity depends on the activity and stability of the biocatalyst, whereas molar conversion can be modulated by applying the appropriate residence time in the bioreactor. Continuous stirred tank membrane reactor (CST-MR) has been scarcely used for the continuous enzymatic synthesis of esters, albeit they may offer several advantages (enhanced mass transfer, improved liquid mixing and reduced clogging) over packed bed reactors. [27, 28, 29, 30] Moreover, the use of a suited membrane allows for continuous removal of the organic liquid phase containing the product, while retaining the biocatalyst inside the reactor.[31]

Mycelium-bound fungal carboxylesterases have been often used for flavour ester production by direct esterification.[28] Dry mycelium of moulds (eg *Aspergillus oryzae*) can be effectively used for different biotransformations,[33, 34] including direct esterification of different alcohols with remarkable advantages, such as easy preparation of the biocatalyst, high stability in organic solvents and high resistance to the inactivation caused by carboxylic acids (including acetic acid); moreover, this biocatalyst allows direct esterification with high molar conversions, enabled by favourable water partition.[35, 36, 37]

In this work, we have investigated the continuous preparation of flavour-active acetate esters (isoamyl and cinnamyl) using mycelium-bound lipase(s) from *Aspergillus oryzae* in a continuous stirred tank membrane reactor (CST-MR).

2. MATERIALS AND METHODS

2.1. Chemicals and biocatalyst

All chemicals were purchased from Sigma-Aldrich and were used without any further purification. *Aspergillus oryzae* MIM[36] (Microbiologia Industriale Milano) was cultured in 500-mL Erlenmeyer flasks containing 125 mL of liquid medium (Tween 80 0.5%v/v, Difco™ yeast extract 1 g/L, (NH₄)₂SO₄ 5 g/L, K₂HPO₄ 1 g/L, MgSO₄·7H₂O

0.2 g/L, pH 5.8) for 48 h at 28°C on a reciprocal shaker (100 rpm). The mycelium suspension was recovered by vacuum filtration using a Buchner funnel and paper filter and washed with distilled water, and lyophilised.

2.2. Batch biotransformations

Different amounts of dry (lyophilised) mycelium of *A. oryzae* were suspended in 2.5 mL of *n*-heptane for 30 minutes and the reactions started by adding different concentrations of isoamyl alcohol and acetic acid; the reaction mixture was magnetically stirred at the desired temperature. Samples (0.25 mL) were taken at different intervals, added to an equal volume of an internal standard solution (*n*-octanol) in *n*-heptane and analysed by GC. Recovery and purification were carried out as previously reported.[7]

2.3. Continuous reaction procedure

The vessel used in this study was a glass-made membrane reactor constructed by the glass workshop of University of Milan. The working volume of the reactor was 200 mL (a picture and a cross section of the membrane reactor are given in Figure 1). The membrane bioreactor was composed with a thermostated water-jacketed glass vessel and an ultrafiltration module (0.1 µm) mounted on the bottle cap. The final suspension was maintained under agitation using a magnetic stirrer.




Fig. 1 Schematic representation and picture of the membrane reactor

The temperature in the reactor was kept constant by circulating water in the jacket. A Gilson Miniplus 2 peristaltic pump controlled incoming liquid flow.

The molar balance for a continuous stirred tank reactor (CSTR) has the following expression, where F_A stands for the molar flow rate of substrate:

$$V = \frac{F_{A,in} - F_{A,out}}{(-r_a)} \quad (1)$$

One can rewrite as:

$$\tau = \frac{C_{A0} - C_A}{(-r_A)} \quad \text{or} \quad \tau = \frac{C_{A0} X_A}{(-r_A)} \quad (2)$$

Finally, rate of the CSTR was calculated using the algebraic Equation 3:

$$r = \frac{C_{A0} X_A}{\tau} \quad (3)$$

The reactor was run for 24-30 hours depending on the residence time of each operation to reach a steady state before collecting samples for analysis.

2.4. Analysis

Isoamyl alcohol and isoamyl acetate concentrations were determined by gas-chromatographic (GC) analysis on a Carlo Erba Fractovap GC equipped with a fused-silica capillary column MEGA-DEX DMP-Beta (dimethyl pentyl- β -cyclodextrin, 25 m \times 0.25 mm i.d.), with the injector temperature at 200°C. Oven temperature was a 5°C/min gradient from 40 to 180°C. The retention times were isoamyl alcohol, 6.7 minutes and isoamyl acetate, 7.1 minutes. Cinnamyl alcohol and cinnamyl acetate concentrations were determined with a fused-silica capillary column MEGA-SE30 (100% methyl polysiloxane; 25 m \times 0.25 mm i.d.), with the injector temperature at 200°C. Oven temperature was 80°C (2 minutes) and then from 80 to 180°C with a 5°C/min gradient. The retention times were cinnamyl alcohol, 9.9 min and cinnamyl acetate, 13.2 min.

3. RESULTS AND DISCUSSION

3.1. Isoamyl acetate preparation: optimisation of batch reaction

Direct esterification of isoamyl alcohol with acetic acid in *n*-heptane for the preparation of isoamyl acetate was firstly carried out in batch mode using dry mycelium of *Aspergillus oryzae*. Isoamyl acetate is used as a flavouring compound, having a characteristic banana flavour property.[1] Optimisation of the biotransformation was accomplished by sequential experimental trials using a

Multisimplex[®] 2.0 software, previously employed for the optimisation of biotransformations.[37] Response variables were the productivity of the biotransformation (defined as amount of product per amount of biocatalyst per unit of time) and the molar conversion of the alcohol, both determined after 24 hours; the control variables were alcohol concentration, biocatalyst concentration, molar ratio (acid/alcohol) and temperature. The initial levels considered for the optimisation are listed in Table 1.

Table 1 Control variables and initial levels considered for the optimisation of the esterification between isoamyl alcohol and acetic acid catalysed by mycelium-bound lipase(s) of *Aspergillus oryzae*

	Control variables			
	Alcohol (mmol/L)	Dry mycelium (mg/mL)	Molar ratio (acid/alcohol)	T (°C)
Reference value	25	20	1.5	40
Step size	5	5	0.2	5
Minimum	10	5	1.0	20
Maximum	200	40	3.0	70

The results of the sequential experiments aimed at the optimisation of the bioconversion are shown in Figure 2. Each trial was performed in triplicate, and the mean value was introduced as response for further optimisation.




Fig. 2 Sequential optimisation of the esterification between isoamyl alcohol and acetic acid catalysed by mycelium-bound lipase(s) of *Aspergillus oryzae*. Experimental responses (Δ productivity; \bullet molar conversion) are reported as function of sequential trials. Values represent the mean of separate assays in triplicate. Molar ratio (acid/alcohol) was always kept >1 for maintaining an excess of acetic acid; therefore, molar conversions are referred to the amount of alcohol (limiting reagent) converted

The highest molar conversion (98%) and overall productivity ($88.2 \mu\text{mol g}^{-1} \text{h}^{-1}$) were simultaneously met in trials 20, 23 and 25; conditions corresponding to the trial 25 (isoamyl alcohol = 54 mmol/L , dry mycelium = 25 mg mL^{-1} , molar ratio acid/alcohol 1.2/1 at 50°C) were chosen for further experiments, since the lowest amount of biocatalyst still gave the highest amount of isoamyl acetate. Critical parameters were acetic acid concentration and temperature, since significant decreases of the productivity were observed in any trial performed at concentrations above 70 mmol/L and at temperature above 60°C . Figure 3 reports the time-course of the direct esterification of isoamyl alcohol with acetic acid under the optimised conditions.




Fig. 3 Esterification between isoamyl alcohol and acetic acid catalysed by mycelium-bound lipase(s) from *A oryzae*. Reaction conditions: initial alcohol concentration 54 mmol/L; initial acetic acid concentration 65 mmol/L in *n*-heptane containing 25 g L⁻¹ of biocatalyst at 50°C

The reaction occurred with high yield and completion of the reaction in 24 hours with a reaction rate quite constant in the first 3 hours (0.089 μmol/mL min).

3.2. Isoamyl acetate preparation: optimisation of continuous reactor

The encouraging results obtained in batch reactions led us to explore the possibility to carry out continuous biotransformations in a continuous stirred tank membrane reactor (CST-MR), where mixing of the heterogeneous system should be optimal. The continuous stirred tank membrane reactor (CST-MR) described in the Material section (50 mL working volume) was employed for continuous operations; the biocatalyst was retained in the reactor by an ultrafiltration membrane, which allowed for permeation of the outflow solution.

The CST-MR was packed with a suspension of dry mycelium (1.25 g, corresponding to 25 g L⁻¹) in *n*-heptane and the solvent was flown in a recirculation mode for 72 hours at a flow rate (Q) of 0.2 mL min⁻¹, corresponding to a hydraulic residence time of 250 min (residence time: $\tau = V/Q$, where V is the reactor volume and Q is the flow rate). No filter plugging or release of enzymatic activity was observed during the time of the operation.

Direct esterification of isoamyl alcohol with acetic acid in CST-MR was then studied; substrates were continuously added in a *n*-heptane solution and the composition of the outflow solution monitored by GC. Experiments were initially focused to reach the best compromise between reaction rate and degree of conversion (Table 2).

Table 2 Optimisation of the continuous esterification of isoamyl alcohol with acetic acid catalysed by mycelium-bound lipase(s) from *A oryzae* carried out in CST-MR. Initial stoichiometric ratio between acid/alcohol was always 1.2

Q (mL/min)	τ (min)	C_{alcohol} (mmol/L) ^a	F_{alcohol} ($\mu\text{mol}/\text{min}$)	Conversion (%)	Rate ($\mu\text{mol}/\text{min}$)
0.10	500	50	5.0	83	0.083
0.10	500	55	5.5	82	0.090
0.10	500	60	6.0	78	0.094
0.14	350	50	7.0	56	0.090

^aInitial alcohol concentration.

Experiments carried out with a

τ

= 500 min allowed for 78%-83% molar conversion, depending on the F_{alcohol} in the inlet flow; the best compromise between conversion and rate was found with a F_{alcohol} of 5.5 $\mu\text{mol}/\text{min}$ (corresponding to a C_{alcohol} = 55 mmol/L and $C_{\text{acetic acid}}$ = 66 mmol/L) corresponding to a rate of 0.092 $\mu\text{mol}/\text{min}$. The rate was slightly better than the one observed in batch reactor, showing the good performance in terms of mixing of the CST-MR. Experiments were also carried out at lower residence time ($t = 350$ min), leading to a significant decrease in the conversion.

3.3.

Isoamyl acetate preparation: stability of the biocatalyst in the continuous reactor

The continuous operation aimed at preparing isoamyl acetate was carried out for 10 days under optimised conditions (inflow solution containing 55 mmol/L isoamyl alcohol and 65 mmol/L acetic acid,

τ

= 500 min at 50°C); no decrease of the molar conversion was observed at the end of the process, thus indicating a notable operational stability of the catalyst. Table 3 summarises the results of the continuous biotransformation, which allowed for a remarkable overall reactor productivity (defined here as the amount of isoamyl acetate produced for volume of reactor) of 169 mg mL⁻¹.

Table 3 Continuous esterification of isoamyl alcohol with acetic acid catalysed by mycelium-bound lipase(s) from *A oryzae* carried out in CST-MR for 10 days

Q	τ	Conv. ^a	Space time	Catalyst	Reactor
-----	--------	--------------------	------------	----------	---------

(mL min ⁻¹)	(min)	(%)	yield (mmol d ⁻¹)	productivity ^b (mmol g ⁻¹)	productivity ^b (mg mL ⁻¹)
0.10	500	82	6.49	51.95	169

^aConversion detected in the exiting flow after 10 days of operation.

^bProductivity is referred to the amount of ester formed after an operation time of 10 days.

A closed-loop reactor was also set up by recirculating the outflowing solution; after 16 hours of operation, the mycelium-bound lipase(s) was able to catalyse a conversion of 98% of isoamyl alcohol into the corresponding acetate ester.

3.4. Cinnamyl acetate preparation: batch and continuous reactor

Biocatalytic preparation of cinnamyl acetate was also studied; cinnamyl acetate is characterised by a typical pineapple flavour.[1] Direct acetylation of cinnamyl alcohol was accomplished applying the optimised conditions employed for the direct esterification of isoamyl alcohol. Figure 4 shows the time-course of the batch biotransformation.




Fig. 4 Esterification between cinnamyl alcohol and acetic acid catalysed by mycelium-bound lipase(s) from *A. oryzae*. Reaction conditions: initial alcohol concentration 54 mmol/L; initial acetic acid concentration 65 mmol/L in *n*-heptane containing 25 g L⁻¹ of biocatalyst at 50°C

The acetylation rate of cinnamyl alcohol was slightly higher (steady reaction rate in the first 3 hours of 0.095 $\mu\text{mol}/\text{mL min}$) than what observed with isoamyl alcohol, whereas final conversion after 24 h was 95%-96%. Since acetylation of cinnamyl alcohol occurred with a time-course similar to what observed for isoamyl alcohol, continuous operation was carried out under the same conditions ($C_{\text{alcohol}} = 55 \text{ mmol}/\text{L}$, $F_{\text{alcohol}} = 5.5 \mu\text{mol}/\text{min}$, $C_{\text{acid}} = 66 \text{ mmol}/\text{L}$, $Q = 0.1 \text{ mL min}^{-1}$,

$\tau =$

500 min at 50°C) and in the same CST-MR. Also, in this case, the biocatalyst was stable over 10 days of reaction; Table 4 summarises the data of the continuous bioprocess.

Table 4 Continuous esterification of cinnamyl alcohol with acetic acid catalysed by mycelium-bound lipase(s) from *A. oryzae* carried out in CST-MR for 10 days

Q (mL min^{-1})	τ (min)	Conv. ^a (%)	Space time yield (mmol d^{-1})	Catalyst productivity ^b (mmol g^{-1})	Reactor productivity ^b (mg mL^{-1})
0.10	500	80	6.34	50.69	223

^aConversion detected in the exiting flow after 10 days of operation.

^bProductivity is referred to the amount of ester formed after an operation time of 10 days.

3.5. Conclusion

The mycelium-bound lipase(s) activity of *Aspergillus oryzae* was exploited for the direct esterification of isoamyl and cinnamyl alcohol with acetic acid for the preparation of the corresponding flavour-active esters. Remarkably, similar acetylation rates and molar conversions were observed, although two alcohols with quite different structure were employed. It is worthy to underline that the biotransformation occurred in *n*-heptane without any particular system for the removal of the water produced; it has previously suggested that the mycelia supply a hydrophobic micro-environment, thus disfavouring water access to the enzymes which catalyses the reaction.[38] A stirred tank membrane reactor (CST-MR) was set up for carrying out continuous biotransformations; reusability of the biocatalyst reduces the cost of fermentation, necessary for its production. Indeed, the CST-MR guaranteed 10 days of operation without any significant loss of the biocatalyst activity; reactor productivity (defined as the amount of ester formed per volume of reactor) was 169 mg/mL for isoamyl acetate and 223 mg/mL for cinnamyl acetate. In conclusion, a highly effective continuous production of isoamyl and cinnamyl acetate was obtained with dry mycelium, without any necessity of costly or laborious enzymatic purification, and without the need of water sorption during the prolonged bioprocess.

CONFLICT OF INTEREST

The authors have no conflicts of interest to declare.

REFERENCES

- 1 Burdock GA. *Fenaroli's handbook of flavor ingredients*, 6th edn. Boca Raton: CRC Press; 2010.
- 2 Dhake KP, Thakare DD, Bhanage BM. Lipase: a potential biocatalyst for the synthesis of valuable flavour and fragrance ester compounds. *Flavour Fragr. J.* 2013;28:71-83.
- 3 Ferreira-Dias S, Sandoval G, Plou F, Valero F. The potential use of lipases in the production of fatty acid derivatives for the food and nutraceutical **AQ5** industries. *Electron J Biotechnol.* 2013;16(3).
- 4 Sá AGA, de Meneses AC, de Araújo PHH, Oliveira D. A review on enzymatic synthesis of aromatic esters used as flavor ingredients for food, cosmetics and pharmaceuticals industries. *Trends Food Sci Technol.* 2017;69:95-105.
- 5 Mathews I, Soltis M, Saldajeno M, et al. Structure of a novel enzyme that catalyzes acyl transfer to alcohols in aqueous conditions. *Biochemistry.* 2007;46:8969-8979.
- 6 Wiermans L, Hofzumahaus S, Schotten C, et al. Transesterifications and peracid-assisted oxidations in aqueous media catalyzed by *Mycobacterium smegmatis* acyl transferase. *Chem Cat Chem.* 2013;5:3719-3724.
- 7 Chiarelli Perdomo I, Gianolio S, Pinto A, et al. Efficient enzymatic preparation of flavor esters in water. *J Agric Food Chem.* 2019;67:6517-6522.
- 8 Kazemi M, Sheng X, Kroutil W, Himo F. Computational study of *Mycobacterium smegmatis* acyl transferase reaction mechanism and specificity. *ACS Catal.* 2018;8:10698-10706.
- 9 Contente ML, Pinto A, Molinari F, Paradisi F. Biocatalytic N-acylation of amines in water using an acyltransferase from *Mycobacterium smegmatis*. *Adv Synth Catal.* 2018;360:4814-4819.
- 10 Mestrom L, Claessen JGR, Hanefeld U. Enzyme-catalyzed synthesis of esters in water. *ChemCatChem.* 2019;11:2004-2010.
- 11 Halling PJ. Solvent selection for biocatalysis in mainly organic systems: Predictions of effects on equilibrium position. *Biotechnol Bioeng.* 1990;35:691-701.
- 12 Klibanov AM. Improving enzymes by using them in organic solvents. *Nature.* 2001;409:241-246.
- 13 Svensson I, Wehtje E, Adlercreutz P, Mattiasson B. Effects of water activity on reaction rates and equilibrium positions in enzymatic esterifications. *Biotechnol Bioeng.* 1994;44:549-556.
- 14 Spizzo P, Basso A, Ebert C, et al. Resolution of (R, S)-flurbiprofen catalysed by dry mycelia in organic solvent. *Tetrahedron.* 2007;63:11005-11010.

- 15 Gubicza L, Kabiri-Badr A, Keoves E, Belafi-Bako K. Large-scale enzymatic production of natural flavour esters in organic solvent with continuous water removal. *J Biotechnol.* 2000;84:193-196.
- 16 De Castro HF, De Oliveira PC, Pereira EB. Evaluation of different approaches for lipase catalysed synthesis of citronellyl acetate. *Biotechnol Lett.* 1997;19:229-232.
- 17 Güvenç A, Kapucu N, Mehmetoglu U. The production of isoamyl acetate using immobilized lipases in a solvent-free system. *Proc Biochem.* 2002;38:379-386.
- 18 Melo LLMM, Pastore GM, Macedo GA. Optimized synthesis of citronellyl flavour esters using free and immobilized lipase from *Rhizopus* sp. *Proc Biochem.* 2005;40:3181-3185.
- 19 Romero MD, Calvo L, Alba C, Daneshfar A, Ghaziaskar HS. Enzymatic synthesis of isoamyl acetate with immobilized *Candida antarctica* lipase in *n*-hexane. *Enzyme Microb Technol.* 2005;37:42-48.
- 20 Karra-Châabouni M, Ghamgui H, Bezzine S, Rekik A, Gargouri Y. Production of flavour esters by immobilized *Staphylococcus simulans* lipase in a solvent-free system. *Proc Biochem.* 2006;41:1692-1698.
- 21 Ozyilmaz G, Gezer E. Production of aroma esters by immobilized *Candida rugosa* and porcine pancreatic lipase into calcium alginate gel. *J Mol Catal B: Enzymatic.* 2010;64:140-145.
- 22 Martins AB, Graebin NG, Lorenzoni ASG, Fernandez-Lafuente R, Ayub MAZ, Rodrigues RC. Rapid and high yields of synthesis of butyl acetate catalyzed by Novozym 435: Reaction optimization by response surface methodology. *Proc Biochem.* 2011;46:2311-2316.
- 23 Matte CR, Bordinhão C, Poppe JK, Rodrigues RC, Hertz PF, Ayub MAZ. Synthesis of butyl butyrate in batch and continuous enzymatic reactors using *Thermomyces lanuginosus* lipase immobilized in Immobead 150. *J Mol Catal B: Enzymatic.* 2016;127:67-75.
- 24 Almeida Gomes Sá A, de Meneses AC, Lerin LA, de Araújo PHH, Sayer C, de Oliveira D. Biocatalysis of aromatic benzyl-propionate ester by different immobilized lipases. *Bioproc Biosyst Eng.* 2018;41:585-591..
- 25 Tamborini L, Romano D, Pinto A, Bertolani A, Molinari F, Conti P. An efficient method for the lipase-catalysed resolution and in-line purification of racemic flurbiprofen in a continuous-flow reactor. *J Mol Catal B: Enzym.* 2012;84:78-82.
- 26 Itabaiana I, de MarizeMiranda LS de Souza ROMA. Towards a continuous flow environment for lipase-catalyzed reactions. *J Mol Catal B: Enzym.* 2013;86:1-9.

- 27 Carvalho CML, Aires-Barros MR, Cabral JMS. A continuous membrane bioreactor for ester synthesis in organic media: II. modeling of MBR continuous operation. *Biotechnol Bioeng.* 2001;72:136-143.
- 28 Trusek-Holownia A, Noworyta A. An integrated process: ester synthesis in an enzymatic membrane reactor and water sorption. *J Biotechnol.* 2007;130:47-56.
- 29 Damjanovic JJ, Zuza MG, Savanovic JK, et al. Covalently immobilized lipase catalyzing high-yielding optimized geranyl butyrate synthesis in a batch and fluidized bed reactor. *J Mol Catal B: Enzymatic.* 2012;75:50-59.
- 30 Giovannini PP, Catani M, Massi A, et al. Continuous production of eugenol esters using enzymatic packed-bed microreactors and an evaluation of the products as antifungal agents. *Flavour Fragr J.* 2019;34:201-210.
- 31 Novak U, Lavric D, Žnidaršič-Plaz P. Continuous Lipase B-catalyzed isoamyl acetate synthesis in a two-liquid phase system using Corning® AFRTM module coupled with a membrane separator enabling biocatalyst recycle. *J Flow Chem.* 2016;6:33-38.
- 32 Molinari F, Marianelli G, Aragozzini F. Production of flavour esters by *Rhizopus oryzae*. *Appl. Microbiol. Biotechnol.* 1995;43:967-973. *Proc Biochem.* 2005;40:3181-3185.
- 33 Zambelli P, Fernandez-Arrojo L, Romano D, et al. Production of fructooligosaccharides by mycelium-bound transfructosylation activity present in *Cladosporium cladosporioides* and *Penicillium sizovae*. *Proc Biochem.* 2014;49:2174-2180.
- 34 Zambelli P, Serra I, Fernandez-Arrojo L, et al. Sweet-and-salty biocatalysis: Fructooligosaccharides production using *Cladosporium cladosporioides* in seawater. *Proc Biochem.* 2015;50:1086-1090.
- 35 Gandolfi R, Converti A, Pirozzi D, Molinari F. Efficient and selective microbial esterification with dry mycelium of *Rhizopus oryzae*. *J Biotechnol.* 2001;92:21-26.
- 36 Tamborini L, Romano D, Pinto A, et al. Biotransformation with whole microbial systems in a continuous flow reactor: resolution of (RS)-flurbiprofen using *Aspergillus oryzae* by direct esterification with ethanol in organic solvent. *Tetrahedron Lett.* 2013;54:6090-6093.
- 37 Contente ML, Molinari F, Zambelli P, et al. Biotransformation of aromatic ketones and ketoesters with the non-conventional yeast *Pichia glucozyma*. *Tetrahedron Lett.* 2014;55:7051-7053.
- 38 Converti A, Gandolfi R, Zilli M, et al. Synthesis of ethyl phenylacetate by lyophilized mycelium of *Aspergillus oryzae*. *Appl Microbiol Biotechnol.* 2005;67:637-640.

AQ4