

# SERRIN'S TYPE OVERDETERMINED PROBLEMS IN CONVEX CONES

GIULIO CIRAOLO AND ALBERTO RONCORONI

ABSTRACT. We consider overdetermined problems of Serrin's type in convex cones for (possibly) degenerate operators in the Euclidean space as well as for a suitable generalization to space forms. We prove rigidity results by showing that the existence of a solution implies that the domain is a spherical sector.

## 1. INTRODUCTION

Given a bounded domain  $E \subset \mathbb{R}^N$ ,  $N \geq 2$ , the classical Serrin's overdetermined problem [40] asserts that there exists a solution to

$$\begin{cases} \Delta u = -1 & \text{in } E, \\ u = 0 & \text{on } \partial E, \\ \partial_\nu u = -c & \text{on } \partial E, \end{cases} \quad (1)$$

for some constant  $c > 0$ , if and only if  $E = B_R(x_0)$  is a ball of radius  $R$  centered at some point  $x_0$ . Moreover, the solution  $u$  is radial and it is given by

$$u(x) = \frac{R^2 - |x - x_0|^2}{2N}, \quad (2)$$

with  $R = Nc$ . Here,  $\nu$  denotes the outward normal to  $\partial\Omega$ .

The starting observation of this manuscript is the following. Let  $\Sigma$  be an open cone in  $\mathbb{R}^N$  with vertex at the origin  $O$ , i.e.

$$\Sigma = \{tx : x \in \omega, t \in (0, +\infty)\}$$

for some open domain  $\omega \subset \mathbb{S}^{N-1}$ . We notice that if  $x_0$  is chosen appropriately then  $u$  given by (2) is still the solution to

$$\begin{cases} \Delta u = -1 & \text{in } B_R(x_0) \cap \Sigma, \\ u = 0 \text{ and } \partial_\nu u = -c & \text{on } \partial B_R(x_0) \setminus \bar{\Sigma}, \\ \partial_\nu u = 0 & \text{on } B_R(x_0) \cap \partial\Sigma. \end{cases}$$

More precisely,  $x_0$  may coincide with  $O$  or it may be just a point of  $\partial\Sigma \setminus \{O\}$  and, in this case,  $B_R(x_0) \cap \Sigma$  is half a sphere lying over a flat portion of  $\partial\Sigma$ . Hence, it is natural to look for a characterization of symmetry in this direction, as done in [35] (see below for a more detailed description).

In order to properly describe the results, we introduce some notation. Given an open cone  $\Sigma$  such that  $\partial\Sigma \setminus \{O\}$  is smooth, we consider a bounded domain  $\Omega \subset \Sigma$  and denote by  $\Gamma_0$  its relative boundary, i.e.

$$\Gamma_0 = \partial\Omega \cap \Sigma,$$

and we set

$$\Gamma_1 = \partial\Omega \setminus \bar{\Gamma}_0.$$

We assume that  $\mathcal{H}_{N-1}(\Gamma_1) > 0$ ,  $\mathcal{H}_{N-1}(\Gamma_0) > 0$  and that  $\Gamma_0$  is a smooth  $(N-1)$ -dimensional manifold, while  $\partial\Gamma_0 = \partial\Gamma_1 \subset \partial\Omega \setminus \{O\}$  is a smooth  $(N-2)$ -dimensional manifold. Following

---

*Date:* December 5, 2019.

*2010 Mathematics Subject Classification.* 35N25, 35B06, 53C24, 35R01.

*Key words and phrases.* Overdetermined problems, rigidity, torsion problem, convex cones, mixed boundary conditions.

[35], such a domain  $\Omega$  is called a *sector-like domain*. In the following, we shall write  $\nu = \nu_x$  to denote the exterior unit normal to  $\partial\Omega$  wherever is defined (that is for  $x \in \Gamma_0 \cup \Gamma_1 \setminus \{O\}$ ).

Under the assumption that  $\Sigma$  is a convex cone, in [35] it is proved that if  $\Omega$  is a sector-like domain and there exists a classical solution  $u \in C^2(\Omega) \cap C^1(\Omega \cup \Gamma_0 \cup \Gamma_1 \setminus \{O\})$  to

$$\begin{cases} \Delta u = -1 & \text{in } \Omega, \\ u = 0 \text{ and } \partial_\nu u = -c & \text{on } \Gamma_0, \\ \partial_\nu u = 0 & \text{on } \Gamma_1 \setminus \{O\}, \end{cases} \quad (3)$$

and such that

$$u \in W^{1,\infty}(\Omega) \cap W^{2,2}(\Omega),$$

then

$$\Omega = B_R(x_0) \cap \Sigma$$

for some  $x_0 \in \mathbb{R}^N$  and  $u$  is given by (2). Differently from the original paper of Serrin [40], the method of moving planes is not helpful (at least when applied in a standard way) and the rigidity result in [35] is proved by using two alternative approaches. One is based on integral identities and it is inspired from [5], the other one uses a  $P$ -function approach as in [42].

In this paper, we generalize the rigidity result for Serrin's problem in [35] in two directions. The former is by considering more general operators than the Laplacian in the Euclidean space, where the operators may be of degenerate type. Here, the generalization is not trivial due to the lack of regularity of the solution (the operator may be degenerate) as well as to other technical details which are not present in the linear case.

The latter is by considering an analogous problem in space forms, i.e. the hyperbolic space and the (hemi)sphere. The operator that we consider is linear and it is interesting since it has been shown that it is a helpful generalization of the torsion problem to space forms ([15], [36], [37]).

Overdetermined problems for quasilinear and possible degenerate operators have attracted a lot of interest in the last decades, see for instance [26, 25, 19, 27, 38]. As Fosdick and Serrin noticed in [40] and [24], Serrin's overdetermined problem for quasilinear elliptic operators is also interesting for possible applications to the study of steady rectilinear motion of viscous incompressible fluids and incompressible non-Newtonian fluids (see also [26]), and in the theory of torsion of a solid straight bar. Roughly speaking, a rigidity result as the one given by Serrin proves that *the tangential stress on the pipe wall is the same at all points of the wall if and only if the pipe has a circular cross section* or that *when a solid straight bar is subject to torsion, the magnitude of the resulting traction which occurs at the surface of the bar is independent of position if and only if the bar has a circular cross section*. There are other possible applications for Serrin's type rigidity results, and we refer to [25, Introduction] for connections to capillarity theory, torsional creep, Born-Infeld theory and other applications to quantum-physics.

As explained in [35], the study of Serrin's overdetermined problem in convex cones is related to relative isoperimetric inequality and Alexandrov soap bubble theorem. In this manuscript we extend this study to non-Euclidean manifolds, in particular to space forms. The study of isoperimetric inequality and Alexandrov theorem in non-Euclidean manifolds has recently attracted a lot of interest in the geometric analysis community (see [31, 36, 7] and references therein). We believe that, by taking inspiration from our results and the ones in [31, 36], one can study Alexandrov theorem and relative isoperimetric inequalities for sector-like domains in more general Riemannian settings.

The study of rigidity problems in convex cones appears also in the context of critical points for Sobolev inequality (which in turns can be related to Yamabe problem), see [12, 32]. Indeed, the study started in this manuscript served as inspiration for [12], where we characterized, together with A. Figalli, the solutions of critical anisotropic  $p$ -Laplace type equations in convex cones.

We also mention that the approach used in this paper originated from [5], which in turns has been later used for proving quantitative estimates for Serrin's overdetermined problem in [6]. As for the symmetry result, this approach is also useful when considering quantitative versions of Alexandrov soap bubble theorem, in particular to describe the appearance of bubbling [13].

**More general operators in the Euclidean space.** Let  $\Omega$  be a sector like domain in  $\mathbb{R}^N$  and let  $f : [0, +\infty) \rightarrow [0, +\infty)$  be such that

$$f \in C^1([0, \infty)) \cap C^3((0, \infty)) \text{ with } f(0) = f'(0) = 0, \quad f''(s) > 0 \text{ for } s > 0$$

$$\text{and } \lim_{s \rightarrow +\infty} \frac{f(s)}{s} = +\infty. \quad (4)$$

We consider the following mixed boundary value problem

$$\begin{cases} L_f u = -1 & \text{in } \Omega, \\ u = 0 & \text{on } \Gamma_0 \\ \partial_\nu u = 0 & \text{on } \Gamma_1 \setminus \{O\}, \end{cases} \quad (5)$$

where the operator  $L_f$  is given by

$$L_f u = \operatorname{div} \left( f'(|\nabla u|) \frac{\nabla u}{|\nabla u|} \right), \quad (6)$$

and the equation  $L_f u = -1$  is understood in the sense of distributions

$$\int_{\Omega} \frac{f'(|\nabla u|)}{|\nabla u|} \nabla u \cdot \nabla \varphi \, dx = \int_{\Omega} \varphi \, dx$$

for any

$$\varphi \in T(\Omega) := \{\varphi \in C^1(\Omega) : \varphi \equiv 0 \text{ on } \Gamma_0\}.$$

Notice that the operator  $L_f$  may be of degenerate type.

We notice that the solution to  $L_f u = -1$  in  $B_R(x_0)$  (a ball of radius  $R$  centered at  $x_0$ ) such that  $u = 0$  on  $\partial B_R(x_0)$  is radial and it is given by

$$u(x) = \int_{|x-x_0|}^R g' \left( \frac{s}{N} \right) ds, \quad (7)$$

where  $g$  denotes the Fenchel conjugate of  $f$  (see for instance [17] or [25]), i.e.

$$g = \sup\{st - f(s) : s \geq 0\}$$

(hence for us  $g'$  is the inverse function of  $f'$ ). Our first main result is the following.

**Theorem 1.** *Let  $f$  satisfy (4). Let  $\Sigma$  be a convex cone such that  $\Sigma \setminus \{O\}$  is smooth and let  $\Omega$  be a sector-like domain in  $\Sigma$ . If there exists a solution  $u \in C^1(\Omega \cup \Gamma_0 \cup \Gamma_1 \setminus \{O\}) \cap W^{1,\infty}(\Omega)$  to (5) such that*

$$\partial_\nu u = -c \text{ on } \Gamma_0 \quad (8)$$

for some constant  $c$ , and satisfying

$$\frac{f'(|\nabla u|)}{|\nabla u|} \nabla u \in W^{1,2}(\Omega, \mathbb{R}^N), \quad (9)$$

then there exists  $x_0 \in \mathbb{R}^N$  such that  $\Omega = \Sigma \cap B_R(x_0)$  with  $c = g'(|\Omega|/|\Gamma_0|)$ ,  $R = N|\Omega|/|\Gamma_0|$ . Moreover  $u$  is given by (7), where  $x_0$  is the origin or, if  $\partial\Sigma$  contains flat regions, it is a point on  $\partial\Sigma$ .

When  $L_f = \Delta$  (i.e.  $f(t) = t^2/2$ ), Theorem 1 is essentially Theorem 1.1 in [35]. Condition (9) holds locally in  $\Omega$  for a large class of elliptic operators, such as the mean curvature operator ( $f(t) = \sqrt{1+t^2}$ ), and for the  $p$ -Laplace operator ( $f(t) = t^p/p$ ,  $p > 1$ ), see [2, Theorem ] and [10, Theorem 2.1]. We stress that the validity of (9) up to the boundary is more subtle, since it depends strongly on how  $\Gamma_0$  and  $\Gamma_1$  intersect. Some global results may be obtained by following the approach in [10], where (9) is proved for Dirichlet or Neumann boundary value problems of  $p$ -Laplace type in domains which are convex or satisfying minimal regularity assumptions on the boundary.

We observe that the overdetermined problem (5) with the condition (8) can be seen as a partially overdetermined problem (see for instance [20] and [21]), since we impose both Dirichlet and Neumann conditions only on a part of the boundary, namely  $\Gamma_0$ , while a sole homogeneous

Neumann boundary condition is assigned on  $\Gamma_1$  (where, however, there is the strong assumption that it is contained in the boundary of a cone).

We notice that the proof of Theorem 1 still works when  $\Gamma_1 = \emptyset$  (hence  $\partial\Omega = \Gamma_0$ ). In this case we obtain the celebrated result of Serrin [40] for the operator  $L_f$  (see also [4], [5], [14], [25], [19], [26], [39], [42]). Moreover, the proof is also suitable to be adapted to the anisotropic counterpart of the overdetermined problem (5) and (8) by following the approach used in this manuscript and in [4] (see also [11] and [41]). We also mention that rigidity theorems in cones are related to the study of relative isoperimetric and Sobolev inequalities in cones, and we refer to [35] for a more detailed discussion (see also [3, 9, 23, 29, 32, 33]).

**Serrin's problem in cones in space forms.** A space form is a complete simply-connected Riemannian manifold  $(M, g)$  with constant sectional curvature  $K$ . Up to homotheties we may assume  $K = 0, 1, -1$ : the case  $K = 0$  corresponds to the Euclidean space  $\mathbb{R}^N$ ,  $K = -1$  is the hyperbolic space  $\mathbb{H}^N$  and  $K = 1$  is the unitary sphere with the round metric  $\mathbb{S}^N$ . More precisely, in the case  $K = 1$  we consider the hemisphere  $\mathbb{S}_+^N$ . These three models can be described as warped product spaces  $M = I \times \mathbb{S}^{N-1}$  equipped with the rotationally symmetric metric

$$g = dr^2 + h(r)^2 g_{\mathbb{S}^{N-1}},$$

where  $g_{\mathbb{S}^{N-1}}$  is the round metric on the  $(N-1)$ -dimensional sphere  $\mathbb{S}^{N-1}$  and

- $h(r) = r$  in the Euclidean case ( $K = 0$ ), with  $I = [0, \infty)$ ;
- $h(r) = \sinh(r)$  in the hyperbolic case ( $K = -1$ ), with  $I = [0, \infty)$ ;
- $h(r) = \sin(r)$  in the spherical case ( $K = 1$ ), with  $I = [0, \pi/2)$  for  $\mathbb{S}_+^N$ .

By using the warping structure of the manifold, we denote by  $O$  the pole of the model and it is natural to define a *cone*  $\Sigma$  with vertex at  $\{O\}$  as the set

$$\Sigma = \{tx : x \in \omega, t \in I\}$$

for some open domain  $\omega \subset \mathbb{S}^{N-1}$ . Moreover, we say that  $\Sigma$  is a *convex cone* if the second fundamental form II is nonnegative defined at every  $p \in \partial\Sigma$ .

Serrin's overdetermined problem for semilinear equations  $\Delta u + f(u) = 0$  in space forms has been studied in [30] and [34] by using the method of moving planes. If one considers the corresponding problem for sector-like domains in space forms, the method of moving planes can not be used and one has to look for alternative approaches. As already mentioned, in the Euclidean space these approaches typically use integral identities and  $P$ -functions (see [5, 42]) and have the common feature that at a crucial step of the proof they use the fact that the radial solution attains the equality sign in a Cauchy-Schwartz inequality, which implies that the Hessian matrix  $\nabla^2 u$  is proportional to the identity. Since the equivalent crucial step in space forms is to prove that the Hessian matrix of the solution is proportional to the metric, then the equation  $\Delta u = -1$  is no more suitable (one can easily verify that in the radial case the Hessian matrix of the solution is not proportional to the metric) and a suitable equation to be considered is

$$\Delta u + NKu = -1 \tag{10}$$

as done in [15] and [36], [37]. It is clear that for  $K = 0$ , i.e. in the Euclidean case, the equation reduces to  $\Delta u = -1$ . For this reason, we believe that, in this setting, (10) is the natural generalization of the Euclidean  $\Delta u = -1$  to space forms.

A Serrin's type rigidity result for (10) can be proved following Weinberger's approach by using a suitable  $P$ -function associated to (10) (see [15] and [37]). This approach is helpful for proving the following Serrin's type rigidity result for convex cones in space forms, which is the second main result of this paper.

**Theorem 2.** *Let  $(M, g)$  be the Euclidean space, hyperbolic space or the hemisphere. Let  $\Sigma \subset M$  be a convex cone such that  $\Sigma \setminus \{O\}$  is smooth and let  $\Omega$  be a sector-like domain in  $\Sigma$ . Assume*

that there exists a solution  $u \in C^1(\Omega \cup \Gamma_0 \cup \Gamma_1 \setminus \{O\}) \cap W^{1,\infty}(\Omega) \cap W^{2,2}(\Omega)$  to

$$\begin{cases} \Delta u + NKu = -1 & \text{in } \Omega, \\ u = 0 & \text{on } \Gamma_0 \\ \partial_\nu u = 0 & \text{on } \Gamma_1 \setminus \{O\}, \end{cases} \quad (11)$$

such that

$$\partial_\nu u = -c \quad \text{on } \Gamma_0 \quad (12)$$

for some constant  $c$ . Then  $\Omega = \Sigma \cap B_R(x_0)$  where  $B_R(x_0)$  is a geodesic ball of radius  $R$  centered at  $x_0$  and  $u$  is given by

$$u(x) = \frac{H(R) - H(d(x, x_0))}{nh(R)},$$

with

$$H(r) = \int_0^r h(s) ds$$

and where  $d(x, x_0)$  denotes the distance between  $x$  and  $x_0$ .

**Organization of the paper.** The paper is organized as follows: in Section 2 we introduce some notation, we recall some basic facts about elementary symmetric function of a matrix and prove some preliminary result needed to prove Theorem 1. Theorems 1 and 2 are proved in Sections 3 and 4, respectively.

## 2. PRELIMINARY RESULTS FOR THEOREM 1

In this section we collect some preliminary results which are needed in the proof of Theorem 1. Let  $f$  satisfy (4) and consider problem (5)

$$\begin{cases} L_f u = -1 & \text{in } \Omega, \\ u = 0 & \text{on } \Gamma_0 \\ \partial_\nu u = 0 & \text{on } \Gamma_1 \setminus \{O\}, \end{cases}$$

where the operator  $L_f$  is given by

$$L_f u = \operatorname{div} \left( f'(|\nabla u|) \frac{\nabla u}{|\nabla u|} \right).$$

**Definition 3.**  $u \in C^1(\Omega \cup \Gamma_0 \cup \Gamma_1 \setminus \{O\})$  is a solution to Problem (5) if

$$\int_\Omega \frac{f'(|\nabla u|)}{|\nabla u|} \nabla u \cdot \nabla \varphi \, dx = \int_\Omega \varphi \, dx \quad (13)$$

for any

$$\varphi \in T(\Omega) := \{\varphi \in C^1(\Omega) : \varphi \equiv 0 \text{ on } \Gamma_0\}. \quad (14)$$

We observe some facts that will be useful in the following. Since the outward normal  $\nu$  to  $\Gamma_0$  is given by

$$\nu = -\frac{\nabla u}{|\nabla u|} \Big|_{\Gamma_0}, \quad (15)$$

then (8) implies that

$$|\nabla u| = c \quad \text{on } \Gamma_0. \quad (16)$$

Moreover we observe that the constant  $c$  in the statement is given by

$$c = g' \left( \frac{|\Omega|}{|\Gamma_0|} \right), \quad (17)$$

as it follows by integrating the equation  $L_f u = -1$ , by using the divergence theorem, formula (16) and the fact that  $\partial_\nu u = 0$  on  $\Gamma_1 \setminus \{O\}$ . We also notice that

$$x \cdot \nu = 0 \quad \text{on } \Gamma_1. \quad (18)$$

It will be useful to write the operator  $L_f$  as the trace of a matrix. Let  $V : \mathbb{R}^N \rightarrow \mathbb{R}$  be given by

$$V(\xi) = f(|\xi|) \quad \text{for } \xi \in \mathbb{R}^N, \quad (19)$$

and notice that

$$\begin{aligned} V_{\xi_i}(\xi) &:= \partial_{\xi_i} V(\xi) = f'(|\xi|) \frac{\xi_i}{|\xi|}, \\ V_{\xi_i \xi_j}(\xi) &:= \partial_{\xi_i \xi_j} V(\xi) = f''(|\xi|) \frac{\xi_i \xi_j}{|\xi|^2} - f'(|\xi|) \frac{\xi_i \xi_j}{|\xi|^3} + f'(|\xi|) \frac{\delta_{ij}}{|\xi|}. \end{aligned} \quad (20)$$

Hence, by setting

$$W = (w_{ij})_{i,j=1,\dots,N}$$

where

$$w_{ij}(x) = \partial_j V_{\xi_i}(\nabla u(x)), \quad (21)$$

we have

$$L_f(u) = \text{Tr}(W). \quad (22)$$

Notice that at regular points, where  $\nabla u \neq 0$ , it holds that

$$W = \nabla_{\xi}^2 V(\nabla u) \nabla^2 u. \quad (23)$$

Our approach to prove Theorem 1 is to write several integral identities and just one pointwise inequality, involving the matrix  $W$ . Writing the operator  $L_f$  as trace of  $W$  has the advantage that we can use the generalization of the so-called Newton's inequalities, as explained in the following subsection.

We mention that, unless otherwise specified, we adopt the Einstein convention of summation over repeated indices.

**2.1. Elementary symmetric functions of a matrix.** Given a matrix  $A = (a_{ij}) \in \mathbb{R}^{N \times N}$ , for any  $k = 1, \dots, N$  we denote by  $S_k(A)$  the sum of all the principal minors of  $A$  of order  $k$ . In particular,  $S_1(A) = \text{Tr}(A)$  is the trace of  $A$ , and  $S_n(A) = \det(A)$  is the determinant of  $A$ . We consider the case  $k = 2$ . By setting

$$S_{ij}^2(A) = -a_{ji} + \delta_{ij} \text{Tr}(A), \quad (24)$$

we can write

$$S_2(A) = \frac{1}{2} \sum_{i,j} S_{ij}^2(A) a_{ij} = \frac{1}{2} ((\text{Tr}(A))^2 - \text{Tr}(A^2)). \quad (25)$$

The elementary symmetric functions of a symmetric matrix  $A$  satisfy the so called Newton's inequalities. In particular,  $S_1$  and  $S_2$  are related by

$$S_2(A) \leq \frac{N-1}{2N} (S_1(A))^2. \quad (26)$$

When the matrix  $A = W$ , with  $W$  given by (23), we have

$$S_{ij}^2(W) = -V_{\xi_j \xi_k}(\nabla u) u_{ki} + \delta_{ij} L_f u, \quad (27)$$

and  $S_{ij}^2(W)$  is divergence free in the following (weak) sense

$$\frac{\partial}{\partial x_j} S_{ij}^2(W) = 0. \quad (28)$$

If  $V$  and  $u$  are sufficiently smooth, (28) was proved in [11, Equation (4.14)]. In Lemma 8 below we will prove (28) under weaker regularity assumptions on  $V$  and  $u$  by approximation (notice that (28) is implicitly written in (43), as follows from (25)).

We will need a generalization of (26) to not necessarily symmetric matrices, which is given by the following lemma.

**Lemma 4** ([11], Lemma 3.2). *Let  $B$  and  $C$  be symmetric matrices in  $\mathbb{R}^{N \times N}$ , and let  $B$  be positive semidefinite. Set  $A = BC$ . Then the following inequality holds:*

$$S_2(A) \leq \frac{N-1}{2N} \text{Tr}(A)^2. \quad (29)$$

Moreover, if  $\text{Tr}(A) \neq 0$  and equality holds in (29), then

$$A = \frac{\text{Tr}(A)}{N} Id,$$

and  $B$  is, in fact, positive definite.

**2.2. Some properties of solutions to (5).** In this subsection we collect some properties of the solutions to (5). We assume that the solution is of class  $C^1(\Omega) \cap W^{1,\infty}(\Omega)$ . From standard regularity elliptic estimates one has that  $u$  is of class  $C^{1,\alpha}(\Omega)$  and  $C^{2,\alpha}$  where  $\nabla u \neq 0$ .

In the following two lemmas we show that  $u > 0$  in  $\Omega \cup \Gamma_1 \setminus \{O\}$  and we prove a Pohozaev-type identity.

**Lemma 5.** *Let  $f$  satisfy (4) and let  $u$  be a solution of (5). Then*

$$u > 0 \quad \text{in} \quad \Omega \cup \Gamma_1 \setminus \{O\}. \quad (30)$$

*Proof.* We write  $u = u^+ - u^-$  and use  $\varphi = u^-$  as test function in (13):

$$0 \geq - \int_{\Omega \cap \{u < 0\}} \frac{f'(|\nabla u|)}{|\nabla u|} |\nabla u^-|^2 dx = \int_{\Omega \cap \{u < 0\}} u^- dx \geq 0,$$

which implies that  $u \geq 0$  in  $\Omega$ . Moreover, if one assumes that  $u(x_0) = 0$  at some point  $x_0 \in \Omega \cup \Gamma_1 \setminus \{O\}$ , then  $\nabla u(x_0) = 0$ . Since  $x_0 \in \Omega \cup \Gamma_1 \setminus \{O\}$  and  $\Gamma_1 \setminus \{O\}$  is smooth, there exists a ball  $B_r \subset \Omega$  such that  $x_0 \in \partial B_r$ . Let  $v$  be the solution of

$$\begin{cases} L_f v = -1 & \text{in } B_r, \\ v = 0 & \text{on } \partial B_r. \end{cases}$$

By comparison principle we have that  $v \leq u$  in  $\overline{B_r}$ ; from  $\nabla u(x_0) = 0$  and since  $\nabla v(x_0) \neq 0$  we get a contradiction.  $\square$

The following Pohozaev-type identity is a typical tool to prove symmetry results. In a similar setting as the one in this paper, a Pohozaev identity was proved in [26].

**Lemma 6** (Pohozaev-type identity). *Let  $\Omega$  be a sector-like domain and assume that  $f$  satisfies (4). Let  $u \in C^1(\Omega \cup \Gamma_0 \cup \Gamma_1 \setminus \{O\}) \cap W^{1,\infty}(\Omega)$  be a solution to (5). Then the following integral identity*

$$\int_{\Omega} [(N+1)u - Nf(|\nabla u|)] dx = \int_{\Gamma_0} [f'(|\nabla u|)|\nabla u| - f(|\nabla u|)] x \cdot \nu d\sigma \quad (31)$$

holds.

*Proof.* We argue by approximation. We first approximate  $f$  with functions  $f_\varepsilon$  such that

$$f_\varepsilon \in C^\infty([0, \infty)) \text{ with } f_\varepsilon(0) = f'_\varepsilon(0) = 0, \quad f''_\varepsilon(s) > 0 \text{ for } s \geq 0, \quad (32)$$

and

$$f_\varepsilon \rightarrow f \quad \text{and} \quad f'_\varepsilon \rightarrow f' \quad \text{uniformly on compact sets of } [0, +\infty). \quad (33)$$

We notice that such an approximation exists as shown in [26, Section 3].

We recall that  $V(\xi) = f(|\xi|)$  (see (19)) for  $\xi \in \mathbb{R}^N$ , and we define  $V^\varepsilon : \mathbb{R}^N \rightarrow \mathbb{R}$  as

$$V^\varepsilon(\xi) := f_\varepsilon(|\xi|).$$

We notice that  $\nabla_\xi V^\varepsilon$  and  $\nabla_\xi V$  can be continuously extended to 0 at  $\xi = 0$ .

We approximate  $\Omega$  by domains  $\Omega_\delta$  obtained by chopping off a  $\delta$ -tubular neighborhood of  $\partial\Gamma_0$  and a  $\delta$ -neighborhood of  $O$ . For  $n \in \mathbb{N}$ , we consider  $u_\delta^n \in C^\infty(\Omega_\delta) \cap C^1(\overline{\Omega}_\delta)$  such that

$$u_\delta^n \rightarrow u \text{ in } C^1(\overline{\Omega}_\delta),$$

as  $n$  goes to infinity (see for instance [8, Section 2.6]).

Since

$$\operatorname{div}(x \cdot \nabla u_\delta^n \nabla_\xi V^\varepsilon(\nabla u_\delta^n)) = x \cdot \nabla u_\delta^n \operatorname{div}(\nabla_\xi V^\varepsilon(\nabla u_\delta^n)) + \nabla(x \cdot \nabla u_\delta^n) \cdot \nabla_\xi V^\varepsilon(\nabla u_\delta^n)$$

and from

$$\begin{aligned} \nabla(x \cdot \nabla u_\delta^n) \cdot \nabla_\xi V^\varepsilon(\nabla u_\delta^n) &= \nabla u_\delta^n \cdot \nabla_\xi V^\varepsilon(\nabla u_\delta^n) + x \nabla^2(u_\delta^n) \cdot \nabla_\xi V^\varepsilon(\nabla u_\delta^n) \\ &= \operatorname{div}(u_\delta^n \nabla_\xi V^\varepsilon(\nabla u_\delta^n)) - u_\delta^n \operatorname{div}(\nabla_\xi V^\varepsilon(\nabla u_\delta^n)) \\ &\quad + \operatorname{div}(x V^\varepsilon(\nabla u_\delta^n)) - N V^\varepsilon(\nabla u_\delta^n), \end{aligned}$$

we obtain

$$\operatorname{div}(\varphi_n \nabla_\xi V^\varepsilon(\nabla u_\delta^n) - x V^\varepsilon(\nabla u_\delta^n)) = \varphi_n \operatorname{div}(\nabla_\xi V^\varepsilon(\nabla u_\delta^n)) - N V^\varepsilon(\nabla u_\delta^n), \quad (34)$$

where

$$\varphi_n(x) = x \cdot \nabla u_\delta^n(x) - u_\delta^n(x).$$

Moreover, from the divergence theorem we have

$$\int_{\Omega_\delta} \nabla_\xi V^\varepsilon(\nabla u_\delta^n) \cdot \nabla \varphi_n \, dx = - \int_{\Omega_\delta} \varphi_n \operatorname{div}(\nabla_\xi V^\varepsilon(\nabla u_\delta^n)) \, dx + \int_{\partial \Omega_\delta} \varphi_n \nabla_\xi V^\varepsilon(\nabla u_\delta^n) \cdot \nu \, d\sigma. \quad (35)$$

We are going to apply the divergence theorem in  $\Omega_\delta$ ; to this end we set

$$\Gamma_{0,\delta} = \Gamma_0 \cap \partial \Omega_\delta, \quad \Gamma_{1,\delta} = \Gamma_1 \cap \partial \Omega_\delta \quad \text{and} \quad \Gamma_\delta = \partial \Omega_\delta \setminus (\Gamma_{0,\delta} \cup \Gamma_{1,\delta}).$$

From (35) and by integrating (34) in  $\Omega_\delta$  we obtain

$$\begin{aligned} \int_{\Omega_\delta} \nabla_\xi V^\varepsilon(\nabla u_\delta^n) \cdot \nabla \varphi_n \, dx &= -N \int_{\Omega_\delta} V^\varepsilon(\nabla u_\delta^n) \, dx - \int_{\Omega_\delta} \operatorname{div}(\varphi_n \nabla_\xi V^\varepsilon(\nabla u_\delta^n)) \, dx \\ &\quad + \int_{\Omega_\delta} \operatorname{div}(x V^\varepsilon(\nabla u_\delta^n)) \, dx, \end{aligned}$$

and from  $x \cdot \nu = 0$  on  $\Gamma_{1,\delta}$ , we find

$$\begin{aligned} \int_{\Omega_\delta} \nabla_\xi V^\varepsilon(\nabla u_\delta^n) \cdot \nabla \varphi_n \, dx &= -N \int_{\Omega_\delta} V^\varepsilon(\nabla u_\delta^n) \, dx - \int_{\Gamma_{0,\delta} \cup \Gamma_{1,\delta}} \varphi_n \nabla_\xi V^\varepsilon(\nabla u_\delta^n) \cdot \nu \, d\sigma \\ &\quad + \int_{\Gamma_{0,\delta}} V^\varepsilon(\nabla u_\delta^n) x \cdot \nu \, d\sigma \\ &\quad - \int_{\Gamma_\delta} [\varphi_n \nabla_\xi V^\varepsilon(\nabla u_\delta^n) - x V^\varepsilon(\nabla u_\delta^n)] \cdot \nu \, d\sigma. \end{aligned}$$

By taking the limit as  $\varepsilon \rightarrow 0$  and then as  $n \rightarrow \infty$ , using that  $\nabla u \cdot \nu = 0$  on  $\Gamma_{1,\delta}$  (since  $\partial_\nu u = 0$  on  $\Gamma_1$ ), we obtain

$$\begin{aligned} \int_{\Omega_\delta} \nabla_\xi V(\nabla u) \cdot \nabla \varphi \, dx &= -N \int_{\Omega_\delta} V(\nabla u) \, dx - \int_{\Gamma_{0,\delta}} \varphi \nabla_\xi V(\nabla u) \cdot \nu \, d\sigma + \int_{\Gamma_{0,\delta}} V(\nabla u) x \cdot \nu \, d\sigma \\ &\quad - \int_{\Gamma_\delta} [\varphi \nabla_\xi V(\nabla u) - x V(\nabla u)] \cdot \nu \, d\sigma \end{aligned} \quad (36)$$

where we let

$$\varphi(x) = x \cdot \nabla u(x) - u(x). \quad (37)$$

Now, we take the limit as  $\delta \rightarrow 0$ . Since  $u \in W^{1,\infty}(\Omega)$  and  $\mathcal{H}_{N-1}(\Gamma_\delta)$  goes to 0 as  $\delta \rightarrow 0$ , we have that the last term in (36) vanishes and we obtain

$$\int_{\Omega} \nabla_\xi V(\nabla u) \cdot \nabla \varphi \, dx = -N \int_{\Omega} V(\nabla u) \, dx - \int_{\Gamma_0} \varphi \nabla_\xi V(\nabla u) \cdot \nu \, d\sigma + \int_{\Gamma_0} V(\nabla u) x \cdot \nu \, d\sigma,$$

i.e. (in terms of  $f$ )

$$\int_{\Omega} \frac{f'(|\nabla u|)}{|\nabla u|} \nabla u \cdot \nabla \varphi \, dx = -N \int_{\Omega} f(|\nabla u|) \, dx - \int_{\Gamma_0} \varphi \frac{f'(|\nabla u|)}{|\nabla u|} \partial_\nu u \, d\sigma + \int_{\Gamma_0} f(|\nabla u|) x \cdot \nu \, d\sigma.$$

Since  $u$  satisfies (13), we get

$$\int_{\Omega} \varphi \, dx = -N \int_{\Omega} f(|\nabla u|) \, dx - \int_{\Gamma_0} \varphi \frac{f'(|\nabla u|)}{|\nabla u|} \partial_{\nu} u \, d\sigma + \int_{\Gamma_0} f(|\nabla u|) x \cdot \nu \, d\sigma. \quad (38)$$

From (37) and since  $u = 0$  on  $\Gamma_0$  and  $\partial_{\nu} u = 0$  on  $\Gamma_1$ , we have

$$\int_{\Omega} \varphi \, dx = -(N+1) \int_{\Omega} u \, dx$$

and

$$\int_{\Gamma_0} \varphi \frac{f'(|\nabla u|)}{|\nabla u|} \partial_{\nu} u \, d\sigma = \int_{\Gamma_0} f'(|\nabla u|) |\nabla u| x \cdot \nu \, d\sigma, \quad (39)$$

where we used the expression of the unit exterior normal on  $\Gamma_0$  given by (15). From (39) and (38) we obtain

$$-(N+1) \int_{\Omega} u \, dx + N \int_{\Omega} f(|\nabla u|) \, dx = - \int_{\Gamma_0} f'(|\nabla u|) |\nabla u| x \cdot \nu \, d\sigma + \int_{\Gamma_0} f(|\nabla u|) x \cdot \nu \, d\sigma.$$

which is (31), and the proof is complete.  $\square$

We conclude this subsection by exploiting the boundary condition  $\partial_{\nu} u = 0$  on  $\Gamma_1$ . Before doing this, we need to recall some notation from differential geometry (see also [18, Appendix A]). We denote by  $D$  the standard Levi-Civita connection. Recall that, given an  $(N-1)$ -dimensional smooth orientable submanifold  $M$  of  $\mathbb{R}^N$  we define the *tangential gradient* of a smooth function  $f : M \rightarrow \mathbb{R}$  with respect to  $M$  as

$$\nabla^T f(x) = \nabla f(x) - \nu \cdot \nabla f(x) \nu$$

for  $x \in M$ , where  $\nabla f$  denotes the usual gradient of  $f$  in  $\mathbb{R}^N$  and  $\nu$  is the outward unit normal at  $x$  to  $M$ . Moreover, we recall that the *second fundamental form* of  $M$  is the bilinear and symmetric form defined on  $TM \times TM$  as

$$\mathbb{I}\mathbb{I}(v, w) = D\nu(v)w \cdot \nu;$$

a submanifold is called *convex* if the second fundamental form is non-negative definite.

**Lemma 7.** *Let  $u$  be the solution to (5). Then*

$$\nabla_{\xi} V(\nabla u) \cdot \nu = 0 \quad \text{on } \Gamma_1, \quad (40)$$

and

$$\nabla(\nabla_{\xi} V(\nabla u) \cdot \nu) \cdot \nabla u = 0 \quad \text{on } \Gamma_1.^1 \quad (41)$$

*Proof.* Since  $\partial_{\nu} u = 0$  on  $\Gamma_1$ , we immediately find (40). By taking the tangential derivative in (40) we get

$$0 = \nabla^T(\nabla_{\xi} V(\nabla u) \cdot \nu) = \nabla(\nabla_{\xi} V(\nabla u) \cdot \nu) - \nu \cdot \nabla(\nabla_{\xi} V(\nabla u) \cdot \nu) \nu \quad \text{on } \Gamma_1.$$

By taking the scalar product with  $\nabla u$  we obtain

$$0 = \nabla(\nabla_{\xi} V(\nabla u) \cdot \nu) \cdot \nabla u - \nu \cdot \nabla(\nabla_{\xi} V(\nabla u) \cdot \nu) \partial_{\nu} u,$$

and since  $\partial_{\nu} u = 0$  on  $\Gamma_1$ , we find (41).  $\square$

<sup>1</sup>We remark that (41) is understood to be zero at points where  $\nabla u = 0$ .

**2.3. Integral Identities for  $S_2$ .** In this Subsection we prove some integral inequalities involving  $S_2(W)$  and the solution to problem (5).

**Lemma 8.** *Let  $\Omega \subset \mathbb{R}^N$  be a sector-like domain and assume that  $f$  satisfies (4). Let  $u \in W^{1,\infty}(\Omega)$  be a solution of (5) such that (9) holds. Then the following inequality*

$$2 \int_{\Omega} S_2(W)u \, dx \geq - \int_{\Omega} S_{ij}^2(W)V_{\xi_i}(\nabla u)\partial_j u \, dx \quad (42)$$

holds. Moreover the equality sign holds in (42) if and only if  $\Pi(\nabla^T u, \nabla^T u) = 0$  on  $\Gamma_1$ .

*Proof.* We split the proof in two steps.

*Step 1: the following identity*

$$2 \int_{\Omega} S_2(W)\phi \, dx = - \int_{\Omega} S_{ij}^2(W)V_{\xi_i}(\nabla u)\partial_j \phi \, dx, \quad (43)$$

holds for every  $\phi \in C_0^1(\Omega)$ .

For  $t > 0$  we set  $\Omega_t = \{x \in \Omega : \text{dist}(x, \partial\Omega) > t\}$ . Let  $\phi \in C_0^1(\Omega)$  be a test function and let  $\varepsilon_0 > 0$  be such that  $\Omega_{\varepsilon_0} \subset \Omega$  and  $\text{supp}(\phi) \subset \Omega_{\varepsilon_0}$ . For  $\varepsilon < \varepsilon_0$  sufficiently small, we set

$$a^i(x) = V_{\xi_i}(\nabla u(x)) \quad \text{for every } i = 1, \dots, N, x \in \Omega.$$

From (9) we have that  $a^i \in W^{1,2}(\Omega)$ ,  $i = 1, \dots, N$ . With this notation, the elements  $w_{ij} = \partial_j V_{\xi_i}(\nabla u)$  of the matrix  $W$  are given by

$$w_{ij} = \partial_j a^i.$$

Let  $\{\rho_\varepsilon\}$  be a family of mollifiers and define  $a_\varepsilon^i = a^i * \rho_\varepsilon$ . Let  $W^\varepsilon = (w_{ij}^\varepsilon)_{i,j=1,\dots,N}$  where  $w_{ij}^\varepsilon = \partial_j a_\varepsilon^i$ , and notice that

$$a_\varepsilon^i \rightarrow a^i \quad \text{in } W^{1,2}(\Omega_{\varepsilon_0}) \quad \text{and} \quad W^\varepsilon \rightarrow W \quad \text{in } L^2(\Omega_{\varepsilon_0}),$$

as  $\varepsilon \rightarrow 0$ . Moreover, since

$$\text{Tr } W^\varepsilon(x) = \int_{\mathbb{R}^N} \rho_\varepsilon(y) \text{Tr } W(x-y) \, dy$$

and  $\text{Tr } W = -1$ , we have that

$$\text{Tr } W^\varepsilon(x) = -1 \quad (44)$$

for every  $x \in \Omega_\varepsilon$ .

Let  $i, j = 1, \dots, N$  be fixed. We have

$$\begin{aligned} w_{ji}^\varepsilon w_{ij}^\varepsilon &= \partial_j (a_\varepsilon^i \partial_i a_\varepsilon^j) - a_\varepsilon^i \partial_j \partial_i a_\varepsilon^j \\ &= \partial_j (a_\varepsilon^i \partial_i a_\varepsilon^j) - a_\varepsilon^i \partial_i \partial_j a_\varepsilon^j \\ &= \partial_j (a_\varepsilon^i \partial_i a_\varepsilon^j) - a_\varepsilon^i \partial_i w_{jj}^\varepsilon, \end{aligned}$$

for every  $x \in \Omega_\varepsilon$ , and by summing up over  $j = 1, \dots, N$ , using (44) (hence  $\partial_i \sum_j w_{jj}^\varepsilon = 0$ ), we obtain

$$\begin{aligned} \sum_j w_{ji}^\varepsilon w_{ij}^\varepsilon &= \sum_j \partial_j (a_\varepsilon^i \partial_i a_\varepsilon^j) \\ &= w_{ii}^\varepsilon \text{Tr } W^\varepsilon - \sum_j \partial_j (S_{ij}^2(W^\varepsilon) a_\varepsilon^i), \quad x \in \Omega_\varepsilon. \end{aligned}$$

By summing over  $i = 1, \dots, N$ , from (25) and (28) we have

$$2S_2(W^\varepsilon) = \sum_{i,j} \partial_j (S_{ij}^2(W^\varepsilon) a_\varepsilon^i), \quad x \in \Omega_\varepsilon. \quad (45)$$

Since

$$\int_{\Omega_{\varepsilon_0}} \partial_j (S_{ij}^2(W^\varepsilon) a_\varepsilon^i) \phi \, dx + \int_{\Omega_{\varepsilon_0}} S_{ij}^2(W^\varepsilon) a_\varepsilon^i \partial_j \phi \, dx = \int_{\partial\Omega_{\varepsilon_0}} S_{ij}^2(W^\varepsilon) a_\varepsilon^i \nu_j \phi \, d\sigma = 0,$$

from (45) and by letting  $\varepsilon$  to zero, we obtain (43).

*Step 2.* Let  $\delta > 0$  and consider a cut-off function  $\eta^\delta \in C_c^\infty(\Omega)$  such that  $\eta^\delta = 1$  in  $\Omega_\delta$  and  $|\nabla \eta^\delta| \leq \frac{C}{\delta}$  in  $\Omega \setminus \Omega_\delta$  for some constant  $C$  not depending on  $\delta$ . By taking  $\phi(x) = u(x)\eta^\delta(x)$  for  $x \in \Omega$  in (43) we obtain

$$2 \int_{\Omega} S_2(W)u\eta^\delta dx = - \int_{\Omega} S_{ij}^2(W)V_{\xi_i}(\nabla u)\partial_j u\eta^\delta dx - \int_{\Omega} S_{ij}^2(W)V_{\xi_i}(\nabla u)u\partial_j(\eta^\delta) dx. \quad (46)$$

From (9) we have that  $W \in L^2(\Omega)$  and the dominated convergence Theorem yields

$$2 \int_{\Omega} S_2(W)u\eta^\delta dx \rightarrow 2 \int_{\Omega} S_2(W)u dx, \quad (47)$$

as  $\delta \rightarrow 0$ . Analogously,

$$\int_{\Omega} S_{ij}^2(W)V_{\xi_i}(\nabla u)\partial_j u\eta^\delta dx \rightarrow \int_{\Omega} S_{ij}^2(W)V_{\xi_i}(\nabla u)\partial_j u dx, \quad (48)$$

as  $\delta \rightarrow 0$ .

Now, we consider the last term in (46). We write  $\Omega$  in the following way:

$$\Omega = A_0^\delta \cup A_1^\delta, \quad (49)$$

where

$$A_0^\delta = \{x \in \Omega : \text{dist}(x, \Gamma_0) \leq \delta\} \quad \text{and} \quad A_1^\delta = \Omega \setminus A_0^\delta.$$

Since  $u = 0$  on  $\Gamma_0$ , we get that

$$u(x) \leq \|u\|_{W^{1,\infty}(\Omega)} \text{dist}(x, \Gamma_0) \leq \|u\|_{W^{1,\infty}(\Omega)} \delta$$

for every  $x \in A_0^\delta$  and we obtain

$$\left| \int_{A_0^\delta} S_{ij}^2(W)V_{\xi_i}(\nabla u)u\partial_j(\eta^\delta) dx \right| \leq C_1 |A_0^\delta|,$$

where  $C_1$  is a constant depending on  $\|u\|_{W^{1,\infty}(\Omega)}$  and  $\|W\|_{L^2(\Omega)}$ , which implies that

$$\lim_{\delta \rightarrow 0} \int_{A_0^\delta} S_{ij}^2(W)V_{\xi_i}(\nabla u)u\partial_j(\eta^\delta) dx = 0. \quad (50)$$

Now we show that

$$\lim_{\delta \rightarrow 0} \int_{A_1^\delta} S_{ij}^2(W(x))V_{\xi_i}(\nabla u(x))u(x)\partial_j(\eta^\delta)(x) dx \geq 0. \quad (51)$$

By choosing  $\delta$  small enough, a point  $x \in A_1^\delta$  can be written in the following way:  $x = \bar{x} + t\nu(\bar{x})$  where  $\bar{x} = \bar{x}(x) \in \Gamma_1$  and  $t = |x - \bar{x}|$  with  $0 < t < \delta$ . Moreover, by using a standard approximation argument,  $\eta^\delta$  can be chosen in such a way that  $\eta^\delta(x) = \frac{1}{\delta} \text{dist}(x, \Gamma_1)$  for any  $x \in A_1^\delta$ , so that

$$\nabla \eta^\delta(x) = -\frac{1}{\delta} \nu(\bar{x}), \quad (52)$$

for every  $x \in A_1^\delta \setminus \Omega_\delta$ . For simplicity of notation we set  $F = (F_1, \dots, F_N)$ , where

$$F_j(x) = u(x)S_{ij}^2(W(x))V_{\xi_i}(\nabla u(x)) \quad (53)$$

for  $j = 1, \dots, N$ , and hence

$$\int_{A_1^\delta} S_{ij}^2(W)V_{\xi_i}(\nabla u)u\partial_j(\eta^\delta) dx = \int_{A_1^\delta} F(x) \cdot \nabla \eta^\delta(x) dx. \quad (54)$$

Since  $\nabla \eta^\delta = 0$  in  $\Omega_\delta$  and  $\nabla \eta^\delta(x) = -\frac{1}{\delta} \nu(\bar{x})$ , for every  $x \in A_1^\delta \setminus \Omega_\delta$ , we have

$$\begin{aligned} \int_{A_1^\delta} F(x) \cdot \nabla \eta^\delta(x) dx &= -\frac{1}{\delta} \int_{A_1^\delta \setminus \Omega_\delta} F(x) \cdot \nu(\bar{x}) dx \\ &= -\frac{1}{\delta} \int_0^\delta dt \int_{\{x \in A_1^\delta : \text{dist}(x, \Gamma_1) = t\}} F(x) \cdot \nu(\bar{x}) d\sigma \end{aligned}$$

where we used coarea formula. Since we are in a *small*  $\delta$ -tubular neighborhood of (part of)  $\Gamma_1$ , we can parametrize  $A_1^\delta \setminus \Omega_\delta$  over (part of)  $\Gamma_1$  as from [28, Formula 14.98] we obtain that

$$\int_{A_1^\delta} F(x) \cdot \nabla \eta^\delta(x) dx = -\frac{1}{\delta} \int_0^\delta dt \int_{\Gamma_1} F(\bar{x} + t\nu(\bar{x})) \cdot \nu(\bar{x}) |\det(Dg)| d\sigma. \quad (55)$$

We notice that, by using this notation, proving (51) is equivalent to prove

$$\lim_{\delta \rightarrow 0} \int_{A_1^\delta} F(x) \cdot \nabla \eta^\delta(x) dx \geq 0, \quad (56)$$

for  $\delta > 0$  sufficiently small.

From (52), (53) and the definition of  $S_{ij}^2$  (24), we have

$$\begin{aligned} F(x) \cdot \nu(\bar{x}) &= -\delta_{ij} V_{\xi_i}(\nabla u(x)) u(x) \nu_j(\bar{x}) - w_{ji}(x) V_{\xi_i}(\nabla u(x)) u(x) \nu_j(\bar{x}) \\ &= -\left\{ \delta_{ij} V_{\xi_i}(\nabla u(x)) u(x) \nu_j(\bar{x}) + u(x) \frac{f'(|\nabla u(x)|)}{|\nabla u(x)|} w_{ji}(x) \partial_i u(x) \nu_j(\bar{x}) \right\} \end{aligned}$$

for almost every  $x = \bar{x} + t\nu(\bar{x}) \in A_1^\delta \setminus \Omega_\delta$ , with  $0 \leq t \leq \delta$ . Since

$$w_{ij} \nu_i \partial_j u = \partial_j (V_{\xi_i}(\nabla u) \nu_i) \partial_j u - V_{\xi_i}(\nabla u) \partial_j \nu_i \partial_j u,$$

we have

$$\begin{aligned} F(x) \cdot \nu(\bar{x}) &= -u(x) \nabla_\xi V(\nabla u(x)) \cdot \nu(\bar{x}) - u(x) \frac{f'(|\nabla u(x)|)}{|\nabla u(x)|} \times \\ &\quad \left\{ \nabla(\nabla_\xi V(\nabla u(x)) \cdot \nu(\bar{x})) \cdot \nabla u(x) - \frac{f'(|\nabla u(x)|)}{|\nabla u(x)|} \partial_j \nu_i(\bar{x}) \partial_j u(x) \partial_i u(x) \right\} \end{aligned} \quad (57)$$

for almost every  $x = \bar{x} + t\nu(\bar{x}) \in A_1^\delta \setminus \Omega_\delta$ , with  $0 \leq t \leq \delta$ . Let

$$\Gamma_1^{\delta,t} = \{x \in A_1^\delta : \text{dist}(x, \Gamma_1) = t\}.$$

We notice that if  $x \in \Gamma_1^{\delta,t}$  then  $\nu(\bar{x}) = \nu^t(x)$  where  $\nu^t(x)$  is the outward normal to  $\Gamma_1^{\delta,t}$  at  $x$ . Hence

$$\partial_j \nu_i(\bar{x}) \partial_j u(x) \partial_i u(x) = \Pi_x^{\delta,t}(\nabla^T u(x), \nabla^T u(x)) \quad (58)$$

where  $\Pi_x^{\delta,t}$  is the second fundamental form of  $\Gamma_1^{\delta,t}$  at  $x$ . Since  $\Sigma$  is a convex cone then the second fundamental form of  $\Gamma_1 \setminus \{O\}$  is non-negative definite. This implies that the second fundamental form of  $\Gamma_1^{\delta,t}$  is non-negative definite for  $t$  sufficiently small [28, Appendix 14.6] and hence

$$\partial_j \nu_i(\bar{x}) \partial_j u(x) \partial_i u(x) \geq 0. \quad (59)$$

From (59) and (57) we obtain

$$F(x) \cdot \nu(\bar{x}) \geq -u(x) \nabla_\xi V(\nabla u(x)) \cdot \nu(\bar{x}) - u(x) \frac{f'(|\nabla u(x)|)}{|\nabla u(x)|} \nabla(\nabla_\xi V(\nabla u(x)) \cdot \nu(\bar{x})) \cdot \nabla u(x) \quad (60)$$

for almost every  $x = \bar{x} + t\nu(\bar{x}) \in A_1^\delta \setminus \Omega_\delta$ , with  $0 \leq t \leq \delta$ . We use (60) in the right-hand side of (55) and, by taking the limit as  $\delta \rightarrow 0$ , we obtain

$$\lim_{\delta \rightarrow 0} \int_{A_1^\delta} F(x) \cdot \nabla \eta^\delta(x) dx \geq - \int_{\Gamma_1} u \left( \nabla_\xi V(\nabla u) \cdot \nu + \frac{f'(|\nabla u|)}{|\nabla u|} \nabla(\nabla_\xi V(\nabla u) \cdot \nu) \cdot \nabla u \right) d\sigma.$$

From (40) and (41) we find (56), and hence (51). From (46), (47), (48), (49), (50) and (51), we obtain (42).  $\square$

## 3. PROOF OF THEOREM 1

*Proof of Theorem 1.* We divide the proof in two steps. We first show that

$$W = -\frac{1}{N}Id \quad \text{a.e. in } \Omega. \quad (61)$$

and

$$\Pi(\nabla^T u, \nabla^T u) = 0 \quad \text{on } \Gamma_1, \quad (62)$$

and then we exploit (61) in order to prove that  $u$  is indeed radial.

*Step 1.* Let  $g$  be the Fenchel conjugate of  $f$  (in our case  $g' = (f')^{-1}$ ), using (20) we get that

$$\begin{aligned} \operatorname{div}(g(|\nabla_\xi V(\nabla u)|)\nabla_\xi V(\nabla u)) &= g'(|\nabla_\xi V(\nabla u)|)\nabla|\nabla_\xi V(\nabla u)|V_\xi(\nabla u) + g(|\nabla_\xi V(\nabla u)|)\operatorname{Tr}(W) \\ &= g'(f'(|\nabla u|))\frac{V_{\xi_i}(\nabla u)}{|\nabla_\xi V(\nabla u)|}\partial_j(V_{\xi_i}(\nabla u))V_{\xi_j}(\nabla u) \\ &\quad + g(f'(|\nabla u|))\operatorname{Tr}(W), \end{aligned}$$

a.e. in  $\Omega$ , where we used (20). Since  $\partial_j V_{\xi_i}(\nabla u) = w_{ij}$  and  $g' = (f')^{-1}$ , we obtain

$$\operatorname{div}(g(|\nabla_\xi V(\nabla u)|)\nabla_\xi V(\nabla u)) = \partial_i w w_{ij} V_{\xi_j}(\nabla u) + g(f'(|\nabla u|))\operatorname{Tr}(W)$$

a.e. in  $\Omega$ , and using again (20) we find

$$\operatorname{div}(g(|\nabla_\xi V(\nabla u)|)\nabla_\xi V(\nabla u)) = \frac{f'(|\nabla u|)}{|\nabla u|}\partial_i w w_{ij}\partial_j u + g(f'(|\nabla u|))\operatorname{Tr}(W)$$

a.e. in  $\Omega$ . Since

$$g(f'(t)) = t f'(t) - f(t) \quad (63)$$

and  $\operatorname{Tr}(W) = -1$ , we obtain

$$\operatorname{div}(g(|\nabla_\xi V(\nabla u)|)\nabla_\xi V(\nabla u)) = \frac{f'(|\nabla u|)}{|\nabla u|}\partial_i w w_{ij}\partial_j u + f(|\nabla u|) - |\nabla u|f'(|\nabla u|) \quad (64)$$

a.e. in  $\Omega$ .

Since (27), (20) and (22) yield

$$-S_{ij}^2(W)V_{\xi_i}(\nabla u)\partial_j u = \frac{f'(|\nabla u|)}{|\nabla u|}w_{ji}\partial_i u\partial_j u + f'(|\nabla u|)|\nabla u|,$$

a.e. in  $\Omega$ , from (64) we obtain

$$-S_{ij}^2(W)V_{\xi_i}(\nabla u)\partial_j u = \operatorname{div}(g(|\nabla_\xi V(\nabla u)|)\nabla_\xi V(\nabla u)) + 2f'(|\nabla u|)|\nabla u| - f(|\nabla u|), \quad (65)$$

a.e. in  $\Omega$ .

From Lemma 8 and (65), we obtain

$$\begin{aligned} 2 \int_{\Omega} S_2(W)u \, dx &\geq - \int_{\Omega} S_{ij}^2(W)V_{\xi_i}(\nabla u)\partial_j u \, dx \\ &= \int_{\partial\Omega} g(|\nabla_\xi V(\nabla u)|)\nabla_\xi V(\nabla u) \cdot \nu \, d\sigma + \int_{\Omega} [2f'(|\nabla u|)|\nabla u| - f(|\nabla u|)] \, dx. \end{aligned}$$

From (20) and (40) we find

$$2 \int_{\Omega} S_2(W)u \, dx \geq \int_{\Gamma_0} g(|\nabla_\xi V(\nabla u)|)\frac{f'(|\nabla u|)}{|\nabla u|}\partial_\nu u \, d\sigma + \int_{\Omega} [2f'(|\nabla u|)|\nabla u| - f(|\nabla u|)] \, dx.$$

From (20) and (8) we have

$$2 \int_{\Omega} S_2(W)u \, dx \geq -g(f'(c))f'(c)|\Gamma_0| + \int_{\Omega} [2f'(|\nabla u|)|\nabla u| - f(|\nabla u|)] \, dx$$

and from (63) we obtain

$$2 \int_{\Omega} S_2(W)u \, dx \geq -[cf'(c) - f(c)]f'(c)|\Gamma_0| + \int_{\Omega} [2f'(|\nabla u|)|\nabla u| - f(|\nabla u|)] \, dx. \quad (66)$$

From the Pohozaev identity (31) and (16) we get

$$(N+1) \int_{\Omega} u \, dx - N \int_{\Omega} f(|\nabla u|) \, dx = (f'(c)c - f(c))N|\Omega|;$$

which we use in (66) to obtain

$$2 \int_{\Omega} S_2(W)u \, dx \geq -\frac{f'(c)|\Gamma_0|}{N|\Omega|} \int_{\Omega} [(N+1)u - Nf(|\nabla u|)] \, dx \\ + \int_{\Omega} [2f'(|\nabla u|)|\nabla u| - f(|\nabla u|)] \, dx. \quad (67)$$

We notice that from (17) we have

$$|\Omega| = f'(c)|\Gamma_0|,$$

and from (67) we obtain

$$2 \int_{\Omega} S_2(W)u \, dx \geq -\frac{N+1}{N} \int_{\Omega} u \, dx + 2 \int_{\Omega} f'(|\nabla u|)|\nabla u| \, dx. \quad (68)$$

By using  $u$  as a test function in (13) we have that

$$\int_{\Omega} u \, dx = \int_{\Omega} f'(|\nabla u|)|\nabla u| \, dx,$$

and from (68) we find

$$2 \int_{\Omega} S_2(W)u \, dx \geq \frac{N-1}{N} \int_{\Omega} u \, dx. \quad (69)$$

From (29) and using the fact that  $\text{Tr}(W) = L_f u = -1$ , we get that also the reverse inequality

$$\frac{N-1}{N} \int_{\Omega} u \, dx \geq \int_{\Omega} 2S_2(W)u \, dx \quad (70)$$

holds. From (69) and (70), we conclude that the equality sign must hold in (69) and (70). From Lemma 4 we have that

$$W = \frac{\text{Tr}(W)}{N} Id$$

a.e. in  $\Omega$ , and since  $\text{Tr}(W) = -1$  we obtain (61). Moreover, Lemma 8 yields (62).

*Step 2:  $u$  is a radial function.* From (61) we have that

$$-\frac{1}{N}\delta_{ij} = \partial_j V_{\xi_i}(\nabla u(x)),$$

for every  $i, j = 1, \dots, N$ , which implies that there exists  $x_0 \in \mathbb{R}^N$  such that

$$\nabla_{\xi} V(\nabla u(x)) = -\frac{1}{N}(x - x_0),$$

i.e. according to (20)

$$\frac{f'(|\nabla u(x)|)}{|\nabla u(x)|} \nabla u(x) = -\frac{1}{N}(x - x_0).$$

Hence

$$\nabla u(x) = -g' \left( \frac{|x - x_0|}{N} \right) \frac{x - x_0}{|x - x_0|} \quad \text{in } \Omega.$$

Since  $u = 0$  on  $\Gamma_0$ , we obtain (7) and in particular  $u$  is radial with respect to  $x_0$ . Moreover, from (62) we find that  $x_0$  must be the origin or, if  $\partial\Sigma$  contains flat regions, a point on  $\partial\Sigma$ .  $\square$

## 4. CONES IN SPACE FORMS: PROOF OF THEOREM 2

The goal of this section is to give an easily readable proof of Theorem 2. More precisely we assume more regularity on the solution than the one actually assumed in Theorem 2 in order to give a concise and clear idea of the proof in this setting, and we omit the technical details which are, in fact, needed. A rigorous treatment of the argument described below can be done by adapting the (technical) details in Section 3 and in [35].

Before starting the proof we declare some notations we use in the statement of Theorem 2 and we are going to adopt in the following. Given a  $N$ -dimensional Riemannian manifold  $(M, g)$ , we denote by  $D$  the Levi-Civita connection of  $g$ . Moreover given a  $C^2$ -map  $u : M \rightarrow \mathbb{R}$ , we denote by  $\nabla u$  the gradient of  $u$ , i.e. the dual field of the differential of  $u$  with respect to  $g$ , and by  $\nabla^2 u = Ddu$  the Hessian of  $u$ . We denote by  $\Delta$  the Laplace-Betrami operator induced by  $g$ ;  $\Delta u$  can be defined as the trace of  $\nabla^2 u$  with respect to  $g$ . Given a vector field  $X$  on an oriented Riemannian manifold  $(M, g)$ , we denote by  $\operatorname{div} X$  the divergence of  $X$  with respect to  $g$ . If  $\{e_k\}$  is a local orthonormal frame on  $(M, g)$ , then

$$\operatorname{div} X = \sum_{k=1}^N g(D_{e_k} X, e_k);$$

notice that, if  $u$  is a  $C^1$ -map and if  $X$  is a  $C^1$  vector field on  $M$ , we have the following *integration by parts* formula

$$\int_{\Omega} g(\nabla u, \nu) dx = - \int_{\Omega} u \operatorname{div} X dx + \int_{\partial\Omega} u g(X, \nu) d\sigma,$$

where  $\nu$  is the outward normal to  $\partial\Omega$  and  $\Omega$  is a bounded domain which is regular enough. Here and in the following,  $dx$  and  $d\sigma$  denote the volume form of  $g$  and the induced  $(N-1)$ -dimensional Hausdorff measure, respectively.

*Proof of Theorem 2.* We divide the proof in four steps.

*Step 1: the P-function.* Let  $u$  be the solution to problem (11) and, as in [15], we consider the  $P$ -function defined by

$$P = |\nabla u|^2 + \frac{2}{N}u + Ku^2.$$

Following [15, Lemma 2.1],  $P$  is a subharmonic function and, since  $u = 0$  on  $\Gamma_0$  and from (16), we have that  $P = c^2$  on  $\Gamma_0$ . Moreover,

$$\nabla P = 2\nabla^2 u \nabla u + \frac{2}{n} \nabla u + 2Ku \nabla u. \quad (71)$$

From the convexity assumption of the cone  $\Sigma$ , we have that

$$g(\nabla^2 u \nabla u, \nu) \leq 0. \quad (72)$$

Indeed, since  $u_\nu = 0$  on  $\Gamma_1$  and by arguing as done for (41), we obtain that

$$0 = g(\nabla u_\nu, \nabla u) = g(\nabla^2 u \nabla u, \nu) + \operatorname{II}(\nabla u, \nabla u) \geq g(\nabla^2 u \nabla u, \nu) \quad \text{on } \Gamma_1,$$

which is (72). From (71) and (72) we obtain

$$\partial_\nu P = 2g(\nabla^2 u \nabla u, \nu) + \frac{2}{n} \partial_\nu u + 2Ku \partial_\nu u \leq 0 \quad \text{on } \Gamma_1 \setminus \{O\}.$$

Hence, the function  $P$  satisfies:

$$\begin{cases} \Delta P \geq 0 & \text{in } \Omega, \\ P = c^2 & \text{on } \Gamma_0 \\ \partial_\nu P \leq 0 & \text{on } \Gamma_1 \setminus \{O\}. \end{cases}$$

Moreover, again from [15, Lemma 2.1], we have that

$$\Delta P = 0 \quad \text{if and only if} \quad \nabla^2 u = \left(-\frac{1}{N} - Ku\right) g. \quad (73)$$

*Step 2: we have*

$$P \leq c^2 \quad \text{in } \Omega. \quad (74)$$

Indeed, we multiply  $\Delta P \geq 0$  by  $(P - c^2)^+$  and by integrating by parts we obtain

$$0 \geq \int_{\Omega \cap \{P > c^2\}} |\nabla P|^2 dx - \int_{\partial\Omega} (P - c^2)^+ \partial_\nu P d\sigma.$$

Since  $P = c^2$  on  $\Gamma_0$  and  $\partial_\nu P \leq 0$  on  $\Gamma_1$  we obtain that

$$0 \geq \int_{\Omega \cap \{P > c^2\}} |\nabla P|^2 dx \geq 0$$

and hence  $P \leq c^2$ .

*Step 3:  $P = c^2$ .* By contradiction, we assume that  $P < c^2$  in  $\Omega$ . Since  $\dot{h} > 0$ , we have

$$c^2 \int_{\Omega} \dot{h} dx > \int_{\Omega} \dot{h} |\nabla u|^2 dx + \frac{2}{N} \int_{\Omega} \dot{h} u dx + K \int_{\Omega} \dot{h} u^2 dx.$$

Since

$$\operatorname{div}(\dot{h} u \nabla u) = \dot{h} |\nabla u|^2 + \dot{h} u \Delta u + \ddot{h} u \partial_r u$$

and

$$\ddot{h} = -Kh,$$

and from  $u = 0$  on  $\Gamma_0$  and  $\partial_\nu u = 0$  on  $\Gamma_1 \setminus \{O\}$ , we have that

$$\begin{aligned} c^2 \int_{\Omega} \dot{h} dx &> - \int_{\Omega} \dot{h} u \Delta u dx - \int_{\Omega} \ddot{h} u \partial_r u dx + \frac{2}{N} \int_{\Omega} \dot{h} u dx + K \int_{\Omega} \dot{h} u^2 dx \\ &= (N+1)K \int_{\Omega} \dot{h} u^2 dx + \left(1 + \frac{2}{N}\right) \int_{\Omega} \dot{h} u dx + K \int_{\Omega} \dot{h} u \partial_r u dx. \end{aligned}$$

From  $\operatorname{div}(h \partial_r) = N\dot{h}$  we have

$$\operatorname{div}(u^2 h \partial_r) = N\dot{h} u^2 + 2hu \partial_r u,$$

and from  $u = 0$  on  $\Gamma_0$  and  $\partial_\nu u = 0$  on  $\Gamma_1 \setminus \{O\}$  we obtain

$$c^2 \int_{\Omega} \dot{h} dx > \left(1 + \frac{2}{N}\right) \left( \int_{\Omega} \dot{h} u dx - K \int_{\Omega} \dot{h} u \partial_r u dx \right). \quad (75)$$

Now we show that if  $u$  is a solution of (11) satisfying (12) then the equality sign holds in (75). Indeed, let  $X = h \partial_r$  be the radial vector field and, by integrating formula (2.8) in [15], we get

$$-\frac{c^2}{N} \int_{\partial\Omega} g(X, \nu) d\sigma + \frac{N+2}{N} \int_{\Omega} \dot{h} u dx - (N-2)K \int_{\Omega} \dot{h} u^2 dx + \left(\frac{2}{N} - 3\right) K \int_{\Omega} u g(X, \nabla u) dx = 0.$$

Since  $\operatorname{div} X = N\dot{h}$  we obtain

$$c^2 \int_{\Omega} \dot{h} dx = \frac{N+2}{N} \int_{\Omega} \dot{h} u dx - (N-2)K \int_{\Omega} \dot{h} u^2 dx + \left(\frac{2}{N} - 3\right) K \int_{\Omega} u g(X, \nabla u) dx,$$

i.e.

$$c^2 \int_{\Omega} \dot{h} dx = \left(1 + \frac{2}{N}\right) \left( \int_{\Omega} \dot{h} u dx - K \int_{\Omega} \dot{h} u \partial_r u dx \right),$$

where we used that  $u = 0$  on  $\Gamma_0$ ,  $\partial_\nu u = 0$  on  $\Gamma_1 \setminus \{O\}$  and  $g(X, \nu) = 0$  on  $\Gamma_1$ . From (75) we find a contradiction and hence  $P \equiv c^2$  in  $\Omega$ .

*Step 4:  $u$  is radial.* Since  $P$  is constant, then  $\Delta P = 0$  and from (73) we find that  $u$  satisfies the following Obata-type problem

$$\begin{cases} \nabla^2 u = \left(-\frac{1}{N} - Ku\right)g & \text{in } \Omega, \\ u = 0 & \text{on } \Gamma_0, \\ \partial_\nu u = 0 & \text{on } \Gamma_1 \setminus \{O\}. \end{cases} \quad (76)$$

We notice that the maximum and the minimum of  $u$  can not be both achieved on  $\Gamma_0$  since otherwise we would have that  $u \equiv 0$ . Hence, at least one between the maximum and the

minimum of  $u$  is achieved at a point  $p \in \Omega \cup \Gamma_1$ . Let  $\gamma : I \rightarrow M$  be a unit speed maximal geodesic satisfying  $\gamma(0) = p$  and let  $f(s) = u(\gamma(s))$ . From the first equation of (76) it follows

$$f''(s) = -\frac{1}{N} - Kf(s).$$

Moreover, the definition of  $f$  and the fact that  $\nabla u(p) = 0$  yield

$$f'(0) = 0 \quad \text{and} \quad f(0) = u(p),$$

and therefore

$$f(s) = \left(u(p) - \frac{1}{N}\right)H(s) - \frac{1}{N}.$$

This implies that  $u$  has the same expression along any geodesic strating from  $p$ , and hence  $u$  depends only on the distance from  $p$ . This means that  $\Omega = \Sigma \cap B_R$  where  $B_R$  is a geodesic ball and  $u$  depends only on the distance from the center of  $B_R$ .  $\square$

**Acknowledgement.** The authors wish to thank Luigi Vezzoni for suggesting useful remarks regarding Section 4. The authors have been partially supported by the ‘‘Gruppo Nazionale per l’Analisi Matematica, la Probabilit  e le loro Applicazioni’’ (GNAMPA) of the ‘‘Istituto Nazionale di Alta Matematica’’ (INdAM, Italy).

#### REFERENCES

- [1] A. D. Alexandrov. *Uniqueness theorem for surfaces in the large I*. Vestnik Leningrad Univ. **11** (1956). (English translation: Amer. Math. Soc. Translations Ser. 2 **21** (1962) 341-354).
- [2] B. Avelin, T. Kuusi, G. Mingione. *Nonlinear Calder n-Zygmund theory in the limiting case*. Arch. Rational Mech. Anal. **227** (2018) 227-663.
- [3] E. Baer, A. Figalli. *Characterization of isoperimetric sets inside almost-convex cones*. Discrete Contin. Dyn. Syst. **37** (2017) 1-14.
- [4] C. Bianchini, G. Ciraolo. *Wulff shape characterizations in overdetermined anisotropic elliptic problems*. Comm. Partial Differential Equations. Vol. 43 (2018) 790-820.
- [5] B. Brandolini, C. Nitsch, P. Salani, C. Trombetti. *Serrin type overdetermined problems: an alternative proof*. Arch. Rational Mech. Anal. **190** (2008) 267-280.
- [6] B. Brandolini, C. Nitsch, P. Salani, C. Trombetti. *On the stability of the Serrin problem*. J. Differential Equations **245**, no. 6 (2008) 1566-1583.
- [7] S. Brendle. *Constant mean curvature surfaces in warped product manifolds*. Publ. Math. Inst. Hautes  tudes Sci. **117** (2013) 247-269.
- [8] V.I. Burenkov. Sobolev Spaces on Domains. Teubner, Stuttgart (1998).
- [9] X. Cabr , X. Ros-Oton, J. Serra. *Sharp isoperimetric inequalities via the ABP method*. J. Eur. Math. Soc. (JEMS) **18** (2016) 2971-2998.
- [10] A. Cianchi, V. Maz’ya. *Second-Order Two-Sided Estimates in Nonlinear Elliptic Problems*. Arch. Rational Mech. Anal. **229** (2018) 569-599.
- [11] A. Cianchi, P. Salani. *Overdetermined anisotropic elliptic problems*. Math. Ann. **345** (2009) 859-881.
- [12] G. Ciraolo, A. Figalli, A. Roncoroni. *Symmetry results for critical anisotropic  $p$ -Laplacian equations in convex cones*. Preprint. [arXiv:1906.00622](https://arxiv.org/abs/1906.00622).
- [13] G. Ciraolo, F. Maggi. *On the shape of compact hypersurfaces with almost constant mean curvature*. Comm. Pure Appl. Math. **70** (2017) 665-716.
- [14] G. Ciraolo, R. Magnanini, S. Sakaguchi. *Symmetry of minimizers with a level surface parallel to the boundary*. J. Eur. Math. Soc. (JEMS) **17** (2015) 2789-2804.
- [15] G. Ciraolo, L. Vezzoni. *On Serrin’s overdetermined problem in space forms*. Manuscripta Math. **159** (2019) 445-452.
- [16] M. Cozzi, A. Farina, E. Valdinoci. *Monotonicity formulae and classification results for singular, degenerate, anisotropic PDEs*. Adv. Math. **293** (2016) 343-381.
- [17] G. Crasta. *Existence, uniqueness and qualitative properties of minima to radially symmetric non-coercive non-convex variational problem*. Math. Z. **235** (2000) 569-589.
- [18] K. Ecker. Regularity Theory for Mean Curvature Flow. Progress in Nonlinear Differential Equations and Their Applications, Birkh user Basel, 2014.
- [19] A. Farina, B. Kawohl. *Remarks on an overdetermined boundary value problem*. Calc. Var. Partial Differential Equations **31** (2008) 351-357.
- [20] A. Farina, E. Valdinoci. *Partially and globally overdetermined problems of elliptic type*. Adv. Nonlinear Anal. **1**, no. 1 (2012) 27-45.

- [21] A. Farina, E. Valdinoci. *On partially and globally overdetermined problems of elliptic type*. Amer. J. Math. **135** (2013) 1699-1726.
- [22] A. Farina, L. Mari, E. Valdinoci. *Splitting theorems, symmetry results and overdetermined problems for Riemannian manifolds*. Comm. Partial Differential Equations **38**, no. 10 (2013) 1818-1862.
- [23] A. Figalli, E. Indrei. *A sharp stability result for the relative isoperimetric inequality inside convex cones*. J. Geom. Anal. **23** (2013) 938-969.
- [24] R. L. Fosdick, J. Serrin. *Rectilinear steady flow of simple fluids*. Proc. Roy. Soc. (London) Ser. A **332** (1973) 311-333.
- [25] I. Fragalà, F. Gazzola, B. Kawohl. *Overdetermined problems with possibly degenerate ellipticity, a geometric approach*. Math. Z. **254** (2006) 117-132.
- [26] N. Garofalo, J.L. Lewis. *A symmetry result related to some overdetermined boundary value problems*. Amer. J. Math. **111** (1989) 9-33.
- [27] N. Garofalo, E. Sartori. *Symmetry in exterior boundary value problems for quasilinear elliptic equations via blow-up and a priori estimates*. Adv. Differential Equations **4** (1999) 137-161.
- [28] D. Gilbarg, N. S. Trudinger. *Elliptic partial differential equations of second order*. Springer-Verlag, Berlin-New York, 1977.
- [29] M. Grossi, F. Pacella. *Positive solutions of nonlinear elliptic equations with critical Sobolev exponent and mixed boundary conditions*. Proc. Roy. Soc. Edinburgh Sect. A **116** (1990) 23-43.
- [30] S. Kumaresan, J. Prajapat. *Serrin's result for hyperbolic space and sphere*. Duke Math. J. **91** (1998) 17-28.
- [31] J. Li, C. Xia. *An integral formula and its applications on sub-static manifolds*. J. Differ. Geom. (2018).
- [32] P.-L. Lions, F. Pacella, M. Tricarico. *Best constants in Sobolev inequalities for functions vanishing on some part of the boundary and related questions*. Indiana Univ. Math. J. **37** (1998) 301-324.
- [33] P.-L. Lions, F. Pacella. *Isoperimetric inequalities for convex cones*. Proc. Amer. Math. Soc. **109** (1990), 477-485.
- [34] R. Molzon. *Symmetry and overdetermined boundary value problems*. Forum Math **3** (1991) 143-156.
- [35] F. Pacella, G. Tralli. *Overdetermined problems and constant mean curvature surfaces in cones*. To appear in Rev. Mat. Iberoam. [arXiv:1802.03197](https://arxiv.org/abs/1802.03197).
- [36] G. Qiu, C. Xia. *A generalization of Reilly's formula and its applications to a new Heintze-Karcher type inequality*. Int. Math. Res. Not. IMRN no. 17 (2015) 7608-7619.
- [37] G. Qiu, C. Xia. *Overdetermined boundary value problems in  $\mathbb{S}^n$* . J. Math. Study **50**, no. 2 (2017) 165-173.
- [38] W. Reichel. *Radial Symmetry for Elliptic Boundary-Value Problems on Exterior Domains*. Arch. Rational Mech. Anal. **137**, no. 4 (1997) 381-394.
- [39] A. Roncoroni. *A Serrin-type symmetry result on model manifolds: an extension of the Weinberger argument*. C. R. Math. Acad. Sci. Paris. **356** (2018) 648-656.
- [40] J. Serrin. *A symmetry problem in potential theory*. Arch. Rational Mech. Anal. **43** (1971) 304-318.
- [41] G. Wang, C. Xia. *A characterization of the Wulff shape by an overdetermined anisotropic PDE*. Arch. Ration. Mech. Anal. **199**, no.1 (2011) 99-115.
- [42] H. Weinberger. *Remark on the preceding paper of Serrin*. Arch. Rational Mech. Anal. **43** (1971), 319-320.

GIULIO CIRAULO, DIPARTIMENTO DI MATEMATICA "FEDERIGO ENRIQUES", UNIVERSITÀ DEGLI STUDI DI MILANO, VIA CESARE SALDINI 50, 20133, MILANO, ITALY

*Email address:* [giulio.ciraolo@unimi.it](mailto:giulio.ciraolo@unimi.it)

ALBERTO RONCORONI, DIPARTIMENTO DI MATEMATICA "F. CASORATI", UNIVERSITÀ DEGLI STUDI DI PAVIA, VIA FERRATA 5, 27100 PAVIA, ITALY

*Email address:* [alberto.roncoroni01@universitadipavia.it](mailto:alberto.roncoroni01@universitadipavia.it)