



GUIDELINES

Evidence based position paper on physical and rehabilitation medicine (PRM) practice for people with spinal deformities during growth. The European PRM position (UEMS PRM Section)

Stefano NEGRINI^{1,2*}, Fitnat DINCER³, Carlotta KIEKENS⁴, Liisamari KRUGER⁵,
Enrique VARELA-DONOSO⁶, Nicolas CHRISTODOULOU⁷

¹Clinical and Experimental Sciences Department, University of Brescia, Brescia, Italy; ²Don Gnocchi Foundation, Milan, Italy; ³Division of Internal Medicine, Department of Physical and Rehabilitation Medicine Hacettepe University, University Hospital, Ankara, Turkey; ⁴Physical and Rehabilitation Medicine Universitair Ziekenhuis Leuven, Pellenberg, Belgium; ⁵Orton Rehabilitation Centre, Orton Orthopedic Hospital, Helsinki, Finland; ⁶Physical and Rehabilitation Medicine Department, Complutense University, Ciudad Universitaria, Madrid, Spain; ⁷European University Cyprus, School of Medicine, Nicosia, Cyprus

*Corresponding author: Stefano Negrini, Clinical and Experimental Sciences Department, University of Brescia, Brescia, Italy.
E-mail: stefano.negrini@unibs.it

ABSTRACT

INTRODUCTION: Scoliosis and other spinal deformities involve 3-4% of the population during growth. Their so-called conservative treatment is in the field of competence of physical and rehabilitation medicine (PRM) physicians. This evidence based position paper represents the official position of the European Union through the European Union of Medical Specialists (UEMS) - PRM Section. The aim of the paper was to improve PRM specialists' professional practice for patients with spinal deformities during growth.

EVIDENCE ACQUISITION: A systematic review of the literature and a Consensus procedure with 26 recommendations by means of a Delphi method process has been performed involving the delegates of all European countries represented in the UEMS-PRM Section.

EVIDENCE SYNTHESIS: the systematic literature review is reported together with 26 recommendations coming from the Consensus Delphi procedure.

CONCLUSIONS: The professional role of PRM physicians in spinal deformities during growth is to propose a complete PRM treatment for the patients considering all the concurring diseases and pathologies, impairments, activity limitations and participation restrictions. The PRM physician's role is to coordinate the individual PRM project developed in team with other health professionals and medical specialists, in agreement with the patient and his family, according to the specific medical diagnoses.

(Cite this article as: Negrini S, Dincer F, Kiekens C, Kruger L, Varela-Donoso E, Christodoulou N. Evidence based position paper on physical and rehabilitation medicine (PRM) practice for people with spinal deformities during growth. The European PRM position (UEMS PRM Section). Eur J Phys Rehabil Med 2017;53:125-31. DOI: 10.23736/S1973-9087.16.04406-3)

Key words: Scoliosis - Physical and rehabilitation medicine - Guidelines.

Introduction

The main spinal deformities during growth include scoliosis, sagittal curves changes (increased or reduced kyphosis and/or lordosis), and spondylolisis-

thesis.¹ They involve 3-4% of the population during growth.^{1,2} They can be secondary to another pathology (neurological, syndromic, congenital bone malformation, and others).^{2,3} The so-called conservative treatment (rehabilitation and orthopedic not surgical) are in

the field of competence of physical and rehabilitation medicine (PRM) physicians. This is even more true in patients with pathologies that give rise to a secondary spinal deformity, that are usually followed-up by PRM specialists during growth. Nowadays there is no uniformity among different countries across Europe and in the world in the PRM approach.⁴⁻⁶ For these reasons the European Union of Medical Specialists (UEMS) - PRM Section decided to develop one of its evidence based position papers, representing the official position of the European Union. The aim of the paper was to improve PRM specialists' professional practice for patients with spinal deformities during growth.

Evidence acquisition

This paper has been developed according to the Methodology defined by the Professional Practice Committee of the UEMS-PRM Section.⁷ The systematic review of the literature has been performed in PubMed the 24th of November 2014. The string used for the first selection has been “(“scoliosis”[MeSH Terms]) OR (“kyphosis”[MeSH Terms]) OR (“spondylolisthesis”[MeSH Terms])”, activating the filters: “Filters activated: Systematic Reviews, Meta-Analysis, Randomized Controlled Trial, Child: birth-18 years”. The selection process is reported in the Supplementary Figure 1, online content only. The only criterion for including the studies has been the professional relevance for PRM physicians as judged by at least two of the authors, with the main author resolving conflicts. The strength of evidence (SoE) and the strength of recommendation (SoR) are given according to the methodology paper. The Consensus with Delphi procedure has followed the 4 steps proposed by the methodology paper.⁷

Evidence synthesis

Systematic review

All paper found in the systematic review are listed in the Supplementary Appendix I, online content only.

The professional role of the PRM physician in the diagnosis and assessment of patients with idiopathic deformities during growth as a member of a multiprofessional working-group is essential.⁸ Historically, these pathologies have been followed mainly by orthopedic

surgeons, but the evolution of that specialty toward surgery reduced the attention to the so called conservative treatment.⁴ In 2004 started an international effort of the conservative community that gave rise to a new scientific society, SOSORT (international Society On Scoliosis Orthopedic and Rehabilitation Treatment), whose name clearly states the importance of PRM in this field. Moreover, treatment of spinal deformities during growth is based on a team approach,^{9, 10} and most of the allied professionals involved are in the field of PRM: physiotherapists and orthotists mainly, but also psychologist and, according to the country, chiropractors, trainers etc.^{10, 11}

The goal of PRM physicians should be in the diagnosis, prognosis, treatment and management of the team, including patients and parents.⁸ They should accurately determine which curves are at risk of worsening and which of these can be influenced by treatment, versus those curves that are at high risk of requiring surgery. Today possible treatments include:¹¹

- physiotherapeutic scoliosis-specific exercises (PSSE);
- bracing;
- surgery.

It is today accepted that these treatments are not alternative one to the other, but complimentary and have to be offered to different patients, or to the same patient in different moments.¹¹ The alternative used for years in some countries, the so-called “wait and see” strategy (*i.e.*, observation and eventual surgery), has now been mostly abandoned due to the new strong evidence on the effectiveness of braces.^{5, 12} Electrostimulation, traction and manual therapy have also been used, but there is no evidence of the effectiveness of those methods.¹¹ Once accepted that evidence-based clinical practice comes from the best evidence, combined with clinical expertise and patient preferences,^{13, 14} the patient should be made aware of the possible options. The final choice should come from a multidisciplinary shared decision making discussion, because both surgery and bracing require specific clinical expertise.

Most of the papers we found focus on adolescent idiopathic scoliosis (Supplementary Appendix I), with one exception on secondary scoliosis,¹⁵ two on kyphosis^{16, 17} and one on spondylolisthesis.¹⁸ Consequently research in these understudied fields is to be supported.

BRACING

Scientific evidence support bracing, but even if all good quality papers are in favor, the epidemiological quality of evidence is low.^{1, 5, 12} Historically two kind of braces have been used to curb progression of the pathological curves: the rigid spinal orthosis and the elastic orthosis.^{1, 5} One RCT showed the efficacy of rigid,¹² and another of elastic bracing over natural history;¹¹ another demonstrated that rigid braces are more effective than elastic ones.¹⁹ A new super-rigid generation of braces has been developed, with promising results also in high degree surgical curves.²⁰ Nevertheless there is no evidence in favor of any particular rigid brace over another.⁵ Brace treatment for adolescent idiopathic scoliosis continues to be frequently used, and the number of brace types has increased. Predicting progressive curves and refining indications requires additional investigation. Choosing the best possible intervention for each patient demands good knowledge of the alternatives and discussion about the possible effects that can be reached. In-depth discussion with the patient and possibly with the family is unconditional. Also very important is to follow up the benefit of the intervention and to identify the need of change of intervention, especially the need of surgery in case of worsening of the situation.

PSSE

The aim of PSSE is to affect Cobb angles and also secondary outcomes, such as strength, mobility, and balance.^{6, 21} The present evidence is enough to consider PSSE as an appropriate intervention for AIS, even if it is not possible to support a specific exercise regimen.^{6, 21} Experts²² and two good quality prospective studies²³⁻²⁵ support the importance of Scoliosis-Specific Exercises based on autocorrection. Nevertheless, we presently do not know which approach is more effective between inpatient or outpatient, intrinsic or extrinsic autocorrection exercise, and symmetric or asymmetric exercise. Three RCTs showed effectiveness of PSSE at 6 months^{26, 27} and at the end of growth.^{24, 25} Other studies have confirmed the efficacy of exercises in reducing the progression rate (mainly in early puberty) and/or improving the Cobb angles (around the end of growth).²⁸⁻³¹ Exercises have been shown to be effective in reducing brace prescription.²³

SURGERY

No strong evidence has been found in terms of prospective controlled studies to support surgical intervention from the medical point of view,^{32, 33} even if the methodological and ethical difficulty to perform these studies must be recognized. Surgery has been defined as the failure of conservative treatment,³⁴ meaning that it should be applied only when everything before fails: in fact it creates a problem (loss of movement function due to fusion of the spine) to avoid another one (progression of the curve). Actual indications during growth include very high degree curves (mainly secondary) with possible impact on cardiorespiratory functions, and cosmetic reasons, only if the patient and the family agree.

Recommendations

The results of the Consensus procedure are reported in Supplementary Tables I, II, online content only.

Overall general recommendation

1. The professional role of PRM physicians in spinal deformities during growth is to propose a complete PRM treatment for the patients considering all concurring diseases and pathologies, impairments, activity limitations and participation restrictions. The PRM physician's role is to coordinate the individual PRM project developed in team with other health professionals and medical specialists, in agreement with the patient and family, according to the specific medical diagnoses. (SoE: IV; SoR: A)

Idiopathic spinal deformities

Recommendations on PRM physicians' role in medical diagnosis according to ICD

2. It is recommended that PRM physicians dealing with spinal deformities gain specific and wide expertise in the specific medical diagnosis and treatment approaches of these patients. If this is not possible, they should work in team with spinal deformities experts of other specialties to develop the required PRM treatments. (SoE: IV; SoR: B)
3. It is recommended that PRM physicians start the PRM process with a definite expert medical diagnosis of the spinal deformity. (SoE: IV; SoR: B)

Recommendations on PRM physicians' role in PRM diagnosis according to ICF

4. It is recommended that spinal deformities patients requiring a PRM intervention (so-called conservative treatment: mainly bracing and/or exercises) are evaluated by PRM physicians to diagnose their impairments, activity limitations and participation restrictions so to better focus a complete PRM approach. (SoE: IV; SoR: A)

Recommendations on PRM physicians' role in PRM assessment according to ICF

5. It is recommended that PRM treatment is proposed and regularly monitored through a complete PRM assessment including evaluation of disease and impairment (classical radiographic and clinical parameters), but also activity limitations and participation restrictions due to health condition and/or treatments. (SoE: IV; SoR: B)

Recommendations on PRM management and process

INCLUSION CRITERIA (E.G. WHEN AND WHY PRESCRIBE PRM INTERVENTIONS)

6. It is recommended that PRM interventions are proposed to spinal deformity patients with an actual impairment, activity limitation or participation restriction, or with a possible or proven progression that could drive to a future activity limitation or participation restriction in adulthood. (SoE: IV; SoR: A)

PROJECT DEFINITION (DEFINITION OF THE OVERALL AIMS AND STRATEGY OF PRM INTERVENTIONS)

7. It is recommended that the PRM projects are proposed in PRM centres specialised in spinal deformities treatment where all the team is present (PRM doctor, physiotherapist, orthotist, trainer, eventually psychologist and others) and an adequate expertise is granted to patients (PRM structure project). (SoE: IV; SoR: B)
8. It is recommended that the project is defined on a totally individualised basis in agreement with patient and family. (SoE: IV; SoR: A)

TEAM WORK (PROFESSIONALS INVOLVED AND SPECIFIC MODALITIES OF TEAM WORK)

9. It is recommended that a coordinated and organized team follow the patient, including at least a PRM physician, a medical specialist expert in spinal deformities (usually an orthopaedic surgeon, or another PRM physician consultant) in case the PRM physician has not a specific expertise, an orthotist, and a physiotherapist. The patient and his family are part of the team. For the possible surgical choices, an orthopaedic surgeon must be part of the team as well. (SoE: IV; SoR: B)

PRM INTERVENTIONS

10. It is recommended that PRM interventions are developed according to the actual Guidelines produced by the international Society On Scoliosis Orthopaedic and Rehabilitation Treatment (SOSORT),¹ including mainly specific exercises and/or bracing, as well as recommendations on activities of daily living. (SoE: IV; SoR: A)

OUTCOME CRITERIA

11. It is recommended to use the classical main outcome criteria, including Cobb degrees thresholds as described in the literature, aesthetics and overall spinal balance on the frontal and sagittal plane. It is recommended that also patient-centred outcomes (quality of life, activity limitations, participation restrictions, impact of treatments) are monitored regularly together with the usual impairment and disease-centred parameters. (SoE: IV; SoR: B)

LENGTH/DURATION/INTENSITY OF TREATMENT (OVERALL PRACTICAL PRM APPROACH)

12. It is recommended that treatment (bracing and/or exercises according to the actual SOSORT Guidelines) is continued until the impairments (pain, aesthetic impact...) are resolved and/or the risk of progression is finished (enough bone maturity according to the quantity of deformity), so to grant to the patient the best possible future in terms of activity and participation according to the actual prognostic instruments. (SoE: IV; SoR: A)

DISCHARGE CRITERIA (E.G. WHEN AND WHY END PRM INTERVENTIONS)

13. It is recommended that patients are followed up until any further progression of the deformity is not expected anymore (stabilization) and/or impairments (pain, aesthetic impact...) have been resolved as far as possible. (SoE: IV; SoR: A)

Recommendations on future research on PRM professional practice

14. It is recommended to develop ICF compatible evaluation instruments for spinal deformities during growth. (SoE: IV; SoR: B)

Secondary spinal deformities

Literature search yielded a few research related with Secondary Deformities. So most of the recommendations in Secondary Spinal Deformities are based on evidence extrapolated from Evidence, Research and Recommendations of Idiopathic Spinal deformities and is therefore supported by expert opinion in Secondary Spinal Deformities.

Recommendations on PRM physicians' role in Medical Diagnosis according to ICD

15. It is recommended that PRM physicians dealing with pathologies that can give rise to secondary spinal deformities monitor regularly patients to immediately screen any spinal pathology. (SoE: IV; SoR: A)
16. It is recommended that PRM physicians dealing with pathologies that can give rise to secondary spinal deformities gain specific expertise also in the diagnosis of spinal secondary disorders and treatment approached to integrate them in the overall approach to the main pathology. If this is not possible, they should work in team with another medical specialist expert in secondary spinal deformities (usually an orthopaedic surgeon or a neurosurgeon) to develop the required PRM treatments. (SoE: IV; SoR: A)
17. It is recommended that PRM physicians dealing with pathologies that can give rise to secondary spinal deformities plan the PRM process approaching also the spinal deformity eventually present according to a definite expert medical diagnosis. (SoE: IV; SoR: A)

Recommendations on PRM physicians' role in PRM diagnosis according to ICF

18. It is recommended that patients with secondary spinal deformities requiring a PRM intervention are evaluated by PRM physicians to diagnose their impairments, activity limitations and participation restrictions so as to have a complete PRM approach. (SoE: IV; SoR: A)

Recommendations on PRM physicians' role in PRM assessment according to ICF

19. It is recommended that, before and during PRM treatment of patients with secondary spinal deformities, the PRM assessment includes the evaluation of the spinal disease and impairment (classical radiographic and clinical parameters) together with those of the main pathology. (SoE: IV; SoR: A)

*Recommendations on PRM management and process***INCLUSION CRITERIA** (E.G. WHEN AND WHY PRESCRIBE PRM INTERVENTIONS)

20. It is recommended that PRM interventions include an approach to the spinal deformity as soon as it is discovered, due to the high risk of progression, possibly leading with time to a big impairment of trunk statics (and consequently gait and/or posture) and even cardiopulmonary dysfunctions. (SoE: IV; SoR: A)

PROJECT DEFINITION (DEFINITION OF THE OVERALL AIMS AND STRATEGY OF PRM INTERVENTIONS)

21. It is recommended that, in case of spinal deformities, the PRM projects include also specific treatments of the spine (such as bracing, and/or exercises to improve spinal posture and control) according to individual needs. (SoE: IV; SoR: A)

TEAM WORK (PROFESSIONALS INVOLVED AND SPECIFIC MODALITIES OF TEAM WORK)

22. It is recommended that a coordinated and organized team follow the patient, including at least: a PRM physician; a medical specialist expert in spinal de-

formities in case the PRM physician is not an expert (usually an orthopaedic surgeon, a neurosurgeon, or another PRM physician consultant); an expert in the original pathology (*e.g.* paediatrician, neuropaediatrician, neurologist, syndromes expert, orthopaedic surgeon, neurosurgeon etc) causing the spinal deformity; an orthotist; a physiotherapist. The patient and his family are part of the team. (SoE: IV; SoR: B)

PRM INTERVENTIONS, OUTCOME CRITERIA AND OVERALL PRACTICAL PRM APPROACH

23. It is recommended that PRM interventions are developed according to the actual Guidelines produced by the international Society On Scoliosis Orthopaedic and Rehabilitation Treatment (SOSORT),¹ including mainly specific exercises and/or bracing, as well as recommendations on activities of daily living, but they have to be modified according to the underlying cause of secondary spinal deformity. (SoE: IV; SoR: A)

DISCHARGE CRITERIA (*E.G.* WHEN AND WHY TO END PRM INTERVENTIONS)

24. It is recommended that patients with secondary spinal deformities are followed up until any further progression of the deformity is not expected anymore (stabilization). (SoE: IV; SoR: B)

Recommendations on future research on PRM professional practice

25. It is recommended to start systematic research on the epidemiology of secondary spinal deformities and possible conservative rehabilitation approach to patients with secondary spinal deformities during growth. (SoE: IV; SoR: B)

26. It is recommended to start a systematic research to identify the best PRM interventions, their length/duration/intensity, and the correct outcome criteria for secondary spinal deformities during growth. (SoE: IV; SoR: B)

Conclusions

The professional role of PRM physicians in spinal deformities during growth is to propose a complete PRM

treatment for patients considering all the concurring diseases and pathologies, impairments, activity limitations and participation restrictions. The PRM physician's role is to coordinate the individual PRM projects developed in team with other health professionals and medical specialists, in agreement with the patient and family, according to the specific medical diagnoses.

References

- Negrini S, Aulisa AG, Aulisa L, Circo AB, de Mauroy JC, Durmala J, *et al.* 2011 SOSORT guidelines: Orthopaedic and Rehabilitation treatment of idiopathic scoliosis during growth. *Scoliosis* 2012;7:3.
- Hresko MT. Clinical practice. Idiopathic scoliosis in adolescents. *N Engl J Med* 2013;368:834-41.
- Weinstein SL, Dolan LA, Cheng JCY, Danielsson A, Morcuende JA. Adolescent idiopathic scoliosis. *Lancet* 2008;371:1527-37.
- Negrini S. Approach to scoliosis changed due to causes other than evidence: patients call for conservative (rehabilitation) experts to join in team orthopedic surgeons. *Disabil Rehabil* 2008;30:731-41.
- Negrini S, Minozzi S, Bettany-Saltikov J, Chockalingam N, Grivas TB, Kotwicki T, *et al.* Braces for idiopathic scoliosis in adolescents. *Cochrane Database Syst Rev.* 2015;6:CD006850.
- Romano M, Minozzi S, Bettany-Saltikov J, Zaina F, Chockalingam N, Kotwicki T, *et al.* Exercises for adolescent idiopathic scoliosis. *Cochrane Database Syst Rev.* 2012;8:CD007837.
- Negrini S, Kiekens C, Zampolini M, Wever D, Varela Donoso E, Christodoulou N. Methodology of "Physical and Rehabilitation Medicine practice, Evidence Based Position Papers: the European position" produced by the UEMS-PRM Section. *Eur J Phys Rehabil Med* 2016;52:134-41.
- Negrini S, Zaina F. Physical and rehabilitation medicine specialists in the medical approach to idiopathic scoliosis. *Eur J Phys Rehabil Med* 2014;50:83-6.
- Tavernaro M, Pellegrini A, Tessadri F, Zaina F, Zonta A, Negrini S. Team care to cure adolescents with braces (avoiding low quality of life, pain and bad compliance): a case-control retrospective study. 2011 SOSORT Award winner. *Scoliosis* 2012;7:17.
- Negrini S, Grivas TB, Kotwicki T, Rigo M, Zaina F, international Society on Scoliosis Orthopaedic and Rehabilitation Treatment (SOSORT). Guidelines on "Standards of management of idiopathic scoliosis with corrective braces in everyday clinics and in clinical research": SOSORT Consensus 2008. *Scoliosis* 2009;4:2.
- Coillard C, Circo AB, Rivard CH. A Prospective Randomized Controlled Trial of the Natural History of Idiopathic Scoliosis versus treatment with the Spinecor brace. *Sosort Award 2011 Winner.* *Eur J Phys Rehabil Med* 2014;50:479-87.
- Weinstein SL, Dolan LA, Wright JG, Dobbs MB. Effects of Bracing in Adolescents with Idiopathic Scoliosis. *N Engl J Med* 2013;369:1512-21.
- Sackett DL, Rosenberg WM, Gray JA, Haynes RB, Richardson WS. Evidence based medicine: what it is and what it isn't. *BMJ* 1996;312:71-2.
- Negrini S. Why evidence-based medicine is a good approach in physical and rehabilitation medicine - Thesis. *Eur J Phys Rehabil Med* 2014;50:585-91.
- Liptak GS. Tethered spinal cord: update of an analysis of published articles. *Eur J Pediatr Surg* 1995;5 Suppl 1:21-3.
- Seidi F, Rajabi R, Ebrahimi I, Alizadeh MH, Minoonejad H. The efficiency of corrective exercise interventions on thoracic hyper-kyphosis angle. *J Back Musculoskelet Rehabil* 2014;27:7-16.
- Lowe TG, Line BG. Evidence based medicine: analysis of Scheuermann kyphosis. *Spine* 2007;32(19 Suppl):S115-119.

18. Klein G, Mehlman CT, McCarty M. Nonoperative treatment of spondylolysis and grade I spondylolisthesis in children and young adults: a meta-analysis of observational studies. *J Pediatr Orthop* 2009;29:146-56.
19. Wong MS, Cheng JCY, Lam TP, Ng BKW, Sin SW, Lee-Shum SLF, *et al.* The effect of rigid versus flexible spinal orthosis on the clinical efficacy and acceptance of the patients with adolescent idiopathic scoliosis. *Spine* 2008;33:1360-5.
20. Lusini M, Donzelli S, Minnella S, Zaina F, Negrini S. Brace treatment is effective in idiopathic scoliosis over 45°: an observational prospective cohort controlled study; 2013 [Internet]. Available from: <http://linkinghub.elsevier.com/retrieve/pii/S1529943013019359> [cited 2014, Jul 10]
21. Bettany-Saltikov J, Parent E, Romano M, Villagrasa M, Negrini S. Physiotherapeutic scoliosis-specific exercises for adolescents with idiopathic scoliosis. *Eur J Phys Rehabil Med* 2014;50:111-21.
22. Weiss H-R, Negrini S, Hawes MC, Rigo M, Kotwicki T, Grivas TB, *et al.* Physical exercises in the treatment of idiopathic scoliosis at risk of brace treatment -- SOSORT consensus paper 2005. *Scoliosis* 2006;1:6.
23. Negrini S, Zaina F, Romano M, Negrini A, Parzini S. Specific exercises reduce brace prescription in adolescent idiopathic scoliosis: a prospective controlled cohort study with worst-case analysis. *J Rehabil Med* 2008;40:451-5.
24. Monticone M, Ambrosini E, Cazzaniga D, Rocca B, Ferrante S. Active self-correction and task-oriented exercises reduce spinal deformity and improve quality of life in subjects with mild adolescent idiopathic scoliosis. Results of a randomised controlled trial. *Eur Spine J* 2014;23:1204-14.
25. Negrini S, Bettany-Saltikov J, De Mauroy JC, Durmala J, Grivas TB, Knott P, *et al.* Letter to the Editor concerning: "Active self-correction and task-oriented exercises reduce spinal deformity and improve quality of life in subjects with mild adolescent idiopathic scoliosis. Results of a randomised controlled trial" by Monticone M, Ambrosini E, Cazzaniga D, Rocca B, Ferrante S (2014). *Eur Spine J*; DOI:10.1007/s00586-014-3241-y. *Eur Spine J* 2014;23:2218-20
26. Wan L, Wang GX, Bian R. [Exercise therapy in treatment of essential S-shaped scoliosis: Evaluation of Cobb angle in breast and lumbar segment through a follow-up of half a year]. *Chin J Clin Rehabil* 2005;9:82-4.
27. Kuru T, Yeldan İ, Dereli EE, Özdiñçler AR, Dikici F, Çolak İ. The efficacy of three-dimensional Schroth exercises in adolescent idiopathic scoliosis: A randomised controlled clinical trial. *Clin Rehabil* 2016;30:181-90.
28. Fusco C, Zaina F, Atanasio S, Romano M, Negrini A, Negrini S. Physical exercises in the treatment of adolescent idiopathic scoliosis: An updated systematic review. *Physiother Theory Pract* 2011;27:80-114.
29. Negrini S, Fusco C, Minozzi S, Atanasio S, Zaina F, Romano M. Exercises reduce the progression rate of adolescent idiopathic scoliosis: results of a comprehensive systematic review of the literature. *Disabil Rehabil* 2008;30:772-85.
30. Negrini S, Antonini G, Carabalona R, Minozzi S. Physical exercises as a treatment for adolescent idiopathic scoliosis. A systematic review. *Pediatr Rehabil* 2003;6:227-35.
31. Bettany-Saltikov J, Weiss H-R, Chockalingam N, Taranu R, Srinivas S, Hogg J, *et al.* Surgical versus non-surgical interventions in people with adolescent idiopathic scoliosis. *Cochrane Database Syst Rev*. 2015;4:CD010663.
32. Weiss H-R. Adolescent idiopathic scoliosis (AIS) - an indication for surgery? A systematic review of the literature. *Disabil Rehabil* 2008;30:799-807.
33. Weiss H-R, Goodall D. Rate of complications in scoliosis surgery - a systematic review of the Pub Med literature. *Scoliosis* 2008;3:9.
34. Stagnara P. Les deformations du rachis. Paris: Masson; 1985.

Conflicts of interest.—The authors certify that there is no conflict of interest with any financial organization regarding the material discussed in the manuscript. Article first published online: July 14, 2016. - Manuscript accepted: July 11, 2016. - Manuscript received: July 7, 2016.

For supplementary materials, please see the online version of this article.


SUPPLEMENTARY MATERIALS

Supplementary Appendix I.—Paper found with the systematic review

- Weinstein SL, Dolan LA, Wright JG, Dobbs MB. Effects of bracing in adolescents with idiopathic scoliosis. *N Engl J Med* 2013;369:1512-21.
- Seidi F, Rajabi R, Ebrahimi I, Alizadeh MH, Minoonejad H. The efficiency of corrective exercise interventions on thoracic hyper-kypnosis angle. *J Back Musculoskelet Rehabil* 2014;27:7-16.
- Romano M, Minozzi S, Zaina F, Saltikov JB, Chockalingam N, Kotwicki T, *et al*. Exercises for adolescent idiopathic scoliosis: a Cochrane systematic review. *Spine (Phila Pa 1976)* 2013;38:E883-93.
- Pinquart M. Body image of children and adolescents with chronic illness: a meta-analytic comparison with healthy peers. *Body Image* 2013;10:141-8.
- Rushton PR, Grevitt MP. Comparison of untreated adolescent idiopathic scoliosis with normal controls: a review and statistical analysis of the literature. *Spine (Phila Pa 1976)* 2013;38:778-85.
- Tis JE, Karlin LI, Akbarina BA, Blakemore LC, Thompson GH, McCarthy RE, *et al*; Growing Spine Committee of the Scoliosis Research Society. Early onset scoliosis: modern treatment and results. *J Pediatr Orthop* 2012;32:647-57.
- Sanders JO, Newton PO, Browne RH, Herring AJ. Bracing in adolescent idiopathic scoliosis, surrogate outcomes, and the number needed to treat. *J Pediatr Orthop* 2012;32 Suppl 2:S153-7.
- Romano M, Minozzi S, Bettany-Saltikov J, Zaina F, Chockalingam N, Kotwicki T, *et al*. Exercises for adolescent idiopathic scoliosis. *Cochrane Database Syst Rev*. 2012 Aug 15;8:CD007837.
- Diab AA. The role of forward head correction in management of adolescent idiopathic scoliotic patients: a randomized controlled trial. *Clin Rehabil* 2012;26:1123-32.
- Miller DJ, Franzone JM, Matsumoto H, Gomez JA, Avendaño J, Hyman JE, *et al*. **Electronic monitoring improves brace-wearing compliance** in patients with adolescent idiopathic scoliosis: a randomized clinical trial. *Spine (Phila Pa 1976)* 2012;37:717-21.
- Negrini S, Aulisa AG, Aulisa L, Circo AB, de Mauroy JC, Durmala J, *et al*. 2011 SOSORT guidelines: Orthopaedic and Rehabilitation treatment of idiopathic scoliosis during growth. *Scoliosis* 2012;7:3.
- Mordecai SC, Dabke HV. Efficacy of exercise therapy for the treatment of adolescent idiopathic scoliosis: a review of the literature. *Eur Spine J* 2012;21:382-9.
- Fusco C, Zaina F, Atanasio S, Romano M, Negrini A, Negrini S. Physical exercises in the treatment of adolescent idiopathic scoliosis: an updated systematic review. *Physiother Theory Pract* 2011;27:80-114.
- Maruyama T, Grivas TB, Kaspiris A. Effectiveness and outcomes of brace treatment: a systematic review. *Physiother Theory Pract* 2011;27:26-42.
- Sponseller PD. Bracing for adolescent idiopathic scoliosis in practice today. *J Pediatr Orthop* 2011;31(1 Suppl):S53-60.
- Negrini S, Minozzi S, Bettany-Saltikov J, Zaina F, Chockalingam N, Grivas TB, *et al*. Braces for idiopathic scoliosis in adolescents. *Spine (Phila Pa 1976)* 2010;35:1285-93.
- Fong DY, Lee CF, Cheung KM, Cheng JC, Ng BK, Lam TP, *et al*. A meta-analysis of the clinical effectiveness of school scoliosis screening. *Spine (Phila Pa 1976)* 2010;35:1061-71.
- Negrini S, Minozzi S, Bettany-Saltikov J, Zaina F, Chockalingam N, Grivas TB, *et al*. Braces for idiopathic scoliosis in adolescents. *Cochrane Database Syst Rev*. 2010 Jan 20;(1):CD006850. doi:10.1002/14651858.CD006850.pub2. Review. Update in: *Cochrane Database Syst Rev*. 2015;6:CD006850.
- Kotwicki T, Durmala J, Czaprowski D, Glowacki M, Kolban M, Snela S, *et al*; SOSORT. Conservative management of idiopathic scoliosis--guidelines based on SOSORT 2006 Consensus. *Ortop Traumatol Rehabil* 2009;11:379-95.
- Klein G, Mehlman CT, McCarty M. Nonoperative treatment of spondylolysis and grade I spondylolisthesis in children and young adults: a meta-analysis of observational studies. *J Pediatr Orthop* 2009;29:146-56.
- Negrini S, Grivas TB, Kotwicki T, Rigo M, Zaina F; international Society on Scoliosis Orthopaedic and Rehabilitation Treatment (SOSORT). Guidelines on "Standards of management of idiopathic scoliosis with corrective braces in everyday clinics and in clinical research": SOSORT Consensus 2008. *Scoliosis* 2009;4:2.
- Li XF, Li H, Liu ZD, Dai LY. Low bone mineral status in adolescent idiopathic scoliosis. *Eur Spine J* 2008;17:1431-40.
- Wong MS, Cheng JC, Lam TP, Ng BK, Sin SW, Lee-Shum SL, *et al*. The effect of rigid versus flexible spinal orthosis on the clinical efficacy and acceptance of the patients with adolescent idiopathic scoliosis. *Spine (Phila Pa 1976)* 2008;33:1360-5.
- Weiss HR. Adolescent idiopathic scoliosis (AIS) - an indication for surgery? A systematic review of the literature. *Disabil Rehabil* 2008;30:799-807.
- Maruyama T. Bracing adolescent idiopathic scoliosis: a systematic review of the literature of effective conservative treatment looking for end results 5 years after weaning. *Disabil Rehabil* 2008;30:786-91.
- Negrini S, Fusco C, Minozzi S, Atanasio S, Zaina F, Romano M. Exercises reduce the progression rate of adolescent idiopathic scoliosis: results of a comprehensive systematic review of the literature. *Disabil Rehabil* 2008;30:772-85.
- Bunge EM, de Koning HJ; brace trial group. Bracing patients with idiopathic scoliosis: design of the Dutch randomized controlled treatment trial. *BMC Musculoskelet Disord* 2008;9:57.
- Weiss HR, Goodall D. The treatment of adolescent idiopathic scoliosis (AIS) according to present evidence. A systematic review. *Eur J Phys Rehabil Med* 2008;44:177-93.
- Dolan LA, Weinstein SL. Surgical rates after observation and bracing for adolescent idiopathic scoliosis: an evidence-based review. *Spine (Phila Pa 1976)* 2007;32(19 Suppl):S91-S100.
- Lowe TG, Line BG. Evidence based medicine: analysis of Scheuermann kyphosis. *Spine (Phila Pa 1976)* 2007;32(19 Suppl):S115-9.
- Wong MS, Cheng CY, Ng BK, Lam TP, Sin SW, Lee-Shum LF, *et al*. The effect of rigid versus flexible spinal orthosis on the gait pattern of patients with adolescent idiopathic scoliosis. *Gait Posture* 2008;27:189-95.
- Weiss HR, Negrini S, Rigo M, Kotwicki T, Hawes MC, Grivas TB, *et al*; (SOSORT guideline committee). Indications for conservative management of scoliosis (guidelines). *Scoliosis* 2006 May 8;1:5.
- Lensinck ML, Frijlink AC, Berger MY, Bierman-Zeinstra SM, Verkerk K, Verhagen AP. Effect of bracing and other conservative interventions in the treatment of idiopathic scoliosis in adolescents: a systematic review of clinical trials. *Phys Ther* 2005;85:1329-39.
- Negrini S, Aulisa L, Ferraro C, Frascini P, Masiero S, Simonazzi P, *et al*. Italian guidelines on rehabilitation treatment of adolescents with scoliosis or other spinal deformities. *Eura Medicophys* 2005;41:183-201.
- Richards BS, Bernstein RM, D'Amato CR, Thompson GH. Standardization of criteria for adolescent idiopathic scoliosis brace studies: SRS Committee on Bracing and Nonoperative Management. *Spine (Phila Pa 1976)* 2005;30:2068-75; discussion 2076-7.
- Weiss HR, Heckel I, Stephan C. Application of passive transverse forces in the rehabilitation of spinal deformities: a randomized controlled study. *Stud Health Technol Inform* 2002;88:304-8.
- Negrini S, Antonini G, Carabalona R, Minozzi S. Physical exercises as a treatment for adolescent idiopathic scoliosis. A systematic review. *Pediatr Rehabil* 2003;6:227-35.
- Hawes MC. The use of exercises in the treatment of scoliosis: an

- evidence-based critical review of the literature. *Pediatr Rehabil* 2003;6:171-82.
39. Athanasopoulos S, Paxinos T, Tsafantakis E, Zachariou K, Chatziconstantinou S. The effect of aerobic training in girls with idiopathic scoliosis. *Scand J Med Sci Sports* 1999;9:36-40.
40. Rowe DE, Bernstein SM, Riddick MF, Adler F, Emans JB, Gardner-Bonneau D. A meta-analysis of the efficacy of non-operative treat-

- ments for idiopathic scoliosis. *J Bone Joint Surg Am* 1997;79:664-74.
41. Liptak GS. Tethered spinal cord: update of an analysis of published articles. *Eur J Pediatr Surg* 1995;5 Suppl 1:21-3.
42. Focarile FA, Bonaldi A, Giarolo MA, Ferrari U, Zilioli E, Ottaviani C. Effectiveness of nonsurgical treatment for idiopathic scoliosis. Overview of available evidence. *Spine (Phila Pa 1976)* 1991;16:395-401.



Supplementary Figure 1.—Flow Chart of papers selection.

SUPPLEMENTARY TABLE I.—Results of the Consensus procedure.

Round	Number of recommendations	Accept	Accept with suggestions	Reject
1	26	57.7%	42.3%	0
2	26	92.3%	7.7%	0
3	26	100%	0	0
4	26	100%	0	0

SUPPLEMENTARY TABLE II.—Overall view of the recommendations.

Content	Number of recommendations		Strength of Recommendations				Strength of Evidence			
	Number		A	B	C	D	I	II	III	IV
Overall recommendation	1		100%	0	0	0	0	0	0	100%
PRM physicians' role in medical diagnosis - ICD	5		60%	40%	0	0	0	0	0	100%
PRM diagnosis and assessment- ICF	4		75%	25%	0	0	0	0	0	100%
PRM management and process	13		61.5%	38.5%	0	0	0	0	0	100%
Future research on PRM professional practice	3		0	100%	0	0	0	0	0	100%
Total	26		57.7%	42.3%	0	0	0	0	0	100%

This document is protected by international copyright laws. No additional reproduction is authorized. It is permitted for personal use to download and save only one file and print only one copy of this Article. It is not permitted to make additional copies (either sporadically or systematically, either printed or electronic) of the Article for any purpose. It is not permitted to distribute the electronic copy of the article through online internet and/or intranet file sharing systems, electronic mailing or any other means which may allow access to the Article. The use of all or any part of the Article for any Commercial Use is not permitted. The production of derivative works from the Article is not permitted. The production of reprints for personal or commercial use is not permitted. It is not permitted to remove, cover, overlay, obscure, block, or change any copyright notices or terms of use which the Publisher may post on the Article. It is not permitted to frame or use framing techniques to enclose any trademark, logo or other proprietary information of the Publisher.