



**21ST INTERNATIONAL SYMPOSIUM  
ON  
HOMOGENEOUS CATALYSIS**

July 8th - 13th, 2018

Beurs van Berlage, Amsterdam, The Netherlands

<http://www.ishc21.amsterdam/>



# ISHC 2018

Book of Abstracts

## O16 Palladium-Catalyzed Oxidative Carbonylation of Aniline: the Effect of Iron Cocatalysts


by Fabio Ragaini, Edoardo Barraco, Claudia Gatti, Francesco Ferrettia | University of Milan, Dipartimento di Chimica, Università degli Studi di Milano, via Golgi 19, 20133 Milano (Italy)

Isocyanates are key intermediates for the chemical industry, but their production is actually based on the use of toxic phosgene (path A in the Figure). Among suitable alternatives, the reductive carbonylation of nitroarenes (Path B) and the oxidative carbonylation of amines (Path C) are the most investigated.

Research in our group mostly focused on the reductive carbonylation of nitroarenes, but the connections we could identify between this reaction and the oxidative carbonylation of amines<sup>1</sup> recently pushed us to investigate the latter reaction. The most active catalytic system reported to date is based on palladium catalysts with iodide as promoter<sup>2</sup>. We decided to investigate this system in more detail and found some surprising results. The most important new findings are:

- When the reaction is performed directly in a steel autoclave, the reported data are fully reproducible. However, if the reaction is performed in a glass liner or a Teflon lined autoclave, the activity of the catalytic system drops to about 1/3 of the original one or lesser.
- One essential role of iodide is to etch some iron from the autoclave walls. The high activity can be restored even in a Teflon coated autoclave if small amounts of iron salts are added.
- Iodide and iron have complex interactions and can even deactivate each other when their molar ratio falls within a certain range.
- If iron is present, iodide is no longer necessary, but a halide (chloride) is anyway required.
- The solvent is not completely innocent in the reaction.

These and other trends will be discussed.



Alternative synthetic procedure for aromatic isocyanates

**References:** [1] Ragaini, F. *Dalton Trans.* **2009**, 6251; [2] a) Fukuoka, S.; Chono, M.; Kohno, M. *J. Chem. Soc., Chem. Commun.* **1984**, 399; b) Gabriele, B.; Salerno, G.; Mancuso, R.; Costa, M. *J. Org. Chem.* **2004**, 69, 4741.