

KdV and FPU: asymptotic integrability of quasi unidirectional waves

Matteo Gallone*, Antonio Ponno† and Bob Rink‡

October 7, 2020

Abstract

In this paper we construct a higher order expansion of the manifold of quasi unidirectional waves in the Fermi-Pasta-Ulam (FPU) chain. We also approximate the dynamics on this manifold. As perturbation parameter we use $h^2 = 1/n^2$, where n is the number of particles of the chain. It is well known that the dynamics of quasi unidirectional waves is described to first order by the Korteweg-de Vries (KdV) equation. Here we show that the dynamics to second order is governed by a combination of the first two nontrivial equations in the KdV hierarchy – for any choice of parameters in the FPU potential. On the other hand, we find that only if the parameters of the FPU potential satisfy a condition, then a combination of the first three nontrivial equations in the KdV hierarchy determines the dynamics of quasi unidirectional waves to third order. The required condition is satisfied by the Toda chain. Our results suggest why the close-to-integrable behavior of the FPU chain (the FPU paradox) persists on a time scale longer than explained by the KdV approximation, and also how a breakdown of integrability (detachment from the KdV hierarchy) is responsible for the eventual thermalization of the system.

1 Introduction

In the early 1950s, Fermi, Pasta and Ulam (FPU) set up a series of numerical experiments, with the purpose to measure the time of approach to statistical equilibrium in non-integrable Hamiltonian systems. Their experiments were motivated by the conviction of Fermi that generic, large size Hamiltonian systems are ergodic. FPU studied a simple family of models: chains of particles interacting with their nearest neighbors through a force of simple polynomial type. The unexpected outcome of their numerical study [15], namely the observed lack of ergodicity and mixing over the computational time then available, was named after them – the FPU problem, or FPU paradox. Although a complete mathematical understanding of the FPU problem is still lacking, some deep insights into specific features have been obtained. It is not the aim of this paper to review the history of the FPU problem; the interested reader is referred to the existing reviews [2, 11, 12, 16, 27].

Nowadays, we know that the FPU paradox is a matter of quasi-integrability. By this we mean that the dynamics of the FPU chain over short timescales resembles that of a closely integrable system, while the approach to statistical equilibrium, in the long run, is due to the perturbation. Which integrable system should be considered “closest” to the FPU chain, depends both on the model and on initial conditions. In this paper, we focus on the generic FPU chain, or FPU α -model, in which the inter-particle forces display quadratic nonlinearities to leading order – see equation (2) below. For this model, the first explanation of the FPU paradox in terms of integrability goes back to the pioneering work of Zabusky and Kruskal [30]. These authors show that the dynamics in the chain can be described, for short times and for smooth low energy initial conditions close to a unidirectional wave, by the Korteweg-de Vries (KdV) equation. This partial differential equation was later proved to be integrable [18, 22, 24, 31]. The approach in [30] was later enforced by Zakharov [32], who proved the integrability of the quadratic Boussinesq equation, which can be regarded a continuum approximation of the full FPU system. On the other hand, it is also well understood now that the low energy short term dynamics of the FPU chain follows that of the integrable Toda chain, for *any* initial condition (even extracted in measure), as first pointed out in [14, 23] and more

*Università degli Studi di Milano, Dipartimento di Matematica “F. Enriques”, Via Saldini 50, 20133 Milano, Italy; matteo.gallone@unimi.it

†Università degli Studi di Padova, Dipartimento di Matematica “T. Levi-Civita”, Via Trieste 63, 35121 Padova, Italy; ponno@math.unipd.it

‡Vrije Universiteit Amsterdam, Department of Mathematics, De Boelelaan 1111, 1081 HV Amsterdam, The Netherlands; b.w.rink@vu.nl

recently in [6, 8, 9, 10, 19]. Of course, when smooth initial data are considered, the two points of view connect, since the continuum limit of the Toda chain consists of two KdV equations [3].

It was shown in recent studies [5, 7, 25] that the approximate description of the FPU dynamics in terms of PDEs can be cast within the theory of normal forms. In particular it was proved that, in a certain regime of smooth low energy initial conditions, the leading order resonant normal form of the FPU system consists of two KdV equations – one describing almost-right traveling waves and the other almost-left traveling waves [5]. This result agrees with the perspective sketched above: the dynamics of the FPU chain is integrable in the short term. On the other hand, the numerically observed FPU energy spectrum turns out to be stable over time scales much longer than that of the validity of the KdV approximation. This in turn suggests that the normal form of the FPU system may actually be integrable even beyond the leading order KdV approximation.

We partially investigate this issue in the present paper. In fact, we will consider the FPU chain with periodic boundary conditions, and we will introduce smooth functions interpolating the positions of the particles at any time. As small parameter we choose $h^2 := 1/n^2$, where n is the number of particles in the chain. In a forthcoming paper [17] we will consider this problem for arbitrary smooth initial data. In this paper, on the other hand, we shall investigate solutions that lie inside an invariant sub-manifold consisting of quasi unidirectional waves. We construct this invariant manifold to high perturbative orders. This method is among the standard techniques for finding invariant manifolds in hydrodynamics [28], and was first introduced by Whitham [29], and extended e.g. in [13], to derive the KdV equation in the theory of shallow water waves.

The evolution equation that we initially find for the dynamics of quasi unidirectional waves, is not integrable beyond the KdV equation found at order h^2 . To investigate the asymptotic integrability of the equation in more detail, we shall apply normal form transformations. The type of transformations that we consider were introduced by Hiraoka and Kodama [20]; in this paper we adapt them for continuous systems with periodic boundary conditions. Our results are the following. First of all, we find that to order h^4 the dynamics of quasi unidirectional waves is *always* governed by an integrable equation from the KdV hierarchy. In contrast, to order h^6 the dynamics is governed by a member of the KdV hierarchy only for particular values of the parameters defining the nearest-neighbor interaction in the chain. The Toda chain is an example of such a system that to order h^6 possesses an integrable normal form. Thus,

*the dynamics of quasi unidirectional waves in generic FPU chains is
integrable over the timescale corresponding to the second order normal form;
the breakdown of integrability generally takes place at third order.*

This agrees, qualitatively, with what is observed in numerical simulations. We also point out the recent work [1], in which a somewhat similar result was proven in the theory of shallow water waves.

The remainder of this paper is organised as follows. In Section 2 we introduce the FPU chain, as well as an exact continuous system interpolating it. In this section we also informally present the main result of this paper as Theorem 1. In Section 3 we provide higher order asymptotic expansions for the interpolating system, in the form of partial differential equations (PDEs). In Section 4 we construct and approximate the manifold of quasi unidirectional waves, and provide an asymptotic expansion of the dynamics on this manifold, see Theorem 2. In Section 5 we prove Theorem 3, which provides the normal form for the dynamics on the manifold of quasi unidirectional waves. Theorem 3 is a direct consequence of Theorem 4, which is a spatially periodic version of a theorem by Hiraoka and Kodama. We prove Theorem 4 in Section 6.

2 Formulation of the main result

In this section we give a brief description of the main result of this paper. We start by recalling that the periodic FPU chain with n particles is the n -degrees of freedom Hamiltonian dynamical system with equations of motion

$$\begin{cases} \frac{dq_j}{dt} = p_j, \\ \frac{dp_j}{dt} = W'(q_{j+1} - q_j) - W'(q_j - q_{j-1}), \end{cases} \quad (1)$$

satisfying the periodic boundary conditions

$$q_{j+n} = q_j, \quad p_{j+n} = p_j \quad \text{for all } j \in \mathbb{Z}.$$

The function W in (1) is a potential energy determining the interaction between neighboring particles in the chain. As usual, we assume that it admits the Taylor expansion

$$W(z) = \frac{1}{2}z^2 + \frac{\alpha}{3}z^3 + \frac{\beta}{4}z^4 + \frac{\gamma}{5}z^5 + \mathcal{O}(z^6). \quad (2)$$

The non-linearity in the forces between neighboring particles is thus determined by the parameters α, β, γ , etc. We assume $\alpha \neq 0$ throughout this paper. As discussed above, a special role is played by the Toda chain (see also Section 5), where the potential is taken from a specific one-parameter family, satisfying

$$\beta = \beta_T := \frac{2}{3}\alpha^2, \quad \gamma = \gamma_T := \frac{1}{3}\alpha^3, \quad \text{etc.}$$

To transform the FPU equations of motion (1) into a continuous system of PDEs, we first replace (1) by an exact evolution equation for an interpolating profile. To this end, let us denote $h := 1/n$ and write $\mathbb{T} := \mathbb{R}/\mathbb{Z}$. Now consider a pair of smooth scalar functions

$$(u, v) = (u, v)(x, t) : \mathbb{T} \times \mathbb{R} \rightarrow \mathbb{R}^2$$

and assume that these functions satisfy the evolution equations

$$\begin{cases} u_t(x, t) = v(x, t), \\ v_t(x, t) = h^{-3} [W'(hu(x+h, t) - hu(x, t)) - W'(hu(x, t) - hu(x-h, t))]. \end{cases} \quad (3)$$

We informally think of (3) as an ODE on the space $C^\infty(\mathbb{T}, \mathbb{R}^2)$ of pairs of smooth scalar functions on \mathbb{T} . We correspondingly think of solutions to (3) as curves $t \mapsto (u, v)(\cdot)(t)$ in $C^\infty(\mathbb{T}, \mathbb{R}^2)$.

Given any solution of (3), one can define, for $j \in \mathbb{Z}$, the functions

$$\begin{cases} q_j(t) := hu(hj, ht), \\ p_j(t) := h^2 v(hj, ht). \end{cases} \quad (4)$$

It is not hard to check that these $q_j(t), p_j(t)$ then form solutions of the FPU equations of motion (1). This motivates us to study (3) instead of (1), the advantage being that (3) is defined on the same phase space for each value of h ; in fact, we may simply think of h as a continuous small parameter.

As long as u is a smooth function of x , we can Taylor expand $u(x \pm h) = u(x) \pm hu_x(x) + \frac{h^2}{2}u_{xx}(x) + \dots$, and write (3) as the perturbed Boussinesq equation

$$\begin{cases} u_t = v, \\ v_t = u_{xx} + h^2 \left(\frac{1}{12}u_{xxxx} + 2\alpha u_x u_{xx} \right) + \mathcal{O}(h^4). \end{cases} \quad (5)$$

Next, instead of studying (3) or (5) directly, we introduce a change of variables $(u, v) \mapsto (U, V)$ in $C^\infty(\mathbb{T}, \mathbb{R}^2)$, that maps the functions $(u, v)(x)$ to the *discrete Riemann invariants* $(U, V)(x)$ defined by

$$U := 2\alpha(D_h u + v), \quad V := 2\alpha(D_h u - v). \quad (6)$$

Here D_h denotes the finite difference operator

$$(D_h u)(x) := \frac{u(x+h/2) - u(x-h/2)}{h}. \quad (7)$$

This D_h is a discrete approximation of the derivative ∂_x , because $D_h u = u_x + \frac{h^2}{24}u_{3x} + \mathcal{O}(h^5)$ if $u : \mathbb{T} \rightarrow \mathbb{R}$ is sufficiently smooth - again by Taylor's theorem. The change of variables $(u, v) \mapsto (U, V)$ transforms the continuum equations (3) into a system of the form

$$\begin{cases} U_t = \mathcal{F}(U, V, h), \\ V_t = -\mathcal{F}(V, U, h). \end{cases} \quad (8)$$

We again think of (8) as an ODE on $C^\infty(\mathbb{T}, \mathbb{R}^2)$. An exact expression for $\mathcal{F}(U, V, h)$ will be given in Section 3. Here it suffices to know that $\mathcal{F}(U, V, h)$ admits an expansion

$$\mathcal{F}(U, V, h) = U_x + \frac{h^2}{24}(U_{xxx} + 6UU_x + 6UV_x + 6U_xV + 6VV_x) + \mathcal{O}(h^4),$$

as long as (U, V) are sufficiently smooth functions of x .

By setting $V \equiv 0$ in the evolution equation for U in (8) we recover the KdV equation

$$U_t = \mathcal{F}(U, 0, h) = U_x + \frac{h^2}{24}(U_{xxx} + 6UU_x) + \mathcal{O}(h^4).$$

Nevertheless, it should be clear that if $V(x) \equiv 0$ while $U(x) \not\equiv 0$, then

$$V_t = -\mathcal{F}(V, U, h) = -\mathcal{F}(0, U, h) = -\frac{1}{4}h^2UU_x \neq 0.$$

This means that, already to order h^2 , the subspace

$$\{(U, V) \in C^\infty(\mathbb{T}, \mathbb{R}^2) \mid V \equiv 0\} \subset C^\infty(\mathbb{T}, \mathbb{R}^2)$$

of “unidirectional waves” is not invariant under the flow of (8). This may cast some doubt on the validity of the above derivation of the KdV approximation, but we will show that nearby the subspace of unidirectional waves one can formally construct a submanifold of quasi unidirectional waves of the form

$$\{(U, V) \in C^\infty(\mathbb{T}, \mathbb{R}^2) \mid V = c(U, h) = \mathcal{O}(h^2)\} \subset C^\infty(\mathbb{T}, \mathbb{R}^2).$$

This submanifold is invariant under (8) to high asymptotic orders. Moreover, the dynamics on the manifold of quasi unidirectional waves is indeed governed to order h^2 by the KdV equation.

More interestingly, we shall also derive higher order equations for the evolution of quasi unidirectional waves. It turns out that, after a suitable change of variables, this evolution is determined to a large extent by the “higher order versions” of the KdV equation, i.e., by the integrable KdV hierarchy. More precisely, the main results of this paper are Theorems 2 and 3, which can be summarised as follows.

Theorem 1. *Inside $C^\infty(\mathbb{T}, \mathbb{R}^2)$ there is a formal invariant manifold for the dynamics (8) of the discrete Riemann invariants, consisting of quasi unidirectional waves, and defined by a slaving relation of the form*

$$V = c(U, h) = h^2 c_2(U) + h^4 c_4(U) + \mathcal{O}(h^6).$$

There also exists a formal near-identity transformation in $C^\infty(\mathbb{T}, \mathbb{R})$ of the form

$$U \mapsto U + h^2 \tilde{G}_2(U) + h^4 \tilde{G}_4(U) + h^6 \tilde{G}_6(U, t) + \mathcal{O}(h^8),$$

bringing the dynamics on the manifold of quasi unidirectional waves into the form

$$\begin{aligned} U_t &= \mathcal{C}_1(U, h) K_1(U) + \\ &+ h^2 \mathcal{C}_3(U, h) K_3(U) + \\ &+ h^4 \mathcal{C}_5(U, h) K_5(U) + \\ &+ h^6 \mathcal{C}_7(U, h) [K_7(U) + R(U)] + \mathcal{O}(h^8). \end{aligned} \quad (9)$$

Here, $K_1(U), K_3(U), K_5(U), K_7(U)$ are the first four commuting vector fields in the KdV hierarchy, and the scalars $\mathcal{C}_1(U, h), \mathcal{C}_3(U, h), \mathcal{C}_5(U, h), \mathcal{C}_7(U, h)$ are constants of motion of the KdV hierarchy (see also Remark 1). The term $R(U)$ can be chosen equal to zero in case

$$14\alpha^3 - 27\beta\alpha + 12\gamma = 0. \quad (10)$$

Theorem 1 shows that the evolution of quasi unidirectional waves is always integrable to order h^4 , and is integrable to order h^6 if the relation (10) holds. This relation holds in particular when $\beta = \beta_T = 2\alpha^2/3$ and $\gamma = \gamma_T = \alpha^3/3$. The one-parameter family of Toda chains is thus part of a co-dimension one family of FPU chains whose dynamics is “more integrable” than that of generic FPU chains.

Remark 1. *The first four commuting vector fields in the KdV hierarchy are given by*

$$\begin{aligned} K_1(U) &= U_x, \\ K_3(U) &= U_{3x} + 6UU_x, \\ K_5(U) &= U_{5x} + 20U_x U_{2x} + 10UU_{3x} + 30U^2 U_x, \\ K_7(U) &= U_{7x} + 70U_{2x} U_{3x} + 42U_x U_{4x} + 14UU_{5x} + \\ &+ 70U_x^3 + 280UU_x U_{2x} + 70U^2 U_{3x} + 140U^3 U_x. \end{aligned} \quad (11)$$

See for instance [24]. We used the short-hand notation $U_{mx} := \partial_{mx} U = \partial_x^m U$ for the m -derivative of U .

The constants of motion $\mathcal{C}_1(U, h), \mathcal{C}_3(U, h), \mathcal{C}_5(U, h), \mathcal{C}_7(U, h)$ in Theorem 3 turn out to be functions of the first three integrals of the KdV hierarchy,

$$\int_{\mathbb{T}} U dx, \quad \int_{\mathbb{T}} U^2 dx \quad \text{and} \quad \int_{\mathbb{T}} U_x^2 - 2U^3 dx. \quad (12)$$

Remark 2. *If (4) holds for smooth functions $u, v : \mathbb{T} \rightarrow \mathbb{R}$, then the total energy*

$$E = \sum_{j=1}^n \frac{1}{2} p_j^2 + W(q_{j+1} - q_j)$$

of the FPU system (1) satisfies

$$E = h^3 \int_{\mathbb{T}} \frac{1}{2}v(x)^2 + \frac{1}{2}u_x(x)^2 dx + \mathcal{O}(h^4).$$

In turn this implies that the “specific energy” (the energy per particle) $\frac{1}{n}E$ is of the order $h^4 = 1/n^4$. This is far from the thermodynamic limit in which $\frac{1}{n}E$ would be of order 1. However, (4) is the unique scaling limit in which dispersion and nonlinearity are of the same order, thus leading at lowest order to a KdV equation for the evolution of quasi unidirectional waves. It is also the scaling limit in which the Toda chain can be smoothly connected to the KdV equation [3] and topologically to the harmonic chain [4].

3 PDE approximations of the FPU dynamics

In this section we provide more details on the derivation of the evolution equations (8) for the discrete Riemann invariants (U, V) defined in (6). Our starting point is the evolution equation for the interpolating profiles (u, v) given in (3):

$$\begin{cases} u_t(x, t) = v(x, t), \\ v_t(x, t) = h^{-3} [W'(hu(x+h, t) - hu(x, t)) - W'(hu(x, t) - hu(x-h, t))] . \end{cases}$$

Assuming that $u(x, t)$ and $v(x, t)$ are smooth functions of x , one may Taylor expand $u(x \pm h, t)$ with respect to h in the right hand side of (3). This yields a perturbed nonlinear wave equation which we present here to high order:

$$\begin{cases} u_t = v, \\ v_t = u_{xx} + h^2 \left(\frac{1}{12}u_{4x} + 2\alpha u_x u_{xx} \right) + h^4 \left(\frac{1}{360}u_{6x} + \frac{\alpha}{3}u_{xx}u_{3x} + \frac{\alpha}{6}u_x u_{4x} + 3\beta u_x^2 u_{xx} \right) \\ + h^6 \left(\frac{1}{20160}u_{8x} + \frac{\alpha}{36}u_{3x}u_{4x} + \frac{\alpha}{60}u_{2x}u_{5x} + \frac{\alpha}{180}u_x u_{6x} + \right. \\ \left. + \frac{\beta}{4}u_{2x}^3 + \beta u_x u_{xx} u_{3x} + \frac{\beta}{4}u_x^2 u_{4x} + 4\gamma u_x^3 u_{2x} \right) + h^8 R(u, h) . \end{cases} \quad (13)$$

Note however that the assumption that u and v are smooth has no a priori justification: it is not guaranteed that (3) admits nontrivial initial data leading to solutions that remain smooth over long times. In contrast to this, we recall here that the dynamics of the KdV equation (and of its hierarchy) on \mathbb{T} preserves the Sobolev smoothness and analyticity of initial data for all times [21].

Remark 3. Equation (13) reveals that the FPU equations (3) may be thought of as a weakly dispersive and weakly nonlinear perturbation of the wave equation. The perturbation to order h^2 is the so-called Boussinesq equation. This equation was proven to be integrable by Zakharov [32] in order to explain the FPU paradox.

Remark 4. Using Taylor’s theorem, one may determine explicit bounds for the remainder $h^8 R(u, h)$ in terms of u . For example, one may easily get a conditional estimate of the form: for every integer $k \geq 0$, there exists a constant C_k independent of u and h , such that

$$\|R(u, h)\|_{C^k(\mathbb{T})} \leq C_k \|u\|_{C^{k+10}(\mathbb{T})}$$

as long as $\|u\|_{C^0(\mathbb{T})} < 1$ and $|h| < 1$. Similar estimates can be obtained for Sobolev norms. We will not pursue such explicit bounds any further. Instead, from now on we shall simply write $\mathcal{O}(h^m)$ for expressions of the form $h^m R(u, h)$.

For the exact wave equation $u_t = v$, $v_t = u_{xx}$ it is common to make the change of variables $(u, v) \mapsto (U, V)$ defined by $U := 2a(u_x + v)$ and $V := 2a(u_x - v)$, where $a \neq 0$ is any constant. These functions U and V are called *Riemann invariants*, and their evolution is determined by the equations of motion

$$\begin{cases} U_t = U_x, \\ V_t = -V_x. \end{cases}$$

of which the solutions

$$U(x, t) = U_0(x+t), \quad V(x, t) = V_0(x-t)$$

are unidirectional traveling waves. This motivates our definition (6) of the discrete Riemann invariants for (3), with the particular choice $a = \alpha$ and u_x replaced by $D_h u$:

$$U := 2\alpha(D_h u + v), \quad V := 2\alpha(D_h u - v).$$

We stress that our definition of the discrete Riemann invariants – choosing $a = \alpha$ and using $D_h u$ instead of u_x – is such that the structure of their evolution equations is relatively simple. To see this, note that (3) can also be written as

$$\begin{cases} u_t = v, \\ v_t = h^{-2} D_h W'(h^2 D_h u). \end{cases} \quad (14)$$

Remark 5. *This follows from the identity*

$$F(G(x+h) - G(x)) - F(G(x) - G(x-h)) = h D_h F(h D_h G(x)), \quad (15)$$

which holds for any pair of functions F and G , applied to $F = W'$ and $G = u$.

Taking the time derivative of U and V as defined in (6), and using (14), we find

$$\begin{cases} U_t = \mathcal{F}(U, V, h) := D_h [U + f(U + V, h)], \\ V_t = -\mathcal{F}(V, U, h) := -D_h [V + f(U + V, h)]. \end{cases} \quad (16)$$

Here

$$f(z, h) := 2\alpha h^{-2} \left[W' \left(\frac{h^2 z}{4\alpha} \right) - \frac{h^2 z}{4\alpha} \right] = \frac{h^2}{8} z^2 + \frac{\beta h^4}{32\alpha^2} z^3 + \frac{\gamma h^6}{128\alpha^3} z^4 + \mathcal{O}(h^8) \quad (17)$$

is the (rescaled) nonlinear part of the inter-particle force. Using that

$$D_h = \partial_x + \frac{h^2}{24} \partial_{3x} + \frac{h^4}{1920} \partial_{5x} + \frac{h^6}{322560} \partial_{7x} + \mathcal{O}(h^8), \quad (18)$$

one may expand $\mathcal{F}(U, V, h)$ in (16) to high order. This gives (when U, V are smooth):

$$\begin{aligned} U_t = \mathcal{F}(U, V, h) &= U_x + h^2 \left(\frac{1}{24} U_{3x} + \frac{1}{8} (U + V)_x^2 \right) \\ &+ h^4 \left(\frac{1}{1920} U_{5x} + \frac{1}{192} (U + V)_{3x}^2 + \frac{\beta}{32\alpha^2} (U + V)_x^3 \right) \\ &+ h^6 \left(\frac{1}{322560} U_{7x} + \frac{1}{15360} (U + V)_{5x}^2 + \frac{\beta}{768\alpha^2} (U + V)_{3x}^3 + \frac{\gamma}{128\alpha^3} (U + V)_x^4 \right) \\ &+ \mathcal{O}(h^8). \end{aligned} \quad (19)$$

Remark 6. *Here and in the sequel, it is understood that*

$$F_{m,x}^n := (F^n)_{m,x}$$

is short hand notation for the m -th derivative of the n -th power of the function F (and not the n -th power of the m -th derivative).

Remark 7. *Note that the evolution equation for V in (16) can be obtained from that for U in (16) by exchanging the roles of U and V and adding a minus sign. This expresses the fact that (16) possesses the time reversal symmetry*

$$(U, V, t) \mapsto (V, U, -t),$$

corresponding in turn to the time reversal symmetry

$$(u, v, t) \mapsto (u, -v, -t)$$

of the original continuous FPU system (3). It is therefore not restrictive, in what follows, to focus our analysis on the evolution of U , since analogous results automatically hold for V .

4 Quasi unidirectional waves

The subspace

$$\{(U, V) \in C^\infty(\mathbb{T}, \mathbb{R}^2) \mid V \equiv 0\} \subset C^\infty(\mathbb{T}, \mathbb{R}^2)$$

is invariant under the evolution of the wave equation

$$U_t = U_x, \quad V_t = -V_x,$$

and consists entirely of unidirectional traveling waves

$$U(x, t) = U_0(x + t), \quad V(x, t) = 0.$$

The evolution of the discrete Riemann invariants of the FPU chain is accurately described by the wave equation to order h^0 because $U_t = \mathcal{F}(U, V, h) = U_x + \mathcal{O}(h^2)$ and $V_t = -\mathcal{F}(V, U, h) = -V_x + \mathcal{O}(h^2)$. Nevertheless, the subspace of unidirectional waves is already not invariant anymore to order h^2 – see Section 2.

It is therefore quite natural to search for an (at least asymptotically) invariant manifold for the dynamics of (16) lying close to the subspace of unidirectional waves. We will try to find such a submanifold in the form of the graph of a function $c = c(U, h) = \mathcal{O}(h^2)$ over the subspace of unidirectional waves:

$$\{(U, V) \in C^\infty(\mathbb{T}, \mathbb{R}^2) \mid V = c(U, h) = \mathcal{O}(h^2)\}.$$

We think of this graph $\{(U, V) \mid V = c(U, h)\}$ as a manifold of quasi unidirectional waves.

Remark 8. *By Definition (6) we have that $V = \alpha(u_x - v)$. The assumption that $V = \mathcal{O}(h^2)$ thus means that $u_x = v + \mathcal{O}(h^2)$. This in turn implies, by the equations of motion (3), that $u_t = u_x + \mathcal{O}(h^2)$ and $v_t = v_x + \mathcal{O}(h^2)$. Thus we recover an approximate left traveling wave in the original variables.*

Inserting the ansatz

$$V = c(U, h) \tag{20}$$

into system (16) yields the following two equations:

$$\begin{cases} U_t = \mathcal{F}(U, c(U, h), h), \\ c(U, h)_t = -\mathcal{F}(c(U, h), U, h). \end{cases} \tag{21}$$

The first of these equations states that solutions $(U, V) = (U, c(U, h))$ inside the manifold of quasi unidirectional waves are governed by a closed evolution equation

$$U_t = \mathcal{F}(U, h) := \mathcal{F}(U, c(U, h), h). \tag{22}$$

Once a solution U of (22) has been found, the solution V is completely determined by (20). We therefore think of the functional relation (20) between U and V as a slaving relation: the free variable U completely determines the behaviour of the slave variable V .

Using the chain rule at its left hand side, the second equation in (21) can be rewritten as

$$c'(U, h)\mathcal{F}(U, c(U, h), h) = -\mathcal{F}(c(U, h), U, h). \tag{23}$$

This equation can be thought of as an invariance equation. It states that a certain relation must hold between the U -component and the V -component of the vector field in (16) restricted to the manifold of quasi unidirectional waves. This relation guarantees the manifold to be invariant.

Remark 9. *We denote by $F'(U)$ the Gateaux derivative (or directional derivative) of an operator F at $U \in C^\infty(\mathbb{T}, \mathbb{R})$. In other words, given a smooth increment function $H \in C^\infty(\mathbb{T}, \mathbb{R})$, we have*

$$F'(U)H := \lim_{\varepsilon \rightarrow 0} \frac{F(U + \varepsilon H) - F(U)}{\varepsilon}.$$

The operators $F = F(U)$ that we consider in this paper generally are maps from $C^\infty(\mathbb{T}, \mathbb{R})$ to $C^\infty(\mathbb{T}, \mathbb{R})$ and may depend for example on U, U_x, U_{xx} , etc. but also on certain averages or even primitives of U .

The following result gives a formula for $c(U, h)$ to order h^4 , and for the vector field $\mathcal{F}(U, h)$ to order h^6 . We use the notation $\langle F \rangle := \int_{\mathbb{T}} F(x) dx$ for $F = F(x) : \mathbb{T} \rightarrow \mathbb{R}$, see also Remark 10.

Theorem 2. *A formal invariant manifold of quasi unidirectional waves, defined by the invariance equation (23), is given to order h^4 as the graph of the function*

$$\begin{aligned} c(U, h) := & h^2 \left\{ \frac{1}{16} (\langle U^2 \rangle - U^2) \right\} + \\ & + h^4 \left\{ \left(\frac{5}{384} - \frac{\beta}{64\alpha^2} \right) (U^3 - \langle U^3 \rangle) - \frac{1}{128} \langle U^2 \rangle (U - \langle U \rangle) - \frac{1}{256} [(U_x)^2 - \langle (U_x)^2 \rangle] \right\} + \\ & + \mathcal{O}(h^6). \end{aligned} \quad (24)$$

The vector field defined in (22) that determines the evolution of quasi unidirectional waves, is given by

$$\begin{aligned} \mathcal{F}(U, h) = & U_x + \frac{h^2}{24} \{U_{3x} + 6UU_x\} + \\ & + \frac{h^4}{1920} \left\{ U_{5x} + 60U_x U_{2x} + 20UU_{3x} + 90 \left(\frac{2\beta}{\alpha^2} - 1 \right) U^2 U_x + 30 \langle U^2 \rangle U_x \right\} + \\ & + \frac{h^6}{322560} \left\{ U_{7x} + 420U_{2x} U_{3x} + 210U_x U_{4x} + 42UU_{5x} + 315 \left(\frac{8\beta}{\alpha^2} - 5 \right) (U_x)^3 + \right. \\ & + 630 \left(\frac{12\beta}{\alpha^2} - 7 \right) UU_x U_{xx} + 630 \left(\frac{2\beta}{\alpha^2} - 1 \right) U^2 U_{3x} + 210 \left(\frac{48\gamma}{\alpha^3} - \frac{60\beta}{\alpha^2} + 23 \right) U^3 U_x + \\ & + 210 \langle U^2 \rangle \left(U_{3x} + \left(\frac{18\beta}{\alpha^2} - 9 \right) UU_x \right) + 105 \left(3 \langle U_x^2 \rangle + \left(\frac{12\beta}{\alpha^2} - 10 \right) \langle U^3 \rangle + 6 \langle U^2 \rangle \langle U \rangle \right) U_x \left. \right\} + \\ & + \mathcal{O}(h^8). \end{aligned} \quad (25)$$

Proof. We shall look for an approximate solution of the invariance equation (23) of the form

$$c(U, h) = h^2 c_2(U) + h^4 c_4(U) + \mathcal{O}(h^6). \quad (26)$$

Combining (23) with the explicit expression for \mathcal{F} given in (19), we get

$$\begin{aligned} & [h^2 c_2(U) + h^4 c_4(U) + \mathcal{O}(h^6)]' \left[U_x + h^2 \left(\frac{1}{24} U_{3x} + (U_x)^2 \right) + \mathcal{O}(h^4) \right] = \\ & - [h^2 c_2(U) + h^4 c_4(U) + \mathcal{O}(h^6)]_x - h^2 \left[\frac{1}{8} (U_x)^2 + \frac{h^2}{24} (c_2(U))_{3x} + \frac{h^2}{4} (U c_2)_x + \mathcal{O}(h^4) \right] \\ & - h^4 \left[\frac{1}{192} (U_{3x})^2 + \frac{\beta}{32\alpha^2} (U_x)^3 \right] + \mathcal{O}(h^6). \end{aligned}$$

Collecting terms of orders h^2 and h^4 produces two equations:

$$c_2'(U) U_x = -\frac{1}{8} U U_x, \quad (27)$$

$$c_4'(U) U_x = -c_2'(U) \left(\frac{1}{24} U_{3x} + \frac{1}{8} (U_x)^2 \right) - \frac{1}{24} (c_2(U))_{3x} - \frac{1}{4} (U c_2(U))_x - \frac{1}{192} (U_{3x})^2 - \frac{\beta}{32\alpha^2} (U_x)^3. \quad (28)$$

A solution of equation (27) is easily found:

$$c_2(U) = \frac{1}{16} (\langle U^2 \rangle - U^2). \quad (29)$$

Here, $\langle U^2 \rangle := \int_{\mathbb{T}} U^2(x) dx$ denotes the average of the function U^2 . We could have omitted the term $\frac{1}{16} \langle U^2 \rangle$ in (29), but including this term in $c_2(U)$ guarantees that $\langle c_2(U) \rangle = 0$. See also Remark 10.

Inserting (29) into equation (28) for $c_4(U)$ gives

$$c_4'(U) U_x = \left(\frac{5}{128} - \frac{3\beta}{64\alpha^2} \right) U^2 U_x - \frac{1}{128} \langle U^2 \rangle U_x - \frac{1}{128} U_{xx} U_x,$$

which admits as a solution

$$c_4(U) = \left(\frac{5}{384} - \frac{\beta}{64\alpha^2} \right) U^3 - \frac{1}{128} \langle U^2 \rangle (U - \langle U \rangle) - \frac{1}{256} [(U_x)^2 - \langle (U_x)^2 \rangle]. \quad (30)$$

Again, we made sure that $\langle c_4(U) \rangle = 0$ by choosing an appropriate “integration constant”. Together, (26), (29) and (30) produce (24).

The evolution equation for U is obtained from (22) by inserting the expansion (26) into (19), giving

$$\begin{aligned} U_t &= \mathcal{F}(U, h) := \mathcal{F}(U, h^2 c_2(U) + h^4 c_4(U) + \mathcal{O}(h^6), h) = \\ &= U_x + h^2 \left(\frac{1}{24} U_{3x} + \frac{1}{8} (U_x)^2 \right) + h^4 \left[\frac{1}{1920} U_{5x} + \frac{1}{192} (U_{3x})^2 + \frac{\beta}{32\alpha^2} (U_x)^3 + \frac{1}{4} (U c_2(U))_x \right] + \\ &+ h^6 \left[\frac{1}{322560} U_{7x} + \frac{1}{15360} (U_{5x})^2 + \frac{\beta}{768\alpha^2} (U_{3x})^3 + \frac{\gamma}{128\alpha^3} (U_x)^4 + \right. \\ &\left. + \frac{1}{8} (c_2(U)^2)_x + \frac{1}{96} (U c_2(U))_{3x} + \frac{3\beta}{32\alpha^2} (U^2 c_2(U))_x + \frac{1}{4} (U c_4(U))_x \right] + \mathcal{O}(h^8). \end{aligned}$$

Substituting the expression (29) and (30) that we found for $c_2(U)$ and $c_4(U)$ gives (25). This completes the proof of the theorem. \square

Remark 10. For any (smooth) function $F = F(x) : \mathbb{T} \rightarrow \mathbb{R}$ we denote by

$$\langle F \rangle := \int_{\mathbb{T}} F(x) dx,$$

the average of F over \mathbb{T} . In Theorem 2 we encounter averages of the functions $F = U, U^2, U^3$ and $(U_x)^2$. By including such average terms as “integration constants” in our choice for c_2 and c_4 , we make sure that $\langle c_2(U) \rangle = \langle c_4(U) \rangle = 0$. This choice is of course somewhat arbitrary, but it makes that the solutions c_2 and c_4 of (27) and (28) are unique.

5 A normal form theorem

In (6) we defined the discrete Riemann invariants U and V in terms of the interpolating profiles u and v that were introduced in Section 2. Our choice to define them as $U = 2\alpha(D_h u + v)$ and $V = 2\alpha(D_h u - v)$ helped us in obtaining a closed expression for the equations that govern their evolution, see (16), but was otherwise somewhat arbitrary. For example, we could have equally well defined U as $2\alpha(u_x + v)$ and V as $2\alpha(u_x - v)$, the difference between the former and the latter choices being only of the order h^2 .

Another definition of the variables U and V would have resulted in another expression for the slaving relation $V = c(U, h)$. More importantly, another choice of U would have given another – and perhaps simpler – expression for the reduced vector field $\mathcal{F}(U, h)$ that determines the dynamics on the invariant manifold of quasi unidirectional waves, see (22). This motivates us to look for a near-to-identity transformation $U \mapsto U + \mathcal{O}(h^2)$, i.e., a small change in the definition of U , that simplifies the evolution equation $U_t = \mathcal{F}(U, h)$ as much as possible. It turns out that we can prove the following normal form theorem.

Theorem 3. Recall that $\mathcal{F} = \mathcal{F}(U, h) : C^\infty(\mathbb{T}, \mathbb{R}) \rightarrow C^\infty(\mathbb{T}, \mathbb{R})$ admits the asymptotic expansion (25). There exists a formal change of variables inside the space $C^\infty(\mathbb{T}, \mathbb{R})$ of the form

$$U \mapsto U + h^2 \tilde{G}_2(U) + h^4 \tilde{G}_4(U) + h^6 \tilde{G}_6(U, t) + \mathcal{O}(h^8), \quad (31)$$

that transforms the evolution equation $U_t = \mathcal{F}(U, h)$ into

$$\begin{aligned} U_t = \tilde{\mathcal{F}}(U, h, t) = & \mathcal{C}_1(U, h) K_1(U) + \\ & + h^2 \mathcal{C}_3(U, h) K_3(U) + \\ & + h^4 \mathcal{C}_5(U, h) K_5(U) + \\ & + h^6 \mathcal{C}_7(U, h) [K_7(U) + R(U)] + \mathcal{O}(h^8). \end{aligned} \quad (32)$$

Here, $K_1(U), K_3(U), K_5(U), K_7(U)$ are the first four commuting vector fields in the KdV hierarchy given in Remark 1. The scalars $\mathcal{C}_1(U, h), \mathcal{C}_3(U, h), \mathcal{C}_5(U, h), \mathcal{C}_7(U, h)$ are constants of motion of the KdV hierarchy. They are given explicitly in Remark 13 below. The term $R = R(U)$ can be chosen equal to zero when

$$14\alpha^3 - 27\alpha\beta + 12\gamma = 0.$$

Remark 11. A Toda chain is an FPU chain with a potential energy of the form

$$W(z) = W_{\mathbb{T}}(z) := \frac{e^{2\alpha z} - (1 + 2\alpha z)}{4\alpha^2} = \frac{1}{2} z^2 + \frac{\alpha}{3} z^3 + \frac{\alpha^2}{6} z^4 + \frac{\alpha^3}{15} z^5 + \dots \quad (33)$$

Toda chains thus define a specific one-parameter family of FPU chains, for which

$$\beta = \beta_T := \frac{2}{3}\alpha^2, \quad \gamma = \gamma_T := \frac{1}{3}\alpha^3, \quad \text{etc.}$$

One readily checks that $14\alpha^3 - 27\alpha\beta_T + 12\gamma_T = 0$. This means that for the Toda chains, the normal form to order h^6 given in Theorem 3 lies in the KdV hierarchy.

Theorem 3 follows from a more general result that we will state below as Theorem 4. A result comparable to Theorem 4 was originally formulated by Hiraoka and Kodama in [20]. These authors consider evolution equations similar to (25) for functions $U \in C^\infty(\mathbb{R}, \mathbb{R})$. In this paper we work with functions $U \in C^\infty(\mathbb{T}, \mathbb{R})$. We thus need to adapt the proof of Hiraoka and Kodama.

Remark 12. *The main new feature in our adaptation concerns the use of primitives F_{-x} of functions $F : \mathbb{T} \rightarrow \mathbb{R}$. Primitives of functions $F \in C^\infty(\mathbb{R}, \mathbb{R})$ are always defined. In contrast, a function $F \in C^\infty(\mathbb{T}, \mathbb{R})$ only possesses a well defined primitive $F_{-x} \in C^\infty(\mathbb{T}, \mathbb{R})$ if $\langle F \rangle = \int_{\mathbb{T}} F(x) dx = 0$. To deal with this complication we shall only make use of transformations in $C^\infty(\mathbb{T}, \mathbb{R})$ that map the space of zero-average functions on \mathbb{T} into itself. This in turn forces us to add some extra ‘‘average’’ terms to the transformations that were originally considered by Hiraoka and Kodama.*

In particular, we shall agree that, if F has zero average, then F_{-x} is the unique primitive of F that has zero average itself. For a general $F \in C^\infty(\mathbb{T}, \mathbb{R})$, we will then have the formulas

$$\langle (F - \langle F \rangle)_{-x} \rangle = 0 \quad \text{and} \quad ((F - \langle F \rangle)_{-x})_x = F - \langle F \rangle.$$

Examples of transformations involving primitives occur in Theorem 4 below, which makes use of maps $G_2, G_4 : C^\infty(\mathbb{T}, \mathbb{R}) \rightarrow C^\infty(\mathbb{T}, \mathbb{R})$ given by formulas of the form

$$G_2(U) := \frac{C_5}{C_3} (a_1 U_{2x} + a_2 (U^2 - \langle U^2 \rangle) + a_3 (U_x (U - \langle U \rangle)_{-x} + \langle U^2 \rangle - \langle U \rangle^2) + a_4 \langle U \rangle (U - \langle U \rangle)) \quad (34)$$

and

$$\begin{aligned} G_4(U) := & \frac{C_7}{C_3} (b_1 U_{4x} + b_2 ((U_x)^2 - \langle (U_x)^2 \rangle) + b_3 (UU_{2x} + \langle (U_x)^2 \rangle) + b_4 (U^3 - \langle U^3 \rangle) \\ & + b_5 (U_x (U^2 - \langle U^2 \rangle)_{-x} + \langle U^3 \rangle - \langle U \rangle \langle U^2 \rangle) \\ & + b_6 ((U_{3x} + 6UU_x)(U - \langle U \rangle)_{-x} + 3\langle U^3 \rangle - \langle (U_x)^2 \rangle - 3\langle U^2 \rangle \langle U \rangle) \\ & + b_7 \langle U \rangle U_{xx} + b_8 \langle U \rangle (U^2 - \langle U^2 \rangle) \\ & + b_9 \langle U \rangle (U_x (U - \langle U \rangle)_{-x} + \langle U^2 \rangle - \langle U \rangle^2) + b_{10} \langle U \rangle^2 (U - \langle U \rangle) + b_{11} \langle U^2 \rangle (U - \langle U \rangle) \\ & + b_{12} \langle (U_x)^2 \rangle + b_{13} \langle U^3 \rangle). \end{aligned} \quad (35)$$

Here, $a_1, \dots, a_4, b_1, \dots, b_{13}, C_3, \dots, C_7 \in \mathbb{R}$ are constants that will be specified later. Formulas (34) and (35) are such that $\langle G_2(U) \rangle = \langle G_4(U) \rangle = 0$ for every $U \in C^\infty(\mathbb{T}, \mathbb{R})$.

The following theorem is our version of the result of Hiraoka and Kodama.

Theorem 4 (Hiraoka–Kodama on \mathbb{T}). *Consider an evolution equation for $U \in C^\infty(\mathbb{T}, \mathbb{R})$ of the form*

$$\begin{aligned} U_t = F(U, h) = & C_1 U_x \\ & + h^2 C_3 (U_{3x} + 6UU_x) \\ & + h^4 C_5 (U_{5x} + A_1 U_x U_{2x} + A_2 UU_{3x} + A_3 U^2 U_x + A_4 \langle U^2 \rangle U_x) \\ & + h^6 C_7 (U_{7x} + B_1 U_{2x} U_{3x} + B_2 U_x U_{4x} + B_3 UU_{5x} + B_4 (U_x)^3 + B_5 UU_x U_{2x} + \\ & \quad + B_6 U^2 U_{3x} + B_7 U^3 U_x + \langle U \rangle (B_8 U_{5x} + B_9 U_x U_{2x} + B_{10} UU_{3x} + B_{11} U^2 U_x) \\ & \quad + \langle U^2 \rangle (B_{12} U_{3x} + B_{13} UU_x) + \langle U \rangle^2 (B_{14} U_{3x} + B_{15} UU_x) \\ & \quad + (B_{16} \langle U^3 \rangle + B_{17} \langle (U_x)^2 \rangle + B_{18} \langle U \rangle \langle U^2 \rangle + B_{19} \langle U \rangle^3) U_x + B_{20} \langle (U_x)^3 \rangle) \\ & + \mathcal{O}(h^8). \end{aligned} \quad (36)$$

Here $A_1, \dots, A_4, B_1, \dots, B_{20}, C_1, \dots, C_7$ are scalar coefficients.

i) *By a normal form transformation of the form*

$$U \mapsto U + h^2 G_2(U) + \mathcal{O}(h^4), \quad (37)$$

with G_2 of the form (34), one can transform equation (36) into the form

$$U_t = \mathcal{C}_1(U, h)K_1(U) + h^2\mathcal{C}_3(U, h)K_3(U) + h^4\mathcal{C}_5(U, h)K_5(U) + \mathcal{O}(h^6). \quad (38)$$

The scalars $\mathcal{C}_1(U, h), \dots, \mathcal{C}_5(U, h)$ are constants of motion of the KdV hierarchy, explicitly given by

$$\begin{aligned} \mathcal{C}_1(U, h) &= C_1(1 + h^4(\tilde{A}_4\langle U^2 \rangle + \tilde{A}_5\langle U \rangle^2)), \\ \mathcal{C}_3(U, h) &= C_3(1 + h^2\tilde{A}_6\langle U \rangle), \\ \mathcal{C}_5(U, h) &= C_5. \end{aligned} \quad (39)$$

The scalars $\tilde{A}_4, \tilde{A}_5, \tilde{A}_6$ in (44) depend on A_1, A_2, A_3 and A_4 as follows:

$$\begin{aligned} \tilde{A}_4 &= A_3 + A_4 - 4A_2 + 10, \\ \tilde{A}_5 &= 20 - 2A_2, \\ \tilde{A}_6 &= A_2 - 10. \end{aligned} \quad (40)$$

ii) By a further normal form transformation of the form

$$U \mapsto U + h^4G_4(U) + \mathcal{O}(h^8), \quad (41)$$

with G_4 of the form (35), one can subsequently transform equation (38) into the form

$$\begin{aligned} U_t &= \mathcal{C}_1(U, h)K_1(U) + h^2\mathcal{C}_3(U, h)K_3(U) + h^4\mathcal{C}_5(U, h)K_5(U) \\ &\quad + h^6\mathcal{C}_7(U, h)[K_7(U) + R(U) + \lambda_7\langle (U_x)^3 \rangle] + \mathcal{O}(h^8). \end{aligned} \quad (42)$$

The term $R(U)$ can be chosen equal to zero when $r = r(\mathbf{A}, \mathbf{B}, \mathbf{C}) = 0$, where

$$\begin{aligned} r &:= 1680 - 72B_1 + 180B_2 - 510B_3 - 72B_3 + 27B_5 - 72B_4 + 27B_5 + 24B_6 - 9B_7 \\ &\quad + \frac{C_5^2}{C_3C_7} \{-2400 + 6A_1^2 + 670A_2 - 30A_1A_2 - 4A_2^2 - 60A_3 + 3A_1A_3 - A_2A_3\}. \end{aligned} \quad (43)$$

The scalars $\mathcal{C}_1(U, h), \dots, \mathcal{C}_7(U, h)$ are constants of motion of the KdV hierarchy, explicitly given by

$$\begin{aligned} \mathcal{C}_1(U, h) &= C_1(1 + h^4(\tilde{A}_4\langle U^2 \rangle + \tilde{A}_5\langle U \rangle^2) + h^6(\lambda_4\langle (U_x)^2 \rangle - 2U^3) + \lambda_5\langle U \rangle\langle U^2 \rangle + \lambda_6\langle U \rangle^3), \\ \mathcal{C}_3(U, h) &= C_3(1 + h^2\tilde{A}_6\langle U \rangle + h^4(\lambda_2\langle U^2 \rangle + \lambda_3\langle U \rangle^2)), \\ \mathcal{C}_5(U, h) &= C_5(1 + h^2\lambda_1\langle U \rangle), \\ \mathcal{C}_7(U, h) &= C_7. \end{aligned} \quad (44)$$

Here, the $\tilde{A}_4, \tilde{A}_5, \tilde{A}_6$ are as in (40), the scalar λ_4 is a free parameter, and

$$\begin{aligned} \lambda_1 &= -14 + B_3 + B_8, \\ \lambda_2 &= 42 + B_{12} - 8B_3 + B_6 - \frac{2C_5^2}{3C_3C_7}(A_2 - 10)^2, \\ \lambda_3 &= 28 + B_{10} + B_{14} - 2B_3 - 10B_8 + \frac{2C_5^2}{3C_3C_7}(100 - 20A_2 + A_2^2), \\ \lambda_5 &= 10B_3 - 2B_6 + 10B_8 - 4B_{10} + B_{11} - 6B_{12} + B_{13} + B_{18} \\ &\quad + \frac{C_5^2}{C_3C_7}(-100 + 40A_2 - A_2^2 - 10A_3 - \frac{2}{3}A_2A_3 + \frac{1}{3}A_3^2 - 10A_4 + \frac{1}{3}A_3A_4), \\ \lambda_6 &= -56 + 4B_3 + 20B_8 - 2B_{10} - 6B_{14} + B_{15} + B_{19} \\ &\quad + \frac{2C_5^2}{3C_3C_7}(-100 + A_2^2 + 10A_3 - A_2A_3 + 10A_4 - A_2A_4), \\ \lambda_7 &= \frac{28}{3} + \frac{B_{16}}{2} + B_{17} + B_{20} - \frac{7B_3}{3} + \frac{B_6}{3} \\ &\quad + \frac{C_5^2}{18C_3C_7}(-300 + 70A_2 - A_2^2 - A_2A_3 - 3A_1A_4 + 6A_2A_4). \end{aligned} \quad (45)$$

We will postpone the proof of Theorem 4 to the next section, and conclude the current section by proving Theorem 3 from Theorem 4.

Proof of Theorem 3. The vector field $\mathcal{F}(U, h)$ given in equation (25) is of the form of the vector field $F(U, h)$ in equation (36) with

$$\begin{aligned} \mathbf{A} &= (A_1, A_2, A_3, A_4) = \left(60, 20, 90 \left(\frac{2\beta}{\alpha^2} - 1 \right), 30 \right), \\ \mathbf{B} &= (B_1, \dots, B_{20}) = \left(420, 210, 42, 315 \left(\frac{8\beta}{\alpha^2} - 5 \right), 630 \left(\frac{12\beta}{\alpha^2} - 7 \right), 630 \left(\frac{2\beta}{\alpha^2} - 1 \right), \right. \\ &\quad \left. 210 \left(\frac{48\gamma}{\alpha^3} - \frac{60\beta}{\alpha^2} + 23 \right), 0, 0, 0, 0, 210, 210 \left(\frac{18\beta}{\alpha^2} - 9 \right), 0, 0, 105 \left(\frac{12\beta}{\alpha^2} - 10 \right), 315, 0, 0, 0 \right), \\ \mathbf{C} &= (C_1, C_3, C_5, C_7) = \left(1, \frac{1}{24}, \frac{1}{1920}, \frac{1}{322560} \right). \end{aligned} \quad (46)$$

(NB: for clarity, we denote vectors with bold characters.) According to Theorem 4 it is thus possible to transform (25) into (42) with two consecutive near-to-identity transformations of the form (37) and (41) respectively. To remove the term $h^6 \mathcal{C}_7(U, h) \lambda_7 \langle (U_x)^3 \rangle = h^6 C_7 \lambda_7 \langle (U_x)^3 \rangle$ in (42), it suffices to apply one more time-dependent change of variables

$$U \mapsto U + h^6 \tilde{G}_6(U, t) := U - h^6 C_7 \lambda_7 \int_0^t \langle U_x^3(s) \rangle ds,$$

which cancels this last term in (42), yielding precisely (32).

The quantity that determines whether the term $R(U)$ in (32) and (42) can be chosen equal to zero, is the constant r defined in (43). Substituting (46) in (43) gives

$$r = -\frac{7560}{\alpha^3} (14\alpha^3 - 27\alpha\beta + 12\gamma).$$

This finishes the proof of Theorem 3. \square

Remark 13. *Explicit expressions for the constants of motion $\mathcal{C}_1(U, h), \dots, \mathcal{C}_7(U, h)$ in Theorem 3 can be obtained by combining (44) with (40), (45) and (46). In the setting of Theorem 3 we have*

$$\tilde{A}_4 = -130 + \frac{180\beta}{\alpha^2}, \quad \tilde{A}_5 = -20, \quad \tilde{A}_6 = 10,$$

so that (39) becomes

$$\mathcal{C}_1(U, h) = 1 + h^4 \left(\left(\frac{180\beta}{\alpha^2} - 130 \right) \langle U^2 \rangle - 20 \langle U \rangle^2 \right), \quad \mathcal{C}_3(U, h) = \frac{1}{24} (1 + 10 h^2 \langle U \rangle), \quad \mathcal{C}_5(U, h) = \frac{1}{1920}. \quad (47)$$

By combining (45) and (46) we obtain

$$\begin{aligned} \lambda_1 &= 28, & \lambda_2 &= -854 + \frac{1260\beta}{\alpha^2}, & \lambda_3 &= 84, \\ \lambda_5 &= 420 \left(16 - \frac{63\beta}{\alpha^2} + \frac{54\beta^2}{\alpha^4} \right), & \lambda_6 &= 28 \left(49 - \frac{90\beta}{\alpha^2} \right), & \lambda_7 &= -427 + \frac{630\beta}{\alpha^2}. \end{aligned}$$

Recall that the scalar λ_4 may be chosen freely, so let us choose $\lambda_4 = 0$. Then (44) becomes

$$\begin{aligned} \mathcal{C}_1(U, h) &= 1 + h^4 \left(\left(\frac{180\beta}{\alpha^2} - 130 \right) \langle U^2 \rangle - 20 \langle U \rangle^2 \right) + \\ &\quad + h^6 \left(420 \left(16 - \frac{63\beta}{\alpha^2} + \frac{54\beta^2}{\alpha^4} \right) \langle U \rangle \langle U^2 \rangle + 28 \left(49 - \frac{90\beta}{\alpha^2} \right) \langle U \rangle^3 \right), \\ \mathcal{C}_3(U, h) &= \frac{1}{24} \left(1 + 10 h^2 \langle U \rangle + h^4 \left(\left(\frac{1260\beta}{\alpha^2} - 854 \right) \langle U^2 \rangle + 84 \langle U \rangle^2 \right) \right), \\ \mathcal{C}_5(U, h) &= \frac{1}{1920} (1 + 28 h^2 \langle U \rangle), \\ \mathcal{C}_7(U, h) &= \frac{1}{322560}. \end{aligned} \quad (48)$$

6 Proof of Theorem 4

In the proof of Theorem 4 that we provide in this section, we shall explicitly compute how coordinate changes of the form $U \mapsto U + h^2 G_2(U) + \mathcal{O}(h^4)$ and $U \mapsto U + h^4 G_4(U) + \mathcal{O}(h^8)$, with $G_2(U)$ given by

(34) and $G_4(U)$ by (35), transform an evolution equation $U_t = F(U, h)$ of the form (36). In particular, we shall compute when exactly it is possible to transform (36) into a member of the KdV hierarchy, to order h^4 and to order h^6 , and which choices of $G_2(U)$ and $G_4(U)$ realise this transformation.

Our proof is divided in three steps: first we transform equation (36) into a normal form to order h^4 . Next, we compute how this first transformation affects the evolution equation to order h^6 . And finally, we normalise this new equation to order h^6 . Before we give a more concise outline of this procedure, we recall the definition of the Lie bracket of two operators f and g :

$$[f, g](U) := f'(U)g(U) - g'(U)f(U). \quad (49)$$

Here $f'(U)g(U)$ denotes the Gateaux derivative (or directional derivative) of an operator f (evaluated at U) in the direction of $g(U)$. See also Remark 9. More explicitly,

$$f'(U)g(U) := \left. \frac{d}{d\varepsilon} \right|_{\varepsilon=0} f(U + \varepsilon g(U)).$$

In this paper, f and g will always be operators from $C^\infty(\mathbb{T}, \mathbb{R})$ to $C^\infty(\mathbb{T}, \mathbb{R})$.

Outline of the normal form procedure. Below we sketch the procedure by which we bring equation (36) in Theorem 4 into normal form. For other settings in which normal form transformations are applied to the FPU chain, we refer to [5, 16, 20]. Recall that equation (36) is of the form

$$U_t = F_1(U) + h^2 F_3(U) + h^4 F_5(U) + h^6 F_7(U) + \mathcal{O}(h^8), \quad (50)$$

in which

$$\begin{aligned} F_1(U) &= C_1 U_x, \\ F_3(U) &= C_3 (U_{3x} + 6UU_x), \\ F_5(U) &= C_5 (U_{5x} + A_1 U_x U_{2x} + A_2 UU_{3x} + A_3 U^2 U_x + A_4 \langle U^2 \rangle U_x), \\ F_7(U) &= C_7 (U_{7x} + B_1 U_{2x} U_{3x} + B_2 U_x U_{4x} + B_3 UU_{5x} + B_4 (U_x)^3 + B_5 UU_x U_{2x} + B_6 U^2 U_{3x} \\ &\quad + B_7 U^3 U_x + \langle U \rangle (B_8 U_{5x} + B_9 U_x U_{2x} + B_{10} UU_{3x} + B_{11} U^2 U_x) \\ &\quad + \langle U^2 \rangle (B_{12} U_{3x} + B_{13} UU_x) + \langle U \rangle^2 (B_{14} U_{3x} + B_{15} UU_x) \\ &\quad + (B_{16} \langle U^3 \rangle + B_{17} \langle (U_x)^2 \rangle + B_{18} \langle U \rangle \langle U^2 \rangle + B_{19} \langle U \rangle^3) U_x + B_{20} \langle (U_x)^3 \rangle). \end{aligned} \quad (51)$$

We will transform this equation by a normal form procedure that consists of two separate transformation steps. First we make a transformation of the form $U \mapsto e^{h^2 G_2}(U) := U + h^2 G_2(U) + \frac{h^4}{2} G_2'(U) G_2(U) + \mathcal{O}(h^6)$ where G_2 is chosen in such a way that $[G_2, F_1] = 0$. This transforms the vector field $F = F(U)$ into

$$e^{h^2[G_2, \cdot]}(F) := 1 + h^2[G_2, F] + \frac{1}{2}h^4[G_2, [G_2, F]] + \frac{1}{6}h^6[G_2, [G_2, [G_2, F]]] + \mathcal{O}(h^8).$$

Expanding F in powers of h in this equation, we obtain the expansion

$$\begin{aligned} U_t = & F_1(U) + h^2 F_3(U) + \\ & + h^4 \underbrace{\{F_5(U) + [G_2, F_3](U)\}}_{=: N_5(U)} + h^6 \underbrace{\{F_7(U) + [G_2, F_5](U) + \frac{1}{2}[G_2, [G_2, F_3]](U)\}}_{=: \frac{1}{2}[G_2, F_5 + N_5](U) =: R_6(U)} + \mathcal{O}(h^8), \end{aligned} \quad (52)$$

for the transformed evolution equation. Note that the term $h^2 F_3(U)$ is unaffected because $[G_2, F_1] = 0$. We now want to choose G_2 in such a way that $N_5 = F_5 + [G_2, F_3]$ is in the KdV hierarchy, see Remark 1. It turns out that this can always be arranged.

The next step is to make a further coordinate change $U \mapsto e^{h^4 G_4}(U) = U + h^4 G_4(U) + \mathcal{O}(h^8)$, again choosing G_4 so that $[G_4, F_1] = 0$. This transforms our evolution equation further into

$$U_t = F_1(U) + h^2 F_3(U) + h^4 N_5(U) + h^6 \underbrace{\{F_7(U) + \frac{1}{2}[G_2, F_5 + N_5](U) + [G_4, F_3](U)\}}_{=: F_7(U) + R_6(U) + [G_4, F_3](U) =: N_7(U)} + \mathcal{O}(h^8). \quad (53)$$

The goal is to choose G_4 so that $N_7 = F_7 + \frac{1}{2}[G_2, F_5 + N_5] + [G_4, F_3]$ is in the KdV hierarchy as well. It turns out that this can only be arranged (within the class of transformations that we consider) if a certain relation among the coefficients of F_5 and F_7 is satisfied.

X	$[X, U_{3x} + 6UU_x]$
U_{2x}	$12U_x U_{2x}$
$U^2 - \langle U^2 \rangle$	$-6U_x U_{2x} - 6U^2 U_x + 6\langle U^2 \rangle U_x$
$U_x(U - \langle U \rangle)_{-x} + \langle U^2 \rangle - \langle U \rangle^2$	$-3U_x U_{2x} - 3UU_{3x} - 3U^2 U_x - 9\langle U^2 \rangle U_x + 6\langle U \rangle^2 U_x + 6\langle U \rangle UU_x + 3\langle U \rangle U_{3x}$
$\langle U \rangle (U - \langle U \rangle)$	$-6\langle U \rangle UU_x + 6\langle U \rangle^2 U_x$

Table 1: Lie brackets determining the first normalisation step.

Normalisation at lowest order. Inspired by Lemma 5.1 in [20] we choose G_2 of the form (34). We have that $[G_2, F_1] = C_1[G_2, \partial_x] = 0$ because $G_2 = G_2(U)$ does not explicitly depend on x . We also remark that the authors of [20] use a simpler class of transformations of the form $(C_3/C_5)G_2(U) = a_1 U_{2x} + a_2 U^2 + a_3 U_x U_{-x}$. This is not sufficient for us, on the one hand because equation (36) that we try to bring into normal form, is more general than the equations considered in [20]. On the other hand, we like to make sure that $\langle G_2(U) \rangle = 0$, so that $e^{h^2 G_2}$ maps the space of zero-average functions into itself. To compute the transformed vector field $N_5(U)$, we use the bracket relations in Table 1.

Remark 14. *The brackets in Table 1 can all be computed by hand, using formula (49). To illustrate,*

$$\begin{aligned}
[U^2 - \langle U^2 \rangle, U_{3x} + 6UU_x] &= \left. \frac{d}{d\varepsilon} \right|_{\varepsilon=0} \left\{ (U + \varepsilon(U_{3x} + 6UU_x))^2 - \langle (U + \varepsilon(U_{3x} + 6UU_x))^2 \rangle \right. \\
&\quad \left. - (U + \varepsilon(U^2 - \langle U^2 \rangle))_{3x} - 6(U + \varepsilon(U^2 - \langle U^2 \rangle))(U + \varepsilon(U^2 - \langle U^2 \rangle))_x \right\} \\
&= 2U(U_{3x} + 6UU_x) - \langle 2U(U_{3x} + 6UU_x) \rangle \\
&\quad - (U^2 - \langle U^2 \rangle)_{3x} - 6U(U^2 - \langle U^2 \rangle)_x - 6(U^2 - \langle U^2 \rangle)U_x.
\end{aligned}$$

Now observe that $\langle UU_{3x} \rangle = -\langle U_x U_{2x} \rangle = -\langle (\frac{1}{2}(U_x^2)_x) \rangle = 0$ (using integration by parts and the fundamental theorem of calculus), that $\langle U^2 U_x \rangle = \langle (\frac{1}{3}U^3)_x \rangle = 0$ (again by the fundamental theorem of calculus), and that $\langle U^2 \rangle_{3x} = \langle U^2 \rangle_x = 0$ (because the average is not a function of x). Using these identities, as well as the chain rule to rewrite the terms $(U^2)_{3x}$ and $U(U^2)_x$, we can simplify our expression for the bracket to $-6U_x U_{2x} - 6U^2 U_x + 6\langle U^2 \rangle U_x$. The other brackets are computed in an analogous fashion.

Using (34), (51) and the table above, we compute that

$$\begin{aligned}
[G_2, F_3](U) &= [G_2, C_3(U_{3x} + 6UU_x)] \\
&= C_5 \left\{ (12a_1 - 6a_2 - 3a_3)U_x U_{2x} - 3a_3 UU_{3x} - (6a_2 + 3a_3)U^2 U_x + \right. \\
&\quad \left. (6a_3 - 6a_4)\langle U \rangle UU_x + 3a_3 \langle U \rangle U_{3x} + (6a_2 - 9a_3)\langle U^2 \rangle U_x + (6a_3 + 6a_4)\langle U \rangle^2 U_x \right\}.
\end{aligned} \tag{54}$$

Recall that we want to make our lowest order normal form a member of the KdV hierarchy, of the form

$$\begin{aligned}
N_5(U) &= F_5(U) + [G_2, F_3](U) = \\
&C_5(U_{5x} + 20U_x U_{2x} + 10UU_{3x} + 30U^2 U_x + \widetilde{A}_4 \langle U^2 \rangle U_x + \widetilde{A}_5 \langle U \rangle^2 U_x + \widetilde{A}_6 \langle U \rangle (U_{3x} + 6UU_x)).
\end{aligned} \tag{55}$$

This can be arranged if we can solve the system of linear equations

$$\begin{cases} A_1 + 12a_1 - 6a_2 - 3a_3 = 20, \\ A_2 - 3a_3 = 10, \\ A_3 - 6a_2 - 3a_3 = 30, \\ 6a_3 - 6a_4 = 18a_3. \end{cases}$$

for the coefficients a_1, \dots, a_4 that define G_2 . It turns out that this system admits the unique solution

$$\begin{aligned}
a_1 &= \frac{1}{12}(A_3 - A_1 - 10), \\
a_2 &= \frac{1}{6}(A_3 - A_2 - 20), \\
a_3 &= \frac{1}{3}(A_2 - 10), \\
a_4 &= \frac{2}{3}(10 - A_2).
\end{aligned} \tag{56}$$

Inserting (56) back into (54) and comparing with (55), we in fact obtain

$$\begin{aligned}\tilde{A}_4 &= A_4 + 6a_2 - 9a_3 = A_3 + A_4 - 4A_2 + 10, \\ \tilde{A}_5 &= 6a_3 + 6a_4 = 20 - 2A_2, \\ \tilde{A}_6 &= 3a_3 = A_2 - 10.\end{aligned}$$

This coincides with (40). Equations (39) follow from (55) combined with (40). This completes the proof of part *i*) of Theorem 4.

A new term higher order term. The normal form transformation generated by $h^2G_2(U)$ simplifies our evolution equation to order h^4 , but it adds new additional terms at order h^6 . From (52) we see that these new terms are

$$R_6(U) = \frac{1}{2}[G_2, F_5 + N_5](U).$$

Here G_2 is as determined as in the previous paragraph, and we in fact have

$$F_5(U) + N_5(U) = C_5^2 \left\{ 2U_{5x} + (A_1 + 20)U_x U_{2x} + (A_2 + 10)UU_{3x} + (A_3 + 30)U^2U_x + (A_3 + 2A_4 - 4A_2 + 10)\langle U^2 \rangle U_x + (20 - 2A_2)\langle U \rangle^2 U_x + (A_2 - 10)\langle U \rangle(U_{3x} + 6UU_x) \right\}.$$

To compute $R_6(U)$ explicitly, we use the bracket relations listed in Table 2. Using Table 2, it is straightforward to compute that

$$\begin{aligned}R_6(U) &= \frac{C_5^2}{C_3} \left(\tilde{B}_1 U_{2x} U_{3x} + \tilde{B}_2 U_x U_{4x} + \tilde{B}_3 U U_{5x} + \tilde{B}_4 (U_x)^3 + \tilde{B}_5 U U_x U_{2x} + \tilde{B}_6 U^2 U_{3x} + \tilde{B}_7 U^3 U_x + \right. \\ &\quad \left. \langle U \rangle \left(\tilde{B}_8 U_{5x} + \tilde{B}_9 U_x U_{2x} + \tilde{B}_{10} U U_{3x} + \tilde{B}_{11} U^2 U_x \right) + \right. \\ &\quad \left. \langle U^2 \rangle \left(\tilde{B}_{12} U_{3x} + \tilde{B}_{13} U U_x \right) + \langle U \rangle^2 \left(\tilde{B}_{14} U_{3x} + \tilde{B}_{15} U U_x \right) + \right. \\ &\quad \left. \left(\tilde{B}_{16} \langle U^3 \rangle + \tilde{B}_{17} \langle (U_x)^2 \rangle + \tilde{B}_{18} \langle U \rangle \langle U^2 \rangle + \tilde{B}_{19} \langle U \rangle^3 \right) U_x + \tilde{B}_{20} \langle (U_x)^3 \rangle \right),\end{aligned}\tag{57}$$

in which

$$\begin{aligned}\tilde{B}_1 &= a_1(A_1 + 20) - 20a_2 - 15a_3, \\ \tilde{B}_2 &= a_1(A_2 + 10) - 10a_2 - 10a_3, \\ \tilde{B}_3 &= -5a_3, \\ \tilde{B}_4 &= a_1(A_3 + 30) - a_2(A_1 + 20) - (a_3/4)(A_1 + A_2 + 30), \\ \tilde{B}_5 &= 2a_1(A_3 + 30) - a_2(A_1 + 3A_2 + 50) - (3a_3/2)(A_1 + A_2 + 30), \\ \tilde{B}_6 &= -(a_2 + 3a_3)(A_2 + 10)/2, \\ \tilde{B}_7 &= -(a_2 + a_3/3)(A_3 + 30), \\ \tilde{B}_8 &= 5a_3, \\ \tilde{B}_9 &= (6a_1 - 3a_2)(A_2 - 10) + 3a_3(A_1 - A_2 + 30)/2 - a_4(A_1 + 20)/2, \\ \tilde{B}_{10} &= 30a_3 - a_4(A_2 + 10)/2, \\ \tilde{B}_{11} &= (6a_2 + 3a_3)(10 - A_2)/2 + (a_3 - 2a_4)(A_3 + 30)/2, \\ \tilde{B}_{12} &= (a_2 - a_3)(A_2 + 10)/2, \\ \tilde{B}_{13} &= (a_2 - a_3)(A_3 + 30), \\ \tilde{B}_{14} &= a_3(2A_2 - 10) + a_4(A_2 + 10)/2, \\ \tilde{B}_{15} &= (a_3 + a_4)(A_3 + 30) + 3(a_3 - a_4)(A_2 - 10), \\ \tilde{B}_{16} &= (a_3/2 - a_2)(A_3 + 2A_4 - 4A_2 + 10) - a_3(A_3 + 30)/6, \\ \tilde{B}_{17} &= a_1(A_3 + 2A_4 - 4A_2 + 10) + a_3(3A_2 - A_1 + 10)/4, \\ \tilde{B}_{18} &= (a_2 - \frac{3}{2}a_3 - a_4)(A_3 + 2A_4 - 4A_2 + 10) + (3a_2 - \frac{9}{2}a_3)(A_2 - 10), \\ \tilde{B}_{19} &= (a_3 + a_4)(A_3 + 2A_4 - A_2 - 20), \\ \tilde{B}_{20} &= (a_2 - a_3)(A_1 - 2A_2)/2.\end{aligned}\tag{58}$$

Here the coefficients a_1, a_2, a_3, a_4 are as defined in (56). Note that $\tilde{B}_4 - \frac{1}{2}\tilde{B}_5 + \tilde{B}_6 + \tilde{B}_{20} = 0$ because $R_6(U)$ has zero average. To summarise, we now have that

$$F_7(U) + R_6(U) = C_7 \left(U_{7x} + \left(B_1 + \frac{C_5^2}{C_3 C_7} \tilde{B}_1 \right) U_{2x} U_{3x} + \left(B_2 + \frac{C_5^2}{C_3 C_7} \tilde{B}_2 \right) U_x U_{4x} + \dots \right)$$

is the new term of order h^6 , after the first normal form transformation.

X	Y	$[X, Y]$
U_{xx}	U_{5x}	0
"	$U_x U_{2x}$	$2U_{2x}U_{3x}$
"	UU_{3x}	$2U_xU_{4x}$
"	U^2U_x	$2(U_x)^3 + 4UU_xU_{2x}$
"	$\langle U^2 \rangle U_x$	$2\langle (U_x)^2 \rangle U_x$
"	$\langle U \rangle^2 U_x$	0
"	$\langle U \rangle (U_{3x} + 6UU_x)$	$12\langle U \rangle U_x U_{2x}$
$U^2 - \langle U^2 \rangle$	U_{5x}	$-20U_{2x}U_{3x} - 10U_xU_{4x}$
"	$U_x U_{2x}$	$-2(U_x)^3 - 2UU_xU_{2x} + \langle (U_x)^3 \rangle$
"	UU_{3x}	$-6UU_xU_{2x} - U^2U_{3x} - 2\langle (U_x)^3 \rangle + \langle U^2 \rangle U_{3x}$
"	U^2U_x	$-2U^3U_x + 2\langle U^2 \rangle UU_x$
"	$\langle U^2 \rangle U_x$	$-2\langle U^3 \rangle U_x + 2\langle U \rangle \langle U^2 \rangle U_x$
"	$\langle U \rangle^2 U_x$	0
"	$\langle U \rangle (U_{3x} + 6UU_x)$	$-6\langle U \rangle U_x U_{2x} - 6\langle U \rangle U^2 U_x + 6\langle U \rangle \langle U^2 \rangle U_x$
$U_x(U - \langle U \rangle)_{-x} + \langle U^2 \rangle - \langle U \rangle^2$	U_{5x}	$-15U_{2x}U_{3x} - 10U_xU_{4x} - 5UU_{5x} + 5\langle U \rangle U_{5x}$
"	$U_x U_{2x}$	$-\frac{1}{2}U_x^3 - 3UU_xU_{2x} - \frac{1}{2}\langle U_x^2 \rangle U_x - \langle U_x^3 \rangle + 3\langle U \rangle U_x U_{2x}$
"	UU_{3x}	$-\frac{1}{2}(U_x)^3 - 3UU_xU_{2x} - 3U^2U_{3x} + \frac{3}{2}\langle (U_x)^2 \rangle U_x + 2\langle (U_x)^3 \rangle - \langle U^2 \rangle U_{3x} + \langle U \rangle^2 U_{3x} + 3\langle U \rangle UU_{3x}$
"	U^2U_x	$-\frac{2}{3}U^3U_x - \frac{1}{3}\langle U^3 \rangle U_x - 2\langle U^2 \rangle UU_x + 2\langle U \rangle^2 UU_x + \langle U \rangle U^2 U_x$
"	$\langle U^2 \rangle U_x$	$\langle U^3 \rangle U_x - 3\langle U \rangle \langle U^2 \rangle U_x + 2\langle U \rangle^3 U_x$
"	$\langle U \rangle^2 U_x$	0
"	$\langle U \rangle (U_{3x} + 6UU_x)$	$-3\langle U \rangle U_x U_{2x} - 3\langle U \rangle UU_{3x} - 3\langle U \rangle U^2 U_x + 3\langle U \rangle^2 U_{3x} + 6\langle U \rangle^2 UU_x + 6\langle U \rangle^3 U_x - 9\langle U \rangle \langle U^2 \rangle U_x$
$\langle U \rangle (U - \langle U \rangle)$	U_{5x}	0
"	$U_x U_{2x}$	$-\langle U \rangle U_x U_{2x}$
"	UU_{3x}	$\langle U \rangle^2 U_{3x} - \langle U \rangle UU_{3x}$
"	U^2U_x	$2\langle U \rangle^2 UU_x - 2\langle U \rangle U^2 U_x$
"	$\langle U^2 \rangle U_x$	$2\langle U \rangle^3 U_x - 2\langle U \rangle \langle U^2 \rangle U_x$
"	$\langle U \rangle^2 U_x$	0
"	$\langle U \rangle (U_{3x} + 6UU_x)$	$6\langle U \rangle^3 U_x - 6\langle U \rangle^2 UU_x$

Table 2: Lie brackets determining the new higher order term.

X	$[X, U_{3x} + 6UU_x]$
U_{4x}	$60U_{2x}U_{3x} + 24U_xU_{4x}$
$\langle U_x \rangle^2 - \langle (U_x)^2 \rangle$	$-6U_{2x}U_{3x} + 6(U_x)^3 - 6\langle (U_x)^3 \rangle + 6\langle (U_x)^2 \rangle U_x$
$UU_{2x} + \langle (U_x)^2 \rangle$	$-3U_{2x}U_{3x} - 3U_xU_{4x} + 12UU_xU_{2x} + 6\langle (U_x)^3 \rangle - 6\langle (U_x)^2 \rangle U_x$
$U^3 - \langle U^3 \rangle$	$-6(U_x)^3 - 18UU_xU_{2x} - 6U^3U_x - 3\langle (U_x)^3 \rangle + 6\langle U^3 \rangle U_x$
$U_x(U^2 - \langle U^2 \rangle)_{-x} + \langle U^3 \rangle - \langle U \rangle \langle U^2 \rangle$	$-3(U_x)^3 - 6UU_xU_{2x} - 3U^2U_{3x} - 2U^3U_x + 3\langle U^2 \rangle U_{3x} + 3\langle (U_x)^2 \rangle U_x + 6\langle U^2 \rangle UU_x - 10\langle U^3 \rangle U_x + 3\langle (U_x)^3 \rangle + 6\langle U \rangle \langle U^2 \rangle U_x$
$(U_{3x} + 6UU_x)(U - \langle U \rangle)_{-x} + 3\langle U^3 \rangle - \langle (U_x)^2 \rangle - 3\langle U^2 \rangle \langle U \rangle$	$-3U_xU_{4x} - 3UU_{5x} - 18(U_x)^3 - 72UU_xU_{2x} - 21U^2U_{3x} - 18U^3U_x - 3\langle U^2 \rangle U_{3x} + 6\langle (U_x)^2 \rangle U_x + 3\langle (U_x)^3 \rangle - 18\langle U^3 \rangle U_x - 18\langle U^2 \rangle UU_x + 18\langle U \rangle \langle U^2 \rangle U_x + 3\langle U \rangle (U_{5x} + 18U_xU_{2x} + 8UU_{3x} + 12U^2U_x)$
$\langle U \rangle U_{xx}$	$12\langle U \rangle U_x U_{2x}$
$\langle U \rangle (U^2 - \langle U^2 \rangle)$	$-6\langle U \rangle U_x U_{2x} - 6\langle U \rangle U^2 U_x + 6\langle U \rangle \langle U^2 \rangle U_x$
$\langle U \rangle (U_x(U - \langle U \rangle)_{-x} + \langle U^2 \rangle - \langle U \rangle^2)$	$-3\langle U \rangle U_x U_{2x} - 3\langle U \rangle UU_{3x} - 3\langle U \rangle U^2 U_x - 9\langle U \rangle \langle U^2 \rangle U_x + 6\langle U \rangle^3 U_x + 6\langle U \rangle^2 UU_x + 3\langle U \rangle^2 U_{3x}$
$\langle U \rangle^2 (U - \langle U \rangle)$	$-6\langle U \rangle^2 UU_x + 6\langle U \rangle^3 U_x$
$\langle U^2 \rangle (U - \langle U \rangle)$	$6\langle U^2 \rangle \langle U \rangle U_x - 6\langle U^2 \rangle UU_x$
$\langle (U_x)^2 \rangle$	$-6\langle (U_x)^2 \rangle U_x + 6\langle (U_x)^3 \rangle$
$\langle U^3 \rangle$	$-6\langle U^3 \rangle U_x + 3\langle (U_x)^3 \rangle$

Table 3: Lie brackets determining the second normalisation step.

Normalisation at second order. In this final paragraph we perform a second change of variables $U \mapsto e^{h^2 G_4}(U) = U + h^4 G_4(U) + \mathcal{O}(h^8)$ to transform our evolution equation into an as simple as possible form at order h^6 . In (53) we see that in this way we add the term $[G_4, F_3]$ to the order h^6 part of the equation. Recall that the transformation generator $G_4(U)$ has the specific form (35). To see what exactly the additional term $[G_4, F_3]$ will look like, we present the relevant Lie brackets in Table 3.

From now on we shall write

$$\mathbf{B} = (B_1, \dots, B_{20}), \tilde{\mathbf{B}} = (\tilde{B}_1, \dots, \tilde{B}_{20}), \tilde{\tilde{\mathbf{B}}} = (\tilde{\tilde{B}}_1, \dots, \tilde{\tilde{B}}_{20}), \text{ and } \mathbf{b} = (b_1, \dots, b_{13}).$$

(Again, we denote vectors with bold characters.) Here, the B_i are as in (51), the \tilde{B}_j as in (58), and the b_k as in (35). From the bracket relations in Table 3, it follows that the new order h^6 part of our evolution equation reads

$$N_7(U) := F_7(U) + R_6(U) + [G_4, F_3](U) = C_7 \left(U_{7x} + \tilde{\tilde{B}}_1 U_{2x} U_{3x} + \tilde{\tilde{B}}_2 U_x U_{4x} + \tilde{\tilde{B}}_3 U U_{5x} + \tilde{\tilde{B}}_4 (U_x)^3 + \tilde{\tilde{B}}_5 U U_x U_{2x} + \tilde{\tilde{B}}_6 U^2 U_{3x} + \tilde{\tilde{B}}_7 U^3 U_x + \dots \right)$$

– we use the notation as in (51) – in which

$$\tilde{\tilde{\mathbf{B}}} = \mathbf{B} + \frac{C_5^2}{C_3 C_7} \tilde{\mathbf{B}} + M \mathbf{b},$$

and M is the 20×13 matrix

$$M = \begin{pmatrix} 60 & -6 & -3 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 24 & 0 & -3 & 0 & 0 & -3 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -3 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 6 & 0 & -6 & -3 & -18 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 12 & -18 & -6 & -72 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -3 & -21 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -6 & -2 & -18 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 3 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 54 & 12 & -6 & -3 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 24 & 0 & 0 & -3 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 36 & 0 & -6 & -3 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 3 & -3 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 6 & -18 & 0 & 0 & 0 & 0 & -6 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 3 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 6 & -6 & 0 & 0 & 0 \\ 0 & 0 & 0 & 6 & -10 & -18 & 0 & 0 & 0 & 0 & 0 & 0 & -6 \\ 0 & 6 & -6 & 0 & 3 & 6 & 0 & 0 & 0 & 0 & 0 & -6 & 0 \\ 0 & 0 & 0 & 0 & 6 & 18 & 0 & 6 & -9 & 0 & 6 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 6 & 6 & 0 & 0 & 0 \\ 0 & -6 & 6 & -3 & 3 & 3 & 0 & 0 & 0 & 0 & 0 & 6 & 3 \end{pmatrix}.$$

Recall that we would like $N_7(U)$ to lie in the KdV hierarchy. More precisely, we want it to be of the form

$$\begin{aligned} & C_7(U_{7x} + 70U_{2x}U_{3x} + \dots) \\ & + \lambda_1 \langle U \rangle (U_{5x} + 20U_x U_{2x} + \dots) + \lambda_2 \langle U^2 \rangle (U_{3x} + 6UU_x) + \lambda_3 \langle U \rangle^2 (U_{3x} + 6UU_x) \\ & + \lambda_4 (\langle (U_x)^2 \rangle - 2\langle U^3 \rangle) U_x + \lambda_5 \langle U \rangle \langle U^2 \rangle U_x + \lambda_6 \langle U \rangle^3 U_x + \lambda_7 \langle (U_x)^3 \rangle, \end{aligned} \quad (59)$$

for certain scalars $\lambda_1, \dots, \lambda_7$. This can be arranged precisely when the system of algebraic equations

$$\tilde{\mathbf{B}} = \mathbf{B} + \frac{C_5^2}{C_3 C_7} \tilde{\mathbf{B}} + M\mathbf{b} = \mathbf{w} \quad (60)$$

can be solved for \mathbf{b} , where $\mathbf{w} = \mathbf{w}(\lambda)$ is defined as

$$\mathbf{w} = (70, 42, 14, 70, 280, 70, 140, \lambda_1, 20\lambda_1, 10\lambda_1, 30\lambda_1, \lambda_2, 6\lambda_2, \lambda_3, 6\lambda_3, -2\lambda_4, \lambda_4, \lambda_5, \lambda_6, \lambda_7). \quad (61)$$

In fact, the system (60) admits a solution \mathbf{b} if and only if

$$\mathbf{w} - \tilde{\mathbf{B}} = \mathbf{w} - \frac{C_5^2}{C_3 C_7} \tilde{\mathbf{B}} - \mathbf{B} \in \text{ran } M.$$

Now note that the orthogonal complement of $\text{ran } M$ (with respect to the Euclidean inner product in \mathbb{R}^{20}) is $(\text{ran } M)^\perp = \text{span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4, \mathbf{v}_5, \mathbf{v}_6, \mathbf{v}_7\}$ with

$$\begin{aligned} \mathbf{v}_1 &= (0, 0, -14, 0, 0, 2, 0, 0, 0, 0, 0, 0, 0, 0, 3, 6, 0, 0, 0, 6), \\ \mathbf{v}_2 &= (0, 0, 32, 0, 0, 0, 0, 0, 0, 4, 0, 0, 0, 0, 1, 0, 0, 0, 1, 0), \\ \mathbf{v}_3 &= (0, 0, -48, 0, 0, 4, 0, 0, 0, -4, 1, 0, 1, 0, 0, 0, 0, 1, 0, 0), \\ \mathbf{v}_4 &= (0, 0, 8, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0), \\ \mathbf{v}_5 &= (0, 0, -8, 0, 0, 1, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0), \\ \mathbf{v}_6 &= (0, 0, 1, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0), \\ \mathbf{v}_7 &= (24, -60, 170, 24, -9, -8, 3, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0). \end{aligned} \quad (62)$$

Indeed, one may check that the \mathbf{v}_i span the kernel of the map $\mathbf{v} \mapsto \mathbf{v}^T M$. It is thus sufficient to require that $\mathbf{w} - \tilde{\mathbf{B}}$ is perpendicular to each of the vectors $\mathbf{v}_1, \dots, \mathbf{v}_7$, i.e., that

$$\left(\mathbf{B} + \frac{C_5^2}{C_3 C_7} \tilde{\mathbf{B}} - \mathbf{w} \right) \cdot \mathbf{v}_j = 0 \quad \text{for } j = 1, \dots, 7. \quad (63)$$

After substituting (56) in (58) to obtain an expression for $\tilde{\mathbf{B}}$ in terms of \mathbf{A} and \mathbf{B} , equations (63) for $j = 1, \dots, 6$ become precisely equations (45). Equation (63) for $j = 7$ is equivalent to the equation $r(\mathbf{A}, \mathbf{B}, \mathbf{C}) = 0$, with r as defined in (43). This last equation thus constitutes the (only) constraint on the parameters A_1, \dots, B_{20} of the vector fields F_5 and F_7 . Equations (44) follow from (55) combined with (40) and (59) combined with (45). This concludes the proof of part *ii*) of Theorem 4.

References

- [1] D. Bambusi, *Hamiltonian Studies on Counter-Propagating Water Waves*, Water Waves, 2020.
- [2] D. Bambusi, A. Carati, A. Maiocchi and A. Maspero, *Some analytic results on the FPU paradox*, Hamiltonian partial differential equations and applications, Fields Inst. Commun. **75**, 235–254 (2015).
- [3] D. Bambusi, T. Kappeler and T. Paul, *From Toda to KdV*, Nonlinearity **28**, no. 7, 2461–2496 (2015).
- [4] D. Bambusi and A. Maspero, *Birkhoff coordinates for the Toda Lattice in the limit of infinitely many particles with an application to FPU*, J. Funct. Anal. **270**, 1818–1887 (2016).
- [5] D. Bambusi and A. Ponno, *On Metastability in FPU*, Commun. Math. Phys. **264**, 539–561 (2006).
- [6] G. Benettin, H. Christodoulidi and A. Ponno, *The Fermi-Pasta-Ulam Problem and Its Underlying Integrable Dynamics*, J. Stat. Phys. **152**, 195–212 (2013).
- [7] G. Benettin, R. Livi and A. Ponno, *The Fermi-Pasta-Ulam Problem: Scaling Laws vs. Initial Conditions*, J. Stat. Phys. **135**, 873–893 (2009).
- [8] G. Benettin, S. Pasquali and A. Ponno, *The Fermi-Pasta-Ulam problem and its underlying integrable dynamics: an approach through Lyapunov Exponents*, J. Stat. Phys. **171**, 521–542 (2018).
- [9] G. Benettin and A. Ponno, *Time-scales to equipartition in the Fermi-Pasta-Ulam problem: finite-size effects and thermodynamic limit*, J. Stat. Phys. **144**, 793–812 (2011).
- [10] G. Benettin and A. Ponno, *Understanding the FPU state in FPU-like models*, to appear in Mathematics in Engineering (2020).
- [11] Chaos focus issue: *The “Fermi–Pasta–Ulam” problem—the first 50 years*. Chaos **15**, 2005.
- [12] T. Dauxois and S. Ruffo, *Fermi-Pasta-Ulam nonlinear lattice oscillations*, Scholarpedia **3**, 5538 (2008).
- [13] H.R. Dullin, G.A. Gottwald and D.D. Holm, *Camassa-Holm, Korteweg-de Vries-5 and other asymptotically equivalent equations for shallow water waves*, Fluid Dynamics Research **33**, 73–95 (2003).
- [14] W.E. Ferguson, H. Flaschka and D.W. McLaughlin, *Nonlinear Normal Modes for the Toda Chain*, J. Comp. Phys. **45**, 157–209 (1982).
- [15] E. Fermi, J. Pasta, and S. Ulam: *Studies of Non Linear Problems*, Los-Alamos Internal Report, Document LA-1940 (1955), first published in: *Enrico Fermi Collected Papers*, Vol. II, The University of Chicago Press, Chicago, and Accademia Nazionale dei Lincei, Roma, 1965, pp. 977-988. The paper was later reprinted in Lect. Appl. Math. **15**, 143–156 (1974) and inside ref. [16].
- [16] G. Gallavotti (Ed.): *The Fermi-Pasta-Ulam Problem: A Status Report*, Lect. Notes Phys. **728**, Springer, Berlin-Heidelberg, 2008.
- [17] M. Gallone, A. Ponno, B. Rink, *Hydrodynamics of the FPU problem and its integrable aspects*; preprint 2018.
- [18] C.S. Gardner, J.M. Greene, M.D. Kruskal and R.M. Miura, *Method for solving the Korteweg-de Vries equation*, Phys. Rev. Lett. **19**, 1095–1097 (1967).
- [19] T. Grava, A. Maspero, G. Mazzuca and A. Ponno, *Adiabatic invariants for the FPUT and Toda chain in the thermodynamic limit*; preprint, submitted (2020); <https://arxiv.org/abs/2001.08070>
- [20] Y. Hiraoka and Y. Kodama, *Normal Form and Solitons*, in A.V. Mikhailov (ed.), Integrability, LNP **767**, Springer 2009, 175–214.
- [21] T. Kappeler and J. Pöschel, *On the periodic KdV equation in weighted Sobolev spaces*, Ann. I.H. Poincaré - AN **26**, 841–853 (2009).
- [22] P.D. Lax, *Integrals of Nonlinear Equations of Evolution and Solitary Waves*, Comm. Pure Appl. Math. **21**, 467–490 (1968).
- [23] S.V. Manakov, *Complete integrability and stochastization of discrete dynamical systems*, Sov. Phys. JEPT **40**, 269–274 (1974).
- [24] R.M. Miura, C.S. Gardner and M.D. Kruskal, *Korteweg-de Vries equation and generalization, II. Existence of conservation laws and constants of motion*, J. Math. Phys. **9**, 1204–1209 (1968).
- [25] A. Ponno and D. Bambusi, *Korteweg - de Vries equation and energy sharing in Fermi-Pasta-Ulam*, Chaos **15**, 015107/1-5 (2005).
- [26] B. Rink, *Proof of Nishida’s conjecture on anharmonic lattices*, Comm. Math. Phys. **261**, 613–627 (2006).

- [27] B. Rink, *Fermi-Pasta-Ulam systems (FPU): mathematical aspects*, Scholarpedia **4**, 9217 (2009).
- [28] R. Temam, *Inertial Manifolds*, The Mathematical Intelligencer **12**, 68–74 (1990).
- [29] G.B. Whitham, *Linear and Nonlinear Waves*, John Wiley & Sons, 1974.
- [30] N.J. Zabusky and M.D. Kruskal, *Interaction of “solitons” in a collisionless plasma and the recurrence of initial states*, Phys. Rev. Lett. **15**, 240–243 (1965).
- [31] V.E. Zakharov and L.D. Feddeev, *Korteweg-de Vries Equation: a completely integrable Hamiltonian system*, Funct. Analysis Appl. **5**, 280–286 (1971).
- [32] V.E. Zakharov, *On stochastization of one dimensional chains of nonlinear oscillators*, Sov. Phys. JETP **38**, 108–110 (1974).