

# Acute Hemorrhagic Edema of Infancy With Associated Hemorrhagic Lacrimation

Gregorio P. **Milani**<sup>1,2</sup>, Mario G. **Bianchetti**<sup>3,4</sup>,  
Alessandra **Ferrarini**<sup>3</sup>, Pietro O. **Rinoldi**<sup>3</sup>,  
Sebastiano A. G. **Lava**<sup>5</sup>

<sup>1</sup> Pediatric Unit, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy;

<sup>2</sup> Department of Clinical Sciences and Community Health, Università degli Studi di Milano, Milan, Italy;

<sup>3</sup> Pediatric Institute of Southern Switzerland, Ospedale San Giovanni, Bellinzona, Switzerland;

<sup>4</sup> Università della Svizzera Italiana, Lugano, Switzerland;

<sup>5</sup> Pediatric Cardiology Unit, Department of Pediatrics, Centre Hospitalier Universitaire Vaudois (CHUV), 1010 Lausanne, Switzerland.

Correspondence: Dr. Sebastiano A. G. Lava, MD MSc, Pediatric Cardiology Unit, Department of Pediatrics, Centre Hospitalier Universitaire Vaudois (CHUV), 1010 Lausanne, Switzerland. Email: [webmaster@sebastianolava.ch](mailto:webmaster@sebastianolava.ch)

Sir:

In the interesting case of acute hemorrhagic edema of young children with associated hemorrhagic lacrimation reported by Sneller H et al. [1], the authors state that no more than one other case [2] of hemorrhagic lacrimation has been seen in infants and children affected by this skin limited small-vessel leukocytoclastic vasculitis [3].

We recently started to develop **AHEBI**, the **acute hemorrhagic edema bibliographic database**, which should progressively integrate all the original articles on acute hemorrhagic edema published since 1913 in Dutch, English, French, German, Italian Portuguese and Spanish. Using the mentioned database, we found 3 further cases of hemorrhagic lacrimation in children affected by acute hemorrhagic edema published in 1972 [4], 2004 [5] and 2018 [6]. It is therefore concluded that in acute hemorrhagic edema, hemorrhagic lacrimation is an uncommon but not exceptional complication that has been so far documented in at least 5 cases [1, 2, 4-6]: 3 girls and 2 boys aged from 5 to 11, median 6 months.

## References

1. Sneller H, Vega C, Zemel L, Chicaiza HP. Acute hemorrhagic edema of infancy with associated hemorrhagic lacrimation. *Pediatr Emerg Care*. doi: 10.1097/PEC.0000000000001542.
2. Mreish S, Al-Tatari H. Hemorrhagic lacrimation and epistaxis in acute hemorrhagic edema of infancy. *Case Rep Pediatr*. 2016;2016:9762185.
3. Lava SAG, Milani GP, Fossali EF, Simonetti GD, Agostoni C, Bianchetti MG. Cutaneous manifestations of small-vessel leukocytoclastic vasculitides in childhood. *Clin Rev Allergy Immunol*. 2017;53:439-451.
4. Verdura G, Lupi G. Porpora vascolare (anafilattoide) acuta in lattante di sei mesi. Confini nosologici della sindrome.

*Minerva Pediatr.* 1972;24:519-525.

5. Di Lernia V, Lombardi M, Lo Scocco G. Infantile acute hemorrhagic edema and rotavirus infection. *Pediatr Dermatol.* 2004;21:548-50.

6. Manzano G, Shantharam R, Webb E, Finelt N, Hengel K. Hemolacria, hematochezia, and hematuria in an 11-month-old boy. *Pediatr Rev.* 2018;39(8):418-420.