

The motivic Hopf map solves the homotopy limit problem for K -theory

Oliver Röndigs, Markus Spitzweck, Paul Arne Østvær

January 20, 2017

Abstract

We solve affirmatively the homotopy limit problem for K -theory over fields of finite virtual cohomological dimension. Our solution employs the motivic slice filtration and the first motivic Hopf map.

1 Introduction

A homotopy limit problem asks for an equivalence between fixed points and homotopy fixed points for a group action [30]. In some contexts, the fixed points are easily described, and one then obtains a description of the otherwise intractable homotopy fixed points. Many distinguished results in algebraic topology take the form of a homotopy limit problem, e.g., the Atiyah-Segal completion theorem linking equivariant K -theory to representation theory, Segal's Burnside ring conjecture on stable cohomotopy, Sullivan's conjecture on the homotopy type of real points of algebraic varieties, and the Quillen-Lichtenbaum conjecture on Galois descent for algebraic K -theory under field extensions.

In this paper we give a surprising solution of the homotopy limit problem for K -theory in the stable motivic homotopy category. This is achieved by analyzing the slice filtration for algebraic and hermitian K -theory [25], [26], [32], and completing with respect to the Hopf element η in the Milnor-Witt K -theory ring [15].

To provide context for our approach, recall that complex conjugation of vector bundles gives rise to the Adams operation Ψ^{-1} and an action of the group C_2 of order two on the complex K -theory spectrum \mathbf{KU} . Atiyah [1] shows there is an isomorphism

$$\mathbf{KO} \xrightarrow{\simeq} \mathbf{KU}^{hC_2} \tag{1.1}$$

between the real K -theory spectrum \mathbf{KO} and the C_2 -homotopy fixed points of \mathbf{KU} . For the corresponding connective K -theory spectra, the homotopy cofiber of

$$\mathbf{ko} \longrightarrow \mathbf{ku}^{hC_2} \tag{1.2}$$

is an infinite sum $\bigvee_{i < 0} \Sigma^{4i} \mathbf{HZ}/2$ of suspensions of the mod-2 Eilenberg-MacLane spectrum.

We are interested in the analogues of (1.1) and (1.2) for algebraic K -theory \mathbf{KGL} with C_2 -action given by the Adams operation Ψ^{-1} and hermitian K -theory \mathbf{KQ} , see for example [25, §3,4]. Throughout we work in the stable motivic homotopy category \mathbf{SH} over a field F of characteristic $\text{char}(F) \neq 2$. Our starting point is, somewhat unexpectedly in view of (1.2), the naturally induced map between fixed points and homotopy fixed points

$$\gamma: \mathbf{kq} \longrightarrow \mathbf{kgl}^{hC_2} \quad (1.3)$$

for the projections of \mathbf{KGL} and \mathbf{KQ} to the effective stable motivic homotopy category \mathbf{SH}^{eff} . The latter is the localizing subcategory of \mathbf{SH} generated by suspension spectra of smooth schemes [32]. Let η be the first motivic Hopf map induced by the natural map of algebraic varieties $\mathbf{A}^2 \setminus \{0\} \longrightarrow \mathbf{P}^1$. Recall that η defines a non-nilpotent element in the homotopy group $\pi_{1,1}\mathbf{1}$ of the motivic sphere spectrum. We let $\pi_*\mathbf{E}$ denote the bigraded coefficients of a generic motivic spectrum \mathbf{E} . The stable cone of η acquires a Bousfield localization functor \mathbf{L}_η defined on all motivic spectra [24, Appendix A]. Let $\text{vcd}_2(F)$ denote the mod-2 cohomological dimension of the absolute Galois group of $F(\sqrt{-1})$ [27, Chapter 1, §3]. We solve the homotopy limit problem (1.3) affirmatively by completing with respect to η .

Theorem 1.1: *Suppose F is a field of $\text{char}(F) \neq 2$ and virtual cohomological dimension $\text{vcd}_2(F) < \infty$. Then (1.3) induces an isomorphism*

$$\mathbf{L}_\eta(\gamma): \mathbf{kq}_\eta^\wedge \xrightarrow{\simeq} \mathbf{kgl}^{hC_2}. \quad (1.4)$$

We show that the homotopy fixed point spectrum \mathbf{kgl}^{hC_2} in (1.4) is η -complete. The proof of Theorem 1.1 invokes the slice filtration

$$\dots \subset \Sigma_T^{q+1}\mathbf{SH}^{\text{eff}} \subset \Sigma_T^q\mathbf{SH}^{\text{eff}} \subset \Sigma_T^{q-1}\mathbf{SH}^{\text{eff}} \subset \dots \quad (1.5)$$

introduced by Voevodsky [32, §2]. Using (1.5) one associates to \mathbf{E} an integrally graded family of slices $\mathfrak{s}_*(\mathbf{E})$ and a trigraded slice spectral sequence

$$\pi_*\mathfrak{s}_*(\mathbf{E}) \implies \pi_*\mathbf{E}. \quad (1.6)$$

We show that (1.6) converges conditionally for the η -completion of \mathbf{kq} and also for the homotopy fixed point spectrum \mathbf{kgl}^{hC_2} . In contrast to the topological situation (1.1), $\pi_*\mathbf{KQ}$ and $\pi_*\mathbf{KGL}$ are unknown over general fields. Nonetheless we obtain a proof of Theorem 1.1 by using the computations of $\mathfrak{s}_*(\mathbf{KQ})$ and $\mathfrak{s}_*(\mathbf{KGL}^{hC_2})$ accomplished in [25].

Next we turn to solving the homotopy limit problem

$$\Upsilon: \mathbf{KQ} \longrightarrow \mathbf{KGL}^{hC_2}. \quad (1.7)$$

Here \mathbf{KGL}^{hC_2} is η -complete essentially due to motivic orientability of algebraic K -theory. We proceed by comparing with the effective cocovers of \mathbf{KQ} and \mathbf{KGL}^{hC_2} . Remarkably, the first motivic Hopf map η turns Υ into an isomorphism without altering the target in (1.7).

Theorem 1.2: *Suppose F is a field of $\text{char}(F) \neq 2$ and virtual cohomological dimension $\text{vcd}_2(F) < \infty$. Then (1.7) induces an isomorphism*

$$\mathbf{L}_\eta(\Upsilon): \mathbf{KQ}_\eta^\wedge \xrightarrow{\cong} \mathbf{KGL}^{hC_2}. \quad (1.8)$$

Supplementing our main results we note that the η -arithmetic square

$$\begin{array}{ccc} \mathbf{KQ} & \longrightarrow & \mathbf{KW} \\ \downarrow & & \downarrow \\ \mathbf{KQ}_\eta^\wedge & \longrightarrow & \mathbf{KQ}_\eta^\wedge[\eta^{-1}] \end{array} \quad (1.9)$$

for \mathbf{KQ} [26, §3.1] coincides up to isomorphism with the Tate diagram [9, (20)]

$$\begin{array}{ccc} \mathbf{KQ} & \longrightarrow & \mathbf{KW} \\ \downarrow & & \downarrow \\ \mathbf{KGL}^{hC_2} & \longrightarrow & \mathbf{KGL}^{tC_2} \end{array} \quad (1.10)$$

for the C_2 -action on \mathbf{KGL} . Here \mathbf{KW} denotes the higher Witt-theory and \mathbf{KGL}^{tC_2} denotes the Tate K -theory spectrum. Moreover, by representability, (1.8) implies that for every $X \in \mathbf{Sm}_F$ — smooth F -schemes of finite type — there is a naturally induced isomorphism

$$\mathbf{L}_\eta(\Upsilon)_\star: \pi_\star \mathbf{KQ}(X)_\eta^\wedge \xrightarrow{\cong} \pi_\star \mathbf{KGL}(X)^{hC_2}.$$

Remark 1.3: *The earlier works [2] and [9] identified the 2-adic completion of the homotopy fixed points by showing an isomorphism $\pi_\star \mathbf{KQ}/2 \xrightarrow{\cong} \pi_\star \mathbf{KGL}^{hC_2}/2$. Explicit calculations are carried out over the complex numbers \mathbb{C} in [10]. However, for the identification of \mathbf{KGL}^{hC_2} in (1.8) it is paramount to work in \mathbf{SH} , so that the η -completion of \mathbf{KQ} makes sense. The commonplace assumption $\text{vcd}_2(F) < \infty$ is also used in [2], [9], and in the context of the Quillen-Lichtenbaum conjecture for étale K -theory [12, §4].*

2 The first motivic Hopf map η

We view $\mathbf{A}^2 \setminus \{0\}$ and \mathbf{P}^1 as motivic spaces pointed at $(1, 1)$ and $[1 : 1]$, respectively. The canonical projection map $\mathbf{A}^2 \setminus \{0\} \rightarrow \mathbf{P}^1$ induces the stable motivic Hopf map $\eta: \mathbf{G}_m \rightarrow \mathbf{1}$ for the motivic sphere spectrum $\mathbf{1}$. Iteration of η yields the cofiber sequence

$$\mathbf{G}_m^{\wedge n} \xrightarrow{\eta^n} \mathbf{1} \rightarrow \mathbf{1}/\eta^n. \quad (2.1)$$

The η -completion \mathbf{E}_η^\wedge of a motivic spectrum \mathbf{E} is defined as the homotopy limit $\text{holim}_{n \rightarrow \infty} \mathbf{E}/\eta^n$ of the canonically induced diagram

$$\dots \rightarrow \mathbf{E}/\eta^{n+1} \rightarrow \mathbf{E}/\eta^n \rightarrow \dots \rightarrow \mathbf{E}/\eta. \quad (2.2)$$

By (2.1) and (2.2) there is a naturally induced map

$$\mathbf{E} \longrightarrow \mathbf{E}_\eta^\wedge. \quad (2.3)$$

We say that \mathbf{E} is η -complete if the map in (2.3) is an isomorphism. The Bousfield localization $\mathbf{L}_\eta \mathbf{E}$ of \mathbf{E} for the cone of η coincides with \mathbf{E}_η^\wedge . Recall that the algebraic cobordism spectrum \mathbf{MGL} is the universal oriented motivic spectrum, see [16], [18].

Lemma 2.1: *Every module over an oriented motivic ring spectrum is η -complete.*

Proof. The unit map for algebraic cobordism $\mathbf{1} \longrightarrow \mathbf{MGL}$ factors through the cone $\mathbf{1}/\eta$, see [26, Lemma 3.24] for an explicit factorization, which implies $\mathbf{MGL} \wedge \eta = 0$. The statement for modules follows readily. \square

3 The slice filtration

In this section we discuss results for the slice filtration [32] which will be applied in the proofs of our main results in Section 5. Throughout we work over a base field F .

To (1.5) one associates distinguished triangles

$$f_{q+1}(\mathbf{E}) \longrightarrow f_q(\mathbf{E}) \longrightarrow s_q(\mathbf{E}), \quad (3.1)$$

for every motivic spectrum \mathbf{E} , see [32, Theorem 2.2]. Here the q th effective cover of \mathbf{E} is the universal map $f_q(\mathbf{E}) \longrightarrow \mathbf{E}$ from $\Sigma_T^q \mathbf{SH}^{\text{eff}}$ to \mathbf{E} . The q th slice $s_q(\mathbf{E}) \in \Sigma_T^q \mathbf{SH}^{\text{eff}}$ is uniquely determined up to isomorphism by (3.1). Every object of $\Sigma_T^{q+1} \mathbf{SH}^{\text{eff}}$ maps trivially to $s_q(\mathbf{E})$. It is technically important for many constructions to have a “strict model” for the slice filtration, e.g., by means of model categories as in [6, §3.1], [20, §3.2]. We note that $f_q s_{q'} \simeq s_{q'} f_q$ follows from [6, (2.2), §6] for all $q, q' \in \mathbb{Z}$.

Lemma 3.1: *The slice filtration is exhaustive in the sense that there is an isomorphism*

$$\text{hocolim}_{q \rightarrow -\infty} f_q(\mathbf{E}) \xrightarrow{\simeq} \mathbf{E}. \quad (3.2)$$

Proof. Each generator $\Sigma^{s,t} X_+$ of \mathbf{SH} is contained in $\Sigma_T^{q'} \mathbf{SH}^{\text{eff}}$ for some $q' \in \mathbb{Z}$. Here $s, t \in \mathbb{Z}$ and $X \in \mathbf{Sm}_F$, see for example [4, Theorem 9.1]. Recall that $f_{q'}$ preserves homotopy colimits [28, Corollary 4.5], [32, Lemma 4.2]. By the universal property of the q' th effective cover it suffices to show there is an isomorphism

$$\mathbf{SH}(\Sigma^{s,t} X_+, \text{hocolim}_{q < q'} f_{q'} f_q(\mathbf{E})) \xrightarrow{\simeq} \mathbf{SH}(\Sigma^{s,t} X_+, f_{q'}(\mathbf{E})).$$

This follows since $f_{q'} f_q \simeq f_{q'}$ for $q < q'$. \square

Lemma 3.2: *The slices of a motivic spectrum are η -complete.*

Proof. Every slice $s_q(\mathbf{E})$ is a module over the motivic ring spectrum $s_0(\mathbf{1})$, cf. [6, §6 (iv),(v)] and [20, Theorem 3.6.13(6)]. If F is a perfect field, then $s_0(\mathbf{1})$ is the motivic cohomology spectrum \mathbf{MZ} by [14, Theorem 10.5.1] and [34, Theorem 6.6]. This extends by base change; every field is essentially smooth over a perfect field [8, Lemma 2.9], and [8, Lemma 2.7(1)] verifies the hypothesis of [21, Theorem 2.12] for an essentially smooth map. To conclude the proof we use Lemma 2.1 and the canonical orientation on \mathbf{MZ} [17, §10]. \square

Corollary 3.3: *Algebraic K-theory \mathbf{KGL} and its effective cover \mathbf{kgl} are η -complete.*

Proof. This follows from Lemma 2.1 by using the orientation map $\mathbf{MGL} \rightarrow \mathbf{KGL}$, see for example [19, Example 2.4] and [29, Examples 2.1, 2.2], and the geometric fact that the algebraic cobordism spectrum is effective [28, Corollary 3.2], [32, §8]. \square

Corollary 3.4: *The homotopy fixed points spectra \mathbf{KGL}^{hC_2} and \mathbf{kgl}^{hC_2} are η -complete.*

Proof. Let \mathbf{E} be short for \mathbf{KGL} or \mathbf{kgl} . We use homotopy limits to model the homotopy fixed points \mathbf{E}^{hC_2} for the C_2 -action given by the Adams operation Ψ^{-1} [7, §18], [25, §3.4]. Corollary 3.3 implies there is an isomorphism

$$\mathrm{holim}_{C_2} \mathbf{E} \xrightarrow{\simeq} \mathrm{holim}_{C_2} \mathrm{holim}_{n \rightarrow \infty} \mathbf{E}/\eta^n.$$

Thus the corollary follows by commuting homotopy limits over small categories, i.e.,

$$\mathrm{holim}_{C_2} \mathrm{holim}_{n \rightarrow \infty} \mathbf{E}/\eta^n \xrightarrow{\simeq} \mathrm{holim}_{n \rightarrow \infty} \mathrm{holim}_{C_2} \mathbf{E}/\eta^n.$$

\square

The q th effective cocover $f^{q-1}(\mathbf{E})$ of \mathbf{E} is uniquely determined up to isomorphism by the distinguished triangle

$$f_q(\mathbf{E}) \longrightarrow \mathbf{E} \longrightarrow f^{q-1}(\mathbf{E}). \quad (3.3)$$

Note that $f^{q-1}(\mathbf{E})$ is a $(q-1)$ -coeffective motivic spectrum, i.e., it is an object of the right orthogonal subcategory of $\Sigma_T^q \mathbf{SH}^{\mathrm{eff}}$. If $q \leq q'$ the isomorphism

$$s_{q'} f_q(\mathbf{E}) \xrightarrow{\simeq} s_{q'}(\mathbf{E}) \quad (3.4)$$

implies $s_{q'} f^{q-1}(\mathbf{E}) \simeq *$. When $q = 0$, (3.3) yields a distinguished triangle for the effective cover $\mathbf{e} \in \mathbf{SH}^{\mathrm{eff}}$ of \mathbf{E} , i.e.,

$$\mathbf{e} \longrightarrow \mathbf{E} \longrightarrow f^{-1}(\mathbf{E}). \quad (3.5)$$

We note that all the nonnegative slices of the coeffective motivic spectrum $f^{-1}(\mathbf{E})$ are trivial.

Lemma 3.5: *If \mathbf{E} has no nontrivial negative slices then $\mathbf{E} \in \mathbf{SH}^{\mathrm{eff}}$.*

Proof. Using (3.1), (3.2) and (3.5) it follows that $f^{-1}(\mathbf{E}) \simeq *$. \square

Lemma 3.6: Suppose $\mathbf{E} \rightarrow \mathbf{F}$ induces an isomorphism $\mathfrak{s}_q \mathfrak{f}^{q-1}(\mathbf{E}) \xrightarrow{\cong} \mathfrak{s}_q \mathfrak{f}^{q-1}(\mathbf{F})$ for all $q \in \mathbb{Z}$. Then there is a naturally induced isomorphism $\mathfrak{f}^{q-1}(\mathbf{E}) \xrightarrow{\cong} \mathfrak{f}^{q-1}(\mathbf{F})$.

Proof. This follows by applying (3.1) and (3.2) to the effective cocovers. \square

Lemma 3.7: For $n > 0$ there is a distinguished triangle

$$\mathfrak{f}_{-n+1} \mathfrak{f}^{-1}(\mathbf{E}) \longrightarrow \mathfrak{f}_{-n} \mathfrak{f}^{-1}(\mathbf{E}) \longrightarrow \mathfrak{s}_{-n}(\mathbf{E}). \quad (3.6)$$

It follows that $\mathfrak{f}_{-n} \mathfrak{f}^{-1}(\mathbf{E})$ is a finite extension of the negative slices of \mathbf{E} .

Proof. This follows from the distinguished triangles:

$$\begin{array}{ccccc} \mathfrak{f}_{-n+1}(\mathbf{e}) & \longrightarrow & \mathfrak{f}_{-n+1}(\mathbf{E}) & \longrightarrow & \mathfrak{f}_{-n+1} \mathfrak{f}^{-1}(\mathbf{E}) \\ \downarrow & & \downarrow & & \downarrow \\ \mathfrak{f}_{-n}(\mathbf{e}) & \longrightarrow & \mathfrak{f}_{-n}(\mathbf{E}) & \longrightarrow & \mathfrak{f}_{-n} \mathfrak{f}^{-1}(\mathbf{E}) \\ \downarrow & & \downarrow & & \downarrow \\ \mathfrak{s}_{-n}(\mathbf{e}) & \longrightarrow & \mathfrak{s}_{-n}(\mathbf{E}) & \longrightarrow & \mathfrak{s}_{-n} \mathfrak{f}^{-1}(\mathbf{E}) \end{array} \quad (3.7)$$

In (3.7) the slice $\mathfrak{s}_{-n}(\mathbf{e}) \simeq *$ by the assumption $n > 0$. This implies (3.6). We note the effective cover $\mathfrak{f}_0 \mathfrak{f}^{-1}(\mathbf{E}) \simeq *$ since the map $\mathfrak{f}_0(\mathbf{e}) \rightarrow \mathfrak{f}_0(\mathbf{E})$ is an isomorphism. It follows that $\mathfrak{f}_{-1} \mathfrak{f}^{-1}(\mathbf{E})$ is isomorphic to $\mathfrak{s}_{-1}(\mathbf{E})$. The conclusion follows from (3.6) by induction on n . \square

Remark 3.8: For $n > 0$, $\mathfrak{f}_{-n} \mathfrak{f}^{-1}(\mathbf{E})$ is η -complete by Lemmas 3.2 and 3.7.

Lemma 3.9: For $n \geq -q > 0$ there are isomorphisms

$$\mathfrak{s}_q(\mathbf{E}) \xrightarrow{\cong} \mathfrak{s}_q \mathfrak{f}^{-1}(\mathbf{E}) \xleftarrow{\cong} \mathfrak{s}_q \mathfrak{f}_{-n} \mathfrak{f}^{-1}(\mathbf{E}). \quad (3.8)$$

Proof. Here we use that $\mathfrak{s}_q(\mathbf{e}) \simeq *$ for $q < 0$. The isomorphism for \mathfrak{f}_{-n} is a special case of (3.4). \square

For every $\mathbf{E} \in \mathbf{SH}$ the distinguished triangle (3.3) yields a commutative diagram:

$$\begin{array}{ccccc} \mathfrak{f}_{q+1}(\mathbf{E}) & \longrightarrow & \mathbf{E} & \longrightarrow & \mathfrak{f}^q(\mathbf{E}) \\ \downarrow & & \parallel & & \downarrow \\ \mathfrak{f}_q(\mathbf{E}) & \longrightarrow & \mathbf{E} & \longrightarrow & \mathfrak{f}^{q-1}(\mathbf{E}) \end{array} \quad (3.9)$$

The slice completion of \mathbf{E} is defined as the homotopy limit

$$\mathrm{sc}(\mathbf{E}) \equiv \mathrm{holim}_{q \rightarrow \infty} \mathfrak{f}^{q-1}(\mathbf{E}). \quad (3.10)$$

Using (3.9) and (3.10) we conclude there is a distinguished triangle

$$\operatorname{holim}_{q \rightarrow \infty} f_q(\mathbf{E}) \longrightarrow \mathbf{E} \longrightarrow \operatorname{sc}(\mathbf{E}). \quad (3.11)$$

We say that \mathbf{E} is *slice complete* if the homotopy limit $\operatorname{holim}_{q \rightarrow \infty} f_q(\mathbf{E})$ in (3.11) is contractible.

Lemma 3.10: *For every $\mathbf{E} \in \mathbf{SH}$, both $s_q(\mathbf{E})$ and $f^q(\mathbf{E})$ are slice complete for all $q \in \mathbb{Z}$.*

Proof. If $q < q'$ there are distinguished triangles

$$f_{q+1}f_{q'}(\mathbf{E}) \xrightarrow{\simeq} f_q f_{q'}(\mathbf{E}) \longrightarrow s_q f_{q'}(\mathbf{E}), \quad f_{q'} f_{q+1}(\mathbf{E}) \xrightarrow{\simeq} f_{q'}(\mathbf{E}) \longrightarrow f_{q'} f^q(\mathbf{E}). \quad (3.12)$$

It follows that $f_{q'} s_q(\mathbf{E}) \simeq s_q f_{q'}(\mathbf{E}) \simeq *$ and $f_{q'} f^q(\mathbf{E}) \simeq *$ for $q < q'$ by (3.12). \square

Lemma 3.11: *Algebraic K-theory \mathbf{KGL} and its effective cover \mathbf{kgl} are slice complete.*

Proof. It suffices to consider \mathbf{KGL} . The associated Nisnevich sheaf of homotopy groups $\pi_{p,q} \mathbf{KGL}$ is trivial when $p < 2q$. Hence $f_q(\mathbf{KGL})$ is q -connected by [26, Lemma 3.17], i.e., for every triple (s, t, d) of integers with $s - t + d < q$ and every $X \in \mathbf{Sm}_F$ of dimension $\leq d$, the group $[\Sigma^{s,t} X_+, f_q(\mathbf{KGL})]$ is trivial. We conclude by letting $q \rightarrow \infty$. \square

From Lemma 3.11 we deduce isomorphisms

$$\mathbf{KGL}^{hC_2} \xrightarrow{\simeq} \operatorname{sc}(\mathbf{KGL})^{hC_2}, \quad \mathbf{kgl}^{hC_2} \xrightarrow{\simeq} \operatorname{sc}(\mathbf{kgl})^{hC_2}.$$

However, it is unclear whether \mathbf{KGL}^{hC_2} and \mathbf{kgl}^{hC_2} are slice complete because homotopy fixed points need not commute with effective cocovers or equivalently with effective covers. To emphasize this issue we construct an example in §6, see also Proposition 3.15.

For $n > 0$ there is a naturally induced distinguished triangle

$$\mathbf{e} \longrightarrow f_{-n}(\mathbf{E}) \longrightarrow f_{-n} f^{-1}(\mathbf{E}). \quad (3.13)$$

Lemma 3.1, (3.5), and (3.13) imply there is a naturally induced isomorphism

$$\operatorname{hocolim}_{n > 0} f_{-n} f^{-1}(\mathbf{E}) \xrightarrow{\simeq} f^{-1}(\mathbf{E}). \quad (3.14)$$

Next we make precise the vagary of identifying the homotopy fixed points $f^{-1}(\mathbf{KGL})^{hC_2}$ with a homotopy colimit. That is, we identify a homotopy limit with a homotopy colimit. Throughout we let \mathbf{E} be a motivic spectrum equipped with a G -action for a finite group G .

Lemma 3.12: *There is a natural isomorphism*

$$\operatorname{hocolim}_{n > 0} f_{-n} f^{-1}(\mathbf{E})^{hG} \xrightarrow{\simeq} f^{-1}(\mathbf{E})^{hG}. \quad (3.15)$$

Proof. For every generator $\Sigma^{s,t}X_+$ of \mathbf{SH} there is a canonically induced map

$$\mathbf{SH}(\Sigma^{s,t}X_+, \operatorname{hocolim}_{n>0} f_{-n}f^{-1}(\mathbf{E})^{hG}) \longrightarrow \mathbf{SH}(\Sigma^{s,t}X_+, f^{-1}(\mathbf{E})^{hG}). \quad (3.16)$$

If $t \geq 0$ the source and target of (3.16) are trivial. If $t < 0$ we show (3.16) is an isomorphism by using the distinguished triangle

$$f_t f^{-1}(\mathbf{E})^{hG} \longrightarrow f^{-1}(\mathbf{E})^{hG} \longrightarrow f^{t-1}(\mathbf{E})^{hG}, \quad (3.17)$$

obtained by applying homotopy fixed points to (3.3) for $f^{-1}(\mathbf{E})$ and identifying $f^{t-1}f^{-1}(\mathbf{E})$ with $f^{t-1}(\mathbf{E})$ by means of the distinguished triangles

$$f_t(\mathbf{e}) \xrightarrow{\simeq} \mathbf{e} \longrightarrow f^{t-1}(\mathbf{e}), \quad f^{t-1}(\mathbf{e}) \longrightarrow f^{t-1}(\mathbf{E}) \xrightarrow{\simeq} f^{t-1}f^{-1}(\mathbf{E}).$$

Since $f^{t-1}(\mathbf{E})^{hG}$ in (3.17) is $(t-1)$ -coeffective there is a canonically induced isomorphism

$$\mathbf{SH}(\Sigma^{s,t}X_+, f_t f^{-1}(\mathbf{E})^{hG}) \xrightarrow{\cong} \mathbf{SH}(\Sigma^{s,t}X_+, f^{-1}(\mathbf{E})^{hG}).$$

On the other hand there are canonical identifications

$$\begin{aligned} \mathbf{SH}(\Sigma^{s,t}X_+, \operatorname{hocolim}_{n>0} f_{-n}f^{-1}(\mathbf{E})^{hG}) &\cong \operatorname{colim}_{n>0} \mathbf{SH}(\Sigma^{s,t}X_+, f_{-n}f^{-1}(\mathbf{E})^{hG}) \\ &\cong \mathbf{SH}(\Sigma^{s,t}X_+, f_t f^{-1}(\mathbf{E})^{hG}). \end{aligned}$$

□

In the following we make the standing assumption that for all $q \in \mathbb{Z}$ there is a naturally induced isomorphism

$$s_q(\mathbf{E}^{hG}) \xrightarrow{\simeq} s_q(\mathbf{E})^{hG}. \quad (3.18)$$

The map in (3.18) arises from the standard adjunction between motivic spectra and “naive” G -motivic spectra. That is, with the trivial G -action on the homotopy fixed points there is a naturally induced G -map $s_q(\mathbf{E}^{hG}) \rightarrow s_q(\mathbf{E})$. Its adjoint is the map in (3.18).

Corollary 3.13: *Assuming (3.18) and $n > 0$ there is a naturally induced isomorphism*

$$f_{-n}f^{-1}(\mathbf{E}^{hG}) \xrightarrow{\simeq} f_{-n}f^{-1}(\mathbf{E})^{hG}. \quad (3.19)$$

Proof. Follows from Lemma 3.7 under the stated assumptions. □

Corollary 3.14: *Assuming (3.18) and $n \geq -q > 0$ there is a naturally induced isomorphism*

$$s_q(\mathbf{E}^{hG}) \xrightarrow{\simeq} s_q(\operatorname{hocolim}_{n>0} f_{-n}f^{-1}(\mathbf{E})^{hG}). \quad (3.20)$$

Proof. From the isomorphisms (3.8) in Lemma 3.9 we obtain

$$s_q(\mathbf{E}^{hG}) \xrightarrow{\simeq} s_q f^{-1}(\mathbf{E}^{hG}) \xleftarrow{\simeq} s_q f_{-n}f^{-1}(\mathbf{E}^{hG}). \quad (3.21)$$

Recall that slices commute with homotopy colimits [28, Corollary 4.5], [32, Lemma 4.2]. Thus the target in (3.20) identifies with the homotopy colimit

$$\operatorname{hocolim}_{n>0} s_q f_{-n} f^{-1}(\mathbf{E})^{hG}. \quad (3.22)$$

With the assumption $n \geq -q > 0$ the q th slice $s_q f_{-n} f^{-1}(\mathbf{E})^{hG}$ maps isomorphically to (3.22). It remains to apply the isomorphism (3.19) in Corollary 3.13 and (3.21). \square

Proposition 3.15: *Assuming (3.18) the slices of \mathbf{e} commute with homotopy fixed points in the sense that there is a naturally induced isomorphism*

$$s_q(\mathbf{e}^{hG}) \xrightarrow{\cong} s_q(\mathbf{e})^{hG} \quad (3.23)$$

for every $q \in \mathbb{Z}$. Moreover, \mathbf{e}^{hG} is an effective motivic spectrum.

Proof. Applying homotopy fixed points to (3.3) yields the distinguished triangle

$$\mathbf{e}^{hG} \longrightarrow \mathbf{E}^{hG} \longrightarrow f^{-1}(\mathbf{E})^{hG}. \quad (3.24)$$

From (3.24) we deduce the commutative diagram of distinguished triangles:

$$\begin{array}{ccccc} s_q(\mathbf{e}^{hG}) & \longrightarrow & s_q(\mathbf{E}^{hG}) & \longrightarrow & s_q(f^{-1}(\mathbf{E})^{hG}) \\ \downarrow & & \downarrow & & \downarrow \\ s_q(\mathbf{e})^{hG} & \longrightarrow & s_q(\mathbf{E})^{hG} & \longrightarrow & s_q(f^{-1}(\mathbf{E}))^{hG} \end{array} \quad (3.25)$$

When $q \geq 0$ it follows that $s_q(f^{-1}(\mathbf{E})) \simeq s_q(f^{-1}(\mathbf{E})^{hG}) \simeq *$ since homotopy limits preserve coeffective motivic spectra. Since the middle vertical map in (3.25) is an isomorphism, see the assumption (3.18), so is (3.23).

When $q < 0$, (3.15) and (3.20) imply that $s_q(\mathbf{E}^{hG}) \longrightarrow s_q(f^{-1}(\mathbf{E})^{hG})$ is an isomorphism. Lemma 3.5 implies $\mathbf{e}^{hG} \in \mathbf{SH}^{\text{eff}}$ and thus $s_q(\mathbf{e}^{hG}) \longrightarrow s_q(\mathbf{e})^{hG}$ is an isomorphism. \square

Lemma 3.16: *Assuming (3.18) there are naturally induced isomorphisms*

$$f^q(\mathbf{e}^{hG}) \xrightarrow{\cong} f^q(\mathbf{e})^{hG}, \quad f_q(\mathbf{e}^{hG}) \xrightarrow{\cong} f_q(\mathbf{e})^{hG}. \quad (3.26)$$

Proof. We show that all the nonnegative effective cocovers of \mathbf{e} commute with homotopy fixed points. With this in hand the assertion for the effective covers of \mathbf{e} follows from (3.3).

We claim there is a commutative diagram:

$$\begin{array}{ccc} s_0(\mathbf{e}^{hG}) & \xrightarrow{\cong} & f^0(\mathbf{e}^{hG}) \\ \cong \downarrow & & \downarrow \\ s_0(\mathbf{e})^{hG} & \xrightarrow{\cong} & f^0(\mathbf{e})^{hG} \end{array} \quad (3.27)$$

Proposition 3.15 shows the left vertical map in (3.27) is an isomorphism and that \mathbf{e}^{hG} is an effective motivic spectrum. Hence $\mathfrak{s}_0(\mathbf{e}) \simeq \mathfrak{f}^0(\mathbf{e})$ and $\mathfrak{s}_0(\mathbf{e}^{hG}) \simeq \mathfrak{f}^0(\mathbf{e}^{hG})$ by comparing (3.1) and (3.3). It follows that the natural map $\mathfrak{f}^0(\mathbf{e}^{hG}) \rightarrow \mathfrak{f}^0(\mathbf{e})^{hG}$ is also an isomorphism.

The cone of the left vertical map in (3.9) is the q th slice. Hence there is a homotopy cofiber sequence $\mathfrak{s}_q(\mathbf{e}) \rightarrow \mathfrak{f}^q(\mathbf{e}) \rightarrow \mathfrak{f}^{q-1}(\mathbf{e})$, and likewise for \mathbf{e}^{hG} . Proposition 3.15 and induction on q implies that $\mathfrak{f}^q(\mathbf{e})$ commutes with homotopy fixed points. \square

Corollary 3.17: *Assuming (3.18) there is a naturally induced isomorphism*

$$\mathrm{sc}(\mathbf{e}^{hG}) \xrightarrow{\simeq} \mathrm{sc}(\mathbf{e})^{hG}. \quad (3.28)$$

If \mathbf{e} is slice complete then so is \mathbf{e}^{hG} .

Proof. Lemma 3.16 and the fact that homotopy limits commute imply there are canonical isomorphisms

$$\mathrm{holim}_{q \rightarrow \infty} \mathfrak{f}^{q-1}(\mathrm{holim}_G \mathbf{e}) \simeq \mathrm{holim}_{q \rightarrow \infty} \mathrm{holim}_G \mathfrak{f}^{q-1}(\mathbf{e}) \simeq \mathrm{holim}_{q \rightarrow \infty} \mathrm{holim}_G \mathfrak{f}^{q-1}(\mathbf{e}) \simeq \mathrm{holim}_G \mathrm{holim}_{q \rightarrow \infty} \mathfrak{f}^{q-1}(\mathbf{e}).$$

For slice completeness of \mathbf{e}^{hG} we use the factorization $\mathbf{e}^{hG} \rightarrow \mathrm{sc}(\mathbf{e}^{hG}) \rightarrow \mathrm{sc}(\mathbf{e})^{hG}$. \square

Corollary 3.18: *Assuming (3.18) there are naturally induced isomorphisms*

$$\mathfrak{f}^q(\mathbf{E}^{hG}) \xrightarrow{\simeq} \mathfrak{f}^q(\mathbf{E})^{hG}, \quad \mathfrak{f}_q(\mathbf{E}^{hG}) \xrightarrow{\simeq} \mathfrak{f}_q(\mathbf{E})^{hG}. \quad (3.29)$$

Proof. There is a naturally induced commutative diagram of distinguished triangles:

$$\begin{array}{ccccc} \mathfrak{f}_0(\mathbf{E}^{hG}) & \longrightarrow & \mathbf{E}^{hG} & \longrightarrow & \mathfrak{f}^{-1}(\mathbf{E}^{hG}) \\ \downarrow & & \parallel & & \downarrow \\ \mathfrak{f}_0(\mathbf{E})^{hG} & \longrightarrow & \mathbf{E}^{hG} & \longrightarrow & \mathfrak{f}^{-1}(\mathbf{E})^{hG} \end{array} \quad (3.30)$$

Proposition 3.15 shows that $\mathfrak{f}_0(\mathbf{E}^{hG}) \rightarrow \mathbf{e}^{hG}$ is a map between effective motivic spectra. It follows that $\mathfrak{f}^{-1}(\mathbf{E}^{hG}) \rightarrow \mathfrak{f}^{-1}(\mathbf{E})^{hG}$ induces an isomorphism on all negative slices. Since it is a map between coeffective spectra, it is in fact an isomorphism according to Lemma 3.6.

The general cases follow by using induction on (3.1) and (3.3). \square

We end this section by discussing G -fixed points in more detail. Let \mathcal{O}_G denote the orbit category of G with objects $\{G/H\}$ and morphisms the G -maps $\mathrm{map}_G(G/H, G/K) \cong (G/K)^H$ [31, §1.8]. Let \mathbf{MSS} be a highly structured model for the stable motivic homotopy category, e.g., motivic functors [5], or motivic symmetric spectra [11]. Let $\mathbf{MSS}^{\mathrm{eff}}$ be the Bousfield colocalization of \mathbf{MSS} with respect to the set of objects $\Sigma^{p,0} \Sigma_{\mathbf{P}^1}^\infty X_+$, where $X \in \mathbf{Sm}_F$ and $p \in \mathbb{Z}$ (it suffices to consider $p \leq 0$). Its homotopy category is $\mathbf{SH}^{\mathrm{eff}}$. As a model for naive G -motivic spectra we use the functor category $[\mathcal{O}_G^{\mathrm{op}}, \mathbf{MSS}]$ with the projective model structure [7, Theorem 11.6.1]. There is a naturally induced Quillen adjunction:

$$i_0^G : [\mathcal{O}_G^{\mathrm{op}}, \mathbf{MSS}^{\mathrm{eff}}] \rightleftarrows [\mathcal{O}_G^{\mathrm{op}}, \mathbf{MSS}] : r_0^G \quad (3.31)$$

Evaluating a naive G -motivic spectrum $\underline{\mathbf{E}}$ at the orbits corresponding to G and the identity element yields the underlying motivic spectrum \mathbf{E} and the G -fixed points \mathbf{E}^G , respectively. Let $\underline{\mathbf{e}}$ be the naive G -motivic spectrum $\mathbf{f}_0^G(\underline{\mathbf{E}})$, where $\mathbf{f}_0^G = \mathbf{L}i_0^G \circ r_0^G$. (Forgetting the G -action, $\underline{\mathbf{e}}$ coincides with \mathbf{e} .) Since evaluating at the identity orbit commutes with (3.31), we obtain an isomorphism

$$\mathbf{f}_0^G(\underline{\mathbf{E}})^G \cong \mathbf{f}_0(\underline{\mathbf{E}}^G). \quad (3.32)$$

In particular, \mathbf{kgl}^{C_2} coincides with the effective hermitian K -theory spectrum \mathbf{kq} , cf. (1.3).

4 The slice spectral sequence

The trigraded slice spectral sequence for \mathbf{E} arising from (1.5) takes the form

$$\pi_* \mathbf{s}_*(\mathbf{E}) \implies \pi_* \mathbf{E}. \quad (4.1)$$

This is an upper half-plane spectral sequence with entering differentials [3, §7] because $\pi_{p,w} \mathbf{s}_q(\mathbf{E}) = 0$ for $q < w$, cf. [32, §7]. A standard argument shows that (4.1) converges conditionally to the motivic homotopy groups of $\mathbf{sc}(\mathbf{E})$ in the sense of [3, Definition 5.10]. For the following result we refer to [26, Lemma 3.14].

Lemma 4.1: *Suppose $\mathbf{e} \in \mathbf{SH}^{\text{eff}}$ and \mathbf{e}/η is slice complete. Then there is a naturally induced isomorphism between \mathbf{e}_η^\wedge to $\mathbf{sc}(\mathbf{e})$.*

Proposition 4.2: *Suppose $\mathbf{e} \in \mathbf{SH}^{\text{eff}}$ and \mathbf{e}/η is slice complete. There is a conditionally convergent slice spectral sequence*

$$\pi_* \mathbf{s}_*(\mathbf{e}) \implies \pi_* \mathbf{e}_\eta^\wedge. \quad (4.2)$$

Proof. This follows from Lemma 4.1 and (4.1). \square

5 Proofs of Theorems 1.1 and 1.2

Corollary 5.1: *For effective hermitian K -theory there is a conditionally convergent slice spectral sequence*

$$\pi_* \mathbf{s}_*(\mathbf{kq}) \implies \pi_* \mathbf{kq}_\eta^\wedge. \quad (5.1)$$

Proof. The only issue is to identify the quotient of \mathbf{kq} by η with a slice complete spectrum. By [25, Theorem 3.4] there is a homotopy cofiber sequence

$$\Sigma^{1,1} \mathbf{KQ} \xrightarrow{\eta} \mathbf{KQ} \longrightarrow \mathbf{KGL} \quad (5.2)$$

relating algebraic and hermitian K -theory via η . Passing to effective covers in (5.2) identifies the cofiber of $\mathbf{f}_0(\eta: \Sigma^{1,1} \mathbf{KQ} \longrightarrow \mathbf{KQ})$ with \mathbf{kgl} . Hence the cofiber of $\eta: \Sigma^{1,1} \mathbf{kq} \longrightarrow \mathbf{kq}$ is an extension of \mathbf{kgl} by $\Sigma^{1,1} \mathbf{s}_{-1}(\mathbf{KQ}) \simeq \mathbf{s}_0(\Sigma^{1,1} \mathbf{KQ})$, cf. [25, Lemma 2.1], so it is slice complete by Lemmas 3.10 and 3.11. This verifies the assumptions in Proposition 4.2 for \mathbf{kq} . \square

Assuming $\text{vcd}_2(F) < \infty$, (3.18) holds for \mathbf{KGL} by [25, Proposition 4.24]. We summarize some useful consequences of the results in §3.

Proposition 5.2: *The following holds when $\text{vcd}_2(F) < \infty$.*

- (1) *The homotopy fixed points spectrum of effective K-theory \mathbf{kgl}^{hC_2} is slice complete. There is a conditionally convergent slice spectral sequence*

$$\pi_* \mathfrak{s}_*(\mathbf{kgl}^{hC_2}) \implies \pi_* \mathbf{kgl}^{hC_2}. \quad (5.3)$$

- (2) *There is a naturally induced isomorphism $f_q(\mathbf{KGL}^{hC_2}) \xrightarrow{\simeq} f_q(\mathbf{KGL})^{hC_2}$.*

Proof. Here (1) follows from Lemma 3.11, Corollary 3.17 and the discussion of (4.1) in §4, while (2) is a special case of Corollary 3.18. \square

Proof of Theorem 1.1. By [25, Theorems 4.18, 4.25, 4.27, Lemma 4.26] the natural map $\Upsilon: \mathbf{KQ} \rightarrow \mathbf{KGL}^{hC_2}$ in (1.7) induces an isomorphism of slices

$$s_q(\Upsilon): s_q(\mathbf{KQ}) \xrightarrow{\simeq} s_q(\mathbf{KGL}^{hC_2}) \simeq \begin{cases} \Sigma^{2q,q}\mathbf{MZ} \vee \bigvee_{i<0} \Sigma^{2q+2i,q}\mathbf{MZ}/2 & q \equiv 0(2), \\ \bigvee_{i<0} \Sigma^{2q+2i+1,q}\mathbf{MZ}/2 & q \equiv 1(2). \end{cases} \quad (5.4)$$

From (5.4) we conclude the natural map $\mathbf{kq} \rightarrow f_0(\mathbf{KGL}^{hC_2})$ induces an isomorphism on slices. By composing with $f_0(\mathbf{KGL}^{hC_2}) \xrightarrow{\simeq} \mathbf{kgl}^{hC_2}$, see Proposition 5.2(2), we conclude there is an isomorphism

$$s_q(\gamma): s_q(\mathbf{kq}) \xrightarrow{\simeq} s_q(\mathbf{kgl}^{hC_2}). \quad (5.5)$$

Thus $\gamma: \mathbf{kq} \rightarrow \mathbf{kgl}^{hC_2}$ in (1.3) induces an isomorphism between the conditionally convergent upper half-plane slice spectral sequences (5.1) and (5.3). The induced map between the filtered target groups is thus an isomorphism [3, Theorem 7.2]. This finishes the proof by passing to Nisnevich sheaves of homotopy groups. \square

Proof of Theorem 1.2. Using (3.5) we obtain the naturally induced commutative diagram of distinguished triangles:

$$\begin{array}{ccccc} \mathbf{kq} & \longrightarrow & \mathbf{KQ} & \longrightarrow & f^{-1}(\mathbf{KQ}) \\ f_0(\Upsilon) \downarrow & & \Upsilon \downarrow & & f^{-1}(\Upsilon) \downarrow \\ f_0(\mathbf{KGL}^{hC_2}) & \longrightarrow & \mathbf{KGL}^{hC_2} & \longrightarrow & f^{-1}(\mathbf{KGL}^{hC_2}) \end{array} \quad (5.6)$$

Lemma 3.6 and (5.4) imply that $f^{-1}(\Upsilon): f^{-1}(\mathbf{KQ}) \rightarrow f^{-1}(\mathbf{KGL}^{hC_2})$ is an isomorphism because it is a map between coeffective spectra and it induces an isomorphism on slices. By composing with the isomorphism $f_0(\mathbf{KGL}^{hC_2}) \xrightarrow{\simeq} \mathbf{kgl}^{hC_2}$ of Proposition 5.2(2) we obtain a commutative diagram of distinguished triangles:

$$\begin{array}{ccccc} \mathbf{kq} & \longrightarrow & \mathbf{KQ} & \longrightarrow & f^{-1}(\mathbf{KQ}) \\ \gamma \downarrow & & \Upsilon \downarrow & & f^{-1}(\Upsilon) \downarrow \simeq \\ \mathbf{kgl}^{hC_2} & \longrightarrow & \mathbf{KGL}^{hC_2} & \longrightarrow & f^{-1}(\mathbf{KGL}^{hC_2}) \end{array} \quad (5.7)$$

This shows that $\mathbf{L}_\eta(\gamma)$ is an isomorphism if and only if $\mathbf{L}_\eta(\Upsilon)$ is an isomorphism. It follows that Theorem 1.1 implies Theorem 1.2. \square

Corollary 5.3: *If $\mathrm{vcd}_2(F) < \infty$ then the η -completion of the Tate K -theory spectrum \mathbf{KGL}^{tC_2} is contractible.*

Proof. Recall that \mathbf{KGL}^{tC_2} is the cone of the norm map from the homotopy orbits \mathbf{KGL}_{hC_2} to the homotopy fixed points \mathbf{KGL}^{hC_2} in the Tate diagram [9, (20)]

$$\begin{array}{ccccc} \mathbf{KGL}_{hC_2} & \longrightarrow & \mathbf{KQ} & \longrightarrow & \mathbf{KW} \\ \parallel & & \downarrow \gamma & & \downarrow \\ \mathbf{KGL}_{hC_2} & \longrightarrow & \mathbf{KGL}^{hC_2} & \longrightarrow & \mathbf{KGL}^{tC_2} \end{array}$$

for the C_2 -action on \mathbf{KGL} . Thus the assertion follows from Theorem 1.2 since the higher Witt-theory spectrum \mathbf{KW} can be identified with $\mathbf{KQ}[\eta^{-1}]$, see e.g., [25, (7)]. \square

6 Appendix

By way of example we show that $\mathbf{SH}^{\mathrm{eff}}$ is not closed under homotopy fixed points. In effect, consider the homotopy fixed points for the trivial C_2 -action on the effective motivic spectrum

$$\bigvee_{i \geq 0} \Sigma^{i,0} \mathbf{MZ}/2 \simeq \prod_{i \geq 0} \Sigma^{i,0} \mathbf{MZ}/2. \quad (6.1)$$

Here the sum and product are isomorphic by [25, Proposition A.5]. Since C_2 -homotopy fixed points commute with products, [25, Lemma 4.22] yields a naturally induced isomorphism

$$\left(\prod_{i \geq 0} \Sigma^{i,0} \mathbf{MZ}/2 \right)^{hC_2} \xrightarrow{\simeq} \prod_{i \geq 0} \prod_{j \geq 0} \Sigma^{i-j,0} \mathbf{MZ}/2. \quad (6.2)$$

Assuming (6.2) is an isomorphism in $\mathbf{SH}^{\mathrm{eff}}$, the countably infinite product $\prod_{n \in \mathbb{N}} \mathbf{MZ}/2$ — corresponding to indices $i = j$ — is effective. However, we show that $\prod_{n \in \mathbb{N}} \mathbf{MZ}/2 \notin \mathbf{SH}^{\mathrm{eff}}$.

Recall from [33, §2] the adjunction between \mathbf{SH} and the stable motivic homotopy category of S^1 -spectra \mathbf{SH}_s :

$$\Sigma_t^\infty : \mathbf{SH}_s \rightleftarrows \mathbf{SH} : \Omega_t^\infty.$$

Now $\Omega_t^\infty \mathbf{MZ}/2$ is the Eilenberg-MacLane S^1 -spectrum $\mathbf{HZ}/2$ associated with the constant presheaf $\mathbf{Z}/2$ by [33, Lemma 5.2]. It follows that $\Omega_t^\infty \prod_{n \in \mathbb{N}} \mathbf{MZ}/2$ is the Eilenberg-MacLane S^1 -spectrum \mathbf{HV} associated with the constant presheaf \mathbf{V} , where \mathbf{V} is a $\mathbf{Z}/2$ -vector space of (uncountable) infinite dimension. If $\prod_{n \in \mathbb{N}} \mathbf{MZ}/2 \in \mathbf{SH}^{\mathrm{eff}}$ we would obtain

$$\Omega_t^\infty \Sigma^{0,1} \prod \mathbf{MZ}/2 \simeq \prod \Omega_t^\infty \Sigma^{0,1} \mathbf{MZ}/2 \in \Sigma_t^1 \mathbf{SH}_s \quad (6.3)$$

since $\Omega_t^\infty \Sigma^{0,1} \mathbf{MZ}/2 \in \Sigma_t^1 \mathbf{SH}_s$ by [14, Theorem 7.4.1] (as conjectured in [33, Conjecture 4]). In (6.3) we use that $\Sigma^{0,1}$, being an equivalence, commutes with products. In \mathbf{SH}_s there is a canonically induced map

$$\alpha: \left(\prod_{n \in \mathbb{N}} \Omega_t^\infty \mathbf{MZ}/2 \right) \otimes_{\Omega_t^\infty \mathbf{MZ}/2} \Omega_t^\infty \Sigma^{0,1} \mathbf{MZ}/2 \longrightarrow \prod_{n \in \mathbb{N}} \Omega_t^\infty \Sigma^{0,1} \mathbf{MZ}/2. \quad (6.4)$$

The tensor product in (6.4) is formed in the module category of the Eilenberg-MacLane S^1 -spectrum $\Omega_t^\infty \mathbf{MZ}/2$ [13], cf. [22], [23], [33, Lemma 5.2]. In particular, the source of α in (6.4) is 1-effective, because $\Omega_t^\infty \Sigma^{0,1} \mathbf{MZ}/2 \in \Sigma_t^1 \mathbf{SH}_s$ by [14, Theorem 7.4.1] and $\prod_{n \in \mathbb{N}} \Omega_t^\infty \mathbf{MZ}/2$ is effective, since it is the Eilenberg-MacLane spectrum associated with \mathbf{V} . We will prove that $\mathbf{SH}_s(\Sigma^{n,0} X_+ \wedge \mathbf{G}_m, \alpha)$ is an isomorphism for every $X \in \mathbf{Sm}_F$ and $n \in \mathbb{Z}$; here \mathbf{G}_m denotes the multiplicative group scheme. In effect, choose an uncountable basis \mathcal{B} of \mathbf{V} and express \mathbf{V} as the filtered colimit of finite dimensional sub- $\mathbf{Z}/2$ -vector spaces $\mathbf{V}' \subset \mathbf{V}$ spanned by finite subsets $\mathcal{F} \subset \mathcal{B}$. Since $\Sigma^{n,0} X_+ \wedge \mathbf{G}_m$ is compact in \mathbf{SH}_s there are isomorphisms

$$\begin{aligned} & \mathbf{SH}_s \left(\Sigma^{n,0} X_+ \wedge \mathbf{G}_m, \left(\prod_{n \in \mathbb{N}} \Omega_t^\infty \mathbf{MZ}/2 \right) \otimes_{\Omega_t^\infty \mathbf{MZ}/2} \Omega_t^\infty \Sigma^{0,1} \mathbf{MZ}/2 \right) \cong \\ & \mathbf{SH}_s \left(\Sigma^{n,0} X_+ \wedge \mathbf{G}_m, \left(\operatorname{colim}_{\mathcal{F} \subset \mathcal{B}} \prod_{f \in \mathcal{F}} \Omega_t^\infty \mathbf{MZ}/2 \right) \otimes_{\Omega_t^\infty \mathbf{MZ}/2} \Omega_t^\infty \Sigma^{0,1} \mathbf{MZ}/2 \right) \cong \\ & \operatorname{colim}_{\mathcal{F} \subset \mathcal{B}} \mathbf{SH}_s \left(\Sigma^{n,0} X_+ \wedge \mathbf{G}_m, \left(\prod_{f \in \mathcal{F}} \Omega_t^\infty \mathbf{MZ}/2 \otimes_{\Omega_t^\infty \mathbf{MZ}/2} \Omega_t^\infty \Sigma^{0,1} \mathbf{MZ}/2 \right) \right) \cong \\ & \operatorname{colim}_{\mathcal{F} \subset \mathcal{B}} \mathbf{SH}_s \left(\Sigma^{n,0} X_+ \wedge \mathbf{G}_m, \Omega_t^\infty \mathbf{MZ}/2 \otimes_{\Omega_t^\infty \mathbf{MZ}/2} \prod_{f \in \mathcal{F}} \Omega_t^\infty \Sigma^{0,1} \mathbf{MZ}/2 \right) \cong \\ & \operatorname{colim}_{\mathcal{F} \subset \mathcal{B}} \mathbf{SH}_s \left(\Sigma^{n,0} X_+ \wedge \mathbf{G}_m, \prod_{f \in \mathcal{F}} \Omega_t^\infty \Sigma^{0,1} \mathbf{MZ}/2 \right) \cong \\ & \operatorname{colim}_{\mathcal{F} \subset \mathcal{B}} \mathbf{SH}_s \left(\Sigma^{n,0} X_+, \Omega_t \left(\prod_{f \in \mathcal{F}} \Omega_t^\infty \Sigma^{0,1} \mathbf{MZ}/2 \right) \right) \cong \\ & \operatorname{colim}_{\mathcal{F} \subset \mathcal{B}} \mathbf{SH}_s \left(\Sigma^{n,0} X_+, \prod_{f \in \mathcal{F}} \Omega_t^\infty \Omega_t \Sigma^{0,1} \mathbf{MZ}/2 \right) \cong \\ & \operatorname{colim}_{\mathcal{F} \subset \mathcal{B}} \mathbf{SH}_s \left(\Sigma^{n,0} X_+, \prod_{f \in \mathcal{F}} \Omega_t^\infty \Sigma^{-1,0} \mathbf{MZ}/2 \right) \cong \\ & \mathbf{SH}_s \left(\Sigma^{n,0} X_+, \operatorname{colim}_{\mathcal{F} \subset \mathcal{B}} \prod_{f \in \mathcal{F}} \Omega_t^\infty \Sigma^{-1,0} \mathbf{MZ}/2 \right) \cong \\ & \mathbf{SH}_s \left(\Sigma^{n,0} X_+, \prod_{n \in \mathbb{N}} \Omega_t^\infty \Sigma^{-1,0} \mathbf{MZ}/2 \right) \cong \mathbf{SH}_s \left(\Sigma^{n,0} X_+, \prod_{n \in \mathbb{N}} \Omega_t^\infty \Omega_t \Sigma^{0,1} \mathbf{MZ}/2 \right) \cong \\ & \mathbf{SH}_s \left(\Sigma^{n,0} X_+, \Omega_t \prod_{n \in \mathbb{N}} \Omega_t^\infty \Sigma^{0,1} \mathbf{MZ}/2 \right) \cong \mathbf{SH}_s \left(\Sigma^{n,0} X_+ \wedge \mathbf{G}_m, \prod_{n \in \mathbb{N}} \Omega_t^\infty \Sigma^{0,1} \mathbf{MZ}/2 \right), \end{aligned}$$

which by canonicity coincides with the map induced by α . If the target in (6.4) is 1-effective, as implied by (6.3), it would follow that α is an isomorphism. One checks that α is not an

isomorphism by choosing a field F such that $F^* \otimes \mathbb{Z}/2$ is an infinitely generated $\mathbb{Z}/2$ -module, e.g., $F = \mathbb{Q}$. The map $\mathbf{SH}_s(\Sigma^{-1,0}\mathrm{Spec}(F)_+, \alpha)$ coincides with the canonical map

$$\left(\prod \mathbb{Z}/2\right) \otimes_{\mathbb{Z}/2} (F^* \otimes \mathbb{Z}/2) \longrightarrow \prod F^* \otimes \mathbb{Z}/2, \quad (6.5)$$

which is not surjective. Hence $\prod_{n \in \mathbb{N}} \Omega_t^\infty \Sigma^{0,1} \mathbf{M}\mathbb{Z}/2$ cannot be 1-effective. As explained above it follows that $(\prod_{i \geq 0} \Sigma^{0,i} \mathbf{M}\mathbb{Z}/2)^{hC_2}$ is noneffective.

Acknowledgments. The authors gratefully acknowledge support from DFG GK 1916, DFG Priority Program 1786, and RCN Frontier Research Group Project no. 250399.

References

- [1] M. F. Atiyah. *K*-theory and reality. *Quart. J. Math. Oxford Ser. (2)*, 17:367–386, 1966.
- [2] A. J. Berrick, M. Karoubi, M. Schlichting, and P. A. Østvær. The Homotopy Fixed Point Theorem and the Quillen-Lichtenbaum conjecture in Hermitian *K*-theory. *Adv. Math.*, 278:34–55, 2015.
- [3] J. M. Boardman. Conditionally convergent spectral sequences. In *Homotopy invariant algebraic structures (Baltimore, MD, 1998)*, volume 239 of *Contemp. Math.*, pages 49–84. Amer. Math. Soc., Providence, RI, 1999.
- [4] D. Dugger and D. C. Isaksen. Motivic cell structures. *Algebr. Geom. Topol.*, 5:615–652, 2005.
- [5] B. I. Dundas, O. Röndigs, and P. A. Østvær. Motivic functors. *Doc. Math.*, 8:489–525 (electronic), 2003.
- [6] J. J. Gutiérrez, O. Röndigs, M. Spitzweck, and P. A. Østvær. Motivic slices and coloured operads. *J. Topol.*, 5(3):727–755, 2012.
- [7] P. S. Hirschhorn. *Model categories and their localizations*, volume 99 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2003.
- [8] M. Hoyois, S. Kelly, and P. A. Østvær. The motivic Steenrod algebra in positive characteristic. [arXiv:1305.5690](https://arxiv.org/abs/1305.5690), to appear in: JEMS.
- [9] P. Hu, I. Kriz, and K. Ormsby. The homotopy limit problem for Hermitian *K*-theory, equivariant motivic homotopy theory and motivic Real cobordism. *Adv. Math.*, 228(1):434–480, 2011.
- [10] D. C. Isaksen and A. Shkemb. Motivic connective *K*-theories and the cohomology of $A(1)$. *J. K-Theory*, 7(3):619–661, 2011.
- [11] J. F. Jardine. Motivic symmetric spectra. *Doc. Math.*, 5:445–553 (electronic), 2000.
- [12] J. F. Jardine. The *K*-theory presheaf of spectra. In *New topological contexts for Galois theory and algebraic geometry (BIRS 2008)*, volume 16 of *Geom. Topol. Monogr.*, pages 151–178. Geom. Topol. Publ., Coventry, 2009.
- [13] J. F. Jardine. *Local homotopy theory*. Springer Monographs in Mathematics. Springer, New York, 2015.
- [14] M. Levine. The homotopy coniveau tower. *J. Topol.*, 1(1):217–267, 2008.
- [15] F. Morel. A^1 -algebraic topology. In *International Congress of Mathematicians. Vol. II*, pages 1035–1059. Eur. Math. Soc., Zürich, 2006.
- [16] N. Naumann, M. Spitzweck, and P. A. Østvær. Chern classes, *K*-theory and Landweber exactness over nonregular base schemes. In *Motives and algebraic cycles*, volume 56 of *Fields Inst. Commun.*, pages 307–317. Amer. Math. Soc., Providence, RI, 2009.
- [17] N. Naumann, M. Spitzweck, and P. A. Østvær. Motivic Landweber exactness. *Doc. Math.*, 14:551–593, 2009.

- [18] I. Panin, K. Pimenov, and O. Röndigs. A universality theorem for Voevodsky’s algebraic cobordism spectrum. *Homology, Homotopy Appl.*, 10(2):211–226, 2008.
- [19] I. Panin, K. Pimenov, and O. Röndigs. On the relation of Voevodsky’s algebraic cobordism to Quillen’s K -theory. *Invent. Math.*, 175(2):435–451, 2009.
- [20] P. Pelaez. Multiplicative properties of the slice filtration. *Astérisque*, (335):xvi+289, 2011.
- [21] P. Pelaez. On the functoriality of the slice filtration. *J. K-Theory*, 11(1):55–71, 2013.
- [22] O. Röndigs and P. A. Østvær. Motives and modules over motivic cohomology. *C. R. Math. Acad. Sci. Paris*, 342(10):751–754, 2006.
- [23] O. Röndigs and P. A. Østvær. Modules over motivic cohomology. *Adv. Math.*, 219(2):689–727, 2008.
- [24] O. Röndigs and P. A. Østvær. Rigidity in motivic homotopy theory. *Math. Ann.*, 341(3):651–675, 2008.
- [25] O. Röndigs and P. A. Østvær. Slices of hermitian K -theory and Milnor’s conjecture on quadratic forms. *Geom. Topol.*, 20(2):1157–1212, 2016.
- [26] O. Röndigs, M. Spitzweck, and P. A. Østvær. The first stable homotopy groups of motivic spheres. [arXiv:1604.00365](https://arxiv.org/abs/1604.00365).
- [27] J.-P. Serre. *Galois cohomology*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, english edition, 2002. Translated from the French by Patrick Ion and revised by the author.
- [28] M. Spitzweck. Relations between slices and quotients of the algebraic cobordism spectrum. *Homology, Homotopy Appl.*, 12(2):335–351, 2010.
- [29] M. Spitzweck and P. A. Østvær. The Bott inverted infinite projective space is homotopy algebraic K -theory. *Bull. Lond. Math. Soc.*, 41(2):281–292, 2009.
- [30] R. W. Thomason. The homotopy limit problem. In *Proceedings of the Northwestern Homotopy Theory Conference (Evanston, Ill., 1982)*, volume 19 of *Contemp. Math.*, pages 407–419. Amer. Math. Soc., Providence, R.I., 1983.
- [31] T. tom Dieck. *Algebraic topology*. EMS Textbooks in Mathematics. European Mathematical Society (EMS), Zürich, 2008.
- [32] V. Voevodsky. Open problems in the motivic stable homotopy theory. I. In *Motives, polylogarithms and Hodge theory, Part I (Irvine, CA, 1998)*, volume 3 of *Int. Press Lect. Ser.*, pages 3–34. Int. Press, Somerville, MA, 2002.
- [33] V. Voevodsky. A possible new approach to the motivic spectral sequence for algebraic K -theory. In *Recent progress in homotopy theory (Baltimore, MD, 2000)*, volume 293 of *Contemp. Math.*, pages 371–379. Amer. Math. Soc., Providence, RI, 2002.
- [34] V. Voevodsky. On the zero slice of the sphere spectrum. *Tr. Mat. Inst. Steklova*, 246(Algebr. Geom. Metody, Svyazi i Prilozh.):106–115, 2004.

Mathematisches Institut, Universität Osnabrück, Germany.
e-mail: oliver.roendigs@uni-osnabrueck.de

Mathematisches Institut, Universität Osnabrück, Germany.
e-mail: markus.spitzweck@uni-osnabrueck.de

Department of Mathematics, University of Oslo, Norway.
e-mail: paularne@math.uio.no