WILEY

Online Proofing System Instructions

The Wiley Online Proofing System allows proof reviewers to review PDF proofs, mark corrections, respond to queries, upload replacement figures, and submit these changes directly from the locally saved PDF proof.

1. For the best experience reviewing your proof in the Wiley Online Proofing System ensure you are connected to the internet. This will allow the PDF proof to connect to the central Wiley Online Proofing System server. If you are connected to the Wiley Online Proofing System server you should see a green check mark icon above in the yellow banner.

Connected

Disconnected

2. Please review the article proof on the following pages and mark any corrections, changes, and query responses using the Annotation Tools outlined on the next 2 pages.

3. Save your proof corrections by clicking the “Publish Comments” button in the yellow banner above. Corrections don’t have to be marked in one sitting. You can publish comments and log back in at a later time to add and publish more comments before you click the “Complete Proof Review” button below.

4. If you need to supply additional or replacement files bigger than 5 Megabytes (MB) do not attach them directly to the PDF Proof, please click the “Upload Files” button to upload files:

5. When your proof review is complete and all corrections have been published to the server by clicking the “Publish Comments” button, please click the “Complete Proof Review” button below:

IMPORTANT: Did you reply to all queries listed on the Author Query Form appearing before your proof?

IMPORTANT: Did you click the “Publish Comments” button to save all your corrections? Any unpublished comments will be lost.

IMPORTANT: Once you click “Complete Proof Review” you will not be able to add or publish additional corrections.

[Click Here](#)

For technical questions about reviewing your proof contact mpsops.wiley@adi-mps.com

You can also check the [OPS Frequently Asked Questions](#) for guidance.

The latest version of Acrobat Reader can be downloaded for free at: <http://get.adobe.com/reader/>

- Export PDF
- Create PDF
- Edit PDF
- Comment

Comment

E-I

How to use it:

- age of nutritional conditions, and landmark events are monitored in populations of relatively homogeneous single cells. *Saccharomyces*, and is initiated after a carbon source [1]. Si are referred to as meiosis-specific genes. *revisiae* depends on the inducer of meiosis) [3]. I functions as a repressor, the genes *pression*) and *RGRI* at rise II mediator subunit osome density [8]. *Si* directly or indirectly

F

How to use it:

1. Small size (35-250 amino acids).
2. Absence of similarity to known proteins.
3. Absence of functional data which could not be the real overlapping gene.
4. Greater than 25% overlap at the N-terminal terminus with another coding feature; over both ends; or ORF containing a tRNA.

How to use it:

- Informal invariance: r
A: Math. Gen., Vol. 12, N

ching fractions for $D0 \rightarrow K+K$
relation in $D0$ decays' Phys

T

How to use it:

- Meiosis has a central role in the sexual reproduction of nearly all eukaryotes. In *Saccharomyces cerevisiae*, analysis of meiosis, especially by a simple change of media, can be conveniently monitored in haploid cells. Sporulation of *Saccharomyces cerevisiae* cell, the a/a cell, and is of a fermentable carbon source. Sporulation and are referred to as (Zb). Transcription of the *IME1* gene, which encodes the activator, in *S. cerevisiae* is induced by the gene *RME1* function. Rme1p to exert repression of GAL1 gene expression) and *HGH1* are required [1, 2, 3, 4]. These genes

5. Attach File Tool – for inserting large amounts of text or replacement figures.

Inserts an icon linking to the attached file in the appropriate place in the text.

How to use it:

- Click on .
- Click on the proof to where you'd like the attached file to be linked.
- Select the file to be attached from your computer or network.
- Select the colour and type of icon that will appear in the proof. Click OK.

The attachment appears in the right-hand panel.

chondrial preparatio
ative damage injury
the extent of membra
malondialdehyde (TBARS) formation.
used by high perform

6. Add stamp Tool – for approving a proof if no corrections are required.

Inserts a selected stamp onto an appropriate place in the proof.

How to use it:

- Click on .
- Select the stamp you want to use. (The **Approved** stamp is usually available directly in the menu that appears. Others are shown under *Dynamic*, *Sign Here*, *Standard Business*).
- Fill in any details and then click on the proof where you'd like the stamp to appear. (Where a proof is to be approved as it is, this would normally be on the first page).

of the business cycle, starting with the
on perfect competition, constant ret
production. In this environment goods
extra...
he...
etermined by the model. The New-Key
otaki (1987), has introduced produ
general equilibrium models with nomin
...
...
...

APPROVED

Drawing tools available on comment ribbon

7. Drawing Markups Tools – for drawing shapes, lines, and freeform annotations on proofs and commenting on these marks.

Allows shapes, lines, and freeform annotations to be drawn on proofs and for comments to be made on these marks.

How to use it:

- Click on one of the shapes in the **Drawing Markups** section.
- Click on the proof at the relevant point and draw the selected shape with the cursor.
- To add a comment to the drawn shape, right-click on shape and select *Open Pop-up Note*.
- Type any text in the red box that appears.

For further information on how to annotate proofs, click on the **Help** menu to reveal a list of further options:

WILEY

COLOR REPRODUCTION IN YOUR ARTICLE

Color figures were included with the final manuscript files that we received for your article. Because of the high cost of color printing, we can only print figures in color if authors cover the expense. **The charge for printing figures in color is \$850 per figure.**

Please indicate if you would like your figures to be printed in color or black and white. Color images will be reproduced online in *Wiley Online Library* at no charge, whether or not you opt for color printing.

Failure to return this form will result in the publication of your figures in black and white.

JOURNAL _____ VOLUME _____ ISSUE _____

TITLE OF MANUSCRIPT _____

MS. NO. _____ NO. OF
COLOR PAGES _____ AUTHOR(S) _____

☐ Please print my figures in black and white

☐ Please print my figures in color

\$ _____

BILL TO:

Name _____

Institution _____

Address _____

Purchase

Order No. _____

Phone _____

Fax _____

E-mail _____

Author Query Form

Journal: JSO
Article: JSO25362

Dear Author,

During the copy-editing of your paper, the following queries arose.

Please refer to the query reference callout numbers in the page proofs and respond to each by marking the necessary comments using the PDF annotation tools.

Please remember illegible or unclear comments and corrections may delay publication.

Many thanks for your assistance.

Query Reference	Query	Remarks/Comments	Query Reviewed
Q1	Author: Please confirm that given names (blue) and surnames/family names (vermilion) have been identified correctly.		
Q2	Author: Please verify that the linked ORCID identifiers are correct for each author.		
Q3	Author: Please provide volume number and page range in Ref. 4.		

Applicability of transoral robotic thyroidectomy: Is it the final solution?

Open thyroidectomy (OT) remains a standard of care management for differentiated thyroid cancer (DTC).¹ However, with the introduction of transoral endoscopic thyroidectomy vestibular approach (TOETVA), the surgical management of DTC has changed.^{2,3} With uptake varying by country, depending on cost, insurance coverage, government healthcare approval, and patient preference, there have been frequent reports assessing transoral robotic thyroidectomy (TORT) safety, perioperative, functional, and oncologic outcomes, as compared with OT and other endoscopic and robotic approaches.^{4,5} TORT is feasible and can be achieved with favorable outcomes.¹⁻⁵ No modifications to the standard TOETVA technique are required, except from an additional and optional fourth robotic arm in the axilla.^{4,5} The fourth arm is a common element in OT. The supplementary arm forms a retractor to enhance and highlighting dissection view and safety.^{4,5} The additional arm, the robotic camera and instruments are working alongside.⁴ The axilla port placement ensure thyroid mobilization, and the robotic arms are not striking procedure itself. The added axilla incision can be used for the removal of the entire thyroid, without fragmenting it.^{4,5}

The debate between alternative approaches to the thyroid gland is unlikely to have any clear resolution, despite the completion of randomized controlled trials.¹ The inherent biases in these trials, essentially comparing surgeons rather than technique, will limit the generalizability of the study results.¹ Setting, patient selection and surgeon candidacy are certainly key elements affecting the outcomes of thyroid cancer surgery.¹

However, it is suggested that the robotic platform may allow a surgeon to obtain the same oncological outcomes as open and endoscopic surgery, but with some advantages.⁵

The robot provides a magnified, three-dimensional surgical view and allows tremor-free articulating instrumentation, has facilitated more precise and complete dissection throughout the surgery, thus overcoming some technical limitations encountered during TOETVA using the conventional endoscopic system. In particular, the robotic system enhanced oncological and radical transoral thyroid surgery in the central compartment and, in turn, extended the indications of TOETVA to most differentiated thyroid cancers.⁵ Therefore, candidates for TORT are different from those for TOETVA and include patients with an ultrasonographically estimated main nodule size of less than or equal to 70 mm, papillary thyroid carcinoma with minimal extrathyroidal extension (T1, T2, and T3), papillary carcinoma with or without evidence of central lymph node metastasis (N0 and N1a), and/or

Graves' disease with an ultrasonographically estimated thyroid volume of less than or equal to 50 mL.⁵ However, patients with differentiated thyroid cancer with posterior extensive extrathyroidal extension (T4) and/or lateral neck metastasis (N1b) are still contraindicated for TORT.⁵

In addition to the advantages of TOETVA, TORT has several specific technical merits. First, complete resection of the pyramidal lobe or upper pole of the thyroid gland is much more feasible with the articulated movement of the endowrist used during TORT than with the rigid endoscopic instruments used during TOETVA.^{4,5} Second, the tremor-filtering robotic system and magnified three-dimensional operative vision provided by TORT enables surgeons to perform fine and safe dissection around critical structures such as the parathyroid glands and recurrent laryngeal nerves. In particular, feasible countertraction of the perineural tissue around the recurrent laryngeal nerve can minimize remnant thyroid tissue around the cricothyroid articulation and facilitate radical central neck dissection.^{4,5}

ORCID

Gianlorenzo Dionigi <http://orcid.org/0000-0003-0864-6087>

Hui Sun MD¹

Gianlorenzo Dionigi MD, FACS²

¹Division of Thyroid Surgery, China-Japan Union Hospital of Jilin University, Jilin Provincial Key Laboratory of Surgical Translational Medicine, Jilin Provincial Precision Medicine Laboratory of Molecular Biology and Translational Medicine on Differentiated Thyroid Carcinoma, Changchun, Jilin, China

²Division for Endocrine and Minimally Invasive Surgery, Department of Human Pathology in Adulthood and Childhood "Gaetano Barresi," University Hospital G. Martino, University of Messina, Messina, Italy

Correspondence

Gianlorenzo Dionigi, Division for Endocrine and Minimally Invasive Surgery, Department of Human Pathology in Adulthood and Childhood "Gaetano Barresi," University Hospital G. Martino, University of Messina, Via C. Valeria 1, Messina, Italy.
Email: gdionigi@unime.it

REFERENCES

- Kim WW, Lee J, Jung JH, Park HY, Tufano RP, Kim HY. A comparison study of the transoral and bilateral axillo-breast approaches in robotic

- thyroidectomy. *J Surg Oncol*. 2018;118(3):381-387. <https://doi.org/10.1002/jso.25175>
2. Bian C, Liu H, Yao XY, et al. Complete endoscopic radical resection of thyroid cancer via an oral vestibule approach. *Oncol Lett*. 2018;16(5):5599-5606. <https://doi.org/10.3892/ol.2018.9369>
3. Razavi CR, Vasiliou E, Tufano RP, Russell JO. Learning curve for transoral endoscopic thyroid lobectomy. *Otolaryngol Head Neck Surg*. 2018;159(4):625-629. <https://doi.org/10.1177/0194599818795881>
4. Cottrill EE, Funk EK, Goldenberg D, Goyal N. Flexible robotic system: a preclinical cadaver feasibility study. *Laryngoscope*. 2018. <https://doi.org/10.1002/lary.27543>
5. Kim WW, Lee J, Jung JH, Park HY, Tufano RP, Kim HY. A comparison study of the transoral and bilateral axillo-breast approaches in robotic thyroidectomy. *J Surg Oncol*. 2018;118(3):381-387. <https://doi.org/10.1002/jso.25175>

UNCORRECTED PROOFS