

Review

# Looking at Alzheimer's Disease Pathogenesis from the Nuclear Side

Laura D'Andrea, Ramona Stringhi, Monica Di Luca and Elena Marcello \*

Department of Pharmacological and Biomolecular Sciences, Università degli Studi di Milano, 20133 Milan, Italy; laura.dandrea@unimi.it (L.D.); ramona.stringhi@unimi.it (R.S.); monica.diluca@unimi.it (M.D.L.)

\* Correspondence: elena.marcello@unimi.it; Tel.: +39-02-50318314

**Abstract:** Alzheimer's disease (AD) is a neurodegenerative disorder representing the most common form of dementia. It is biologically characterized by the deposition of extracellular amyloid- $\beta$  ( $A\beta$ ) senile plaques and intracellular neurofibrillary tangles, constituted by hyperphosphorylated tau protein. The key protein in AD pathogenesis is the amyloid precursor protein (APP), which is cleaved by secretases to produce several metabolites, including  $A\beta$  and APP intracellular domain (AICD). The greatest genetic risk factor associated with AD is represented by the *Apolipoprotein E  $\epsilon 4$*  (*APOE  $\epsilon 4$* ) allele. Importantly, all of the above-mentioned molecules that are strictly related to AD pathogenesis have also been described as playing roles in the cell nucleus. Accordingly, evidence suggests that nuclear functions are compromised in AD. Furthermore, modulation of transcription maintains cellular homeostasis, and alterations in transcriptomic profiles have been found in neurodegenerative diseases. This report reviews recent advancements in the AD players-mediated gene expression.  $A\beta$ , tau, AICD, and *APOE  $\epsilon 4$*  localize in the nucleus and regulate the transcription of several genes, part of which is involved in AD pathogenesis, thus suggesting that targeting nuclear functions might provide new therapeutic tools for the disease.



**Citation:** D'Andrea, L.; Stringhi, R.; Di Luca, M.; Marcello, E. Looking at Alzheimer's Disease Pathogenesis from the Nuclear Side. *Biomolecules* **2021**, *11*, 1261. <https://doi.org/10.3390/biom11091261>

Academic Editors: Sara Merlo, Dmitry Lim and Simona Federica Spampinato

Received: 30 July 2021

Accepted: 20 August 2021

Published: 24 August 2021

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

**Keywords:** Alzheimer's disease; transcription; amyloid- $\beta$ ; AICD; tau; *APOE  $\epsilon 4$* ; nucleus

## 1. Introduction

Alzheimer's disease (AD) is the main cause of dementia and is becoming one of the most burdening diseases of the 21 century. In 2018, a dementia prevalence of about 50 million people worldwide has been estimated, with this number projected to triple by 2050 [1]. The incidence of AD rises exponentially with age; the majority of cases occur after age 65, constituting late-onset AD, while cases occurring earlier than age 65 represent less than 5% of all cases and are termed early-onset AD. Clinical symptoms include gradual loss of cognitive, affective, and behavioral functions, leading to increased impairment in activities of daily life [2].

The progression of AD is called the AD continuum and includes three broad phases: preclinical AD, mild cognitive impairment (MCI) due to AD, and dementia due to AD [3,4]. The AD phase is further classified into the stages of mild, moderate, and severe, which reflect the degree to which symptoms interfere with the ability to carry out everyday activities. The length of each phase in the continuum is variable and influenced by age, genetics, biological sex, and other factors [5]. Although the current diagnosis of AD relies on a combination of neuropsychological evaluations, biomarker measurements, and brain imaging, the diagnosis of AD is defined by the presence of amyloid  $\beta$  ( $A\beta$ ) and phosphorylated tau. Indeed, the disease is mainly characterized by the accumulation of extracellular  $A\beta$  in senile plaques and intracellular neurofibrillary tangles (NFTs) constituted by hyperphosphorylated tau protein [6].

Regarding AD pathogenesis, the amyloid hypothesis has become the dominant model and is guiding the development of potential treatments. According to this hypothesis, an imbalance between production and clearance of  $A\beta$  is an early and initiating factor

that leads to an increase in A $\beta$  levels, and thus to its aggregation in oligomers and later in plaques. Such events trigger synaptic failure and inflammatory responses together with microglial and astrocytic activation, altered neuronal ionic homeostasis, oxidative injury, tau hyperphosphorylation, NFTs formation, and eventually neuronal loss and dementia onset [7].

A $\beta$ , first isolated from the meningeal vessels of AD patients [8], is a small peptide of 40–42 amino acids generated via proteolytic cleavages of the amyloid precursor protein (APP). APP is a transmembrane protein; the enzymes responsible for APP cleavage have been identified in the 1990s [9]. The coordinated activity of the  $\beta$ - and  $\gamma$ -secretases liberates the A $\beta$  peptide from APP [10].  $\beta$ -secretase cleaves at the N-terminus of the A $\beta$  domain, releasing a soluble fragment (sAPP $\beta$ ) into the extracellular space and leaving a C-terminal fragment of 99 amino acids (CTF99) inserted into the membrane [9]. Several studies have identified a membrane-bound aspartyl protease with structural similarities to the pepsin family, named  $\beta$ -site APP cleaving enzyme-1 (BACE1), as the enzyme responsible for the  $\beta$ -cleavage of APP [11–13]. CTF99 is then cleaved by  $\gamma$ -secretase at the C-terminus of the A $\beta$  sequence to liberate the A $\beta$  peptide and the cytoplasmic APP intracellular domain (AICD). A heterotetrameric complex mediates  $\gamma$ -secretase activity and is constituted by Presenilin (PS), nicastrin (NCT), anterior pharynx defective (APH)-1a or APH-1b, and the PS enhancer (PEN)-2 [14]. PS contains the two critical Asp residues within the transmembrane domains 6 and 7 that are part of the aspartyl protease catalytic domain of  $\gamma$ -secretase [15]. The  $\gamma$ -secretase complex can produce A $\beta$  peptides in lengths of 38, 40, and 42 amino acids that are released from the complex [16].

An alternative and nonamyloidogenic physiological pathway implicates the activity of  $\alpha$ -secretase, which prevents A $\beta$  generation and leads to the release of the neuroprotective soluble APP ectodomain (sAPP $\alpha$ ) and the formation of an APP C-terminal fragment of 83 amino acids (CTF83) [17]. The metalloprotease A Disintegrin and Metalloproteinase 10 (ADAM10) is an enzyme capable of cleaving APP within an A $\beta$  sequence and which exerts the main  $\alpha$ -secretase activity in neuronal cells [18,19]. The  $\gamma$ -secretase then cleaves the CTF83 fragment partially inserted in the membrane to produce p3 peptide and the AICD.

The A $\beta$  monomers generated by APP cleavage can aggregate to form oligomers, protofibrils, and fibrils. The aggregation rate of A $\beta$  peptides is strongly correlated with the ratio of A $\beta$ 42/A $\beta$ 40 peptides in cellular and animal models [20,21]. Accumulation and further aggregation of protofibrils and fibrils lead to the formation of insoluble amyloid plaques, the main histopathological signature of AD [7]. However, in the past few decades, several studies have demonstrated that soluble forms of A $\beta$ , rather than the large insoluble fibrils in plaques, are toxic to synapses [22]. Both synthetic and naturally-secreted forms of A $\beta$  oligomers inhibit long-term potentiation (LTP) both ex vivo [23] and in vivo [24] and lead to frank loss of spines [25]. In addition, it has been shown that A $\beta$  oligomers may drive cognitive deficits in animal models of AD [26,27] and potentially also in AD patients [28,29].

As the other classical hallmark of AD, NFTs represent neuronal cytoplasmic aggregates of tau protein that form paired helical filaments [30]. The abnormal phosphorylation of tau is the first step in the formation of these aggregates. Afterward, tau dissociates from axonal microtubules and aberrantly accumulates in the somatic cytoplasm and dendrites. Somatodendritic accumulations of phosphorylated tau form before fibrils and are called pretangles [31]. NFTs consisting of paired helical filaments of tau develop later and correlate with neuronal degeneration. In the human brain, silver-stainable NFTs develop in the transentorhinal cortex as well as in a few subcortical brain stem nuclei [32] and then spread into the entorhinal and other brain regions in a distinct hierarchical sequence that is different from that of A $\beta$ -plaques [33].

Remarkably, in support of the amyloid hypothesis, mutations causing inherited forms of AD lead to increased production of A $\beta$  and the development of AD, while mutations in the *Microtubule-Associated Protein (MAPT)* gene encoding tau do not cause familial AD, although they can lead to other neurodegenerative tauopathies which often include

Parkinsonian symptoms [34]. Indeed, more than 200 mutations in the genes coding for *PS* and about 20 in *APP* genes cause familiar cases of AD in an autosomal dominant manner. Such mutations are localized in A $\beta$  sequences, close to the secretase cleavage sites, and affect the enzyme-substrate complex, leading to an increase in A $\beta$  or to an increase in the A $\beta$ 42/A $\beta$ 40 ratio [14]. These mutations in *APP* or *PS* have also been exploited to generate animal models of the disease [35,36]. These inherited forms of AD represent approximately 1%–2% of AD cases, can present with very early ages of onset and a more rapid rate of progression, and are sometimes associated with other neurologic symptoms seen less frequently in sporadic AD [37].

In the case of late-onset AD, the greatest risk factors are advanced age and the presence of the *APOE*  $\epsilon$ 4 allele [38]. Interestingly, the relatively rare *APOE*  $\epsilon$ 2 allele remains by far the strongest genetic protective factor against sporadic AD, highlighting the importance of *APOE*'s role in AD pathogenesis. Relative to carrying the most common *APOE*  $\epsilon$ 3 homozygous genotype, carrying at least one *APOE*  $\epsilon$ 4 allele increases the risk of developing AD approximately 3.7 times, given *APOE*  $\epsilon$ 4 homozygous increases that risk up to 12 times. Conversely, possessing a single *APOE*  $\epsilon$ 2 allele reduces the risk by approximately 40%, and *APOE*  $\epsilon$ 2 homozygous reduces the risk even further [39,40]. The post-mortem neuropathological factors correlated with the *APOE* genotype are a higher A $\beta$  plaque burden and more severe cerebral amyloid angiopathy in *APOE*  $\epsilon$ 4 carriers [41], also confirmed by A $\beta$  Positron Emission Tomography imaging across preclinical and clinical AD stages [42].

*APOE*  $\epsilon$ 4 affects A $\beta$  pathology because it promotes the seeding of A $\beta$  peptide into A $\beta$  oligomers, protofibrils, and fibrils [43,44] while also inhibiting A $\beta$  clearance from the brain, prolonging its half-life in the interstitial fluid [45,46] and inhibiting its enzymatic degradation [47]. In addition, in the last few years, the comprehension of the role of *APOE* in AD pathogenesis has expanded beyond A $\beta$  peptide-centric mechanisms to NFTs degeneration, microglia, astrocyte responses, and blood–brain barrier disruption [48].

In light of these considerations, AD pathogenesis can be considered multifaceted and, therefore, difficult to pinpoint because it is the result of a complex interplay of crossing pathways. In particular, it has been shown that several elements implicated in AD pathogenesis can translocate to the nucleus and regulate transcription, thereby creating a network of pathways leading to AD dementia. This review summarizes the state-of-the-art understanding of and research into the AD players working as regulators of transcription and their potential contribution to AD pathogenesis. The description of such mechanisms is relevant for the development of therapeutic strategies aimed at modulating transcription in AD.

## 2. APP Metabolites and Their Role as Transcription Regulators

As described in the introduction, APP undergoes a consecutive shedding and intramembrane proteolysis mediated by secretases that can be summarized under the term “regulated intramembrane proteolysis” [49]. This is a cellular process that is frequently implicated in important signaling pathways [50,51], also involves the nucleus. Among APP metabolites, the intracellular domain AICD, which is released into the cytosol, has been extensively studied regarding its function in nuclear signaling [52,53]. However, in the last decade, the nuclear function of A $\beta$ 42 has also been investigated, also showing DNA binding properties of this peptide.

### 2.1. APP Intracellular Domain as a Potential Transcription Factor

A growing number of studies have reported the nuclear role of AICD in transcriptional control. Cytosolic AICD is subjected to rapid degradation [54,55], which leads to a short half-life [56]. Nevertheless, binding with the adaptor protein Fe65 stabilizes it and promotes its nuclear translocation [52,56,57]. Fe65, also known as the Amyloid Beta Precursor Protein Binding Family B Member 1 (APBB1), is a multifunctional protein belonging to the Fe65 family with the related proteins Fe65L1 and Fe65L2. It was recognized as an APP [58] and AICD [52,59] binding partner in yeast two-hybrid screenings. Fe65

cytosolic/nuclear localization is modulated by APP binding [60] and phosphorylation [61], as well as by its own phosphorylation [62]. When associated with APP, Fe65 is anchored to the membrane [60]; the  $\gamma$ -secretase-mediated APP processing releases the AICD-Fe65 complex and allows for its nuclear translocation [57,63]. Beyond its role in the nucleus, Fe65 has been reported to be involved in key cellular processes, such as actin dynamics [64] and calcium homeostasis [65], both impaired in AD.

AICD-Fe65 nuclear functions are modulated by the assembly of ternary complexes with different factors, such as Tip60 or CP2/LSF/LBP1. One well-characterized AICD-Fe65 coactivator is the acetyl transferase Tip60 [52,53,66]. In the nucleus, this complex can be detected as a dot-like structure [53,67]. AICD/Fe65/Tip60 complex is involved in the regulation of actin cytoskeleton, controlling the expression of *Stathmin1* (*STMN1*), a regulator of microtubules dynamics [67]. Nevertheless, the role of AICD in the modulation of *STMN1* transcription requires further elucidation.

It has further been shown that AICD/Fe65/Tip60 complex acts on other promoters. For example, it can activate the transcription of *KAI1* gene, replacing the N-CoR corepressor complex [66]. In the context of AICD-mediated gene expression control, it is important to underline that AICD regulates, both in vitro and in vivo, the expression of *Neprilysin* (*NEP*) [68–70], a protease involved in A $\beta$  degradation. Thus, the AICD-mediated *NEP* regulation suggests a feedback mechanism in the control of APP metabolism. In HEK293 cells, the ternary complex AICD/Fe65/Tip60 works as an activator of *NEP* promoter, triggering a consequent increment in the protein levels [69]. Furthermore, chromatin immunoprecipitation (ChIP) experiments have confirmed that AICD binds *NEP* promoter (contributing to its activation) in NB7 cells and rat primary cortical neurons through an epigenetic mechanism involving a competition between AICD and Histone deacetylases [68]. Although several studies have demonstrated a contribution of AICD in *NEP* transcription, this regulation mechanism remains controversial [71].

AICD/Fe65 complex also associates with CP2/LSF/LBP1 family, inducing the expression of glycogen synthase kinase 3 $\beta$  (GSK3 $\beta$ ) and leading to neurotoxicity as a result of tau phosphorylation increment and  $\beta$ -catenin levels reduction. This subsequently induces apoptosis [72]. Accordingly, other studies in neuroblastoma SHSY5Y cells have proposed the nuclear AICD as an encoder of pro-apoptotic signals through the association with the p53 factor [73].

Protein localization and function are finely tuned by post-translational modifications. Phosphorylation and SUMOylation modulate AICD nuclear activity [74,75]. Thr66 phosphorylation stabilizes AICD and is crucial for its localization and transcriptional activity [74,76], promoting the formation of a ternary complex with Fe65 and CP2 transcription factors and activating GSK3 $\beta$ , thus leading to neurotoxicity [74]. In accordance with these processes, other studies have shown that AICD controls GSK3 $\beta$  expression and activity in different cell types and primary neurons [53,72,77,78]. It has been recently reported that the SUMOylation of AICD Lys43 increases the association with Fe65 and nuclear localization [75]. Furthermore, SUMOylated AICD competes with the histone deacetylase 1 (HDAC1) for interaction with cyclic AMP (cAMP) response element-binding protein (CREB) and p65, controlling *NEP* and *TTR* expression [75].

Although Fe65 is a well-characterized AICD coactivator, a study conducted by Zhou and colleagues showed that AICD can also control transcription independently of Fe65 [79]. Indeed, in neuronal cells, Wnt signaling triggers the accumulation of AICD in the nucleus, where it associates with  $\beta$ -Catenin. This facilitates its activity as a transcriptional transactivator [79]. Wnt and  $\beta$ -catenin constitute a signaling pathway fundamental for cellular homeostasis. When the Wnt signal is off, APC, GSK3 $\beta$ , and CK1 complex promote  $\beta$ -catenin ubiquitination and degradation. Conversely, when Wnt is activated,  $\beta$ -catenin translocates to the nucleus, where it controls gene expression. It is important to note that in brains affected by AD, Wnt signaling is compromised [80]. AICD and Wnt/ $\beta$ -catenin pathways are connected by a negative feedback loop: Wnt signaling promotes AICD- $\beta$ -catenin interaction in the nucleus [79], and in turn, AICD promotes the formation of

GSK3 $\beta$ -Axin complex and  $\beta$ -catenin ubiquitination and degradation [81]. Finally, AICD-mediated Wnt signaling regulation is involved in neuronal development, promoting the differentiation of neuronal cells and blocking their proliferation [81].

Other studies have shown that AICD can negatively regulate neurogenesis in vivo in APP KO mice and AICD mouse models [82–84], and in vitro can inhibit the differentiation of human neural stem cells (hNSC) [85,86]. A recent study showed that AICD cooperates with Forkhead box O3a (FOXO3a) in the control of neurogenesis in neural stem cells (NSCs) and neural progenitor cells (NPCs) derived from the hippocampus of an AICD mouse model [87]. A negative feedback regulation has been proposed between AICD and FOXO3a: AICD promotes FOXO3a transcriptional activity, which in turn negatively regulates AICD transcription. Accordingly, it has been shown that AICD drives FOXO3a transcriptional activity in primary cortical neurons [88].

Mitochondrial structure, function, and dynamics are affected in AD, and an emerging role of AICD has been proposed in AD mitochondrial dysfunction [89]. In collaboration with FOXO3a, AICD activity is implicated in mitochondrial functions; in particular, AICD controls the transcription of *Phosphatase and Tensin Homolog (PTEN)-Induced Kinase 1 (Pink-1)* in a FOXO3a-dependent manner [89]. Pink-1 has a cytosolic and mitochondrial localization, and its AICD-FOXO3a-mediated regulation has repercussions on mitochondrial functions and mitophagy [89]. Taken together, the above-mentioned studies suggest that, in the nucleus, AICD-mediated transcriptional modulation controls key molecular pathways that result in compromised processes leading to AD.

## 2.2. Amyloid- $\beta$ Nuclear Localization and DNA Binding Properties

In the last decade, research has identified and examined the nuclear function of A $\beta$ 42. An important contribution to the study of A $\beta$  as a putative transcriptional regulator arises from the in vitro identification of a specific A $\beta$  binding domain on DNA [90]. An electrophoretic mobility shift assay (EMSA) identified the DNA guanine-rich sequence “KGGRKTGGGG” (where K is guanine or thymine, and R is a purine) as an A $\beta$ -interacting consensus sequence. Such a sequence is recognized in the promoter of genes involved in AD pathogenesis, including *APP*, *BACE1*, and *APOE* [90]. Other works have confirmed that A $\beta$  induces the expression of *BACE1*, suggesting a role in the regulation of its own metabolism [91].

In concert with these studies, ChIP experiments have shown that A $\beta$  binds the *APP* promoter [92], and in SH-SY5Y neuroblastoma cells, A $\beta$  exposure leads to an increase in *APP* transcription [93]. Conversely, in cortical primary neurons, A $\beta$  exerts an opposite effect on *APP* mRNA levels, inducing suppression of its expression [91]. Moreover, such a discrepancy might be dependent on the dose and the time of exposure and on differences in the regulation mechanisms occurring in diverse kinds of cells.

Importantly, confocal imaging and electron transmission microscopy approaches have confirmed that A $\beta$ 42 localizes in the nuclei of neuroblastoma cells and APP/PS1 hippocampal neurons, supporting a nuclear role of A $\beta$ 42 [93]. A genome-wide DNA microarray conducted in neuroblastoma cells led to the identification of genes upregulated on A $\beta$  exposure. Among them, gene encoding was identified for the *Insulin-Like Growth Factor Binding Protein 3 and 5*, whose levels increase even in the transgenic mouse expressing human APP (TgCRND8 mouse line) [94]. It is important to note that A $\beta$  might act even as a transcriptional repressor, as reported for *Irf1* and *KAl1* promoters [29].

Notably, oxidative stress [92] and antibiotic treatment [95] trigger A $\beta$  nuclear translocation, suggesting that A $\beta$ -mediated changes in gene expression are required upon different stimuli in response to cellular stress. Although A $\beta$  localizes in the nucleus [93] and has DNA binding properties [90,92], the mechanisms by which it enters the nucleus are not understood, and A $\beta$  partners in the transcriptional control remain unidentified. Furthermore, whether the exogenous and endogenous intracellular A $\beta$  might reach the nucleus through the same mechanisms and trigger the activation/repression of the same nuclear pathways has not yet been clarified. In this regard, the employment of patient-derived human in-

duced pluripotent stem cells (iPSCs), a potent tool for modeling brain diseases, might provide an important contribution. Although different studies on AD took advantage of such cellular system, A $\beta$  nuclear localization and function have not been explored in iPSCs yet. Moreover, iPSC-derived neurons might represent an important tool for understanding A $\beta$  nuclear trafficking and its relevance in gene expression control. These observations leave many open questions regarding A $\beta$  nuclear functions, but at same time present new perspectives for therapeutic interventions aimed at restoring physiological activity.

### 3. The Relevance of Nuclear Tau in Alzheimer's Disease Pathogenesis

The tau protein was first described as an essential cytosolic factor involved in stabilizing microtubule assembly [96]. As reported above, tau aggregates are a common feature of several neurodegenerative disorders (termed tauopathies), including AD. Tau is expressed in the adult human brain in six different isoforms, which derive from the alternative splicing of exons 2, 3, and 10 of the *MAPT* gene, located on chromosome 17q21.1 [97,98]. The resulting proteins can have 0 (0N tau isoform), 1 (1N tau isoform), or 2 (2N tau isoform) inserts in the N-terminal domain, and 3 (3R tau isoform) or 4 (4R tau isoform) microtubule binding repeats in the C-terminal region [99]. Although the majority of current research on neuronal tau has focused on its role in the cytoplasmic compartment, a nuclear form of tau has also been characterized. In particular, it has been shown that each tau isoform has a preferred subcellular localization and thus a precise function [100]. In addition, the tau protein undergoes post-translational modifications and, therefore, it is important to understand whether these modifications can further influence its localization to the nucleus and its functional relevance.

Various effective studies on nuclear tau have been conducted, but several discrepancies have also emerged. In general, both phosphorylated and dephosphorylated isoforms have been detected in the nuclear compartment, which may vary depending on cell type and intranuclear localization. For example, the nuclear compartment of neuronal cells is characterized by the presence of dephosphorylated forms of tau [101–105]. Moreover, in human brains, dephosphorylated forms of tau have a specific localization in the nucleolus [102,106,107], while in murine neurons, it is diffusely expressed in the whole nuclear compartment [104]. Several in vitro studies have demonstrated the capability of tau to bind DNA [108,109]. These findings have also been confirmed in a cellular environment: 14% of total tau localizes in the chromatin fraction containing DNA, chromatin, and associated proteins [101].

Data obtained from cell cultures as well as mouse and human brains have established the involvement of nuclear tau in mechanisms of nucleoplasmic and ribosomal DNA (rDNA) protection against different cellular stressors, such as thermal denaturation [110] and hydroxyl free radical (–OH)-triggered double-strand DNA breakage [111,112]. Such stresses induce changes in tau phosphorylation states, causing the dephosphorylation of cytoplasmic tau and the consequent nuclear shuttling. The stressor dismissal coincides with a decrease in nuclear tau to basal levels, while the phosphorylated level of cytoplasmic form increases. These findings suggest that tau phosphorylation/dephosphorylation represents a major mechanism of regulation for the trafficking of tau from the nucleus to the cytoplasm [113].

Tau–DNA interaction is mediated through tau's proline-rich domain (PRD) and microtubule binding domain (MBD) with the DNA minor groove in order to form protein–DNA complexes [114]. However, tau has an additional interaction with DNA—more precisely, it co-localizes with dimethylated lysine 9 of histone 3 (H3K9me2)-rich DNA sequences, creating further protein–DNA complexes with the AT-rich  $\alpha$ -satellite DNA sequences, organized as constitutive heterochromatin [107]. In order to highlight the tau protein's physiological role, tau has been overexpressed in *Drosophila melanogaster*, and it has been shown that it relaxes heterochromatin [115]. On the other hand, experiments conducted on KO-tau mice have highlighted its function as an arranger of nucleolar and/or heterochromatinization of a portion of antisense noncoding RNAs [116]. In addition, the depletion of tau results in an

increase in rDNA transcription with an associated decrease in heterochromatin and DNA methylation, suggesting that under physiological conditions, tau is involved in silencing the rDNA. How tau is able to affect chromatin conformation remains unclear. However, it has been described that tau associates with TIP5 in the heterochromatin, a key player in genome stability [117].

Notably, the widespread loss of heterochromatin organization, which could be rescued by overexpressing nuclear human tau protein in KO tau neurons, has also been observed in AD neurons that displayed pathological hyperphosphorylated tau [116]. Heterochromatin loss likely permits increased expression of nonprotein-coding RNA transcripts. Indeed, increased expression of a regulatory RNA has already been implicated in the pathogenesis of AD. In particular, a noncoding antisense transcript for *BACE1* (*BACE1-AS*) is augmented in human AD brains, triggering upregulation of *BACE1* mRNA and protein levels [118]. The loss of heterochromatin, and the consequent aberrant gene expression, are toxic effectors of tau-induced neurodegeneration, underlining the role of chromatin structure as a potential therapeutic target in AD [115].

A further epigenetic mechanism involved in the organization of the chromatin structure and reflecting changes in gene activity is H3K9 acetylation (H3K9ac). Klein and colleagues used H3K9 acetylation (H3K9ac) as a marker in an epigenome-wide association study to reveal a tau-driven disorganization of chromatin in AD affecting 5990 out of 26,384 H3K9ac domains [119]. Together, these findings suggest that AD involves a reconfiguration of the epigenome, and the identification of this process highlights potential epigenetic strategies for early-stage disease treatment [119,120].

Tau is also detectable in a specific region of acrocentric chromosomes of dividing cells, called the nuclear organizer region (NOR). This site is enriched in rRNA genes and represents the sub-nucleolar compartment where the formation of nucleolus takes place [121,122]. A post-translational modification of tau involving phosphorylation plays a key role in regulating its function and interaction. Indeed, NOR-associated tau is dephosphorylated, and its phosphorylation induces tau detachment from the DNA [114]. Since tau hyperphosphorylation is a hallmark of AD, this event can enhance tau-DNA dissociation. Notably, in AD patients, the dentate gyrus (DG) and the CA1 region of the hippocampus display an altered mRNA expression and altered protein levels in rRNA transcription key regulators, such as upstream binding factor (UBF), nucleolin (NLC), and nucleophosmin (NMP1). These factors are decreased in the AD hippocampi from the first stages of pathology, even before neuron loss [123]. These alterations may represent an early sign of AD, which precedes the reduction in dendritic arborization and the synaptic reduction in CA1 and DG neurons, finally resulting in hippocampal atrophy.

A recent study highlighted a further transcriptional pathway aberrantly changed in AD. Through examination of AD post-mortem brain tissues and *Drosophila melanogaster* models, a direct correlation between the increased activation/mobilization of transposable elements (TEs) and the pathogenic activity of tau has been demonstrated [124]. This event represents a key driver of anomalous cell cycle activation in neurons and subsequent neuronal death. Among upregulated transposable elements in human tauopathy, the human endogenous retrovirus (HERV) family (including HERV-K) has emerged [125]. Interestingly, a causal association between HERV-K and neuronal dysfunction has previously been established, as the expression of HERV-K or the retroviral envelope protein that it encodes decrease synaptic activity in mice [126].

Recent research has further described a mechanism of transcription regulation governed by tau, lacking a direct interaction with DNA but depending on the modulation of nuclear calcium ( $\text{Ca}^{2+}$ ) levels. During synaptic plasticity events, an influx of  $\text{Ca}^{2+}$  occurs, resulting in the regulation of transcription through the activation of CREB. Using an animal model of tauopathy (tau<sup>R406W</sup> Transgenic *Drosophila*) and iPSC-derived neurons from AD patients, Mahoney and colleagues highlighted an aberrant depletion of nuclear  $\text{Ca}^{2+}$ . Synapse-to-nucleus communication is mainly governed by  $\text{Ca}^{2+}$  [127], and pathogenic tau directly contributes to CREB and calcium depletion. Indeed, CREB-regulated genes are

significantly over-represented in tau<sup>R406W</sup> transgenic *Drosophila*. In addition, pharmacological activation of big potassium (BK) channels is able to restore nuclear Ca<sup>2+</sup> levels and suppress neurodegeneration in tau<sup>R406W</sup> transgenic *Drosophila* [128]. Although it is not yet evident whether the regulation of such genes is a direct consequence of tau-induced CREB depletion, these findings provide a deeper understanding of the involvement of tau in AD pathogenesis.

#### 4. APOE $\epsilon$ 4 as a Transcriptional Regulator

As described in the introduction, APOE  $\epsilon$ 4 allele represents the strongest genetic risk factor for AD [38]. In the mammal brain, APOE proteins are mainly expressed in astrocytes and microglia; moreover, neurons may produce them under stress and in reaction to toxic events [129–132]. Interestingly, it has been shown that APOE proteins can translocate to the nucleus, where they are able to bind DNA and function as a transcription factor in human glioblastoma cells [133]. Furthermore, in neuroblastoma, APOE  $\epsilon$ 4 negatively regulates the Brain-Derived Neurotrophic Factor (BDNF) through a mechanism involving the enhancement of the nuclear translocation of histone deacetylases [134], suggesting that APOE  $\epsilon$ 4 controls the expression of key genes involved in brain functioning.

In a recent work, the effect of APOE proteins was tested on human neuronal cultures depleted of glia to exclude the contribution of the secreted proteins to the APOE-mediated phenotype in neurons. Neurons were plated on MEFs and transfected with recombinant APOE  $\epsilon$ 2, APOE  $\epsilon$ 3, and APOE  $\epsilon$ 4 [135]. From such experiments, an intriguing mechanism regarding APOE-mediated signaling has emerged. APOE is, in fact, able to activate a noncanonical MAP kinase signaling pathway. In particular, APOE binds the receptors on the cell surface, triggering the activation of map kinase kinase Dual Leucine Zipper Kinase (DLK), which phosphorylates the map kinase Mitogen-Activated Protein Kinase Kinase 7 (MKK7), leading to the phosphorylation of the MAP kinase Extracellular Signal-Related Kinase 1/2 (ERK1/2). Such a signaling cascade results in the stimulation of c-Fos phosphorylation and in the consequent AP-1-dependent APP transcription, which triggers an increment in A $\beta$  production, with an efficacy increasing from APOE  $\epsilon$ 2 to APOE  $\epsilon$ 3 to APOE  $\epsilon$ 4 [135]. Notably, other studies have shown that APOE  $\epsilon$ 4 interacts with APP and controls its metabolism, inducing a decrease in sAPP production [136].

The advent of new technologies allowing for the generation of 2D and 3D cultures derived from iPSCs is a strong in vitro tool and was recently implied in combination with the gene editing technique CRISPR/Cas9 for the study of APOE  $\epsilon$ 4 contribution to AD [137]. Although in this study, a direct APOE  $\epsilon$ 4-mediated regulation of gene expression was not assessed, transcriptomic analysis conducted in APOE  $\epsilon$ 4-expressing neurons showed that the genes that were primarily downregulated were associated with synaptic functions [137]. In addition, APOE  $\epsilon$ 4 neurons show an increment in A $\beta$ 42 with respect to the isogenic line expressing APOE  $\epsilon$ 3 [137], thus confirming the APOE  $\epsilon$ 4 impact on AD-related dysfunctions. Furthermore, ChIP experiments followed by high-throughput DNA sequencing (ChIP-seq) led to the identification of APOE  $\epsilon$ 3 or APOE  $\epsilon$ 4 target genes [133]. A total of 1700 genes were found regulated by APOE  $\epsilon$ 4 and not by APOE  $\epsilon$ 3, 76 of which were related to AD [133].

It is important to underline that APOE is involved in the transcriptional control of APP pathway players. The promoter of *Sirtuin1* (*Sirt1*), a transcriptional activator of the  $\alpha$ -secretase ADAM10 gene [138], has been identified among the promoters bound by APOE. In particular, APOE  $\epsilon$ 4 acts as a transcriptional repressor of *Sirt1* [133]. Accordingly, APOE  $\epsilon$ 4 expression leads to a reduction in *Sirt1* levels in AD brains and primary neurons [136]. The role of APOE  $\epsilon$ 4 as a negative transcriptional regulator was confirmed for at least three other genes, i.e., *MAP Kinase Activating Death Domain* (*MADD*), *Activity Dependent Neuroprotective Protein* (*ADNP*), and *COMM Domain 6* (*COMMD6*) [133]. Moreover, it is not possible to exclude an additional function as a transcriptional activator. These studies provide evidence that APOE proteins play a key role in the modulation of neuronal transcriptome; in particular, they indicate that APOE  $\epsilon$ 4 controls (both directly and indirectly) the expression of genes implicated in AD, confirming its contribution to this pathology. As described



above, the isogenic conversion of *APOE*  $\epsilon 4$  to *APOE*  $\epsilon 3$  ameliorates AD-related phenotypes in AD iPSCs-derived brain cell types [137] and iPSCs-derived cerebral organoids [139]. Furthermore, *APOE*  $\epsilon 4$ , but not *APOE*  $\epsilon 3$ , induces changes in the expression of AD related genes [133]. Such observations suggest that the genetic conversion *APOE*  $\epsilon 4$  allele to *APOE*  $\epsilon 3$  isoform might represent a promising therapeutic tool. The CRISPR/Cas9 approach allows the genetic correction of DNA mutation and constitutes a new frontier for molecular biology research and gene therapy. Once ethical concerns are overcome, this technique might provide a powerful instrument for long-lasting genetic modifications aimed at restoring gene functions as in the case of *APOE*  $\epsilon 4$ -*APOE*  $\epsilon 3$  conversion.


## 5. Conclusions

A growing number of studies have focused on nuclear and transcriptional dysfunctions in AD [140–142]. In the era of omics, high-throughput approaches, and big data generation, many reports have highlighted changes in the transcriptomic profiles of AD for in vitro and in vivo models and patients [143–148]. In addition, gene association studies have identified several putative AD-associated risk factors genes [149]. Such studies shed light on the unexplored mechanisms involved in the pathogenesis of this neurodegenerative disorder. Moreover, other research has provided new insights into the nuclear function of the “canonical” AD-associated molecules, such as the APP metabolites  $A\beta$  and AICD, tau, and *APOE*  $\epsilon 4$  (Figure 1).

$A\beta$ , AICD, and *APOE* are able to bind DNA [66,68,72,79,88,90,92,133,138] and act as transcriptional regulators. Importantly, they modulate the expression of key players of AD pathogenesis. *APOE* regulates *APP* [136] and the transcriptional activator of ADAM10 secretase, *Sirt1* [133].  $A\beta$  can bind *APP*, *BACE1*, and *APOE* promoters [90], while AICD regulates, both in vitro and in vivo, the expression of *Neprilysin* [68–70], a protease involved in  $A\beta$  clearance, just to cite the most relevant regulated genes (Figure 1 and Table 1). Such evidence suggests a feedback mechanism that must be finely tuned in physiological conditions; such equilibrium is disrupted in AD, in a vicious cycle that feeds itself.

Tau has also been described to play nuclear functions, and a nuclear form of tau has been identified. When tau protein is dephosphorylated it is involved in DNA protection against cellular stressors [111–113], as well as in the rearrangement of a heterochromatin portion enriched in noncoding RNAs [117] (Figure 1 and Table 1). Since AD pathogenesis is characterized by abnormal tau hyperphosphorylation, the loss of heterochromatin and the resultant anomalous gene expression could be the toxic effectors of tau-induced neurodegeneration. Effectively, altered levels of rRNA transcription key regulators are reported in post-mortem analysis of AD brains from the first stages of pathology [123]. Overall, these findings provide new insights into tau-induced alterations, suggesting potential strategies for early-stage disease treatment.

All the studies examined in this report underline the importance of proper nuclear signaling in the context of AD and indicate that correction of transcriptional defects might provide a novel therapeutic approach.



**Figure 1.** Schematic representation of how AD effectors contribute to neuronal dysfunction by modulating transcription. APP undergoes intramembrane proteolysis mediated by  $\beta$ - and  $\gamma$ -secretases to generate  $A\beta$  and AICD. The intracellular metabolite AICD binds Fe65, which promotes its nuclear translocation and the consequent interaction with the DNA. The AICD/Fe65/DNA complex controls the expression of *Stathmin1* (*STMN1*), activates *Neprilysin* (*NEP*) promoter, and induces the expression of *Glycogen Synthase Kinase 3 $\beta$*  (*GSK3 $\beta$* ). The  $A\beta$  monomers, generated by APP cleavage, translocate to the nucleus and recognize a sequence in the promoter of genes involved in AD pathogenesis, including *APP*, *BACE1*, and *APOE*. Tau hyperphosphorylated aggregates result in an anomalous activation of transposable elements (TEs) and an aberrant CREB-regulated genes over-representation. When nuclear tau is dephosphorylated, it is involved in DNA protection against cellular stressors and the rearrangement of heterochromatin, enriched in noncoding RNAs. AD pathogenesis is characterized by abnormal tau hyperphosphorylation, causing tau detachment from DNA and the consequent loss of heterochromatin and anomalous noncoding RNAs expression. *APOE  $\epsilon 4$*  binding to its receptor activates a noncanonical signaling pathway which triggers the *APP* transcription and  $A\beta_{40/42}$  production. When *APOE  $\epsilon 4$*  interacts with DNA, it acts as a transcriptional repressor of *Sirtuin1* (*SirT1*), a transcriptional activator of the  $\alpha$ -secretase *ADAM10* gene. (Red and blue arrows indicate the increase and the decrease in gene transcription respectively. Question mark (?) shows an unknown effect (activation or repression) of transcription).

**Table 1.** Nuclear functions and target genes of AD players.

| Molecules | DNA Binding | Nuclear Function        | Target Genes  | References                                    |
|-----------|-------------|-------------------------|---|---|
| AICD      | yes         | Transcription regulator | <i>STMN1</i><br><i>KAI1</i><br><i>NEP</i><br><i>GSK3<math>\beta</math></i><br><i>Pink-1</i> | [67]<br>[66]<br>[68–70,75]<br>[72,74]<br>[89] |

Table 1. Cont.

| Molecules         | DNA Binding | Nuclear Function   | Target Genes     | References                      |
|-------------------|-------------|--|------------------|---------------------------------|
| A $\beta$         | yes         | Transcription regulator                                    | APP              | [90,93]                         |
|                   |             |  | BACE             | [90]                            |
|                   |             |  | APOE             | [90]                            |
|                   |             |  | IRP1             | [29]                            |
|                   |             |  | KAI1             | [29]                            |
| tau               | yes         | DNA protection<br>Chromatin remodeling<br>Genome stability | Unknown genes    | [111,112]<br>[107,115]<br>[117] |
| APOE $\epsilon$ 4 | yes         | Transcription regulator                                    | APP              | [135,136]                       |
|                   |             |  | SirT1            | [133,138]                       |
|                   |             |  | MADD             | [133]                           |
|                   |             |  | ADNP             | [133]                           |
|                   |             |  | COMMD6           | [133]                           |
|                   |             |  | 1700 other genes | [133]                           |

**Author Contributions:** Writing—original draft preparation, L.D., R.S. and E.M.; editing and revision M.D.L. and E.M. All authors have read and agreed to the published version of the manuscript.

**Funding:** This project has received funding from the Italian Ministry of University and Research (PRIN 2017MYJ5TH to M.D.L., PRIN 2017B9NCSX to E.M., MIUR Progetto Eccellenza), from the Fondazione Cariplo to EM (Grant no. 2018-0511), from an intramural grant of the University of Milan to EM (PSR2019\_EMARC). This work was supported by MIUR - PON “Ricerca e Innovazione” PerMedNet id project ARS01\_01226.

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** Not applicable.

**Acknowledgments:** Cartoons used in Figure 1 have been downloaded from Servier Medical Art (smart.servier.com).

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

- Patterson, C. *Alzheimer's Disease International World Alzheimer Report 2018—The State of the Art of Dementia Research*; Alzheimer's Disease International: London, UK, 2018.
- Livingston, G.; Sommerlad, A.; Orgeta, V.; Costafreda, S.G.; Huntley, J.; Ames, D.; Ballard, C.; Banerjee, S.; Burns, A.; Cohen-Mansfield, J.; et al. Dementia prevention, intervention, and care. *Lancet* **2017**, *390*, 2673–2734. [[CrossRef](#)]
- Sperling, R.A.; Aisen, P.S.; Beckett, L.A.; Bennett, D.A.; Craft, S.; Fagan, A.M.; Iwatsubo, T.; Jack, C.R.; Kaye, J.; Montine, T.J.; et al. Toward defining the preclinical stages of Alzheimer's disease: Recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimer's Dement.* **2011**, *7*, 280–292. [[CrossRef](#)] [[PubMed](#)]
- Albert, M.S.; DeKosky, S.T.; Dickson, D.; Dubois, B.; Feldman, H.H.; Fox, N.C.; Gamst, A.; Holtzman, D.M.; Jagust, W.J.; Petersen, R.C.; et al. The diagnosis of mild cognitive impairment due to Alzheimer's disease: Recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimer's Dement.* **2011**, *7*, 270–279. [[CrossRef](#)]
- Vermunt, L.; Sikkes, S.A.M.; van den Hout, A.; Handels, R.; Bos, I.; van der Flier, W.M.; Kern, S.; Ousset, P.J.; Maruff, P.; Skoog, I.; et al. Duration of preclinical, prodromal, and dementia stages of Alzheimer's disease in relation to age, sex, and APOE genotype. *Alzheimer's Dement.* **2019**, *15*, 888–898. [[CrossRef](#)]
- Selkoe, D.J. Alzheimer's disease. *Cold Spring Harb. Perspect. Biol.* **2011**, *3*, a004457. [[CrossRef](#)] [[PubMed](#)]
- Selkoe, D.J.; Hardy, J. The amyloid hypothesis of Alzheimer's disease at 25 years. *EMBO Mol. Med.* **2016**, *8*, 595–608. [[CrossRef](#)]
- Glenner, G.G.; Wong, C.W. Alzheimer's disease: Initial report of the purification and characterization of a novel cerebrovascular amyloid protein. *Biochem. Biophys. Res. Commun.* **1984**, *120*, 885–890. [[CrossRef](#)]

9. Marcello, E.; Epis, R.; Gardoni, F.; Di Luca, M. The amyloid cascade: The old and the new. *J. Nutr. Health Aging* **2008**, *12*, 58S–60S. [[CrossRef](#)]
10. Haass, C.; Hung, A.Y.; Schlossmacher, M.G.; Teplow, D.B.; Selkoe, D.J.  $\beta$ -Amyloid peptide and a 3-kDa fragment are derived by distinct cellular mechanisms. *J. Biol. Chem.* **1993**, *268*, 3021–3024. [[CrossRef](#)]
11. Vassar, R.; Bennett, B.D.; Babu-Khan, S.; Kahn, S.; Mendiaz, E.A.; Denis, P.; Teplow, D.B.; Ross, S.; Amarante, P.; Loeloff, R.; et al.  $\beta$ -Secretase cleavage of Alzheimer's amyloid precursor protein by the transmembrane aspartic protease BACE. *Science* **1999**, *286*, 735–741. [[CrossRef](#)]
12. Yan, R.; Blenkowski, M.J.; Shuck, M.E.; Miao, H.; Tory, M.C.; Pauley, A.M.; Brashler, J.R.; Stratman, N.C.; Mathews, W.R.; Buhl, A.E.; et al. Membrane-anchored aspartyl protease with Alzheimer's disease  $\beta$ -secretase activity. *Nature* **1999**, *402*, 533–537. [[CrossRef](#)]
13. Sinha, S.; Anderson, J.P.; Barbour, R.; Basi, G.S.; Caccaveffo, R.; Davis, D.; Doan, M.; Dovey, H.F.; Frigon, N.; Hong, J.; et al. Purification and cloning of amyloid precursor protein  $\beta$ -secretase from human brain. *Nature* **1999**, *402*, 537–540. [[CrossRef](#)]
14. Voytyuk, I.; De Strooper, B.; Chávez-Gutiérrez, L. Modulation of  $\gamma$ - and  $\beta$ -Secretases as Early Prevention Against Alzheimer's Disease. *Biol. Psychiatry* **2018**, *83*, 320–327. [[CrossRef](#)]
15. Wolfe, M.S.; Xia, W.; Ostaszewski, B.L.; Diehl, T.S.; Kimberly, W.T.; Selkoe, D.J. Two transmembrane aspartates in presenilin-1 required for presenilin endoproteolysis and  $\gamma$ -secretase activity. *Nature* **1999**, *398*, 513–517. [[CrossRef](#)]
16. Zhou, R.; Yang, G.; Guo, X.; Zhou, Q.; Lei, J.; Shi, Y. Recognition of the amyloid precursor protein by human  $\gamma$ -secretase. *Science* **2019**, *363*, eaaw0930. [[CrossRef](#)]
17. Lammich, S.; Kojro, E.; Postina, R.; Gilbert, S.; Pfeiffer, R.; Jasionowski, M.; Haass, C.; Fahrenholz, F. Constitutive and regulated  $\alpha$ -secretase cleavage of Alzheimer's amyloid precursor protein by a disintegrin metalloprotease. *Proc. Natl. Acad. Sci. USA* **1999**, *96*, 3922–3927. [[CrossRef](#)] [[PubMed](#)]
18. Kuhn, P.H.; Wang, H.; Dislich, B.; Colombo, A.; Zeitschel, U.; Ellwart, J.W.; Kremmer, E.; Roßner, S.; Lichtenthaler, S.F. ADAM10 is the physiologically relevant, constitutive  $\alpha$ -secretase of the amyloid precursor protein in primary neurons. *EMBO J.* **2010**, *29*, 3020–3032. [[CrossRef](#)] [[PubMed](#)]
19. Marcello, E.; Borroni, B.; Pelucchi, S.; Gardoni, F.; Di Luca, M. ADAM10 as a therapeutic target for brain diseases: From developmental disorders to Alzheimer's disease. *Expert Opin. Ther. Targets* **2017**, *21*, 1017–1026. [[CrossRef](#)] [[PubMed](#)]
20. Pontrello, C.G.; Sun, M.Y.; Lin, A.; Fiacco, T.A.; DeFea, K.A.; Ethell, I.M. Cofilin under control of  $\beta$ -arrestin-2 in NMDA-dependent dendritic spine plasticity, long-term depression (LTD), and learning. *Proc. Natl. Acad. Sci. USA* **2012**, *109*, E442–E451. [[CrossRef](#)]
21. Terrill-Usery, S.E.; Colvin, B.A.; Davenport, R.E.; Nichols, M.R. A $\beta$ 40 has a subtle effect on A $\beta$ 42 protofibril formation, but to a lesser degree than A $\beta$ 42 concentration, in A $\beta$ 42/A $\beta$ 40 mixtures. *Arch. Biochem. Biophys.* **2016**, *597*, 1–11. [[CrossRef](#)] [[PubMed](#)]
22. Cline, E.N.; Bicca, M.A.; Viola, K.L.; Klein, W.L. The Amyloid- $\beta$  Oligomer Hypothesis: Beginning of the Third Decade. *J. Alzheimer's Dis.* **2018**, *64*, S567–S610. [[CrossRef](#)]
23. Lambert, M.P.; Barlow, A.K.; Chromy, B.A.; Edwards, C.; Freed, R.; Liosatos, M.; Morgan, T.E.; Rozovsky, I.; Trommer, B.; Viola, K.L.; et al. Diffusible, nonfibrillar ligands derived from A $\beta$ 1–42 are potent central nervous system neurotoxins. *Proc. Natl. Acad. Sci. USA* **1998**, *95*, 6448–6453. [[CrossRef](#)]
24. Walsh, D.M.; Klyubin, I.; Fadeeva, J.V.; Cullen, W.K.; Anwyl, R.; Wolfe, M.S.; Rowan, M.J.; Selkoe, D.J. Naturally secreted oligomers of amyloid  $\beta$  protein potently inhibit hippocampal long-term potentiation in vivo. *Nature* **2002**, *416*, 535–539. [[CrossRef](#)] [[PubMed](#)]
25. Shankar, G.M.; Bloodgood, B.L.; Townsend, M.; Walsh, D.M.; Selkoe, D.J.; Sabatini, B.L. Natural oligomers of the Alzheimer amyloid- $\beta$  protein induce reversible synapse loss by modulating an NMDA-type glutamate receptor-dependent signaling pathway. *J. Neurosci.* **2007**, *27*, 2866–2875. [[CrossRef](#)] [[PubMed](#)]
26. Cleary, J.P.; Walsh, D.M.; Hofmeister, J.J.; Shankar, G.M.; Kuskowski, M.A.; Selkoe, D.J.; Ashe, K.H. Natural oligomers of the amyloid- $\beta$  protein specifically disrupt cognitive function. *Nat. Neurosci.* **2005**, *8*, 79–84. [[CrossRef](#)] [[PubMed](#)]
27. Lesné, S.; Koh, M.; Kotilinek, L.; Kaye, R.; Glabe, C.G.; Yang, A.; Gallagher, M.; Ashe, K.H. A Specific Amyloid-beta Protein Assembly in the Brain Impairs Memory. *Nature* **2006**, *440*, 352–357. [[CrossRef](#)] [[PubMed](#)]
28. Esparza, T.J.; Zhao, H.; Cirrito, J.R.; Cairns, N.J.; Bateman, R.J.; Holtzman, D.M.; Brody, D.L. Amyloid-beta oligomerization in Alzheimer dementia versus high-pathology controls. *Ann. Neurol.* **2013**, *73*, 104–119. [[CrossRef](#)] [[PubMed](#)]
29. Perez-Nievas, B.G.; Stein, T.D.; Tai, H.C.; Dols-Icardo, O.; Scotton, T.C.; Barroeta-Espar, I.; Fernandez-Carballo, L.; De Munain, E.L.; Perez, J.; Marquie, M.; et al. Dissecting phenotypic traits linked to human resilience to Alzheimer's pathology. *Brain* **2013**, *136*, 2510–2526. [[CrossRef](#)]
30. Iqbal, K.; Zaidi, T.; Wen, G.Y.; Grundke-Iqbal, I.; Merz, P.A.; Shaikh, S.S.; Wisniewski, H.M.; Alafuzoff, I.; Winblad, B. Defective brain microtubule assembly in Alzheimer's disease. *Lancet* **1986**, *2*, 421–426. [[CrossRef](#)]
31. Spires-Jones, T.L.; Attems, J.; Thal, D.R. Interactions of pathological proteins in neurodegenerative diseases. *Acta Neuropathol.* **2017**, *134*, 187–205. [[CrossRef](#)]
32. Braak, H.; Del Tredici, K. The pathological process underlying Alzheimer's disease in individuals under thirty. *Acta Neuropathol.* **2011**, *121*, 171–181. [[CrossRef](#)]
33. Braak, H.; Braak, E. Neuropathological staging of Alzheimer-related changes. *Acta Neuropathol.* **1991**, *82*, 239–259. [[CrossRef](#)]
34. Hutton, M.; Lendon, C.L.; Rizzu, P.; Baker, M.; Froelich, S.; Houlden, H.H.; Pickering-Brown, S.; Chakraverty, S.; Isaacs, A.; Grover, A.; et al. Association of missense and 5'-splice-site mutations in tau with the inherited dementia FTDP-17. *Nature* **1998**, *393*, 702–705. [[CrossRef](#)]

35. Saraceno, C.; Musardo, S.; Marcello, E.; Pelucchi, S.; Di Luca, M. Modeling Alzheimer's disease: From past to future. *Front. Pharmacol.* **2013**, *19*, 77. [[CrossRef](#)]
36. Epis, R.; Gardoni, F.; Marcello, E.; Genazzani, A.; Canonico, P.L.; Di Luca, M. Searching for new animal models of Alzheimer's disease. *Eur. J. Pharmacol.* **2010**, *626*, 57–63. [[CrossRef](#)] [[PubMed](#)]
37. Bateman, R.J.; Xiong, C.; Benzinger, T.L.S.; Fagan, A.M.; Goate, A.; Fox, N.C.; Marcus, D.S.; Cairns, N.J.; Xie, X.; Blazey, T.M.; et al. Clinical and biomarker changes in dominantly inherited Alzheimer's disease. *N. Engl. J. Med.* **2012**, *367*, 795–804. [[CrossRef](#)]
38. van der Lee, S.J.; Wolters, F.J.; Ikram, M.K.; Hofman, A.; Ikram, M.A.; Amin, N.; van Duijn, C.M. The effect of APOE and other common genetic variants on the onset of Alzheimer's disease and dementia: A community-based cohort study. *Lancet Neurol.* **2018**, *17*, 434–444. [[CrossRef](#)]
39. Reiman, E.M.; Arboleda-Velasquez, J.F.; Quiroz, Y.T.; Huentelman, M.J.; Beach, T.G.; Caselli, R.J.; Chen, Y.; Su, Y.; Myers, A.J.; Hardy, J.; et al. Exceptionally low likelihood of Alzheimer's dementia in APOE2 homozygotes from a 5000-person neuropathological study. *Nat. Commun.* **2020**, *11*, 667. [[CrossRef](#)] [[PubMed](#)]
40. Rebeck, W.G.; Reiter, J.S.; Strickland, D.K.; Hyman, B.T. Apolipoprotein E in sporadic Alzheimer's disease: Allelic variation and receptor interactions. *Neuron* **1993**, *11*, 575–580. [[CrossRef](#)]
41. Serrano-Pozo, A.; Qian, J.; Monsell, S.E.; Betensky, R.A.; Hyman, B.T. APOE $\epsilon$ 2 is associated with milder clinical and pathological Alzheimer disease. *Ann. Neurol.* **2015**, *77*, 917–929. [[CrossRef](#)] [[PubMed](#)]
42. Ossenkoppele, R.; Jansen, W.J.; Rabinovici, G.D.; Knol, D.L.; van der Flier, W.M.; van Berckel, B.N.M.; Scheltens, P.; Visser, P.J.; Verfaillie, S.C.J.; Zwan, M.D.; et al. Prevalence of amyloid PET positivity in dementia syndromes: A meta-analysis. *JAMA* **2015**, *313*, 1939–1949. [[CrossRef](#)]
43. Hashimoto, T.; Serrano-Pozo, A.; Hori, Y.; Adams, K.W.; Takeda, S.; Banerji, A.O.; Mitani, A.; Joyner, D.; Thyssen, D.H.; Bacskai, B.J.; et al. Apolipoprotein e, especially apolipoprotein E4, increases the oligomerization of amyloid  $\beta$  peptide. *J. Neurosci.* **2012**, *32*, 15181–15192. [[CrossRef](#)]
44. Hori, Y.; Hashimoto, T.; Nomoto, H.; Hyman, B.T.; Iwatsubo, T. Role of apolipoprotein E in  $\beta$ -amyloidogenesis: Isoform-specific effects on protofibril to fibril conversion of A $\beta$  in vitro and brain A $\beta$  deposition in vivo. *J. Biol. Chem.* **2015**, *290*, 15163–15174. [[CrossRef](#)] [[PubMed](#)]
45. Liu, C.C.; Zhao, N.; Fu, Y.; Wang, N.; Linares, C.; Tsai, C.W.; Bu, G. ApoE4 Accelerates Early Seeding of Amyloid Pathology. *Neuron* **2017**, *96*, 1024–1032. [[CrossRef](#)] [[PubMed](#)]
46. Castellano, J.M.; Kim, J.; Stewart, F.R.; Jiang, H.; DeMattos, R.B.; Patterson, B.W.; Fagan, A.M.; Morris, J.C.; Mawuenyega, K.G.; Cruchaga, C.; et al. Human APOE isoforms differentially regulate brain amyloid- $\beta$  peptide clearance. *Sci. Transl. Med.* **2011**, *3*, 89. [[CrossRef](#)] [[PubMed](#)]
47. Deane, R.; Sagare, A.; Hamm, K.; Parisi, M.; Lane, S.; Finn, M.B.; Holtzman, D.M.; Zlokovic, B.V. apoE isoform-specific disruption of amyloid  $\beta$  peptide clearance from mouse brain. *J. Clin. Investig.* **2008**, *118*, 4002–4013. [[CrossRef](#)]
48. Serrano-Pozo, A.; Das, S.; Hyman, B.T. APOE and Alzheimer's disease: Advances in genetics, pathophysiology, and therapeutic approaches. *Lancet Neurol.* **2021**, *20*, 68–80. [[CrossRef](#)]
49. Lichtenthaler, S.F.; Haass, C.; Steiner, H. Regulated intramembrane proteolysis—Lessons from amyloid precursor protein processing. *J. Neurochem.* **2011**, *117*, 779–796. [[CrossRef](#)]
50. Selkoe, D.; Kopan, R. Notch and Presenilin: Regulated intramembrane proteolysis links development and degeneration. *Annu. Rev. Neurosci.* **2003**, *26*, 565–597. [[CrossRef](#)]
51. De Strooper, B.; Iwatsubo, T.; Wolfe, M.S. Presenilins and  $\gamma$ -secretase: Structure, function, and role in Alzheimer disease. *Cold Spring Harb. Perspect. Med.* **2012**, *2*, a006304. [[CrossRef](#)]
52. Cao, X.; Sudhof, T.C. A transcriptionally [correction of transcriptively] active complex of APP with Fe65 and histone acetyltransferase Tip60. *Science* **2001**, *293*, 115–120. [[CrossRef](#)]
53. von Rotz, R.C.; Kohli, B.M.; Bosset, J.; Meier, M.; Suzuki, T.; Nitsch, R.M.; Konietzko, U. The APP intracellular domain forms nuclear multiprotein complexes and regulates the transcription of its own precursor. *J. Cell Sci.* **2004**, *117*, 4435–4448. [[CrossRef](#)]
54. Edbauer, D.; Willem, M.; Lammich, S.; Steiner, H.; Haass, C. Insulin-degrading enzyme rapidly removes the  $\beta$ -amyloid precursor protein intracellular domain (AICD). *J. Biol. Chem.* **2002**, *277*, 13389–13393. [[CrossRef](#)]
55. Lu, D.C.; Rabizadeh, S.; Chandra, S.; Shayya, R.F.; Ellerby, L.M.; Ye, X.; Salvesen, G.S.; Koo, E.H.; Bredesen, D.E. A second cytotoxic proteolytic peptide derived from amyloid  $\beta$ -protein precursor. *Nat. Med.* **2000**, *6*, 397–404. [[CrossRef](#)] [[PubMed](#)]
56. Cupers, P.; Orlans, I.; Craessaerts, K.; Annaert, W.; De Strooper, B. The amyloid precursor protein (APP) -cytoplasmic fragment generated by gamma-secretase is rapidly degraded but distributes partially in a nuclear fraction of neurones in culture. *J. Neurochem.* **2001**, *78*, 1168–1178. [[CrossRef](#)] [[PubMed](#)]
57. Kimberly, W.T.; Zheng, J.B.; Gue, S.Y.; Selkoe, D.J. The Intracellular Domain of the betaamyloid precursor protein is stabilized by Fe65 and translocates to the nucleus in a notch-like Manner. *J. Biol Chem.* **2001**. [[CrossRef](#)]
58. Fiore, F.; Zambrano, N.; Minopoli, G.; Donini, V.; Duilio, A.; Russo, T. The regions of the Fe65 protein homologous to the phosphotyrosine interaction/phosphotyrosine binding domain of Shc bind the intracellular domain of the Alzheimer's amyloid precursor protein. *J. Biol. Chem.* **1995**. [[CrossRef](#)]

59. Bressler, S.L.; Gray, M.D.; Sopher, B.L.; Hu, Q.; Hearn, M.G.; Pham, D.G.; Dinulos, M.B.; Fukuchi, K.I.; Sisodia, S.S.; Miller, M.A.; et al. cDNA cloning and chromosome mapping of the human Fe65 gene: Interaction of the conserved cytoplasmic domains of the human  $\beta$ -amyloid precursor protein and its homologues with the mouse Fe65 protein. *Hum. Mol. Genet.* **1996**, *5*, 1589–1598. [[CrossRef](#)] [[PubMed](#)]
60. Minopoli, G.; De Candia, P.; Bonetti, A.; Faraonio, R.; Zambrano, N.; Russo, T. The beta-amyloid precursor protein functions as a cytosolic anchoring site that prevents Fe65 nuclear translocation. *J. Biol. Chem.* **2001**, *276*, 6545–6550. [[CrossRef](#)]
61. Ando, K.; Iijima, K.I.; Elliott, J.I.; Kirino, Y.; Suzuki, T. Phosphorylation-dependent regulation of the interaction of amyloid precursor protein with Fe65 affects the production of beta-amyloid. *J. Biol. Chem.* **2001**, *276*, 40353–40361. [[CrossRef](#)] [[PubMed](#)]
62. Perkinton, M.S.; Standen, C.L.; Lau, K.F.; Kesavapany, S.; Byers, H.L.; Ward, M.; McLoughlin, D.M.; Miller, C.C.J. The c-Abl tyrosine kinase phosphorylates the Fe65 adaptor protein to stimulate Fe65/amyloid precursor protein nuclear signaling. *J. Biol. Chem.* **2004**, *279*, 22084–22091. [[CrossRef](#)]
63. Kinoshita, A.; Whelan, C.M.; Smith, C.J.; Berezovska, O.; Hyman, B.T. Direct visualization of the gamma secretase-generated carboxyl-terminal domain of the amyloid precursor protein: Association with Fe65 and translocation to the nucleus. *J. Neurochem.* **2002**, *82*, 839–847. [[CrossRef](#)] [[PubMed](#)]
64. Augustin, V.; Kins, S. Fe65: A Scaffolding Protein of Actin Regulators. *Cells* **2021**, *25*, 1599. [[CrossRef](#)]
65. Nensa, F.M.; Neumann, M.H.D.; Schrotter, A.; Przyborski, A.; Mastalski, T.; Susdalzew, S.; Looße, C.; Helling, S.; El Magraoui, F.; Erdmann, R.; et al. Amyloid beta a4 precursor protein-binding family b member 1 (FE65) interactomics revealed synaptic vesicle glycoprotein 2a (SV2A) and sarcoplasmic/endoplasmic reticulum calcium ATPase 2 (SERCA2) as new binding proteins in the human brain. *Mol. Cell. Proteom.* **2014**, *13*, 475–488. [[CrossRef](#)] [[PubMed](#)]
66. Baek, S.H.; Ohgi, K.A.; Rose, D.W.; Koo, E.H.; Glass, C.K.; Rosenfeld, M.G. Exchange of N-CoR corepressor and Tip60 coactivator complexes links gene expression by NF-kappaB and beta-Amyloid Precursor Protein. *Cell.* **2002**, *110*, 55–67. [[CrossRef](#)]
67. Müller, T.; Schrötter, A.; Looße, C.; Pfeiffer, K.; Theiss, C.; Kauth, M.; Meyer, H.E.; Marcus, K. A ternary complex consisting of AICD, FE65, and TIP60 down-regulates Stathmin1. *Biochim. Biophys. Acta* **2013**, *1834*, 387–394. [[CrossRef](#)]
68. Belyaev, N.D.; Nalivaeva, N.N.; Makova, N.Z.; Turner, A.J. Nephilysin gene expression requires binding of the amyloid precursor protein intracellular domain to its promoter: Implications for Alzheimer disease. *EMBO Rep.* **2009**, *10*, 94–100. [[CrossRef](#)]
69. Pardossi-piquard, R.; Petit, A.; Kawarai, T.; Sunyach, C.; Alves, C.; Vincent, B.; Ring, S.; Adamio, L.D.; Shen, J.; Müller, U.; et al. Presenilin-dependent transcriptional control of the A $\beta$ -degrading enzyme neprilysin by intracellular domains of beta APP and APLP. *Neuron.* **2005**, *46*, 541–554. [[CrossRef](#)]
70. Grimm, M.O.W.; Mett, J.; Stahlmann, C.P.; Grösgen, S.; Haupenthal, V.J.; Blümel, T.; Hundsdörfer, B.; Zimmer, V.C.; Mylonas, N.T.; Tanila, H.; et al. APP intracellular domain derived from amyloidogenic  $\beta$ - and  $\gamma$ -secretase cleavage regulates neprilysin expression. *Front. Aging Neurosci.* **2015**, *19*, 77. [[CrossRef](#)]
71. Grimm, M.O.W.; Mett, J.; Stahlmann, C.P.; Haupenthal, V.J.; Zimmer, V.C.; Hartmann, T. Nephilysin and A $\beta$  clearance: Impact of the APP Intracellular Domain in NEP Regulation and Implications in Alzheimer's Disease. *Front. Aging Neurosci.* **2013**, *23*, 95–98. [[CrossRef](#)] [[PubMed](#)]
72. Kim, H.S.; Kim, E.; Lee, J.; Park, C.H.; Kim, S. C-terminal fragments of amyloid precursor protein exert neurotoxicity by inducing glycogen synthase kinase-3 $\beta$  expression. *FASEB J.* **2003**, *17*, 1951–1953. [[CrossRef](#)]
73. Ozaki, T.; Li, Y.; Kikuchi, H.; Tomita, T.; Iwatsubo, T.; Nakagawara, A. The intracellular domain of the amyloid precursor protein (AICD) enhances the p53-mediated apoptosis. *Biochem. Biophys. Res. Commun.* **2006**, *351*, 57–63. [[CrossRef](#)] [[PubMed](#)]
74. Chang, K.; Kim, H.; Ha, T.; Ha, J.; Shin, K.Y.; Jeong, Y.H.; Lee, J.; Park, C.; Kim, S.; Baik, T.; et al. Phosphorylation of amyloid precursor protein (APP) at Thr668 regulates the nuclear translocation of the APP intracellular domain and induces neurodegeneration. *Mol. Cell. Biol.* **2006**, *26*, 4327–4338. [[CrossRef](#)]
75. Liu, Y.; Hsu, W.; Ma, Y.; Lee, E.H.Y. Melatonin Induction of APP Intracellular Domain 50 SUMOylation Alleviates AD through Enhanced Transcriptional Activation and A $\beta$ Degradation. *Mol. Ther.* **2021**, *29*, 376–395. [[CrossRef](#)]
76. Pastorino, L.; Sun, A.; Lu, P.; Zhou, X.Z.; Balastik, M.; Finn, G.; Wulf, G.; Lim, J.; Li, S.; Li, X.; et al. The prolyl isomerase Pin1 regulates amyloid precursor protein processing and amyloid-beta production. *Nature* **2006**, *440*, 528–534. [[CrossRef](#)] [[PubMed](#)]
77. Ryan, K.A.; Pimplikar, S.W. Activation of GSK-3 and phosphorylation of CRMP2 in transgenic mice expressing APP intracellular domain. *J. Cell Biol.* **2005**, *171*, 327–335. [[CrossRef](#)] [[PubMed](#)]
78. Kirouac, L.; Rajic, A.J.; Cribbs, D.H.; Padmanabhan, J. Activation of Ras-ERK Signaling and GSK-3 by Amyloid Precursor Protein and Amyloid Beta Facilitates Neurodegeneration in Alzheimer's disease. *eNeuro* **2017**, *4*. [[CrossRef](#)] [[PubMed](#)]
79. Zhou, F.; Gong, K.; Van Laar, T.; Gong, Y.; Zhang, L. Wnt / $\beta$ -catenin signal pathway stabilizes APP intracellular domain (AICD) and promotes its transcriptional activity. *Biochem. Biophys. Res. Commun.* **2011**, *412*, 68–73. [[CrossRef](#)] [[PubMed](#)]
80. Jia, L.; Piña-crespo, J.; Li, Y. Restoring Wnt/ $\beta$ -catenin signaling is a promising therapeutic strategy for Alzheimer's disease. *Mol. Brain* **2019**, *12*, 104. [[CrossRef](#)]
81. Zhou, F.; Gong, K.; Song, B.; Ma, T.; Van Laar, T.; Gong, Y.; Zhang, L. The APP intracellular domain (AICD) inhibits Wnt signalling and promotes neurite outgrowth. *Biochim. Biophys. Acta* **2012**, *1823*, 1233–1241. [[CrossRef](#)]
82. Ghosal, K.; Stathopoulos, A.; Pimplikar, S.W. APP intracellular domain impairs adult neurogenesis in transgenic mice by inducing neuroinflammation. *PLoS ONE* **2010**, *5*, e11866. [[CrossRef](#)]
83. Ghosal, K.; Fan, Q.; Dawson, H.N.; Pimplikar, S.W. Tau Protein Mediates APP Intracellular Domain (AICD) -Induced Alzheimer's-Like Pathological Features in Mice. *PLoS ONE* **2016**, *11*, e0159435. [[CrossRef](#)]

84. Ma, Q.; Futagawa, T.; Yang, W.; Jiang, X.; Zeng, L.; Takeda, Y.; Xu, X.; Bagnard, D.; Schachner, M.; Furley, A.J.; et al. A TAG1-APP signalling pathway through Fe65 negatively modulates neurogenesis. *Nat. Cell. Biol.* **2008**, *10*, 283–294. [[CrossRef](#)]
85. Coronel, R.; Lachgar, M.; Bernabeu-zornoza, A.; Palmer, C.; Domínguez-alvaro, M.; Revilla, A.; Ocaña, I.; Fernández, A.; Martínez-serrano, A.; Cano, E.; et al. Neuronal and Glial Differentiation of Human Neural Stem Cells Is Regulated by Amyloid Precursor Protein (APP) Levels. *Mol. Neurobiol.* **2019**, *56*, 1248–1261. [[CrossRef](#)] [[PubMed](#)]
86. Shu, R.; Wong, W.; Ma, Q.H.; Yang, Z.Z.; Zhu, H.; Liu, F.J.; Wang, P.; Ma, J.; Yan, S.; Polo, J.M.; et al. APP intracellular domain acts as a transcriptional regulator of miR-663 suppressing neuronal differentiation. *Cell Death Dis.* **2015**, *6*, e1651. [[CrossRef](#)] [[PubMed](#)]
87. Jiang, M.; Vanan, S.; Tu, H.; Zhang, W.; Zhang, Z.; Chia, S.; Eun, S.; Zeng, X.; Yu, W.; Xu, J.; et al. Amyloid precursor protein intracellular domain-dependent regulation of FOXO3a inhibits adult hippocampal neurogenesis. *Neurobiol. Aging* **2020**, *95*, 250–263. [[CrossRef](#)] [[PubMed](#)]
88. Law, B.M.; Guest, A.L.; Pullen, M.W.J.; Perkinson, M.S. Increased Foxo3a Nuclear Translocation and Activity is an Early Neuronal Response to  $\beta\gamma$ -Secretase-Mediated Processing of the Amyloid- $\beta$  Protein Precursor: Utility of an APP-GAL4 Reporter Assay. *J. Alzheimer's Dis.* **2018**, *61*, 673–688. [[CrossRef](#)] [[PubMed](#)]
89. Goiran, T.; Duplan, E.; Chami, M.; Bourgeois, A.; El Manaa, W.; Rouland, L.; Dunys, J.; Lauritzen, I.; You, H.; Stambolic, V.; et al.  $\beta$ -Amyloid Precursor Protein Intracellular Domain Controls Mitochondrial Function by Modulating Phosphatase and Tensin Homolog—Induced Kinase 1 Transcription in Cells and in Alzheimer Mice Models. *Biol. Psychiatry* **2018**, *83*, 416–427. [[CrossRef](#)]
90. Maloney, B.; Lahiri, D.K. The Alzheimer's amyloid  $\beta$ -peptide ( $A\beta$ ) binds a specific DNA  $A\beta$ -interacting domain ( $A\beta$ ID) in the APP, BACE1, and APOE promoters in a sequence-specific manner: Characterizing a new regulatory motif. *Gene* **2011**, *488*, 1–12. [[CrossRef](#)]
91. Gezen, D.; İrem, A.; Esin, L.A.; Merve, C.; Erdinç, A. The Transcriptional Regulatory Properties of Amyloid Beta 1–42 may Include Regulation of Genes Related to Neurodegeneration. *Neuromol. Med.* **2018**, *20*, 363–375. [[CrossRef](#)]
92. Bailey, J.A.; Maloney, B.; Ge, Y.; Lahiri, D.K. Functional activity of the novel Alzheimer's amyloid  $\beta$ -peptide interacting domain ( $A\beta$  ID) in the APP and BACE1 promoter sequences and implications in activating apoptotic genes and in amyloidogenesis. *Gene* **2011**, *488*, 13–22. [[CrossRef](#)] [[PubMed](#)]
93. Barucker, C.; Harmeyer, A.; Weiske, J.; Fauler, B.; Albring, K.F.; Prokop, S.; Hildebrand, P.; Lurz, R.; Heppner, F.L.; Huber, O.; et al. Nuclear Translocation Uncovers the Amyloid Peptide  $A\beta$ 42 as a Regulator of Gene Transcription. *J. Biol. Chem.* **2014**, *289*, 20182–20191. [[CrossRef](#)] [[PubMed](#)]
94. Barucker, C.; Sommer, A.; Beckmann, G.; Eravci, M.; Harmeyer, A. Alzheimer Amyloid Peptide  $a\beta$ 42 Regulates Gene Expression of Transcription and Growth Factors. *J. Alzheimers Dis.* **2015**, *44*, 613–624. [[CrossRef](#)] [[PubMed](#)]
95. Gezen, D. Vitamin D Receptor Regulates Amyloid Beta 1–42 Production with Protein Disulfide Isomerase A3. *ACS Chem. Neurosci.* **2017**, *8*, 2335–2346. [[CrossRef](#)]
96. Weingarten, M.D.; Lockwood, A.H.; Hwo, S.-Y.; Kirschner, M.W. A protein factor essential for microtubule assembly. *Proc. Natl. Acad. Sci. USA* **1975**, *72*, 1858–1862. [[CrossRef](#)]
97. Buée, L.; Bussiére, T.; Buée-Scherrer, V.; Delacourte, A.; Hof, P.R. Tau protein isoforms, phosphorylation and role in neurodegenerative disorders. *Brain Res. Brain Res. Rev.* **2000**, *33*, 95–130. [[CrossRef](#)]
98. Martin, L.; Latypova, X.; Terro, F. Post-translational modifications of tau protein: Implications for Alzheimer's disease. *Neurochem. Int.* **2011**, *58*, 458–471. [[CrossRef](#)]
99. Maina, M.B.; Al-hilaly, Y.K.; Serpell, L.C. Nuclear Tau and Its Potential Role in Alzheimer's Disease. *Biomolecules* **2016**, *6*, 9. [[CrossRef](#)]
100. Liu, C.; Götz, J. Profiling murine tau with 0N, 1N and 2N isoform-specific antibodies in brain and peripheral organs reveals distinct subcellular localization, with the 1N isoform being enriched in the nucleus. *PLoS ONE* **2013**, *8*, e84849. [[CrossRef](#)]
101. Greenwood, J.A.; Johnson, G.V. Localization and in situ phosphorylation state of nuclear tau. *Exp. Cell Res.* **1995**, *220*, 332–337. [[CrossRef](#)]
102. Loomis, P.A.; Howardt, T.H.; Castleberry, R.P.; Binder, L. 1 Identification of nuclear X isoforms in human neuroblastoma cells. *Proc. Natl. Acad. Sci. USA* **1990**, *87*, 8422–8426. [[CrossRef](#)]
103. Violet, M.; Delattre, L.; Tardivel, M.; Sultan, A.; Chauderlier, A.; Caillierez, R.; Talahari, S.; Nesslany, F.; Lefebvre, B.; Bonnefoy, E.; et al. A major role for Tau in neuronal DNA and RNA protection in vivo under physiological and hyperthermic conditions. *Front. Cell. Neurosci.* **2014**. [[CrossRef](#)]
104. Sultan, A.; Nesslany, F.; Violet, M.; Begard, S.; Loyens, A.; Talahari, S.; Mansuroglu, Z.; Marzin, D.; Sergeant, N.; Humez, S.; et al. Nuclear tau, a key player in neuronal DNA protection. *J. Biol. Chem.* **2011**. [[CrossRef](#)]
105. Brady, R.M.; Zinkowski, R.P.; Binder, L.I. Presence of tau in isolated nuclei from human brain. *Neurobiol. Aging* **1995**, *16*, 479–486. [[CrossRef](#)]
106. Thurston, V.C.; Zikowski, R.P.; Binder, L.I. Tau as a nucleolar protein in human nonneural cells in vitro and in vivo. *Chromosoma* **1996**, *105*, 20–30. [[CrossRef](#)] [[PubMed](#)]
107. Sjöberg, M.K.; Shestakova, E.; Mansuroglu, Z.; Maccioni, R.B.; Bonnefoy, E. Tau protein binds to pericentromeric DNA: A putative role for nuclear tau in nucleolar organization. *J. Cell Sci.* **2006**, *19*, 2025–2034. [[CrossRef](#)] [[PubMed](#)]
108. Wiche, G.; Corces, V.G.; Avila, J. Preferential binding of hog brain microtubule-associated proteins to mouse satellite versus bulk DNA preparations. *Nature* **1978**, *273*, 403–405. [[CrossRef](#)] [[PubMed](#)]

109. Corces, V.G.; Salas, J.; Salas, M.L.; Avila, J. Binding of microtubule proteins to DNA: Specificity of the interaction. *Eur. J. Biochem.* **1978**, *86*, 473–479. [[CrossRef](#)] [[PubMed](#)]
110. Hua, Q.; He, R.Q.; Haque, N.; Qu, M.H.; del Carmen Alonso, A.I.; Grundke-Iqbal, I.; Iqbal, K. Microtubule associated protein tau binds to double-stranded but not single-stranded DNA. *Cell. Mol. Life Sci.* **2003**, *60*, 413–421. [[CrossRef](#)] [[PubMed](#)]
111. Wei, Y.; Qu, M.H.; Wang, X.S.; Chen, L.; Wang, D.L.; Liu, Y.; Hua, Q.; He, R.Q. Binding to the minor groove of the double-strand, tau protein prevents DNA from damage by peroxidation. *PLoS ONE* **2008**, *3*, e2600. [[CrossRef](#)] [[PubMed](#)]
112. Camero, S.; Benítez, M.J.; Barrantes, A.; Ayuso, J.M.; Cuadros, R.; Avila, J.; Jiménez, J.S. Tau protein provides DNA with thermodynamic and structural features which are similar to those found in histone-DNA complex. *J. Alzheimer's Dis.* **2014**, *39*, 649–660. [[CrossRef](#)]
113. Multhaup, G.; Huber, O.; Galas, M. Amyloid Precursor Protein (APP) Metabolites APP Intracellular Fragment (AICD), A $\beta$ 42, and Tau in Nuclear Roles. *J. Biol. Chem.* **2015**, *290*, 23515–23522. [[CrossRef](#)]
114. Qi, H.; Cantrelle, F.X.; Benhelli-Mokrani, H.; Smet-Nocca, C.; Buée, L.; Lippens, G.; Bonnefoy, E.; Galas, M.C.; Landrieu, I. Nuclear magnetic resonance spectroscopy characterization of interaction of Tau with DNA and its regulation by phosphorylation. *Biochemistry* **2015**, *54*, 1525–1533. [[CrossRef](#)] [[PubMed](#)]
115. Frost, B.; Hemberg, M.; Lewis, J.; Feany, M.B. Tau promotes neurodegeneration through global chromatin relaxation. *Nat. Neurosci.* **2014**, *17*, 357–366. [[CrossRef](#)] [[PubMed](#)]
116. Mansuroglu, Z.; Benhelli-Mokrani, H.; Marcato, V.; Sultan, A.; Violet, M.; Chauderlier, A.; Delattre, L.; Loyens, A.; Talahari, S.; Bégard, S.; et al. Loss of Tau protein affects the structure, transcription and repair of neuronal pericentromeric heterochromatin. *Sci. Rep.* **2016**, *6*, 33047. [[CrossRef](#)] [[PubMed](#)]
117. Maina, M.B.; Bailey, L.J.; Wagih, S.; Biasetti, L.; Pollack, S.J.; Quinn, J.P.; Thorpe, J.R.; Doherty, A.J.; Serpell, L.C. The involvement of tau in nucleolar transcription and the stress response. *Acta Neuropathol. Commun.* **2018**, *6*, 70. [[CrossRef](#)] [[PubMed](#)]
118. Faghihi, M.A.; Modarresi, F.; Khalil, A.M.; Wood, D.E.; Sahagan, B.G.; Morgan, T.E.; Finc, C.E.; St Laurent, G.; Kenny, P.J.; Wahlestedt, C. Expression of a noncoding RNA is elevated in Alzheimer's disease and drives rapid feed-forward regulation of beta-secretase. *Nat. Med.* **2008**, *14*, 723–730. [[CrossRef](#)]
119. Klein, H.U.; McCabe, C.; Gjonasca, E.; Sullivan, S.E.; Kaskow, B.J.; Tang, A.; Smith, R.V.; Xu, J.; Pfenning, A.R.; Bernstein, B.E.; et al. Epigenome-wide study uncovers large-scale changes in histone acetylation driven by tau pathology in aging and Alzheimer's human brains. *Nat. Neurosci.* **2019**, *22*, 37–46. [[CrossRef](#)]
120. Nativio, R.; Lan, Y.; Donauhe, G.; Sidoli, S.; Berson, A.; Srinivasan, A.R.; Shcherbakova, O.; Amlie-Wolf, A.; Nie, J.; Cui, X.; et al. An integrated multi-omics approach identifies epigenetic alterations associated with Alzheimer's disease. *Nat. Genet.* **2020**, *52*, 1024–1035. [[CrossRef](#)]
121. Olson, M.O.J. Nucleolus: Structure and Function. *Encycl. Life Sci.* **2006**. [[CrossRef](#)]
122. Lu, J.; Miao, J.; Su, T.; Liu, Y.; He, R. Formaldehyde induces hyperphosphorylation and polymerization of Tau protein both in vitro and in vivo. *Biochim. Biophys. Acta.* **2013**, *1830*, 4102–4116. [[CrossRef](#)] [[PubMed](#)]
123. Hernández-Ortega, K.; Garcia-Esparcia, P.; Gil, L.; Lucas, J.J.; Ferrer, I. Altered Machinery of Protein Synthesis in Alzheimer's: From the Nucleolus to the Ribosome. *Brain Pathol.* **2016**, *26*, 593–605. [[CrossRef](#)] [[PubMed](#)]
124. De Jager, P.L.; Liu, Z.; Shulman, J.M. Tau Activates Transposable Elements in Alzheimer's Disease. *Cell. Rep.* **2018**, *23*, 2874–2880. [[CrossRef](#)]
125. Sun, W.; Samini, H.; Gamez, M.; Zare, H.; Frost, B. Pathogenic tau-induced piRNA depletion promotes neuronal death through transposable element dysregulation in neurodegenerative tauopathies. *Nat. Neurosci.* **2018**, *21*, 1038–1048. [[CrossRef](#)] [[PubMed](#)]
126. Li, W.; Lee, M.H.; Henderson, L.; Tyagi, R.; Bachani, M.; Steiner, J.; Campanac, E.; Hoffman, D.A.; von Geldern, G.; Jhonson, K.; et al. Human endogenous retrovirus-K contributes to motor neuron disease. *Sci. Transl. Med.* **2015**, *7*, 307ra153. [[CrossRef](#)]
127. Bading, H. Nuclear calcium signalling in the regulation of brain function. *Nat. Rev. Neurosci.* **2013**, *14*, 593–608. [[CrossRef](#)]
128. Mahoney, R.; Thomas, E.O.; Ramirez, P.; Miller, H.E.; Beckmann, A.; Zuniga, G.; Dobrowolski, R.; Frost, B. Pathogenic Tau Causes a Toxic Depletion of Nuclear Calcium II Pathogenic Tau Causes a Toxic Depletion of Nuclear Calcium. *Cell. Rep.* **2020**, *32*, 107900. [[CrossRef](#)]
129. Huang, Y.; Weisgraber, K.H.; Mucke, L.; Mahley, R.W. Apolipoprotein E: Diversity of cellular origin, structural and biophysical properties, and effects in Alzheimer's Disease. *J. Mol. Neurosci.* **2004**, *23*, 189–204. [[CrossRef](#)]
130. Grehan, S.; Tse, E.; Taylor, J.M. Two Distal Downstream Enhancers Direct Expression of the Human Apolipoprotein E Gene to Astrocytes in the Brain. *J. Neurosci.* **2001**, *21*, 812–822. [[CrossRef](#)]
131. Ignatius, M.J.; Gebicke-haerter, P.J.; Pitasb, R.E.; Shootera, E.M. Apolipoprotein E in nerve injury and repair. *Prog. Brain. Res.* **1987**, *71*, 177–178. [[CrossRef](#)]
132. Huang, Y.; Mahley, R.W. Apolipoprotein E: Structure and function in lipid metabolism, neurobiology, and Alzheimer's diseases. *Neurobiol. Dis.* **2014**, *72*. [[CrossRef](#)] [[PubMed](#)]
133. Theendakara, V.; Peters-libeu, C.A.; Spilman, X.P.; Poksay, K.S.; Bredesen, D.E.; Rao, R.V. Direct Transcriptional Effects of Apolipoprotein. *Eur. J. Neurosci.* **2016**, *36*, 685–700. [[CrossRef](#)] [[PubMed](#)]
134. Sen, A.; Nelson, T.J.; Alkon, D.L. ApoE4 and A $\beta$  Oligomers Reduce BDNF Expression via HDAC Nuclear Translocation. *J. Neurosci.* **2015**, *35*, 7538–7551. [[CrossRef](#)]
135. Huang, Y.A.; Zhou, B.; Wernig, M.; Südhof, T.C. APOE2, APOE3, and APOE4 Differentially Stimulate APP Transcription and A $\beta$  Secretion. *Cell* **2017**, *168*, 427–441. [[CrossRef](#)] [[PubMed](#)]



136. Theendakara, V.; Patent, A.; Peters, C.A.; Philpot, B.; Flores, S.; Descamps, O.; Poksay, K.S.; Zhang, Q.; Cailing, G.; Hart, M.; et al. Neuroprotective Sirtuin ratio reversed by APOE4. *Proc. Natl. Acad. Sci. USA* **2013**, *110*, 18303–18308. [[CrossRef](#)] [[PubMed](#)]
137. Lin, Y.T.; Seo, J.; Gao, F.; Feldman, M.H.; Wen, H.; Penney, J.; Cam, H.P.; GJoneska, E.; Raja, W.; Cheng, J.; et al. APOE4 Causes Widespread Molecular and Cellular Alterations Associated with Alzheimer's Disease Phenotypes in Human iPSC-Derived Brain Cell Types. *Neuron* **2018**, *98*, 1141–1154. [[CrossRef](#)]
138. Lee, H.R.; Shin, H.K.; Park, S.Y.; Kim, H.Y.; Lee, W.S.; Rhim, B.Y.; Hong, K.W.; Kim, C.D. Cilostazol Suppresses  $\beta$ -Amyloid Production by Activating A Disintegrin and Metalloproteinase 10 Via the Upregulation of SIRT1-Coupled Retinoic Acid Receptor- $\beta$ . *J. Neurosci. Res.* **2014**, *92*, 1581–1590. [[CrossRef](#)]
139. Zhao, J.; Fu, Y.; Yamazaki, Y.; Ren, Y.; Davis, M.D.; Liu, C.C.; Lu, W.; Wang, X.; Chen, K.; Cherukuri, Y.; et al. APOE4 exacerbates synapse loss and neurodegeneration in Alzheimer's disease patient iPSC-derived cerebral organoids. *Nat. Commun.* **2020**, *11*, 5540. [[CrossRef](#)]
140. Chen, X.; Zhang, Y.; Xu, H.; Bu, G. Transcriptional regulation and its misregulation in Alzheimer's disease. *Mol. Brain* **2013**, *6*, 44. [[CrossRef](#)]
141. Bagyinszky, E.; Giau, V.; An, S.A. Transcriptomics in Alzheimer's Disease: Aspects and Challenges. *Int. J. Mol. Sci.* **2020**, *21*, 3517. [[CrossRef](#)]
142. Verheijen, J.; Sleegers, K. Understanding Alzheimer Disease at the Interface between Genetics and Transcriptomics. *Trends Genet.* **2018**, *34*, 434–447. [[CrossRef](#)] [[PubMed](#)]
143. Chen, W.; Lu, A.; Craessaerts, K.; Pavie, B.; Frigerio, C.S.; Corthout, N.; Qian, K.; Laláková, J.; Kühnemund, M.; Voytyuk, I.; et al. Spatial Transcriptomics and In Situ Sequencing to Study Alzheimer's Disease II Resource Spatial Transcriptomics and In Situ Sequencing to Study Alzheimer's Disease. *Cell* **2020**, *182*, 976–991. [[CrossRef](#)] [[PubMed](#)]
144. Pickett, E.K.; Herrmann, A.G.; McQueen, J.; Abt, K.; Dando, O.; Tulloch, J.; Jain, P.; Dunnett, S.; Sohrabi, S.; Fjeldsta, M.P.; et al. Amyloid Beta and Tau Cooperate to Cause Reversible Behavioral and Transcriptional Deficits in a Model of Alzheimer's Disease. *Cell. Rep.* **2019**, *29*, 3592–3604. [[CrossRef](#)] [[PubMed](#)]
145. Zhong, S.; Wang, M.; Zhan, Y.; Zhang, J.; Yang, X.; Fu, S.; Bi, D.; Gao, F.; Shen, Y.; Chen, Z. Single-nucleus RNA sequencing reveals transcriptional changes of hippocampal neurons in APP23 mouse model of Alzheimer's disease. *Biosci. Biotechnol. Biochem.* **2020**, *84*, 919–926. [[CrossRef](#)] [[PubMed](#)]
146. Magistri, M.; Velmeshev, D.; Makhmutova, M.; Faghihi, M.A. Transcriptomics Profiling of Alzheimer's Disease Reveal Neurovascular Defects, Altered Amyloid- $\beta$  Homeostasis, and Deregulated Expression of Long Noncoding RNAs. *J. Alzheimer's Dis.* **2015**, *48*, 647–665. [[CrossRef](#)] [[PubMed](#)]
147. Mathys, H.; Davila-Velderrain, J.; Peng, Z.; Gao, F.; Mohammadi, S.; Young, J.Z.; Menon, M.; He, L.; Abdurrob, F.; Jiang, X.; et al. Single-cell transcriptomic analysis of Alzheimer's disease. *Nature* **2019**, *570*, 332–337. [[CrossRef](#)]
148. Wan, Y.; Al-ouran, R.; Mangleburg, C.G.; Perumal, T.M.; Lee, T.V.; Allison, K.; Swarup, V.; Funk, C.C.; Gaiteri, C.; Allen, M.; et al. Resource Meta-Analysis of the Alzheimer's Disease Human Brain Transcriptome and Functional Dissection in Mouse Models. *Cell. Rep.* **2020**, *32*, 107908. [[CrossRef](#)]
149. Jansen, I.E.; Savage, J.E.; Watanabe, K.; Bryois, J.; Williams, D.M.; Steinberg, S.; Sealock, J.; Karlsson, I.K.; Hägg, S.; Athanasiu, L.; et al. Genome-wide meta-analysis identifies new loci and functional pathways influencing Alzheimer's disease risk. *Nat. Genet.* **2019**, *51*, 404–413. [[CrossRef](#)]