

Journal de l'École polytechnique

Mathématiques

Andrea ASPRI, Elena BERETTA, & Corrado MASCIA

Analysis of a Mogi-type model describing surface deformations induced by a magma chamber embedded in an elastic half-space

Tome 4 (2017), p. 223-255.

http://jep.cedram.org/item?id=JEP_2017__4__223_0

© Les auteurs, 2017.

Certains droits réservés.



Cet article est mis à disposition selon les termes de la licence
CREATIVE COMMONS ATTRIBUTION – PAS DE MODIFICATION 3.0 FRANCE.
<http://creativecommons.org/licenses/by-nd/3.0/fr/>

L'accès aux articles de la revue « Journal de l'École polytechnique — Mathématiques » (<http://jep.cedram.org/>), implique l'accord avec les conditions générales d'utilisation (<http://jep.cedram.org/legal/>).

Publié avec le soutien
du Centre National de la Recherche Scientifique

cedram

Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>

ANALYSIS OF A MOGI-TYPE MODEL DESCRIBING SURFACE DEFORMATIONS INDUCED BY A MAGMA CHAMBER EMBEDDED IN AN ELASTIC HALF-SPACE

BY ANDREA ASPRI, ELENA BERETTA & CORRADO MASCIA

ABSTRACT. — Motivated by a vulcanological problem, we establish a sound mathematical approach for surface deformation effects generated by a magma chamber embedded into Earth's interior and exerting on it a uniform hydrostatic pressure. Modeling assumptions translate the problem into classical elasto-static system (homogeneous and isotropic) in a half-space with an embedded cavity. The boundary conditions are traction-free for the air/crust boundary and uniformly hydrostatic for the chamber boundary. These are complemented with zero-displacement condition at infinity (with decay rate). After a short presentation of the model and of its geophysical interest, we establish the well-posedness of the problem and provide an appropriate integral formulation for its solution for cavity with general shape. Based on that, assuming that the chamber is centered at some fixed point \mathbf{z} and has diameter $r > 0$, small with respect to the depth d , we derive rigorously the principal term in the asymptotic expansion for the surface deformation as $\varepsilon = r/d \rightarrow 0^+$. Such a formula provides a rigorous proof of the Mogi point source model in the case of spherical cavities generalizing it to the case of cavities of arbitrary shape.

RÉSUMÉ (Analyse d'un modèle du type de Mogi décrivant les déformations de surface induites par une chambre magmatique contenue dans un demi-espace élastique)

Motivés par un problème volcanologique, nous établissons une approche mathématique solide pour les effets de déformation de surface engendrés par une chambre magmatique contenue à l'intérieur de la Terre et soumise à une pression hydrostatique uniforme. Des hypothèses de modélisation traduisent le problème en un système élasto-statique (homogène et isotrope) classique dans un demi-espace avec une cavité incluse. Les conditions au bord sont sans traction au bord air/croûte, et uniformément hydrostatiques au bord de la chambre. Elles sont complétées par une condition de déplacement nul à l'infini (avec taux de décroissance). Après une courte présentation du modèle et de son intérêt géophysique, nous établissons que le problème est bien posé et proposons une formulation intégrale appropriée pour sa solution dans le cas d'une cavité de forme générale. En conséquence, supposant que la chambre est centrée en un point \mathbf{z} et est de diamètre $r > 0$ petit par rapport à la profondeur d , nous en déduisons rigoureusement le terme principal du développement asymptotique de la déformation de surface sous la forme $\varepsilon = r/d \rightarrow 0^+$. Une telle formule permet de donner une preuve rigoureuse du modèle des points sources de Mogi dans le cas des cavités de forme arbitraire, qui généralise celle dans le cas des cavités sphériques.

CONTENTS

1. Introduction.....	224
2. Description of the mathematical model.....	227

MATHEMATICAL SUBJECT CLASSIFICATION (2010). — 35C20, 31B10, 35J25, 86A60.

KEYWORDS. — Lamé operator, asymptotic expansions, single and double layer potentials.

3. Integral representation and well-posedness of the direct problem.....	229
4. Rigorous derivation of the asymptotic expansion.....	241
Appendix. Neumann function for the half-space with zero traction.....	248
References.....	253

1. INTRODUCTION

Measurements of crustal deformations are crucial in studying and monitoring volcanoes activity. In this context, one of the main goals is to investigate the mechanics of volcanic systems in order to obtain models for which the predicted synthetic data best fit the observed ones. Indeed, if the model is accurate it gives good images of the Earth's interior. In particular, concerning volcanoes monitoring, synthetic data might be used to localize underground magma chambers and their volume changes. In this sense, a well-established and widely used model, in the geological literature, is the one proposed by Mogi in [33] where the magma chamber is modeled by a pressurized spherical cavity of radius r , buried in a homogeneous, isotropic, elastic half-space (\mathbb{R}_+^3) at depth $d \ll r$. Specifically, Mogi provides an explicit formula of the leading term of the displacement field on the boundary (\mathbb{R}^2) of the half-space when the ratio r/d goes to zero. This model has been further investigated by McTigue in [30] who extends the analysis by detecting the second order term in the asymptotic formula. For other geometries, we mention [16] where Davies considers the case of a pressurized cavity in the shape of prolate and oblate ellipsoids, restricted to radius ratio to depth sufficiently small, deriving, also in this case, an explicit formula of the leading order term.

We stress that the analysis carried out in [16, 30, 33] and the derivation of the formulas therein are informal. For this reason the principal aim of this article is to contribute with a rigorous mathematical analysis of such modeling with the crucial difference relative to the shape of the cavity; our analysis does not require this shape to be a sphere or an ellipsoid but a general subdomain of the half-space. More precisely, we derive a rigorous asymptotic expansion of the boundary displacement vector field due to the presence of a cavity C , buried in an isotropic, homogeneous and elastic half-space (\mathbb{R}_+^3), of the form

$$C = C_\varepsilon := \mathbf{z} + \varepsilon\Omega,$$

where \mathbf{z} is the center of the cavity, ε is a suitable scaling parameter (see Section 2) and Ω , shape of the cavity, is a bounded Lipschitz domain containing the origin. Furthermore, we follow the assumption in [16, 33] supposing that the boundary of the cavity C is subjected to a constant pressure p . In details, denoting by \mathbf{u}_ε the displacement vector field, we have, for $\mathbf{y} \in \mathbb{R}^2$,

$$(1.1) \quad u_\varepsilon^k(\mathbf{y}) = \varepsilon^3 |\Omega| p \widehat{\nabla}_{\mathbf{z}} \mathbf{N}^{(k)}(\mathbf{z}, \mathbf{y}) : \mathbb{M}\mathbf{I} + O(\varepsilon^4),$$

for $k = 1, 2, 3$, as $\varepsilon \rightarrow 0$ (see Theorem 4.1), where u_ε^k stands for the k -th component of the displacement vector. Here $p\varepsilon^3$ represents the total work exerted by the cavity

on the half-space, $\widehat{\nabla} \mathbf{N}^{(k)}$ denotes the symmetric part of the gradient related to the k -th column vector of the *Neumann function* that is the solution to

$$\operatorname{div}(\mathbb{C} \widehat{\nabla} \mathbf{N}(\cdot, \mathbf{y})) = \delta_{\mathbf{y}} \mathbf{I} \quad \text{in } \mathbb{R}_-^3, \quad \partial \mathbf{N}(\cdot, \mathbf{y}) / \partial \boldsymbol{\nu} = \mathbf{0} \quad \text{on } \mathbb{R}^2,$$

with the decay conditions at infinity

$$\mathbf{N} = O(|\mathbf{x}|^{-1}), \quad |\nabla \mathbf{N}| = O(|\mathbf{x}|^{-2}),$$

where $\partial / \partial \boldsymbol{\nu}$ is the conormal derivative operator. In addition, \mathbf{I} is the identity matrix in \mathbb{R}^3 and \mathbb{M} is the fourth-order moment elastic tensor defined by

$$\mathbb{M} := \mathbb{I} + \frac{1}{|\Omega|} \int_{\partial \Omega} \mathbb{C}(\boldsymbol{\theta}^{qr}(\boldsymbol{\zeta}) \otimes \mathbf{n}(\boldsymbol{\zeta})) d\sigma(\boldsymbol{\zeta}),$$

for $q, r = 1, 2, 3$, where \mathbb{I} is the symmetric identity tensor, \mathbb{C} is the isotropic elasticity tensor and \mathbf{n} is the outward unit normal vector to $\partial \Omega$. Finally, the functions $\boldsymbol{\theta}^{qr}$, with $q, r = 1, 2, 3$, are the solutions to

$$\operatorname{div}(\mathbb{C} \widehat{\nabla} \boldsymbol{\theta}^{qr}) = 0 \quad \text{in } \mathbb{R}^3 \setminus \Omega, \quad \frac{\partial \boldsymbol{\theta}^{qr}}{\partial \boldsymbol{\nu}} = -\frac{1}{3\lambda + 2\mu} \mathbb{C} \mathbf{n} \quad \text{on } \partial \Omega,$$

with the decay conditions at infinity

$$|\boldsymbol{\theta}^{qr}| = O(|\mathbf{x}|^{-1}), \quad |\nabla \boldsymbol{\theta}^{qr}| = O(|\mathbf{x}|^{-2}), \quad \text{as } |\mathbf{x}| \rightarrow \infty.$$

The leading term in (1.1) contains the information on the location of the cavity, \mathbf{z} , on its shape and on the work $p \varepsilon^3$, hence it might be used to detect the unknown cavity if boundary measurements of the displacement field are given.

In order to establish these results we use the approach introduced by Ammari and Kang, based on layer potentials techniques for bounded domains, see for example [4, 5, 6], and following the path outlined in [8]. We highlight that proving the existence and uniqueness results for unbounded domains with unbounded boundary is, in general, much more difficult with respect to the case of bounded or exterior domains. The main obstacle is the control of both solution decay and integrability on the boundary. Indeed, the usual approach is based on the employment of some weighted functional spaces, see for example [7]. Here, taking advantage of the explicit formula of the Neumann function \mathbf{N} and rewriting the problem into an integral formulation via layer potential techniques, we are able to prove the existence and uniqueness of the displacement field generated by an arbitrary finite cavity C in a standard Sobolev setting. These results follow showing the invertibility of the integral operators, some of them with singular kernel, that come from the layer potential technique. After that, we derive the asymptotic expansion (1.1), as ε goes to zero, for a cavity of a generic shape.

The approach based on the determination of asymptotic expansions in the case of small inclusions or cavities in bounded domains, which goes back to Friedman and Vogelius [22], has been extensively and successfully used for the reconstruction, from boundary measurements, of location and geometrical features of unknown conductivity inhomogeneities in the framework of electrical impedance tomography (see, for

example, [11] and [5] for an extended bibliography). In the last decade some of these results have been extended to elasticity in the case of bounded domains, see [4]. Specifically, starting from boundary measurements given by the couple potentials/currents or deformations/tractions, in the case, respectively, of electrical impedance tomography and linear elasticity, information about the conductivity profile and the elastic parameters of the medium have been inferred. It is well known that without any a priori assumptions on the features of the problem, the reconstruction procedures give poor quality results. This is due to the severe ill-posedness of the inverse boundary value problem modeling both the electrical impedance tomography [2] and the elasticity problems [3, 34]. However, in certain situations one has some a priori information about the structure of the medium to be reconstructed. These additional details allow to restore the well-posedness of the problem and, in particular, to gain uniqueness and Lipschitz continuous dependence of inclusions or cavities from the boundary measurements. One way to proceed, for instance, is to consider the medium with a smooth background conductivity or elastic parameters except for a finite number of small inhomogeneities [4, 5]. Therefore, by means of partial or complete asymptotic formulas of solutions to the conductivity/elastic problems and some efficient algorithms, information about the location and size of the inclusions can be reconstructed, see for example [4, 5, 24].

Despite the extensive literature in this field, we remark that the mathematical problem of this work represents an intriguing novelty because we have to deal with a pressurized cavity, that is, a hole with nonzero tractions on its boundary, buried in a half-space. These two peculiarities do not allow to reduce the boundary value problem to a classical one based on cavities (see, for example, problems in [4, 5] and reference therein).

The paper is organized as follows. In Section 2, we illustrate the model of ground deformations and motivate the assumptions at the basis of the simplified linearized version. In Section 3, we recall some arguments about linear elasticity and layer potentials techniques and then we analyze the well-posedness of the direct problem via an integral representation formula for the displacement field. Section 4 is devoted to the proof of the main result regarding the asymptotic formula for the boundary displacement field. In addition, as a consequence of the asymptotic expansion, we obtain the classical Mogi's formula for spherical cavities.

NOTATION. — We denote scalar quantities in italic type, e.g. λ, μ, ν , points and vectors in bold italic type, e.g. $\mathbf{x}, \mathbf{y}, \mathbf{z}$ and $\mathbf{u}, \mathbf{v}, \mathbf{w}$, matrices and second-order tensors in bold type, e.g. $\mathbf{A}, \mathbf{B}, \mathbf{C}$, and fourth-order tensors in blackboard bold type, e.g. $\mathbb{A}, \mathbb{B}, \mathbb{C}$. The half-space $\{\mathbf{x} = (x_1, x_2, x_3) \in \mathbb{R}^3 : x_3 < 0\}$ is indicated by \mathbb{R}_-^3 . The unit outer normal vector to a surface is represented by \mathbf{n} . The transpose of a second-order tensor \mathbf{A} is denoted by \mathbf{A}^T and its symmetric part by $\widehat{\mathbf{A}} = \frac{1}{2}(\mathbf{A} + \mathbf{A}^T)$. To indicate the inner product between two vectors \mathbf{u} and \mathbf{v} we use $\mathbf{u} \cdot \mathbf{v} = \sum_i u_i v_i$ whereas for second-order tensors $\mathbf{A} : \mathbf{B} = \sum_{i,j} a_{ij} b_{ij}$. The cross product of two vectors \mathbf{u} and \mathbf{v} is denoted by $\mathbf{u} \times \mathbf{v}$. With $|\mathbf{A}|$ we mean the norm induced by the inner product between second-order tensors, that is, $|\mathbf{A}| = \sqrt{\mathbf{A} : \mathbf{A}}$.

Acknowledgements. — The authors are indebted with Maurizio Battaglia for drawing the attention to the problem and for some useful discussion on its volcanological significance. The authors thank the New York University in Abu Dhabi (EAU) for its kind hospitality that permitted a further development of the present research.

2. DESCRIPTION OF THE MATHEMATICAL MODEL

Monitoring of volcanoes activity, targeted to the forecasting of volcanic hazards and development of appropriate prevention strategies, is usually performed by combining different types of geophysical measurements. Ground deformations are among the most significant being directly available (in particular, with the development of Global Positioning Systems) and, at the same time, straightly interpretable in term of elastic behaviors of the Earth's crust adjacent to the magma chamber (see [10, 18, 37]). A well-established model has been proposed by Mogi, [33], following previous results (see description in [17, 28, 36]), based on the assumption that ground deformation effects are primarily generated by the presence of an underground magma chamber exerting a uniform pressure on the surrounding medium. Precisely, the model relies on three key founding schematizations.

1. *Geometry of the model.* — The earth's crust is an infinite half-space (with free air/crust surface located on the plane $x_3 = 0$) and the magma chamber, buried in the half-space, is assumed to be spherical with radius r and depth d such that $r \ll d$.

2. *Geophysics of the crust.* — The crust is a perfectly elastic body, isotropic and homogeneous, whose deformations are described by the linearized elastostatic equations, hence are completely characterized by the Lamé parameters μ, λ (or, equivalently, Poisson ratio ν and shear modulus μ). The free air/crust boundary is assumed to be a traction-free surface.

3. *Crust-chamber interaction.* — The cavity describing the magma chamber is assumed to be filled with an ideal incompressible fluid at equilibrium, so that the pressure p exerted on its boundary on the external elastic medium is hydrostatic and uniform.

Assuming that the center of the sphere is located at $\mathbf{z} = (z_1, z_2, z_3)$ with $z_3 < 0$, the displacement $\mathbf{u} = (u_1, u_2, u_3)$ at a surface point $\mathbf{y} = (y_1, y_2, 0)$ is given by

$$(2.1) \quad u^\alpha(\mathbf{y}) = \frac{1-\nu}{\mu} \frac{\varepsilon^3 p (z_\alpha - y_\alpha)}{|\mathbf{z} - \mathbf{y}|^3}, \quad (\alpha = 1, 2) \quad u^3(\mathbf{y}) = \frac{1-\nu}{\mu} \frac{\varepsilon^3 p z_3}{|\mathbf{z} - \mathbf{y}|^3}$$

in the limit $\varepsilon := r/|z_3| \rightarrow 0$ (see Fig. 2.1). A higher-order approximation has been proposed by McTigue [30] with the intent of providing a formal expansion able to cover the case of a spherical body with finite (but small) positive radius.

Being based on the assumption that the ratio radius/depth $\varepsilon := r/|z_3|$ is small, the Mogi model corresponds to the assumption that the magma chamber is well-approximated by a single point producing a uniform pressure in the radial direction; as such, it is sometimes referred to as a *point source model*. However, even if the source




FIGURE 2.1. Normalized Mogi displacement profiles given in (2.1): horizontal components u_α , $\alpha = 1, 2$, dashed line; vertical component u_3 , continuous line.

is reduced to a single point, the model still records the spherical form of the cavity. Different geometrical form may lead to different deformation effects.

The Mogi model has been widely applied to real data of different volcanoes to infer approximate location and strength of the magma chamber. The main benefit of such strategy lies in the fact that it provides a simple formula with an explicit appearance of the basic physical parameters depth and total work (combining pressure and volume) and, thus, that it can be readily compared with real deformation data to provide explicit forecasts.

The simplicity of formulas (2.1) makes the application model viable, but it compensates only partially the intrinsic reductions of the approach. As a consequence, variations on the geometry of the magma chamber have been proposed to provide more realistic frameworks but with the target to furnish explicit formulas for ground deformations; in this way synthetic data can be compared with real data via appropriate inversion algorithms. Such attempt has focused on oblate and prolate ellipsoids [16, 40], rectangular dislocations [35] and horizontal penny-shaped cracks [20]. For completeness, we mention also the attempts to study the heterogeneity of the crust, see [29] and references therein, the case of a non-flat crust surface [39, 14] and the combination of elastic properties with gravitational effects and time-dependent processes modeling of the crust, see [9, 13].

In all cases, refined descriptions have the inherent drawback of requiring a detailed knowledge of the crust elastic properties. In absence of reliable complete data and measurements, the risk of introducing an additional degree of freedom in the parameter choice is substantial. This observation partly supports the approach of the Mogi model which consists in keeping as far as possible the parameters choice limited and, consequently, the model simple.

Now, let us introduce in details the boundary value problem which emerges from the previous assumptions on the geometry of the model, geophysics of the crust and

crust-chamber interaction. Denoting by \mathbb{R}_-^3 the (open) half-space described by the condition $x_3 < 0$, the domain D occupied by the Earth’s crust is $D := \mathbb{R}_-^3 \setminus \overline{C}$, where $C \subset \mathbb{R}_-^3$, describing the magma chamber, is assumed to be an open set with bounded boundary ∂C . Hence, the boundary of D is composed by two disconnected components: the two-dimensional plane $\mathbb{R}^2 := \{\mathbf{x} = (x_1, x_2, x_3) \in \mathbb{R}^3 : x_3 = 0\}$, which constitutes the free air/crust border, and the set ∂C , corresponding to the crust/chamber edge. In this setting, we end up with the following boundary value problem

$$\operatorname{div}(\mathbb{C}\widehat{\nabla}\mathbf{u}) = \mathbf{0} \quad \text{in } D, \quad (\mathbb{C}\widehat{\nabla}\mathbf{u})\mathbf{e}_3 = \mathbf{0} \quad \text{in } \mathbb{R}^2, \quad (\mathbb{C}\widehat{\nabla}\mathbf{u})\mathbf{n} = p\mathbf{n} \quad \text{on } \partial C,$$

which we consider with the asymptotic conditions at infinity

$$\lim_{|\mathbf{x}| \rightarrow \infty} \mathbf{u}(\mathbf{x}) = \lim_{|\mathbf{x}| \rightarrow \infty} |\mathbf{x}| \nabla \mathbf{u}(\mathbf{x}) = \mathbf{0}.$$

The conjunction of conditions on ∂D and at infinity determines a displacement-traction problem.

At this point, the model provides the displacement \mathbf{u} of a generic *finite* cavity C . The next step is to deduce a corresponding point source model, in the spirit of the Mogi spherical one. To this aim, we assume the cavity C of the form

$$C = d\mathbf{z} + r\Omega,$$

where $d, r > 0$ are characteristic length-scales for depth and diameter of the cavity, its center $d\mathbf{z}$ belongs to \mathbb{R}_-^3 and its shape Ω is a bounded domain containing the origin. The Mogi model corresponds to Ω given by a sphere of radius 1.

Introducing the rescaling $(\mathbf{x}, \mathbf{u}) \mapsto (\mathbf{x}/d, \mathbf{u}/r)$ and denoting the new variables again by \mathbf{x} and \mathbf{u} , the above problem takes the form (with unchanged far-field conditions)

$$\operatorname{div}(\mathbb{C}\widehat{\nabla}\mathbf{u}) = \mathbf{0} \quad \text{in } D_\varepsilon, \quad (\mathbb{C}\widehat{\nabla}\mathbf{u})\mathbf{e}_3 = \mathbf{0} \quad \text{in } \mathbb{R}^2, \quad (\mathbb{C}\widehat{\nabla}\mathbf{u})\mathbf{n} = p\mathbf{n} \quad \text{on } \partial C_\varepsilon,$$

where $\varepsilon = r/d$, $C_\varepsilon := \mathbf{z} + \varepsilon\Omega$, $D_\varepsilon := \mathbb{R}_-^3 \setminus \overline{C}_\varepsilon$ and p is a “rescaled” pressure, ratio of the original pressure p and ε .

In the next section we study the well-posedness of the linear elastic model in the case of a finite pressurized cavity C and then we derive the asymptotic expansion of the solution when $C := C_\varepsilon := \mathbf{z} + \varepsilon\Omega$.

3. INTEGRAL REPRESENTATION AND WELL-POSEDNESS OF THE DIRECT PROBLEM

Since $\mathbb{C} := \lambda\mathbf{I} \otimes \mathbf{I} + 2\mu\mathbb{I}$, the elastostatic Lamé operator \mathcal{L} for a homogeneous and isotropic elastic medium is given by

$$\mathcal{L}\mathbf{u} := \operatorname{div}(\mathbb{C}\widehat{\nabla}\mathbf{u}) = \mu\Delta\mathbf{u} + (\lambda + \mu)\nabla \operatorname{div} \mathbf{u},$$

where \mathbf{u} represents the vector of the displacements and $\widehat{\nabla}\mathbf{u} = \frac{1}{2}(\nabla\mathbf{u} + \nabla\mathbf{u}^T)$ the strain tensor. With $\partial\mathbf{u}/\partial\nu$ we depict the conormal derivative on the boundary of a domain, that is, the traction vector, which has the expression

$$\frac{\partial\mathbf{u}}{\partial\nu} := (\mathbb{C}\widehat{\nabla}\mathbf{u})\mathbf{n} = \lambda(\operatorname{div} \mathbf{u})\mathbf{n} + 2\mu(\widehat{\nabla}\mathbf{u})\mathbf{n}$$

or, equivalently,

$$\frac{\partial \mathbf{u}}{\partial \boldsymbol{\nu}} = 2\mu \frac{\partial \mathbf{u}}{\partial \mathbf{n}} + \lambda(\operatorname{div} \mathbf{u})\mathbf{n} + \mu(\mathbf{n} \times \operatorname{rot} \mathbf{u}).$$

Here, we analyze the linear elastostatic boundary value problem

$$(3.1) \quad \begin{cases} \operatorname{div}(\mathbb{C}\widehat{\nabla} \mathbf{u}) = \mathbf{0} & \text{in } \mathbb{R}^3 \setminus C \\ \frac{\partial \mathbf{u}}{\partial \boldsymbol{\nu}} = p\mathbf{n} & \text{on } \partial C \\ \frac{\partial \mathbf{u}}{\partial \boldsymbol{\nu}} = \mathbf{0} & \text{on } \mathbb{R}^2 \\ \mathbf{u} = o(1), \quad \nabla \mathbf{u} = o(|\mathbf{x}|^{-1}) & |\mathbf{x}| \rightarrow \infty, \end{cases}$$

where C is the cavity and p is a constant representing the pressure. For the Lamé parameters, we consider the physical range $3\lambda + 2\mu > 0$ and $\mu > 0$ which ensures positive definiteness of \mathbb{C} . For the sequel, we recall that the positive definiteness of the tensor \mathbb{C} implies the strong ellipticity which corresponds to the request $\mu > 0$ and $\lambda + 2\mu > 0$, see [23].

The aim of this section is to provide an integral representation formula and to establish the well-posedness of the problem. To do that, we consider three steps:

- firstly, we recall Betti's formulas, definition and some properties of single and double layer potentials of linear elasticity;
- then, we give the expression of the fundamental solution \mathbf{N} of the half-space with null traction on the boundary, found by Mindlin in [31, 32];
- finally, we represent the solution to (3.1) by an integral formula through the fundamental solution of the half-space.

All these objects will be used to prove the well-posedness of the problem (3.1).

3.1. PRELIMINARIES. — We recall Betti's formulas for the Lamé system which can be obtained by integration by parts, see for example [4, 27]. Given a bounded Lipschitz domain $C \subset \mathbb{R}^3$ and two vectors $\mathbf{u}, \mathbf{v} \in \mathbb{R}^3$, the *first Betti formula* is

$$(3.2) \quad \int_{\partial C} \mathbf{u} \cdot \frac{\partial \mathbf{v}}{\partial \boldsymbol{\nu}} d\sigma(\mathbf{x}) = \int_C \mathbf{u} \cdot \mathcal{L}\mathbf{v} d\mathbf{x} + \int_C Q(\mathbf{u}, \mathbf{v}) d\mathbf{x},$$

where the quadratic form Q associated to the Lamé system is

$$Q(\mathbf{u}, \mathbf{v}) := \lambda(\operatorname{div} \mathbf{u})(\operatorname{div} \mathbf{v}) + 2\mu\widehat{\nabla} \mathbf{u} : \widehat{\nabla} \mathbf{v}.$$

From (3.2) it is straightforward to find the *second Betti formula*

$$(3.3) \quad \int_C (\mathbf{u} \cdot \mathcal{L}\mathbf{v} - \mathbf{v} \cdot \mathcal{L}\mathbf{u}) d\mathbf{x} = \int_{\partial C} \left(\mathbf{u} \cdot \frac{\partial \mathbf{v}}{\partial \boldsymbol{\nu}} - \mathbf{v} \cdot \frac{\partial \mathbf{u}}{\partial \boldsymbol{\nu}} \right) d\sigma(\mathbf{x}).$$

Formula (3.2) will be used to prove that the solution of (3.1) is unique, and the equality (3.3) to get an integral representation formula for it. To accomplish this second goal, a leading role is played by the fundamental solution of the Lamé system: the *Kelvin matrix* $\boldsymbol{\Gamma}$ (or *Kelvin-Somigliana matrix*) solution to the equation

$$\operatorname{div}(\mathbb{C}\widehat{\nabla} \boldsymbol{\Gamma}) = \delta_{\mathbf{0}} \mathbf{I}, \quad \mathbf{x} \in \mathbb{R}^3 \setminus \{\mathbf{0}\},$$

where $\delta_{\mathbf{0}}$ is the Dirac function centred at $\mathbf{0}$. Setting $C_{\mu,\nu} := 1/\{16\pi\mu(1-\nu)\}$, where ν is the *Poisson ratio* which is related to λ and μ by the identity

$$\nu = \frac{\lambda}{2(\lambda + \mu)},$$

the explicit expression of $\mathbf{\Gamma} = (\Gamma_{ij})$ is

$$(3.4) \quad \Gamma_{ij}(\mathbf{x}) = -C_{\mu,\nu} \left\{ \frac{(3-4\nu)\delta_{ij}}{|\mathbf{x}|} + \frac{x_i x_j}{|\mathbf{x}|^3} \right\}, \quad i, j = 1, 2, 3,$$

where δ_{ij} is the Kronecker symbol and Γ_{ij} stands for the i -th component of the displacement when a force is applied in the j -th direction at the point $\mathbf{0}$. For the reader's convenience, we write also the gradient of $\mathbf{\Gamma}$ to highlight its behaviour at infinity

$$(3.5) \quad \frac{\partial \Gamma_{ij}}{\partial x_k}(\mathbf{x}) = C_{\mu,\nu} \left\{ \frac{(3-4\nu)\delta_{ij}x_k - \delta_{ik}x_j - \delta_{jk}x_i}{|\mathbf{x}|^3} + \frac{3x_i x_j x_k}{|\mathbf{x}|^5} \right\}, \quad i, j, k = 1, 2, 3.$$

Therefore, from (3.4) and (3.5) it is straightforward to see that

$$(3.6) \quad |\mathbf{\Gamma}(\mathbf{x})| = O(1/|\mathbf{x}|) \quad \text{and} \quad |\nabla \mathbf{\Gamma}(\mathbf{x})| = O(1/|\mathbf{x}|^2) \quad \text{as} \quad |\mathbf{x}| \rightarrow \infty.$$

With the Kelvin matrix $\mathbf{\Gamma}$ at hand, we recall the definition of the single and double layer potentials corresponding to the operator \mathcal{L} . Given $\varphi \in L^2(\partial C)$, we set (see [4, 5, 27])

$$(3.7) \quad \begin{aligned} \mathbf{S}^\Gamma \varphi(\mathbf{x}) &:= \int_{\partial C} \mathbf{\Gamma}(\mathbf{x} - \mathbf{y}) \varphi(\mathbf{y}) \, d\sigma(\mathbf{y}), & \mathbf{x} \in \mathbb{R}^3, \\ \mathbf{D}^\Gamma \varphi(\mathbf{x}) &:= \int_{\partial C} \frac{\partial \mathbf{\Gamma}}{\partial \nu(\mathbf{y})}(\mathbf{x} - \mathbf{y}) \varphi(\mathbf{y}) \, d\sigma(\mathbf{y}), & \mathbf{x} \in \mathbb{R}^3 \setminus \partial C, \end{aligned}$$

where $\partial \mathbf{\Gamma} / \partial \nu$ denotes the conormal derivative applied to each column of the matrix $\mathbf{\Gamma}$.

In the sequel the subscripts $+$ and $-$ indicate the limits from outside and inside C , respectively. The double layer potential and the conormal derivative of the single layer potential satisfy the jump relations

$$(3.8) \quad \begin{aligned} \mathbf{D}^\Gamma \varphi \Big|_{\pm}(\mathbf{x}) &= (\mp \frac{1}{2} \mathbf{I} + \mathbf{K}) \varphi(\mathbf{x}) & \text{for almost any } \mathbf{x} \in \partial C, \\ \frac{\partial \mathbf{S}^\Gamma \varphi}{\partial \nu} \Big|_{\pm}(\mathbf{x}) &= (\pm \frac{1}{2} \mathbf{I} + \mathbf{K}^*) \varphi(\mathbf{x}) & \text{for almost any } \mathbf{x} \in \partial C, \end{aligned}$$

where \mathbf{K} and \mathbf{K}^* are the L^2 -adjoint Neumann-Poincaré boundary integral operators defined, in the sense of Cauchy principal value, by

$$\begin{aligned} \mathbf{K} \varphi(\mathbf{x}) &:= \text{p.v.} \int_{\partial C} \frac{\partial \mathbf{\Gamma}}{\partial \nu(\mathbf{y})}(\mathbf{x} - \mathbf{y}) \varphi(\mathbf{y}) \, d\sigma(\mathbf{y}), \\ \mathbf{K}^* \varphi(\mathbf{x}) &:= \text{p.v.} \int_{\partial C} \frac{\partial \mathbf{\Gamma}}{\partial \nu(\mathbf{x})}(\mathbf{x} - \mathbf{y}) \varphi(\mathbf{y}) \, d\sigma(\mathbf{y}). \end{aligned}$$

It is worth noticing that these two operators are not compact even on smooth domains, in contrast with the analogous operators for the Laplace equation (see [5]), due to the presence in their kernels of the terms

$$\frac{n_i(x_j - y_j)}{|\mathbf{x} - \mathbf{y}|^3} - \frac{n_j(x_i - y_i)}{|\mathbf{x} - \mathbf{y}|^3}, \quad i \neq j,$$

which make the kernel not integrable. Indeed, even in the case of smooth domains, we cannot approximate locally the terms $\mathbf{n} \times (\mathbf{x} - \mathbf{y})$ with a smooth function, that is, by power of $|\mathbf{x} - \mathbf{y}|$ via Taylor expansion, in order to obtain an integrable kernel on ∂C . Therefore, the analysis to prove the invertibility of the operators in (3.8) is intricate and usually based on regularizing operators (see [27]) in the case of smooth domains. For the Lipschitz domains the analysis is much more involved and based on Rellich formulas (see [15] and its companion article [19]).

3.2. FUNDAMENTAL SOLUTION OF THE HALF-SPACE. — In this subsection we show the explicit expression of the Neumann function of the half-space presented for the first time in [31] by means of Galerkin vector and nuclei of strain of the theory of linear elasticity, and secondly in [32] using the Papkovitch-Neuber representation of the displacement vector and the potential theory.

We consider the boundary value problem

$$(3.9) \quad \begin{cases} \operatorname{div}(\mathbb{C}\widehat{\nabla}\mathbf{v}) = \mathbf{b} & \text{in } \mathbb{R}_-^3 \\ \frac{\partial \mathbf{v}}{\partial \boldsymbol{\nu}} = \mathbf{0} & \text{on } \mathbb{R}^2 \\ \mathbf{v} = o(\mathbf{1}), \quad \nabla \mathbf{v} = o(|\mathbf{x}|^{-1}) & \text{as } |\mathbf{x}| \rightarrow +\infty. \end{cases}$$

The Neumann function of (3.9) is the kernel \mathbf{N} of the integral operator

$$(3.10) \quad \mathbf{v}(\mathbf{x}) = \int_{\mathbb{R}_-^3} \mathbf{N}(\mathbf{x}, \mathbf{y}) \mathbf{b}(\mathbf{y}) \, d\mathbf{y},$$

giving the solution to the problem.

Given $\mathbf{y} = (y_1, y_2, y_3)$, we set $\tilde{\mathbf{y}} := (y_1, y_2, -y_3)$.

THEOREM 3.1. — *The Neumann function \mathbf{N} of problem (3.9) can be decomposed as*

$$\mathbf{N}(\mathbf{x}, \mathbf{y}) = \boldsymbol{\Gamma}(\mathbf{x} - \mathbf{y}) + \mathbf{R}^1(\mathbf{x} - \tilde{\mathbf{y}}) + y_3 \mathbf{R}^2(\mathbf{x} - \tilde{\mathbf{y}}) + y_3^2 \mathbf{R}^3(\mathbf{x} - \tilde{\mathbf{y}}),$$

where $\boldsymbol{\Gamma}$ is the Kelvin matrix, see (3.4), and \mathbf{R}^k , $k = 1, 2, 3$, have components R_{ij}^k given by

$$\begin{aligned} R_{ij}^1(\boldsymbol{\eta}) &:= C_{\mu,\nu} \{ -(\tilde{f} + c_\nu \tilde{g}) \delta_{ij} - (3 - 4\nu) \eta_i \eta_j \tilde{f}^3 \\ &\quad + c_\nu [\delta_{i3} \eta_j - \delta_{j3} (1 - \delta_{i3}) \eta_i] \tilde{f} \tilde{g} + c_\nu (1 - \delta_{i3}) (1 - \delta_{j3}) \eta_i \eta_j \tilde{f} \tilde{g}^2 \} \\ R_{ij}^2(\boldsymbol{\eta}) &:= 2C_{\mu,\nu} \{ (3 - 4\nu) [\delta_{i3} (1 - \delta_{j3}) \eta_j + \delta_{j3} (1 - \delta_{i3}) \eta_i] \tilde{f}^3 - (1 - 2\delta_{3j}) \delta_{ij} \eta_3 \tilde{f}^3 \\ &\quad + 3(1 - 2\delta_{3j}) \eta_i \eta_j \eta_3 \tilde{f}^5 \} \\ R_{ij}^3(\boldsymbol{\eta}) &:= 2C_{\mu,\nu} (1 - 2\delta_{j3}) \{ \delta_{ij} \tilde{f}^3 - 3\eta_i \eta_j \tilde{f}^5 \}. \end{aligned}$$

for $i, j = 1, 2, 3$, where $c_\nu := 4(1 - \nu)(1 - 2\nu)$ and

$$\tilde{f}(\boldsymbol{\eta}) := \frac{1}{|\boldsymbol{\eta}|}, \quad \tilde{g}(\boldsymbol{\eta}) := \frac{1}{|\boldsymbol{\eta}| - \eta_3}.$$

For the proof of Theorem 3.1 see the appendix. Uniqueness of the solution to (3.9) is similar to the one for problem (3.1) which we present in the following section.

The matrix \mathbf{R} , defined by

$$(3.11) \quad \mathbf{R}(\boldsymbol{\eta}, y_3) := \mathbf{R}^1(\boldsymbol{\eta}) + y_3 \mathbf{R}^2(\boldsymbol{\eta}) + y_3^2 \mathbf{R}^3(\boldsymbol{\eta}),$$

gives the regular part of the Neumann function since the singular point $\boldsymbol{\eta} = \mathbf{0}$ corresponds to $\mathbf{y} = (y_1, y_2, -y_3)$ with $y_3 < 0$, which belongs to \mathbb{R}_+^3 .

To convert the problem (3.1) into an integral form, bounds on the decay at infinity of the Neumann function and its derivative at infinity are needed.

PROPOSITION 3.2. — *For any $M_x, M_y > 0$, there exists $C > 0$ such that*

$$(3.12) \quad |\mathbf{N}(\mathbf{x}, \mathbf{y})| \leq C |\mathbf{x}|^{-1} \quad \text{and} \quad |\nabla \mathbf{N}(\mathbf{x}, \mathbf{y})| \leq C |\mathbf{x}|^{-2}$$

for any $\mathbf{x}, \mathbf{y} \in \mathbb{R}_-^3$ with $|\mathbf{x}| \geq M_x$ and $|\mathbf{y}| \leq M_y$.

The proof of this proposition is a consequence of the properties on homogeneous functions and their derivatives, noticing that \mathbf{R}^k are homogeneous of degree $-k$, for $k = 1, 2, 3$.

3.3. REPRESENTATION FORMULA. — Next, we derive an integral representation formula for \mathbf{u} solution to problem (3.1). For, we make use of single and double layer potentials defined in (3.7) and integral contributions relative to the regular part \mathbf{R} of the Neumann function \mathbf{N} , defined by

$$(3.13) \quad \begin{aligned} \mathbf{S}^R \boldsymbol{\varphi}(\mathbf{x}) &:= \int_{\partial C} (\mathbf{R}(\mathbf{x}, \mathbf{y}))^T \boldsymbol{\varphi}(\mathbf{y}) \, d\sigma(\mathbf{y}), & \mathbf{x} \in \mathbb{R}_-^3, \\ \mathbf{D}^R \boldsymbol{\varphi}(\mathbf{x}) &:= \int_{\partial C} \left(\frac{\partial \mathbf{R}}{\partial \nu(\mathbf{y})}(\mathbf{x}, \mathbf{y}) \right)^T \boldsymbol{\varphi}(\mathbf{y}) \, d\sigma(\mathbf{y}), & \mathbf{x} \in \mathbb{R}_-^3, \end{aligned}$$

where $\boldsymbol{\varphi} \in L^2(\partial C)$.

THEOREM 3.3. — *The solution \mathbf{u} to (3.1) is such that*

$$(3.14) \quad \mathbf{u} = p \mathbf{S}^\Gamma \mathbf{n} - \mathbf{D}^\Gamma \mathbf{f} + p \mathbf{S}^R \mathbf{n} - \mathbf{D}^R \mathbf{f} \quad \text{in } \mathbb{R}_-^3 \setminus \overline{C},$$

where $\mathbf{S}^\Gamma, \mathbf{D}^\Gamma$ are defined in (3.7), $\mathbf{S}^R, \mathbf{D}^R$ in (3.13), $p\mathbf{n}$ is the boundary condition in (3.1) and \mathbf{f} is the trace of \mathbf{u} on ∂C .

Before proving this theorem, we observe that if a solution to (3.1) exists, then \mathbf{f} solves the integral equation

$$(3.15) \quad \left(\frac{1}{2} \mathbf{I} + \mathbf{K} + \mathbf{D}^R \right) \mathbf{f} = p (\mathbf{S}^\Gamma \mathbf{n} + \mathbf{S}^R \mathbf{n}) \quad \text{on } \partial C,$$

obtained by the application of the trace properties of the double layer potential (3.8) in formula (3.14).

Proof of Theorem 3.3. — Given $r, \varepsilon > 0$ such that $C \subset B_r(\mathbf{0})$ and $B_\varepsilon(\mathbf{y}) \subset \mathbb{R}_-^3 \setminus \overline{C}$, let

$$\Omega_{r,\varepsilon} = (\mathbb{R}_-^3 \cap B_r(\mathbf{0})) \setminus (C \cup B_\varepsilon(\mathbf{y}))$$

with r sufficiently large such that to contain the cavity C ; additionally, we define $\partial B_r^h(\mathbf{0})$ as the intersection of the hemisphere with the boundary of the half-space, and with $\partial B_r^b(\mathbf{0})$ the spherical cap (see Figure 3.1). Now, we apply Betti's formula (3.3)




FIGURE 3.1. Domain $\Omega_{r,\varepsilon}$.

to \mathbf{u} and the k -th column vector of \mathbf{N} , indicated by $\mathbf{N}^{(k)}$, for $k = 1, 2, 3$, in $\Omega_{r,\varepsilon}$, hence

$$\begin{aligned} 0 &= \int_{\Omega_{r,\varepsilon}} \left[\mathbf{u}(\mathbf{x}) \cdot \mathcal{L}\mathbf{N}^{(k)}(\mathbf{x}, \mathbf{y}) - \mathbf{N}^{(k)}(\mathbf{x}, \mathbf{y}) \cdot \mathcal{L}\mathbf{u}(\mathbf{x}) \right] d\mathbf{x} \\ &= \int_{\partial B_r^b(\mathbf{0})} \left[\frac{\partial \mathbf{N}^{(k)}}{\partial \boldsymbol{\nu}_\mathbf{x}}(\mathbf{x}, \mathbf{y}) \cdot \mathbf{u}(\mathbf{x}) - \mathbf{N}^{(k)}(\mathbf{x}, \mathbf{y}) \cdot \frac{\partial \mathbf{u}}{\partial \boldsymbol{\nu}_\mathbf{x}}(\mathbf{x}) \right] d\sigma(\mathbf{x}) \\ &\quad - \int_{\partial B_\varepsilon(\mathbf{y})} \left[\frac{\partial \mathbf{N}^{(k)}}{\partial \boldsymbol{\nu}_\mathbf{x}}(\mathbf{x}, \mathbf{y}) \cdot \mathbf{u}(\mathbf{x}) - \mathbf{N}^{(k)}(\mathbf{x}, \mathbf{y}) \cdot \frac{\partial \mathbf{u}}{\partial \boldsymbol{\nu}_\mathbf{x}}(\mathbf{x}) \right] d\sigma(\mathbf{x}) \\ &\quad - \int_{\partial C} \left[\frac{\partial \mathbf{N}^{(k)}}{\partial \boldsymbol{\nu}_\mathbf{x}}(\mathbf{x}, \mathbf{y}) \cdot \mathbf{u}(\mathbf{x}) - \mathbf{N}^{(k)}(\mathbf{x}, \mathbf{y}) \cdot \frac{\partial \mathbf{u}}{\partial \boldsymbol{\nu}_\mathbf{x}}(\mathbf{x}) \right] d\sigma(\mathbf{x}) \\ &:= I_1 + I_2 + I_3, \end{aligned}$$

since, from (3.1) and the boundary condition in (3.9),

$$\int_{\partial B_r^b(\mathbf{0})} \left[\frac{\partial \mathbf{N}^{(k)}}{\partial \boldsymbol{\nu}_\mathbf{x}}(\mathbf{x}, \mathbf{y}) \cdot \mathbf{u}(\mathbf{x}) - \mathbf{N}^{(k)}(\mathbf{x}, \mathbf{y}) \cdot \frac{\partial \mathbf{u}}{\partial \boldsymbol{\nu}_\mathbf{x}}(\mathbf{x}) \right] d\sigma(\mathbf{x}) = 0.$$

We show that the term I_1 goes to zero by using the behaviour at infinity of \mathbf{u} given in (3.1) and of the Neumann function given in (3.12). Indeed, we have

$$\left| \int_{\partial B_r^b(\mathbf{0})} \frac{\partial \mathbf{N}^{(k)}}{\partial \boldsymbol{\nu}_x}(\mathbf{x}, \mathbf{y}) \cdot \mathbf{u}(\mathbf{x}) \, d\sigma(\mathbf{x}) \right| \leq \int_{\partial B_r^b(\mathbf{0})} |\mathbf{u}| \left| \frac{\partial \mathbf{N}^{(k)}}{\partial \boldsymbol{\nu}_x} \right| d\sigma(\mathbf{x}) \leq \frac{C}{r^2} \int_{\partial B_r^b(\mathbf{0})} |\mathbf{u}(\mathbf{x})| d\sigma(\mathbf{x}).$$

This last integral can be estimated by means of the spherical coordinates

$$x_1 = r \sin \varphi \cos \theta, \quad x_2 = r \sin \varphi \sin \theta, \quad x_3 = r \cos \varphi,$$

where $\varphi \in [\pi/2, \pi]$, since $B_r^b(\mathbf{0})$ is a hemisphere in \mathbb{R}^3 , and $\theta \in [0, 2\pi]$, indeed

$$\begin{aligned} \frac{C}{r^2} \int_{\partial B_r^b(\mathbf{0})} |\mathbf{u}| \, d\sigma(\mathbf{x}) &= C \int_{\pi/2}^{\pi} \int_0^{2\pi} |\mathbf{u}(r, \theta, \varphi)| \sin \varphi \, d\theta \, d\varphi \\ &\leq C \sup_{\substack{\theta \in [0, 2\pi] \\ \varphi \in [\pi/2, \pi]}} |\mathbf{u}(r, \theta, \varphi)| \longrightarrow 0, \end{aligned}$$

as $r \rightarrow +\infty$, since $\mathbf{u} = o(\mathbf{1})$. Similarly

$$\left| \int_{\partial B_r^b(\mathbf{0})} \mathbf{N}^{(k)}(\mathbf{x}, \mathbf{y}) \cdot \frac{\partial \mathbf{u}}{\partial \boldsymbol{\nu}_x}(\mathbf{x}) \, d\sigma(\mathbf{x}) \right| \leq \int_{\partial B_r^b(\mathbf{0})} |\mathbf{N}^{(k)}| \left| \frac{\partial \mathbf{u}}{\partial \boldsymbol{\nu}_x} \right| d\sigma(\mathbf{x}) \leq \frac{C}{r} \int_{\partial B_r^b(\mathbf{0})} \left| \frac{\partial \mathbf{u}}{\partial \boldsymbol{\nu}_x} \right| d\sigma(\mathbf{x}).$$

Again, passing through spherical coordinates, we get

$$(3.16) \quad \frac{C}{r} \int_{\partial B_r^b(\mathbf{0})} \left| \frac{\partial \mathbf{u}}{\partial \boldsymbol{\nu}_x} \right| d\sigma(\mathbf{x}) \leq C \sup_{\substack{\theta \in [0, 2\pi] \\ \varphi \in [\pi/2, \pi]}} r \left| \frac{\partial \mathbf{u}}{\partial \boldsymbol{\nu}}(r, \theta, \varphi) \right| \longrightarrow 0,$$

as $r \rightarrow +\infty$, since $|\nabla \mathbf{u}| = o(r^{-1})$.

The integral I_2 gives the value of the function \mathbf{u} in \mathbf{y} as ε goes to zero. Indeed, we have

$$\begin{aligned} \left| \int_{\partial B_\varepsilon(\mathbf{y})} \mathbf{N}^{(k)}(\mathbf{x}, \mathbf{y}) \cdot \frac{\partial \mathbf{u}}{\partial \boldsymbol{\nu}_x}(\mathbf{x}) \, d\sigma(\mathbf{x}) \right| &\leq \int_{\partial B_\varepsilon(\mathbf{y})} |\mathbf{N}^{(k)}| \left| \frac{\partial \mathbf{u}}{\partial \boldsymbol{\nu}_x} \right| d\sigma(\mathbf{x}) \\ &\leq \sup_{\mathbf{x} \in \partial B_\varepsilon(\mathbf{y})} \left| \frac{\partial \mathbf{u}}{\partial \boldsymbol{\nu}_x} \right| \int_{\partial B_\varepsilon(\mathbf{y})} \left[|\boldsymbol{\Gamma}^{(k)}| + |\mathbf{R}^{(k)}| \right] d\sigma(\mathbf{x}) = O(\varepsilon), \end{aligned}$$

since the second integral has a continuous kernel. On the other hand

$$\begin{aligned} & - \int_{\partial B_\varepsilon(\mathbf{y})} \frac{\partial \mathbf{N}^{(k)}}{\partial \boldsymbol{\nu}_x}(\mathbf{x}, \mathbf{y}) \cdot \mathbf{u}(\mathbf{x}) \, d\sigma(\mathbf{x}) \\ &= -\mathbf{u}(\mathbf{y}) \cdot \int_{\partial B_\varepsilon(\mathbf{y})} \frac{\partial \mathbf{N}^{(k)}}{\partial \boldsymbol{\nu}_x}(\mathbf{x}, \mathbf{y}) \, d\sigma(\mathbf{x}) + \int_{\partial B_\varepsilon(\mathbf{y})} [\mathbf{u}(\mathbf{y}) - \mathbf{u}(\mathbf{x})] \cdot \frac{\partial \mathbf{N}^{(k)}}{\partial \boldsymbol{\nu}_x}(\mathbf{x}, \mathbf{y}) \, d\sigma(\mathbf{x}). \end{aligned}$$

The latter integral tends to zero when ε goes to zero because

$$\left| \int_{\partial B_\varepsilon(\mathbf{y})} [\mathbf{u}(\mathbf{y}) - \mathbf{u}(\mathbf{x})] \cdot \frac{\partial \mathbf{N}^{(k)}}{\partial \boldsymbol{\nu}_x}(\mathbf{x}, \mathbf{y}) d\sigma(\mathbf{x}) \right| \leq \sup_{\mathbf{x} \in \partial B_\varepsilon(\mathbf{y})} |\mathbf{u}(\mathbf{y}) - \mathbf{u}(\mathbf{x})| \int_{\partial B_\varepsilon(\mathbf{y})} \left| \frac{\partial \mathbf{N}^{(k)}}{\partial \boldsymbol{\nu}_x} \right| d\sigma(\mathbf{x})$$

and this last integral is bounded when ε goes to zero. Let finally observe that

$$\begin{aligned} & -\mathbf{u}(\mathbf{y}) \cdot \int_{\partial B_\varepsilon(\mathbf{y})} \frac{\partial \mathbf{N}^{(k)}}{\partial \boldsymbol{\nu}_x}(\mathbf{x}, \mathbf{y}) d\sigma(\mathbf{x}) \\ (3.17) \quad & = -\mathbf{u}(\mathbf{y}) \cdot \int_{\partial B_\varepsilon(\mathbf{y})} \frac{\partial(\boldsymbol{\Gamma}^{(k)} + \mathbf{R}^{(k)})}{\partial \boldsymbol{\nu}_x}(\mathbf{x}, \mathbf{y}) d\sigma(\mathbf{x}) \\ & = -\mathbf{u}(\mathbf{y}) \cdot \int_{\partial B_\varepsilon(\mathbf{y})} \frac{\partial \boldsymbol{\Gamma}^{(k)}}{\partial \boldsymbol{\nu}_x}(\mathbf{x} - \mathbf{y}) d\sigma(\mathbf{x}) - \mathbf{u}(\mathbf{y}) \cdot \int_{\partial B_\varepsilon(\mathbf{y})} \frac{\partial \mathbf{R}^{(k)}}{\partial \boldsymbol{\nu}_x}(\mathbf{x}, \mathbf{y}) d\sigma(\mathbf{x}), \end{aligned}$$

where the latter integral tends to zero as $\varepsilon \rightarrow 0$, since $\mathbf{R}^{(k)}$ represents the regular part of the Neumann function. To deal with the first integral, we preliminarily observe that direct differentiation gives

$$\begin{aligned} (3.18) \quad \left(\frac{\partial \boldsymbol{\Gamma}^{(k)}}{\partial \boldsymbol{\nu}_x} \right)_h(\mathbf{x} - \mathbf{y}) & = -c'_\nu \left\{ n_k(\mathbf{x}) \frac{\partial}{\partial x_h} \frac{1}{|\mathbf{x} - \mathbf{y}|} - n_h(\mathbf{x}) \frac{\partial}{\partial x_k} \frac{1}{|\mathbf{x} - \mathbf{y}|} \right. \\ & \quad \left. + \left[\delta_{hk} + \frac{3}{(1-2\nu)} \frac{\partial |\mathbf{x} - \mathbf{y}|}{\partial x_k} \frac{\partial |\mathbf{x} - \mathbf{y}|}{\partial x_h} \right] \frac{\partial}{\partial n(\mathbf{x})} \frac{1}{|\mathbf{x} - \mathbf{y}|} \right\}, \end{aligned}$$

where $c'_\nu := (1-2\nu)/(8\pi(1-\nu))$. We substitute this expression into the integral (3.17) and we take into account that

$$n_h(\mathbf{x}) = \frac{x_h - y_h}{|\mathbf{x} - \mathbf{y}|}, \quad \frac{\partial}{\partial x_k} \frac{1}{|\mathbf{x} - \mathbf{y}|} = -\frac{x_k - y_k}{|\mathbf{x} - \mathbf{y}|^3},$$

hence

$$\int_{\partial B_\varepsilon(\mathbf{y})} n_h(\mathbf{x}) \frac{\partial}{\partial x_k} \frac{1}{|\mathbf{x} - \mathbf{y}|} d\sigma(\mathbf{x}) = - \int_{\partial B_\varepsilon(\mathbf{y})} \frac{(x_h - y_h)(x_k - y_k)}{|\mathbf{x} - \mathbf{y}|^4} d\sigma(\mathbf{x}).$$

To compute this last integral we use spherical coordinates, that is,

$$x_1 - y_1 = \varepsilon \sin \varphi \cos \theta, \quad x_2 - y_2 = \varepsilon \sin \varphi \sin \theta, \quad x_3 - y_3 = \varepsilon \cos \varphi,$$

where $\varphi \in [0, \pi]$ and $\theta \in [0, 2\pi)$. From a simple calculation it follows

$$(3.19) \quad - \int_{\partial B_\varepsilon(\mathbf{y})} \frac{(x_h - y_h)(x_k - y_k)}{|\mathbf{x} - \mathbf{y}|^4} d\sigma(\mathbf{x}) = \begin{cases} 0 & \text{if } h \neq k \\ -\frac{4}{3}\pi & \text{if } h = k. \end{cases}$$

Therefore, from (3.18) and (3.19), we have

$$(3.20) \quad \int_{\partial B_\varepsilon(\mathbf{y})} \left(n_k(\mathbf{x}) \frac{\partial}{\partial x_h} \frac{1}{|\mathbf{x} - \mathbf{y}|} - n_h(\mathbf{x}) \frac{\partial}{\partial x_k} \frac{1}{|\mathbf{x} - \mathbf{y}|} \right) d\sigma(\mathbf{x}) = 0,$$

for any h and k . Hence, (3.17) becomes

$$\begin{aligned}
 -\mathbf{u}(\mathbf{y}) \cdot \int_{\partial B_\varepsilon(\mathbf{y})} \frac{\partial \mathbf{N}^{(k)}}{\partial \boldsymbol{\nu}_{\mathbf{x}}}(\mathbf{x} - \mathbf{y}) d\sigma(\mathbf{x}) &= c'_\nu \sum_{h=1}^3 u_h(\mathbf{y}) \cdot \\
 &\cdot \int_{\partial B_\varepsilon(\mathbf{y})} \left[\left(\delta_{hk} + \frac{3}{(1-2\nu)} \frac{\partial |\mathbf{x} - \mathbf{y}|}{\partial x_k} \frac{\partial |\mathbf{x} - \mathbf{y}|}{\partial x_h} \right) \frac{\partial}{\partial \mathbf{n}_{\mathbf{x}}} \frac{1}{|\mathbf{x} - \mathbf{y}|} \right] d\sigma(\mathbf{x}) + O(\varepsilon).
 \end{aligned}$$

Employing again the spherical coordinates and the definition of c'_ν , we find that

$$(3.21) \quad \frac{1-2\nu}{8\pi(1-\nu)} \int_{\partial B_\varepsilon(\mathbf{y})} \delta_{hk} \frac{\partial}{\partial \mathbf{n}_{\mathbf{x}}} \frac{1}{|\mathbf{x} - \mathbf{y}|} d\sigma(\mathbf{x}) = \begin{cases} -\frac{1-2\nu}{2(1-\nu)} & \text{if } h = k \\ 0 & \text{if } h \neq k. \end{cases}$$

Similarly

$$\begin{aligned}
 (3.22) \quad \frac{3}{8\pi(1-\nu)} \int_{\partial B_\varepsilon(\mathbf{y})} \left(\frac{\partial |\mathbf{x} - \mathbf{y}|}{\partial x_k} \frac{\partial |\mathbf{x} - \mathbf{y}|}{\partial x_h} \right) \frac{\partial}{\partial \mathbf{n}_{\mathbf{x}}} \frac{1}{|\mathbf{x} - \mathbf{y}|} d\sigma(\mathbf{x}) \\
 = \begin{cases} -\frac{1}{2(1-\nu)} & \text{if } h = k \\ 0 & \text{if } h \neq k. \end{cases}
 \end{aligned}$$

Putting together all the results in (3.21) and (3.22), we find that

$$\lim_{\varepsilon \rightarrow 0} \left(-\mathbf{u}(\mathbf{y}) \cdot \int_{\partial B_\varepsilon(\mathbf{y})} \frac{\partial \mathbf{N}^{(k)}}{\partial \boldsymbol{\nu}_{\mathbf{x}}}(\mathbf{x} - \mathbf{y}) d\sigma(\mathbf{x}) \right) = -u_k(\mathbf{x}).$$

Using the definition of single and double layer potentials (3.7), (3.13) and splitting \mathbf{N} as $\boldsymbol{\Gamma} + \mathbf{R}$, Formula (3.14) holds. \square

From the behaviour of the Neumann function given in (3.12) and the representation formula in (3.14), we immediately get

COROLLARY 3.4. — *If \mathbf{u} is a solution to (3.1), then*

$$(3.23) \quad \mathbf{u}(\mathbf{y}) = O(|\mathbf{y}|^{-1}) \quad \text{as } |\mathbf{y}| \rightarrow \infty.$$

3.4. WELL-POSEDNESS. — The well-posedness of the boundary value problem (3.1) reduces to show the invertibility of

$$(3.24) \quad \frac{1}{2}\mathbf{I} + \mathbf{K} + \mathbf{D}^R : L^2(\partial C) \rightarrow L^2(\partial C).$$

In particular, in order to prove the injectivity of the operator (3.24) we show the uniqueness of \mathbf{u} following the classical approach based on the application of the Betti's formula (3.2) and the energy method, see [21, 27]. From the injectivity, it follows the existence of \mathbf{u} proving the surjectivity of (3.24) which is obtained by the application of the index theory regarding bounded and linear operators.

First of all, let us recall the closed range theorem due to Banach (see [25, 41]).

THEOREM 3.5. — *Let X and Y be Banach spaces, and T a bounded linear operator defined in X into Y . Then the following propositions are all equivalent:*

- a. $\text{Im}(T)$ is closed in Y ; b. $\text{Im}(T^*)$ is closed in X^* ;
 c. $\text{Im}(T) = (\text{Ker}(T^*))^\perp$; d. $\text{Im}(T^*) = (\text{Ker}(T))^\perp$.

Through this theorem we can prove

LEMMA 3.6. — *The operator $\frac{1}{2}\mathbf{I} + \mathbf{K} : \mathbf{L}^2(\partial C) \rightarrow \mathbf{L}^2(\partial C)$ is invertible with bounded inverse.*

Proof. — The assertion of this lemma is based on the invertibility of the operator $\frac{1}{2}\mathbf{I} + \mathbf{K}^*$ studied in [15]; it is known that

$$\frac{1}{2}\mathbf{I} + \mathbf{K}^* : \mathbf{L}^2(\partial C) \longrightarrow \mathbf{L}^2(\partial C)$$

is a bounded linear operator, injective and with dense and closed range. Therefore, from Theorem 3.5 we have

$$\text{Ker}\left(\frac{1}{2}\mathbf{I} + \mathbf{K}\right) = \{0\}, \quad \text{Im}\left(\frac{1}{2}\mathbf{I} + \mathbf{K}\right)^\perp = \{0\}$$

and $\text{Im}(1/2\mathbf{I} + \mathbf{K})$ is closed. Then, it follows that the operator $\frac{1}{2}\mathbf{I} + \mathbf{K} : \mathbf{L}^2(\partial C) \rightarrow \mathbf{L}^2(\partial C)$ is bijective and the assertion follows exploiting the bounded inverse theorem. \square

Since \mathbf{D}^R has a continuous kernel we prove its compactness adapting the arguments contained in [26].

LEMMA 3.7. — *The operator $\mathbf{D}^R : \mathbf{L}^2(\partial C) \rightarrow \mathbf{L}^2(\partial C)$ is compact.*

Proof. — For the sake of simplicity, we call

$$\mathbf{H}(\mathbf{x}, \mathbf{y}) := \frac{\partial \mathbf{R}}{\partial \boldsymbol{\nu}}(\mathbf{x}, \mathbf{y}), \quad \mathbf{x}, \mathbf{y} \in \partial C$$

and we denote by $\mathbf{H}^{(k)}$, $k = 1, 2, 3$, the column vectors of the matrix \mathbf{H} .

Let \mathbf{S} be a bounded set such that $\mathbf{S} \subset \mathbf{L}^2(\partial C)$, that is, $\|\varphi\|_{\mathbf{L}^2(\partial C)} \leq K$, for any $\varphi \in \mathbf{S}$. Then, applying Cauchy-Schwarz inequality

$$|(\mathbf{D}^R \varphi(\mathbf{y}))_k|^2 \leq \|\mathbf{H}^{(k)}(\cdot, \mathbf{y})\|_{\mathbf{L}^2(\partial C)}^2 \|\varphi\|_{\mathbf{L}^2(\partial C)}^2 \leq K |\partial C| \max_{\mathbf{x}, \mathbf{y} \in \partial C} |\mathbf{H}^{(k)}|,$$

with $k = 1, 2, 3$, for all $\mathbf{y} \in \partial C$ and $\varphi \in \mathbf{S}$. Hence $|\mathbf{D}^R(\varphi)| \leq K'$, with $K' > 0$, which implies that $\mathbf{D}^R(\mathbf{S})$ is bounded. Moreover, for all $\varepsilon > 0$ there exist $\varphi, \varphi' \in \mathbf{S}$ and $\delta > 0$ such that if $\|\varphi(\mathbf{y}) - \varphi'(\mathbf{y})\|_{\mathbf{L}^2(\partial C)} < \delta$ then, applying again the Cauchy-Schwarz inequality

$$|\mathbf{D}^R(\varphi - \varphi')(\mathbf{y})| < \varepsilon.$$

Thus $\mathbf{D}^R(\mathbf{S}) \subset \mathbf{C}(\partial C)$, where $\mathbf{C}(\partial C)$ indicates the space of continuous function on ∂C . Since each component of the matrix \mathbf{H} is uniformly continuous on the compact set $\partial C \times \partial C$, for every $\varepsilon > 0$ there exists $\delta > 0$ such that

$$|\mathbf{H}^{(k)}(\mathbf{z}, \mathbf{x}) - \mathbf{H}^{(k)}(\mathbf{z}, \mathbf{y})| \leq \frac{\varepsilon}{\sqrt{3K}|\partial C|^{1/2}},$$

for all $\mathbf{x}, \mathbf{y}, \mathbf{z} \in \partial C$ with $|\mathbf{x} - \mathbf{y}| < \delta$. Since

$$\begin{aligned} |(\mathbf{D}^R \boldsymbol{\varphi})_k(\mathbf{x}) - (\mathbf{D}^R \boldsymbol{\varphi})_k(\mathbf{y})| &\leq \int_{\partial C} |\mathbf{H}^{(k)}(\mathbf{z}, \mathbf{x}) - \mathbf{H}^{(k)}(\mathbf{z}, \mathbf{y})| |\boldsymbol{\varphi}(\mathbf{z})| d\sigma(\mathbf{z}) \\ &\leq \|\mathbf{H}^{(k)}(\cdot, \mathbf{x}) - \mathbf{H}^{(k)}(\cdot, \mathbf{y})\|_{L^2(\partial C)} \|\boldsymbol{\varphi}\|_{L^2(\partial C)} \leq \frac{\varepsilon}{\sqrt{3}}, \end{aligned}$$

for $k = 1, 2, 3$, hence

$$|(\mathbf{D}^R \boldsymbol{\varphi})(\mathbf{x}) - (\mathbf{D}^R \boldsymbol{\varphi})(\mathbf{y})| \leq \varepsilon,$$

for all $\mathbf{x}, \mathbf{y} \in \partial C$ and $\boldsymbol{\varphi} \in \mathbf{S}$, that is, $\mathbf{D}^R(\mathbf{S})$ is equicontinuous. The assertion follows from Ascoli-Arzelà Theorem and noticing that $\mathbf{C}(\partial C)$ is dense in $\mathbf{L}^2(\partial C)$. \square

We now prove

THEOREM 3.8 (uniqueness). — *The boundary value problem (3.1) admits a unique solution.*

Proof. — Let \mathbf{u}^1 and \mathbf{u}^2 be solutions to (3.1). Then the difference $\mathbf{v} := \mathbf{u}^1 - \mathbf{u}^2$ solves the homogeneous version of (3.1), that is,

$$(3.25) \quad \operatorname{div}(\mathbb{C} \widehat{\nabla} \mathbf{v}) = \mathbf{0} \quad \text{in } \mathbb{R}_-^3 \setminus C$$

with homogeneous boundary conditions

$$(3.26) \quad \begin{aligned} \frac{\partial \mathbf{v}}{\partial \boldsymbol{\nu}} = \mathbf{0} \quad \text{on } \partial C, \quad \frac{\partial \mathbf{v}}{\partial \boldsymbol{\nu}} = \mathbf{0} \quad \text{on } \mathbb{R}^2 \\ \mathbf{v} = O(|\mathbf{x}|^{-1}) \quad \nabla \mathbf{v} = o(|\mathbf{x}|^{-1}) \quad |\mathbf{x}| \rightarrow \infty, \end{aligned}$$

where we make use of the decay condition at infinity comes from Corollary 3.4. Applying Betti's formula (3.2) to \mathbf{v} in $\Omega_r = (\mathbb{R}_-^3 \cap B_r(\mathbf{0})) \setminus C$, we find

$$\int_{\partial \Omega_r} \mathbf{v} \cdot \frac{\partial \mathbf{v}}{\partial \boldsymbol{\nu}} d\sigma(\mathbf{x}) = \int_{\Omega_r} Q(\mathbf{v}, \mathbf{v}) d\mathbf{x},$$

where Q is the quadratic form $Q(\mathbf{v}, \mathbf{v}) = \lambda(\operatorname{div} \mathbf{v})^2 + 2\mu|\widehat{\nabla} \mathbf{v}|^2$. From the behaviour of \mathbf{v} and the boundary conditions (3.26), we estimate the previous integral defined on the surface of Ω_r ; contributions over the surface of the cavity and the intersection of the hemisphere with the half-space are null by means of (3.26), whereas on the spherical cap

$$\left| \int_{\partial B_r^b(\mathbf{0})} \mathbf{v} \cdot \frac{\partial \mathbf{v}}{\partial \boldsymbol{\nu}} d\sigma(\mathbf{x}) \right| \leq \int_{\partial B_r^b(\mathbf{0})} |\mathbf{v}| \left| \frac{\partial \mathbf{v}}{\partial \boldsymbol{\nu}} \right| d\sigma(\mathbf{x}) \leq \frac{C}{r} \int_{\partial B_r^b(\mathbf{0})} \left| \frac{\partial \mathbf{v}}{\partial \boldsymbol{\nu}} \right| d\sigma(\mathbf{x}).$$

As already done in (3.16) to obtain the representation formula, this integral can be evaluated by spherical coordinates; in particular, it tends to zero when $r \rightarrow +\infty$. Therefore

$$\int_{\mathbb{R}_-^3 \setminus C} \left\{ \lambda(\operatorname{div} \mathbf{v})^2 + 2\mu|\widehat{\nabla} \mathbf{v}|^2 \right\} d\mathbf{x} = 0.$$

Since the quadratic form is positive definite for the parameters range $3\lambda + 2\mu > 0$ and $\mu > 0$, we have that

$$(3.27) \quad \widehat{\nabla} \mathbf{v} = \mathbf{0} \quad \text{and} \quad \operatorname{div} \mathbf{v} = 0 \quad \text{in } \mathbb{R}_-^3 \setminus C.$$

It follows that the rigid displacements $\mathbf{v} = \mathbf{a} + \mathbf{A}\mathbf{x}$, with $\mathbf{a} \in \mathbb{R}^3$ and \mathbf{A} belonging to the space of the anti-symmetric matrices (see [4, 12]), could be the only nonzero solutions which satisfy (3.25), the boundary conditions in (3.26) and (3.27). However, in this case they are excluded thanks to the behaviour of the function \mathbf{v} at infinity. Hence, we obtain that $\mathbf{v} = \mathbf{0}$, that is, $\mathbf{u}^1 = \mathbf{u}^2$ in $\mathbb{R}_-^3 \setminus C$. \square

The uniqueness result for problem (3.1) ensures the injectivity of the operator (3.24). In order to prove the surjectivity of the operator (3.24) we recall, for the reader's convenience, the definition of the index of an operator (see [1, 25]).

DEFINITION 3.1. — Given a bounded operator $T : X \rightarrow Y$ between two Banach spaces, the index of the operator T is the extended real number defined as

$$i(T) = \dim(\operatorname{Ker}(T)) - \dim(Y/\operatorname{Im}(T)),$$

where $\dim(\operatorname{Ker}(T))$ is called the *nullity* and $\dim(Y/\operatorname{Im}(T))$ the *defect* of T . In particular, when the nullity and the defect are both finite the operator T is said to be *Fredholm*.

We recall also an important theorem regarding the index of a bounded linear operator perturbed with a compact operator (see [1]).

THEOREM 3.9. — *Let $T : X \rightarrow Y$ be a Fredholm bounded linear operator and $K : X \rightarrow Y$ a compact operator between two Banach spaces. Then $T + K$ is Fredholm with index $i(T + K) = i(T)$.*

Now all the ingredients are supplied in order to prove the surjectivity of the operator.

THEOREM 3.10. — *The operator $\frac{1}{2}\mathbf{I} + \mathbf{K} + \mathbf{D}^R$ is onto in $L^2(\partial C)$.*

Proof. — From Lemma 3.6 we have that the operator $\frac{1}{2}\mathbf{I} + \mathbf{K} : L^2(\partial C) \rightarrow L^2(\partial C)$ is Fredholm with index $i(\frac{1}{2}\mathbf{I} + \mathbf{K}) = 0$, because both the nullity and the defect of this operator are null. Moreover, since the operator \mathbf{D}^R is compact from Lemma 3.7, it follows by means of Theorem 3.9 that

$$i(\frac{1}{2}\mathbf{I} + \mathbf{K} + \mathbf{D}^R) = 0.$$

Hence

$$\dim(\operatorname{Ker}(\frac{1}{2}\mathbf{I} + \mathbf{K} + \mathbf{D}^R)) = \dim(L^2(\partial C)/\operatorname{Im}(\frac{1}{2}\mathbf{I} + \mathbf{K} + \mathbf{D}^R)).$$

Since the operator $\frac{1}{2}\mathbf{I} + \mathbf{K} + \mathbf{D}^R$ is injective it shows that $\dim(\operatorname{Ker}(\frac{1}{2}\mathbf{I} + \mathbf{K} + \mathbf{D}^R)) = 0$. Finally, $\dim(L^2(\partial C)/\operatorname{Im}(\frac{1}{2}\mathbf{I} + \mathbf{K} + \mathbf{D}^R)) = 0$, that is, $\operatorname{Im}(\frac{1}{2}\mathbf{I} + \mathbf{K} + \mathbf{D}^R) = L^2(\partial C)$. \square

Summing up, it follows:

COROLLARY 3.11. — *There exists a unique solution to (3.1).*

Proof. — Uniqueness follows from Theorem 3.8 and the existence from Theorem 3.10. \square

4. RIGOROUS DERIVATION OF THE ASYMPTOTIC EXPANSION

In this section, with the integral representation formula (3.14) at hand, we consider the hypothesis that the cavity C is small compared to the distance from the boundary of the half-space. The aim is to derive an asymptotic expansion of the solution \mathbf{u} .

In particular, let us take the cavity, that from now on we denote by C_ε to highlight the dependence on ε , as

$$C_\varepsilon = \mathbf{z} + \varepsilon\Omega,$$

where Ω is a bounded Lipschitz domain containing the origin. At the same time, we write the solution of the boundary value problem (3.1) as \mathbf{u}_ε . From (3.14), recalling that $\mathbf{N} = \mathbf{\Gamma} + \mathbf{R}$, we have

$$(4.1) \quad \begin{aligned} u_\varepsilon^k(\mathbf{y}) &= p \int_{\partial C_\varepsilon} \mathbf{N}^{(k)}(\mathbf{x}, \mathbf{y}) \cdot \mathbf{n}(\mathbf{x}) \, d\sigma(\mathbf{x}) - \int_{\partial C_\varepsilon} \frac{\partial \mathbf{N}^{(k)}}{\partial \boldsymbol{\nu}}(\mathbf{x}, \mathbf{y}) \cdot \mathbf{f}(\mathbf{x}) \, d\sigma(\mathbf{x}) \\ &:= I_1^{(k)}(\mathbf{y}) + I_2^{(k)}(\mathbf{y}), \quad \mathbf{y} \in \mathbb{R}^2, \end{aligned}$$

for $k = 1, 2, 3$, where u_ε^k indicates the k -th component of the displacement vector and $\mathbf{f} = \mathbf{f}_\varepsilon$ is the solution of (3.15), that is,

$$(4.2) \quad \left(\frac{1}{2}\mathbb{I} + \mathbf{K}_\varepsilon + \mathbf{D}_\varepsilon^R\right) \mathbf{f}_\varepsilon(\mathbf{x}) = p \left(\mathbf{S}_\varepsilon^\Gamma(\mathbf{n})(\mathbf{x}) + \mathbf{S}_\varepsilon^R(\mathbf{n})(\mathbf{x})\right), \quad \mathbf{x} \in \partial C_\varepsilon,$$

where we add the dependence on ε to all the layer potentials to distinguish them, in the sequel, from the layer potential defined over a domain independent of ε . In what follows, with \mathbb{I} we indicate the fourth-order symmetric tensor such that $\mathbb{I}\mathbf{A} = \widehat{\mathbf{A}}$ and for any fixed value of $\varepsilon > 0$, given $\mathbf{h} : \partial C_\varepsilon \rightarrow \mathbb{R}^3$, we introduce the function $\widehat{\mathbf{h}} : \partial\Omega \rightarrow \mathbb{R}^3$ defined by

$$\widehat{\mathbf{h}}(\boldsymbol{\zeta}) := \mathbf{h}(\mathbf{z} + \varepsilon\boldsymbol{\zeta}), \quad \boldsymbol{\zeta} \in \partial\Omega.$$

Moreover, we consider the functions $\boldsymbol{\theta}^{qr}$, for $q, r = 1, 2, 3$, solutions to

$$(4.3) \quad \operatorname{div}(\mathbb{C}\widehat{\nabla}\boldsymbol{\theta}^{qr}) = 0 \quad \text{in } \mathbb{R}^3 \setminus \Omega, \quad \frac{\partial \boldsymbol{\theta}^{qr}}{\partial \boldsymbol{\nu}} = -\frac{1}{3\lambda + 2\mu} \mathbb{C}\mathbf{n} \quad \text{on } \partial\Omega,$$

with the decay conditions at infinity

$$(4.4) \quad |\boldsymbol{\theta}^{qr}| = O(|\mathbf{x}|^{-1}), \quad |\nabla\boldsymbol{\theta}^{qr}| = O(|\mathbf{x}|^{-2}), \quad \text{as } |\mathbf{x}| \rightarrow \infty,$$

where the condition $\partial\boldsymbol{\theta}^{qr}/\partial\boldsymbol{\nu}$ has to be read as

$$\left(\frac{\partial \boldsymbol{\theta}^{qr}}{\partial \boldsymbol{\nu}}\right)_i = -\frac{1}{3\lambda + 2\mu} C_{ijqr} n_j.$$

We now state our main result.

THEOREM 4.1 (asymptotic expansion). — *There exists $\varepsilon_0 > 0$ such that for all $\varepsilon \in (0, \varepsilon_0)$ the following expansion holds at any $\mathbf{y} \in \mathbb{R}^2$:*

$$(4.5) \quad u_\varepsilon^k(\mathbf{y}) = \varepsilon^3 |\Omega| p \widehat{\nabla}_{\mathbf{z}} \mathbf{N}^{(k)}(\mathbf{z}, \mathbf{y}) : \mathbb{M} \mathbf{I} + O(\varepsilon^4),$$

for $k = 1, 2, 3$, where $O(\varepsilon^4)$ denotes a quantity bounded by $C\varepsilon^4$ for some uniform constant $C > 0$, and \mathbb{M} is the fourth-order elastic moment tensor defined by

$$(4.6) \quad \mathbb{M} := \mathbb{I} + \frac{1}{|\Omega|} \int_{\partial\Omega} \mathbb{C}(\boldsymbol{\theta}^{qr}(\boldsymbol{\zeta}) \otimes \mathbf{n}(\boldsymbol{\zeta})) d\sigma(\boldsymbol{\zeta}),$$

where $\boldsymbol{\theta}^{qr}$, for $q, r = 1, 2, 3$, solve the problem in (4.3) and (4.4).

Before proving the theorem on the asymptotic expansion of \mathbf{u}_ε , we need to present some results.

LEMMA 4.2. — *The integral equation (4.2), when $\mathbf{x} = \mathbf{z} + \varepsilon\boldsymbol{\zeta}$, with $\boldsymbol{\zeta} \in \partial\Omega$, is such that*

$$(4.7) \quad \left(\frac{1}{2}\mathbf{I} + \mathbf{K} + \varepsilon^2 \boldsymbol{\Lambda}_{\Omega, \varepsilon}\right) \widehat{\mathbf{f}}(\boldsymbol{\zeta}) = \varepsilon p \mathbf{S}^\Gamma(\mathbf{n})(\boldsymbol{\zeta}) + O(\varepsilon^2),$$

where

$$\boldsymbol{\Lambda}_{\Omega, \varepsilon} \widehat{\mathbf{f}}(\boldsymbol{\eta}) := \int_{\partial\Omega} \frac{\partial \mathbf{R}}{\partial \boldsymbol{\nu}(\boldsymbol{\eta})}(\mathbf{z} + \varepsilon\boldsymbol{\eta}, \mathbf{z} + \varepsilon\boldsymbol{\zeta}) \widehat{\mathbf{f}}(\boldsymbol{\eta}) d\sigma(\boldsymbol{\eta})$$

is uniformly bounded in ε . Moreover, when ε is sufficiently small, we have

$$\widehat{\mathbf{f}}(\boldsymbol{\zeta}) = \varepsilon p \left(\frac{1}{2}\mathbf{I} + \mathbf{K}\right)^{-1} \mathbf{S}^\Gamma(\mathbf{n})(\boldsymbol{\zeta}) + O(\varepsilon^2), \quad \boldsymbol{\zeta} \in \partial\Omega.$$

Proof. — At the point $\mathbf{z} + \varepsilon\boldsymbol{\zeta}$, where $\boldsymbol{\zeta} \in \partial\Omega$, we obtain

$$\begin{aligned} \mathbf{D}_\varepsilon^R \mathbf{f}(\mathbf{z} + \varepsilon\boldsymbol{\zeta}) &= \int_{\partial C_\varepsilon} \frac{\partial \mathbf{R}}{\partial \boldsymbol{\nu}(\mathbf{t})}(\mathbf{t}, \mathbf{z} + \varepsilon\boldsymbol{\zeta}) \mathbf{f}(\mathbf{t}) d\sigma(\mathbf{t}) \\ &= \varepsilon^2 \int_{\partial\Omega} \frac{\partial \mathbf{R}}{\partial \boldsymbol{\nu}(\boldsymbol{\eta})}(\mathbf{z} + \varepsilon\boldsymbol{\eta}, \mathbf{z} + \varepsilon\boldsymbol{\zeta}) \widehat{\mathbf{f}}(\boldsymbol{\eta}) d\sigma(\boldsymbol{\eta}). \end{aligned}$$

Therefore, recalling that the kernel $\partial \mathbf{R} / \partial \boldsymbol{\nu}(\boldsymbol{\eta})$ is continuous, we get

$$(4.8) \quad \mathbf{D}_\varepsilon^R = \varepsilon^2 \boldsymbol{\Lambda}_{\Omega, \varepsilon}$$

where $\|\boldsymbol{\Lambda}_{\Omega, \varepsilon}\| \leq C'$, with C' independent of ε . For the integral

$$\mathbf{K}_\varepsilon \mathbf{f}(\mathbf{z} + \varepsilon\boldsymbol{\zeta}) = \text{p.v.} \int_{\partial C_\varepsilon} \frac{\partial \mathbf{I}}{\partial \boldsymbol{\nu}(\mathbf{t})}(\mathbf{t} - \mathbf{z} - \varepsilon\boldsymbol{\zeta}) \mathbf{f}(\mathbf{t}) d\sigma(\mathbf{t})$$

we use the explicit expression of the conormal derivative of the fundamental solution of the Lamé operator given in (3.18). In particular, since (3.18) is a homogeneous

function of degree -2 , with the substitution $\mathbf{t} = \mathbf{z} + \varepsilon\boldsymbol{\eta}$, we find

$$\begin{aligned} \left(\frac{\partial\boldsymbol{\Gamma}^{(k)}}{\partial\nu}\right)_h(\varepsilon(\boldsymbol{\eta} - \boldsymbol{\zeta})) &= -\frac{1}{4\pi\varepsilon^2} \left\{ \left[\frac{1-2\nu}{2(1-\nu)}\delta_{hk} + \frac{3}{2(1-\nu)} \frac{\eta_k - \zeta_k}{|\boldsymbol{\eta} - \boldsymbol{\zeta}|} \frac{\eta_h - \zeta_h}{|\boldsymbol{\eta} - \boldsymbol{\zeta}|} \right] \frac{\partial}{\partial\mathbf{n}(\boldsymbol{\eta})} \frac{1}{|\boldsymbol{\eta} - \boldsymbol{\zeta}|} \right. \\ &\quad \left. + \frac{1-2\nu}{2(1-\nu)} n_h(\boldsymbol{\eta}) \frac{\eta_k - \zeta_k}{|\boldsymbol{\eta} - \boldsymbol{\zeta}|^3} - \frac{1-2\nu}{2(1-\nu)} n_k(\boldsymbol{\eta}) \frac{\eta_h - \zeta_h}{|\boldsymbol{\eta} - \boldsymbol{\zeta}|^3} \right\} \\ &= \frac{1}{\varepsilon^2} \left(\frac{\partial\boldsymbol{\Gamma}^{(k)}}{\partial\nu}\right)_h(\boldsymbol{\eta} - \boldsymbol{\zeta}), \end{aligned}$$

for $h, k = 1, 2, 3$. Therefore, we immediately obtain that

$$(4.9) \quad \mathbf{K}_\varepsilon \mathbf{f}(\mathbf{z} + \varepsilon\boldsymbol{\zeta}) = \text{p.v.} \int_{\partial\Omega} \frac{\partial\boldsymbol{\Gamma}}{\partial\nu(\boldsymbol{\eta})}(\boldsymbol{\eta} - \boldsymbol{\zeta}) \widehat{\mathbf{f}}(\boldsymbol{\eta}) d\sigma(\boldsymbol{\eta}) = \mathbf{K}\widehat{\mathbf{f}}(\boldsymbol{\zeta}).$$

Evaluating the other integrals in (4.2) we obtain

$$\mathbf{S}_\varepsilon^\Gamma(\mathbf{n})(\mathbf{z} + \varepsilon\boldsymbol{\zeta}) = \int_{\partial C_\varepsilon} \boldsymbol{\Gamma}(\mathbf{t} - \mathbf{z} - \varepsilon\boldsymbol{\zeta}) \mathbf{n}(\mathbf{t}) d\sigma(\mathbf{t}),$$

hence, choosing $\mathbf{t} = \mathbf{z} + \varepsilon\boldsymbol{\eta}$, with $\boldsymbol{\eta} \in \partial\Omega$, we find

$$(4.10) \quad \mathbf{S}_\varepsilon^\Gamma(\mathbf{n})(\mathbf{z} + \varepsilon\boldsymbol{\zeta}) = \varepsilon^2 \int_{\partial\Omega} \boldsymbol{\Gamma}(\varepsilon(\boldsymbol{\eta} - \boldsymbol{\zeta})) \mathbf{n}(\boldsymbol{\eta}) d\sigma(\boldsymbol{\eta}) = \varepsilon \mathbf{S}^\Gamma(\mathbf{n})(\boldsymbol{\zeta}),$$

where the last equality follows noticing that the fundamental solution is homogeneous of degree -1 . In a similar way

$$\mathbf{S}_\varepsilon^R(\mathbf{n})(\mathbf{z} + \varepsilon\boldsymbol{\zeta}) = \int_{\partial C_\varepsilon} \mathbf{R}(\mathbf{t}, \mathbf{z} + \varepsilon\boldsymbol{\zeta}) \mathbf{n}(\mathbf{t}) d\sigma(\mathbf{t}),$$

hence, taking again $\mathbf{t} = \mathbf{z} + \varepsilon\boldsymbol{\eta}$, we find

$$\mathbf{S}_\varepsilon^R(\mathbf{n})(\mathbf{z} + \varepsilon\boldsymbol{\zeta}) = \varepsilon^2 \int_{\partial\Omega} \mathbf{R}(\mathbf{z} + \varepsilon\boldsymbol{\eta}, \mathbf{z} + \varepsilon\boldsymbol{\zeta}) \mathbf{n}(\boldsymbol{\eta}) d\sigma(\boldsymbol{\eta})$$

and since \mathbf{R} is regular it follows that

$$(4.11) \quad \mathbf{S}_\varepsilon^R(\mathbf{n})(\mathbf{z} + \varepsilon\boldsymbol{\zeta}) = O(\varepsilon^2).$$

Relation (4.7) follows putting together the result in (4.8), (4.9), (4.10) and (4.11).

To conclude, from (4.7) we have

$$\left(\frac{1}{2}\mathbf{I} + \mathbf{K}\right) \left(\mathbf{I} + \varepsilon^2 \left(\frac{1}{2}\mathbf{I} + \mathbf{K}\right)^{-1} \boldsymbol{\Lambda}_{\varepsilon, \Omega}\right) \widehat{\mathbf{f}} = \varepsilon p \mathbf{S}^\Gamma(\mathbf{n}) + O(\varepsilon^2) \quad \text{on } \partial\Omega.$$

From Lemma 3.6 and the continuous property of $\boldsymbol{\Lambda}_{\varepsilon, \Omega}$ described before, we have

$$\left\| \left(\frac{1}{2}\mathbf{I} + \mathbf{K}\right)^{-1} \boldsymbol{\Lambda}_{\varepsilon, \Omega} \right\| \leq C,$$

where $C > 0$ is independent of ε . On the other hand, choosing $\varepsilon_0^2 = 1/2C$, it follows that, for all $\varepsilon \in (0, \varepsilon_0)$,

$$\mathbf{I} + \varepsilon^2 \left(\frac{1}{2}\mathbf{I} + \mathbf{K}\right)^{-1} \boldsymbol{\Lambda}_{\varepsilon, \Omega}$$

is invertible and

$$\left(\mathbf{I} + \varepsilon^2 \left(\frac{1}{2}\mathbf{I} + \mathbf{K}\right)^{-1} \mathbf{\Lambda}_{\varepsilon, \Omega}\right)^{-1} = \mathbf{I} + O(\varepsilon^2).$$

Therefore

$$\widehat{\mathbf{f}} = \varepsilon p \left(\frac{1}{2}\mathbf{I} + \mathbf{K}\right)^{-1} \mathbf{S}^\Gamma(\mathbf{n}) + O(\varepsilon^2) \quad \text{on } \partial\Omega,$$

that is, the assertion. \square

For ease of reading, we define the function $\mathbf{w} : \partial\Omega \rightarrow \partial\Omega$ as

$$(4.12) \quad \mathbf{w}(\boldsymbol{\zeta}) := -\left(\frac{1}{2}\mathbf{I} + \mathbf{K}\right)^{-1} \mathbf{S}^\Gamma(\mathbf{n})(\boldsymbol{\zeta}), \quad \boldsymbol{\zeta} \in \partial\Omega.$$

Taking the problem

$$(4.13) \quad \operatorname{div} \left(\mathbb{C} \widehat{\nabla} \mathbf{v} \right) = 0 \quad \text{in } \mathbb{R}^3 \setminus \Omega, \quad \frac{\partial \mathbf{v}}{\partial \boldsymbol{\nu}} = -\mathbf{n} \quad \text{on } \partial\Omega$$

with decay conditions at infinity

$$(4.14) \quad \mathbf{v} = O(|\mathbf{x}|^{-1}), \quad |\nabla \mathbf{v}| = O(|\mathbf{x}|^{-2}) \quad \text{as } |\mathbf{x}| \rightarrow +\infty,$$

we show that $\mathbf{w}(\mathbf{x})$, for $\mathbf{x} \in \partial\Omega$, is the trace of \mathbf{v} on the boundary of Ω . The well-posedness of this problem is a classical result in the theory of linear elasticity so we remind the reader, for example, to [21, 23, 27].

PROPOSITION 4.3. — *The function \mathbf{w} , defined in (4.12), is such that $\mathbf{w} = \mathbf{v}|_{\mathbf{x} \in \partial\Omega}$, where \mathbf{v} is the solution to (4.13) and (4.14).*

Proof. — Applying the second Betti formula to the fundamental solution $\boldsymbol{\Gamma}$ and the function \mathbf{v} into the domain $B_r(\mathbf{0}) \setminus (\Omega \cup B_\varepsilon(\mathbf{x}))$, with $\varepsilon > 0$ and $r > 0$ sufficiently large so that the domain contains the cavity Ω , we obtain, as done in a similar way in the proof of the Theorem 3.3,

$$\mathbf{v}(\mathbf{x}) = -\mathbf{S}^\Gamma \mathbf{n}(\mathbf{x}) - \mathbf{D}^\Gamma \mathbf{v}(\mathbf{x}), \quad \mathbf{x} \in \mathbb{R}^3 \setminus \Omega.$$

Therefore, from the single and double layer potential properties for the elastostatic equations, we find

$$\mathbf{v}(\mathbf{x}) = -\mathbf{S}^\Gamma \mathbf{n}(\mathbf{x}) - \left(-\frac{1}{2}\mathbf{I} + \mathbf{K}\right) \mathbf{v}(\mathbf{x}), \quad \mathbf{x} \in \partial\Omega,$$

hence

$$\mathbf{v}(\mathbf{x}) = -\left(\frac{1}{2}\mathbf{I} + \mathbf{K}\right)^{-1} \mathbf{S}^\Gamma(\mathbf{n})(\mathbf{x}), \quad \mathbf{x} \in \partial\Omega,$$

that is, the assertion. \square

We note that the function \mathbf{v} , as well as its trace \mathbf{w} on $\partial\Omega$, can be written in terms of the functions $\boldsymbol{\theta}^{qr}$. Indeed, taking

$$\mathbf{v} = \boldsymbol{\theta}^{qr} \delta_{qr}, \quad q, r = 1, 2, 3,$$

and using (4.3) and (4.4), it is straightforward to see that the elastostatic equation and the boundary condition in (4.13) are satisfied.

Proof of Theorem 4.1. — We study separately the two integrals $I_1^{(k)}, I_2^{(k)}$ defined in (4.1). Since $\mathbf{y} \in \mathbb{R}^2$ and $\mathbf{x} \in \partial C_\varepsilon = \mathbf{z} + \varepsilon\boldsymbol{\zeta}$, with $\boldsymbol{\zeta} \in \partial\Omega$, we consider the Taylor expansion for the Neumann function, that is,

$$(4.15) \quad \mathbf{N}^{(k)}(\mathbf{z} + \varepsilon\boldsymbol{\zeta}, \mathbf{y}) = \mathbf{N}^{(k)}(\mathbf{z}, \mathbf{y}) + \varepsilon \nabla \mathbf{N}^{(k)}(\mathbf{z}, \mathbf{y}) \boldsymbol{\zeta} + O(\varepsilon^2),$$

for $k = 1, 2, 3$. By the change of variable $\mathbf{x} = \mathbf{z} + \varepsilon\boldsymbol{\zeta}$ and substituting (4.15) in $I_1^{(k)}$, we find

$$\begin{aligned} I_1^{(k)} &= \varepsilon^2 p \mathbf{N}^{(k)}(\mathbf{z}, \mathbf{y}) \cdot \int_{\partial\Omega} \mathbf{n}(\boldsymbol{\zeta}) \, d\sigma(\boldsymbol{\zeta}) + \varepsilon^3 p \int_{\partial\Omega} \mathbf{n}(\boldsymbol{\zeta}) \cdot \nabla \mathbf{N}^{(k)}(\mathbf{z}, \mathbf{y}) \boldsymbol{\zeta} \, d\sigma(\boldsymbol{\zeta}) + O(\varepsilon^4) \\ &:= p \left(\varepsilon^2 I_{11}^{(k)} + \varepsilon^3 I_{12}^{(k)} \right) + O(\varepsilon^4). \end{aligned}$$

The integral $I_{11}^{(k)}$ is null, in fact, applying the divergence theorem

$$\int_{\partial\Omega} \mathbf{n}(\boldsymbol{\zeta}) \, d\sigma(\boldsymbol{\zeta}) = 0.$$

For the integral $I_{12}^{(k)}$, we use the equality $\mathbf{n} \cdot \nabla \mathbf{N}^{(k)} \boldsymbol{\zeta} = \nabla \mathbf{N}^{(k)} : (\mathbf{n}(\boldsymbol{\zeta}) \otimes \boldsymbol{\zeta})$, therefore

$$(4.16) \quad I_{12}^{(k)} = \varepsilon^3 p \nabla \mathbf{N}^{(k)}(\mathbf{z}, \mathbf{y}) : \int_{\partial\Omega} (\mathbf{n}(\boldsymbol{\zeta}) \otimes \boldsymbol{\zeta}) \, d\sigma(\boldsymbol{\zeta}) + O(\varepsilon^4), \quad k = 1, 2, 3.$$

For the term $I_2^{(k)}$ we use the result in Lemma 4.2 and the Taylor expansion of the conormal derivative of $\mathbf{N}^{(k)}(\mathbf{x}, \mathbf{y})$, for $k = 1, 2, 3$. In particular, for $\mathbf{x} = \mathbf{z} + \varepsilon\boldsymbol{\zeta}$, when $\boldsymbol{\zeta} \in \partial\Omega$ and $\mathbf{y} \in \mathbb{R}^2$, we consider only the first term of the asymptotic expansion, that is,

$$\frac{\partial \mathbf{N}^{(k)}}{\partial \boldsymbol{\nu}(\mathbf{x})}(\mathbf{x}, \mathbf{y}) = \frac{\partial \mathbf{N}^{(k)}}{\partial \boldsymbol{\nu}(\boldsymbol{\zeta})}(\mathbf{z}, \mathbf{y}) + O(\varepsilon), \quad k = 1, 2, 3.$$

Therefore

$$\begin{aligned} I_2^{(k)} &= -\varepsilon^2 \int_{\partial\Omega} \frac{\partial \mathbf{N}^{(k)}}{\partial \boldsymbol{\nu}(\mathbf{x})}(\mathbf{z} + \varepsilon\boldsymbol{\zeta}, \mathbf{y}) \cdot \widehat{\mathbf{f}}(\boldsymbol{\zeta}) \, d\sigma(\boldsymbol{\zeta}) \\ &= -\varepsilon^3 p \int_{\partial\Omega} \frac{\partial \mathbf{N}^{(k)}}{\partial \boldsymbol{\nu}(\boldsymbol{\zeta})}(\mathbf{z}, \mathbf{y}) \cdot \mathbf{w}(\boldsymbol{\zeta}) \, d\sigma(\boldsymbol{\zeta}) + O(\varepsilon^4), \end{aligned}$$

for any k , where \mathbf{w} is defined in (4.12). Since $\partial \mathbf{N}^{(k)} / \partial \boldsymbol{\nu}(\boldsymbol{\zeta}) = \mathbb{C} \widehat{\nabla} \mathbf{N}^{(k)} \mathbf{n}(\boldsymbol{\zeta})$, we have

$$\mathbb{C} \widehat{\nabla} \mathbf{N}^{(k)} \mathbf{n}(\boldsymbol{\zeta}) \cdot \mathbf{w}(\boldsymbol{\zeta}) = \mathbb{C} \widehat{\nabla} \mathbf{N}^{(k)} : (\mathbf{w}(\boldsymbol{\zeta}) \otimes \mathbf{n}(\boldsymbol{\zeta})).$$

Therefore

$$(4.17) \quad I_2^{(k)}(\mathbf{y}) = \varepsilon^3 p \mathbb{C} \widehat{\nabla} \mathbf{N}^{(k)}(\mathbf{z}, \mathbf{y}) : \int_{\partial\Omega} (\mathbf{w}(\boldsymbol{\zeta}) \otimes \mathbf{n}(\boldsymbol{\zeta})) \, d\sigma(\boldsymbol{\zeta}) + O(\varepsilon^4).$$

Collecting the result in (4.16) and (4.17), Equation (4.1) becomes

$$u_\varepsilon^k(\mathbf{y}) = I_1^{(k)}(\mathbf{y}) + I_2^{(k)}(\mathbf{y}) \\ = \varepsilon^3 p \left[\nabla \mathbf{N}^{(k)}(\mathbf{z}, \mathbf{y}) : \int_{\partial\Omega} (\mathbf{n} \otimes \boldsymbol{\zeta}) d\sigma(\boldsymbol{\zeta}) + \mathbb{C} \widehat{\nabla} \mathbf{N}^{(k)}(\mathbf{z}, \mathbf{y}) : \int_{\partial\Omega} (\mathbf{w} \otimes \mathbf{n}) d\sigma(\boldsymbol{\zeta}) \right] + O(\varepsilon^4).$$

Now, handling this expression, we highlight the moment elastic tensor. We have

$$(4.18) \quad \int_{\partial\Omega} (\mathbf{n}(\boldsymbol{\zeta}) \otimes \boldsymbol{\zeta}) d\sigma(\boldsymbol{\zeta}) = |\Omega| \mathbf{I},$$

indeed, for any $i, j = 1, 2, 3$, it follows

$$\int_{\partial\Omega} \zeta_i n_j d\sigma(\boldsymbol{\zeta}) = \int_{\partial\Omega} \mathbf{n} \cdot \zeta_i \mathbf{e}_j d\sigma(\boldsymbol{\zeta}) = \int_{\Omega} \operatorname{div}(\zeta_i \mathbf{e}_j) d\boldsymbol{\zeta} = \int_{\Omega} \mathbf{e}_j \cdot \mathbf{e}_i d\boldsymbol{\zeta} = |\Omega| \delta_{ij},$$

where \mathbf{e}_j is the j -th unit vector of \mathbb{R}^3 . Hence, by (4.18) and taking the symmetric part of $\nabla \mathbf{N}^{(k)}$, for any k , we find

$$u_\varepsilon^k = \varepsilon^3 p \left[\widehat{\nabla} \mathbf{N}^{(k)} : \mathbf{I} |\Omega| + \mathbb{C} \widehat{\nabla} \mathbf{N}^{(k)} : \int_{\partial\Omega} \mathbf{w} \otimes \mathbf{n} d\sigma(\boldsymbol{\zeta}) \right] + O(\varepsilon^4).$$

Using the symmetries of \mathbb{C} , we have

$$u_\varepsilon^k = \varepsilon^3 |\Omega| p \widehat{\nabla} \mathbf{N}^{(k)} : \left[\mathbf{I} + \frac{1}{|\Omega|} \int_{\partial\Omega} \mathbb{C}(\mathbf{w} \otimes \mathbf{n}) d\sigma(\boldsymbol{\zeta}) \right] + O(\varepsilon^4),$$

for $k = 1, 2, 3$. Now, taking into account that $\mathbf{I} = \mathbb{I}$ and using the equality $\mathbf{w} = \boldsymbol{\theta}^{qr} \delta_{qr}$, with $q, r = 1, 2, 3$, we have the assertion. \square

THE MOGI MODEL. — In this subsection, starting from the asymptotic expansion (4.5), that is,

$$u_\varepsilon^k(\mathbf{y}) = \varepsilon^3 |\Omega| p \widehat{\nabla}_z \mathbf{N}^{(k)}(\mathbf{z}, \mathbf{y}) : \mathbb{M} \mathbf{I} + O(\varepsilon^4), \quad k = 1, 2, 3,$$

where \mathbb{M} is the tensor given in (4.6), we recover the Mogi model, presented within the Section 2, related to a spherical cavity. We first recall that

$$\mathbb{M} \mathbf{I} = \left[\mathbb{I} + \frac{1}{|\Omega|} \int_{\partial\Omega} \mathbb{C}(\boldsymbol{\theta}^{qr} \otimes \mathbf{n}) d\sigma(\boldsymbol{\zeta}) \right] \mathbf{I} = \mathbf{I} + \frac{1}{|\Omega|} \int_{\partial\Omega} \mathbb{C}(\mathbf{w} \otimes \mathbf{n}) d\sigma(\boldsymbol{\zeta}),$$

where, in the last equality, we use the link between the functions \mathbf{w} and $\boldsymbol{\theta}^{qr}$, that is, $\mathbf{w} = \boldsymbol{\theta}^{qr} \delta_{qr}$, $q, r = 1, 2, 3$. Therefore, to get the Mogi's formula, we first find the explicit expression of \mathbf{w} when the cavity Ω is the unit sphere and then we calculate the gradient of the Neumann function \mathbf{N} .

We recall that \mathbf{w} is the trace on the boundary of the cavity of the solution to the external problem

$$\operatorname{div}(\mathbb{C} \widehat{\nabla} \mathbf{v}) = 0 \quad \text{in } \mathbb{R}^3 \setminus B_1(\mathbf{0}), \quad \frac{\partial \mathbf{v}}{\partial \nu} = -\mathbf{n} \quad \text{on } \partial B_1(\mathbf{0}),$$

where $B_1(\mathbf{0}) = \{\mathbf{x} \in \mathbb{R}^3 : |\mathbf{x}| \leq 1\}$ with decay at infinity

$$\mathbf{v} = O(|\mathbf{x}|^{-1}), \quad |\nabla \mathbf{v}| = O(|\mathbf{x}|^{-2}) \quad \text{as } |\mathbf{x}| \rightarrow +\infty.$$

We look for a solution of the form

$$\mathbf{v}(\mathbf{x}) = \phi(r) \mathbf{x} \quad \text{with } r := |\mathbf{x}|,$$

so that

$$\Delta v_i = \left\{ \phi'' + \frac{4\phi'}{r} \right\} x_i, \quad \operatorname{div} \mathbf{v} = r\phi' + 3\phi, \quad \nabla \operatorname{div} \mathbf{v} = \left\{ \phi'' + \frac{4\phi'}{r} \right\} \mathbf{x}.$$

By direct substitution, since $\mathbf{n} = \mathbf{x}$ on ∂B , we get

$$\begin{aligned} \operatorname{div}(\mathbb{C}\widehat{\nabla} \mathbf{v}) &= (\lambda + 2\mu) \left(\phi'' + \frac{4\phi'}{r} \right) \mathbf{x}, \\ \frac{\partial \mathbf{v}}{\partial \nu} &= \{ (\lambda + 2\mu)r\phi' + (3\lambda + 2\mu)\phi \} \mathbf{x} \end{aligned}$$

Thus, we need to find a function $\phi : [1, +\infty) \rightarrow \mathbb{R}$ such that

$$\phi'' + \frac{4\phi'}{r} = 0, \quad (\lambda + 2\mu)r\phi' + (3\lambda + 2\mu)\phi|_{r=1} = -1, \quad \phi|_{r=+\infty} = 0.$$

The condition at infinity implies that $B = 0$ and $A = 1/4\mu$. Therefore, the solution is $\mathbf{v}(\mathbf{x}) = \mathbf{x}/4\mu|\mathbf{x}|^3$, which implies that

$$\mathbf{w}(\mathbf{x}) := \mathbf{v}(\mathbf{x}) \Big|_{|\mathbf{x}|=1} = \frac{\mathbf{x}}{4\mu}.$$

With the function \mathbf{w} at hand, we have that

$$\mathbf{I} + \frac{1}{|B_1(\mathbf{0})|} \int_{\partial B_1(\mathbf{0})} \mathbb{C}(\mathbf{w}(\boldsymbol{\zeta}) \otimes \mathbf{n}(\boldsymbol{\zeta})) d\sigma(\boldsymbol{\zeta}) = \mathbf{I} + \frac{3}{16\pi\mu} \int_{\partial B_1(\mathbf{0})} \frac{\mathbb{C}(\boldsymbol{\zeta} \otimes \boldsymbol{\zeta})}{|\boldsymbol{\zeta}|^3} d\sigma(\boldsymbol{\zeta}).$$

Through the use of spherical coordinates and orthogonality relations for the circular functions, it holds

$$\int_{\partial B_1(\mathbf{0})} \frac{\boldsymbol{\zeta} \otimes \boldsymbol{\zeta}}{|\boldsymbol{\zeta}|^3} d\sigma(\boldsymbol{\zeta}) = \frac{4\pi}{3} \mathbf{I},$$

hence the second-order tensor $\mathbb{M}\mathbf{I}$ is given by

$$\mathbb{M}\mathbf{I} = \frac{3(\lambda + 2\mu)}{4\mu} \mathbf{I}.$$

It implies

$$(4.19) \quad u_\varepsilon^k(\mathbf{y}) = \frac{\pi(\lambda + 2\mu)}{\mu} \varepsilon^3 p \operatorname{Tr}(\widehat{\nabla}_{\mathbf{z}} \mathbf{N}^{(k)}(\mathbf{z}, \mathbf{y})) + O(\varepsilon^4), \quad k = 1, 2, 3.$$

For the Neumann's function \mathbf{N} (see the appendix for its explicit expression), we are interested only in the trace of $\nabla_{\mathbf{z}} \mathbf{N}(\mathbf{z}, \mathbf{y})$ computed at $y_3 = 0$.

Evaluating $\mathbf{N} = \mathbf{N}(\mathbf{z}, \mathbf{y})$ at $y_3 = 0$, we get

$$\begin{aligned} \kappa_\mu^{-1} N_{\alpha\alpha} &= -f - (z_\alpha - y_\alpha)^2 f^3 - (1 - 2\nu)g + (1 - 2\nu)(z_\alpha - y_\alpha)^2 f g^2 \\ \kappa_\mu^{-1} N_{\beta\alpha} &= (z_\alpha - y_\alpha)(z_\beta - y_\beta) \{ -f^3 + (1 - 2\nu)fg \} \\ \kappa_\mu^{-1} N_{3\alpha} &= (z_\alpha - y_\alpha) \{ -z_3 f^3 + (1 - 2\nu)fg \} \\ \kappa_\mu^{-1} N_{\alpha 3} &= (z_\alpha - y_\alpha) \{ -z_3 f^3 - (1 - 2\nu)fg \} \\ \kappa_\mu^{-1} N_{33} &= -2(1 - \nu)f - z_3^2 f^3 \end{aligned}$$

where $\alpha, \beta = 1, 2$ and $\kappa_\mu = 1/(4\pi\mu)$, with $f = 1/|\mathbf{z} - \mathbf{y}|$ and $g = 1/\{|\mathbf{z} - \mathbf{y}| - z_3\}$.

Let $\rho^2 := (z_1 - y_1)^2 + (z_2 - y_2)^2$. Using the identities

$$\rho^2 f^2 = 1 - z_3^2 f^2, \quad (1 - z_3 f)g = f$$

and the differentiation formulas

$$\begin{aligned} \partial_{z_\alpha} f &= -(z_\alpha - y_\alpha) f^3, & \partial_{z_3} f &= -z_3 f^3, \\ \partial_{z_\alpha} g &= -(z_\alpha - y_\alpha) f g, & \partial_{z_3} g &= f g, \\ \partial_{z_\alpha} (fg) &= -(z_\alpha - y_\alpha) (f + g) f^2 g, & \partial_{z_3} (fg) &= f^3, \end{aligned}$$

we deduce the following formulas for some of the derivatives of $\kappa_\mu^{-1} N_{ij}$

$$\begin{aligned} \kappa_\mu^{-1} \partial_{z_\alpha} N_{\alpha\alpha} &= (z_\alpha - y_\alpha) \{ -f^3 + 3(z_\alpha - y_\alpha)^2 f^5 \\ &\quad + (1 - 2\nu) [3f - (z_\alpha - y_\alpha)^2 f^2 (f + 2g)] g^2 \} \\ \kappa_\mu^{-1} \partial_{z_\beta} N_{\beta\alpha} &= (z_\alpha - y_\alpha) \{ -f^3 + 3(z_\beta - y_\beta)^2 f^5 \\ &\quad + (1 - 2\nu) [f - (z_\beta - y_\beta)^2 f^2 (f + 2g)] g^2 \} \\ \kappa_\mu^{-1} \partial_{z_3} N_{3\alpha} &= (z_\alpha - y_\alpha) \{ -2\nu f^3 + 3z_3^2 f^5 \} \\ \kappa_\mu^{-1} \partial_{z_\alpha} N_{\alpha 3} &= -z_3 f^3 + 3(z_\alpha - y_\alpha)^2 z_3 f^5 + (1 - 2\nu) [-1 + (z_\alpha - y_\alpha)^2 (f + g) f] f g \\ \kappa_\mu^{-1} \partial_{z_3} N_{33} &= -2\nu z_3 f^3 + 3z_3^3 f^5. \end{aligned}$$

As a consequence, we obtain

$$(4.20) \quad \begin{aligned} \text{Tr}(\widehat{\nabla} \mathbf{N}^{(\alpha)}) &= 2\kappa_\mu (1 - 2\nu) (z_\alpha - y_\alpha) f^3, \quad \text{for } \alpha = 1, 2 \\ \text{Tr}(\widehat{\nabla} \mathbf{N}^{(3)}) &= 2\kappa_\mu (1 - 2\nu) z_3 f^3. \end{aligned}$$

Combining (4.19), (4.20) and using the explicit expression for f , we find

$$\begin{aligned} u_\varepsilon^\alpha(\mathbf{y}) &= \frac{1 - \nu}{\mu} \frac{\varepsilon^3 p(z_\alpha - y_\alpha)}{|\mathbf{z} - \mathbf{y}|^3} + O(\varepsilon^4), \quad \text{for } \alpha = 1, 2 \\ u_\varepsilon^3(\mathbf{y}) &= \frac{1 - \nu}{\mu} \frac{\varepsilon^3 p z_3}{|\mathbf{z} - \mathbf{y}|^3} + O(\varepsilon^4), \end{aligned}$$

that are the components given in (2.1).

APPENDIX. NEUMANN FUNCTION FOR THE HALF-SPACE WITH ZERO TRACTION

For the reader's convenience, we provide here the complete derivation of the explicit formula for the Neumann function \mathbf{N} of the problem

$$\mathcal{L}\mathbf{v} := \text{div}(\mathbb{C}\widehat{\nabla}\mathbf{v}) = \mathbf{b} \quad \text{in } \mathbb{R}_-^3, \quad (\mathbb{C}\widehat{\nabla}\mathbf{v})\mathbf{e}_3 = \mathbf{0} \quad \text{in } \mathbb{R}^2$$

as stated in Theorem 3.1. By definition, \mathbf{N} is the kernel of the integral operator which associates to the forcing term \mathbf{b} the solution \mathbf{v} to the boundary value problem, viz.

$$\mathbf{v}(\mathbf{x}) = \int_{\mathbb{R}_-^3} \mathbf{N}(\mathbf{x}, \mathbf{y}) \mathbf{b}(\mathbf{y}) d\mathbf{y}.$$

The explicit expression for the Neumann function has been determined by Mindlin in [31, 32] using different approaches. Here, we follow the second one which is based on the Papkovitch–Neuber representation of the displacement \mathbf{v} and potential theory.

There exist other mathematical approaches to obtain the Neumann function. See for example [38] for an application of a Fourier method.

Let us introduce the functions

$$\phi_0(\mathbf{x}) := \frac{1}{|\mathbf{x}|} \quad \text{and} \quad \psi_0(\mathbf{x}) := \frac{\phi_0(\mathbf{x})}{1 - x_3\phi_0(\mathbf{x})} = \frac{1}{|\mathbf{x}| - x_3},$$

observing that, apart from $\partial_i\phi_0 = -x_i\phi_0^3$, $i = 1, 2, 3$, the following identities hold true for $\alpha = 1, 2$,

$$\phi_0 - \psi_0 = -x_3\phi_0\psi_0, \quad \partial_\alpha\psi_0 = -x_\alpha\phi_0\psi_0^2, \quad \partial_3\psi_0 = \phi_0\psi_0, \quad \partial_3(\phi_0\psi_0) = \phi_0^3.$$

We denote by ϕ and $\tilde{\phi}$ the values $\phi_0(\mathbf{x} + \mathbf{e}_3)$ and $\phi_0(\mathbf{x} - \mathbf{e}_3)$, respectively, with analogous notation for ψ_0 , that is, ψ and $\tilde{\psi}$ are the values $\psi_0(\mathbf{x} + \mathbf{e}_3)$ and $\psi_0(\mathbf{x} - \mathbf{e}_3)$ respectively.

PROPOSITION A.1. — *Let \mathbf{I} be the identity matrix and δ the Dirac delta concentrated at $-\mathbf{e}_3$. Then, the matrix-valued function $\mathcal{N} = \mathcal{N}(\mathbf{x})$ solution to*

$$\mathcal{L}\mathbf{v} := \operatorname{div}(\mathbb{C}\widehat{\nabla}\mathbf{v}) = \delta\mathbf{I} \quad \text{in } \mathbb{R}_-^3, \quad (\mathbb{C}\widehat{\nabla}\mathbf{v})\mathbf{e}_3 = \mathbf{0} \quad \text{in } \mathbb{R}^2,$$

is given by

$$\begin{aligned} \mathcal{N}_{\alpha\alpha} &= -C_{\mu,\nu} \left\{ (3 - 4\nu)\phi + x_\alpha^2\phi^3 + \tilde{\phi} + [(3 - 4\nu)x_\alpha^2 - 2x_3]\tilde{\phi}^3 \right. \\ &\quad \left. + 6x_\alpha^2x_3\tilde{\phi}^5 + c_\nu(\tilde{\psi} - x_\alpha^2\tilde{\phi}\tilde{\psi}^2) \right\} \\ \mathcal{N}_{\alpha\beta} &= -C_{\mu,\nu}x_\alpha x_\beta \left\{ \phi^3 + (3 - 4\nu)\tilde{\phi}^3 + 6x_3\tilde{\phi}^5 - c_\nu\tilde{\phi}\tilde{\psi}^2 \right\} \\ \text{(A.1)} \quad \mathcal{N}_{3\alpha} &= -C_{\mu,\nu}x_\alpha \left\{ (x_3 + 1)\phi^3 + (3 - 4\nu)(x_3 + 1)\tilde{\phi}^3 + 6x_3(x_3 - 1)\tilde{\phi}^5 - c_\nu\tilde{\phi}\tilde{\psi} \right\} \\ \mathcal{N}_{\alpha 3} &= -C_{\mu,\nu}x_\alpha \left\{ (x_3 + 1)\phi^3 + (3 - 4\nu)(x_3 + 1)\tilde{\phi}^3 - 6x_3(x_3 - 1)\tilde{\phi}^5 + c_\nu\tilde{\phi}\tilde{\psi} \right\}, \\ \mathcal{N}_{33} &= -C_{\mu,\nu} \left\{ (3 - 4\nu)\phi + (x_3 + 1)^2\phi^3 + (1 + c_\nu)\tilde{\phi} \right. \\ &\quad \left. + [(3 - 4\nu)(x_3 - 1)^2 + 2x_3]\tilde{\phi}^3 - 6x_3(x_3 - 1)^2\tilde{\phi}^5 \right\}, \end{aligned}$$

where $C_{\mu,\nu} = 1/\{16\pi\mu(1 - \nu)\}$, $c_\nu := 4(1 - \nu)(1 - 2\nu)$ and $\alpha = 1, 2$.

To establish (A.1), we observe that the columns $\mathcal{N}^{(i)}$ of \mathcal{N} are determined by solving the equation $\mathcal{L}\mathbf{v} = \mathbf{e}_i\delta$ for $i = 1, 2, 3$ and using the Papkovitch–Neuber representation

$$\text{(A.2)} \quad \mathbf{v} = C_{\mu,\nu} \left\{ 4(1 - \nu)\mathbf{h} - \nabla(\mathbf{x} \cdot \mathbf{h} + \beta) \right\} \quad \text{with} \quad \begin{cases} \Delta\mathbf{h} = 4\pi\mathbf{e}_i\delta \\ \Delta\beta = 4\pi\delta_{i3}\delta. \end{cases}$$

where δ_{ij} is the Kronecker symbol. The coupling between \mathbf{h} and β is determined by the boundary conditions on the plane $\{x_3 = 0\}$, which are

$$\text{(A.3)} \quad \begin{aligned} (1 - 2\nu)(\partial_3h_\alpha + \partial_\alpha h_3) - \mathbf{x} \cdot \partial_{\alpha 3}^2\mathbf{h} - \partial_{\alpha 3}^2\beta &= 0, \quad (\alpha = 1, 2), \\ 2\nu \operatorname{div} \mathbf{h} + 2(1 - 2\nu)\partial_3h_3 - \mathbf{x} \cdot \partial_{33}^2\mathbf{h} - \partial_{33}^2\beta &= 0, \end{aligned} \quad \text{for } x_3 = 0.$$

Given $\mathbf{y} := (y_1, y_2, y_3)$, let $\tilde{\mathbf{y}}$ be the reflexed point. Set

$$G(\mathbf{x}, \mathbf{y}) := -\phi_0(\mathbf{x} - \mathbf{y}) + \phi_0(\mathbf{x} - \tilde{\mathbf{y}}).$$

Denoting by $\langle f, g \rangle$ the action of the distribution f on the function g , we determine \mathbf{h} and β taking advantage of the relation (which is deduced from the second Green identity)

$$(A.4) \quad F(\mathbf{x}) = \frac{1}{4\pi} \langle \Delta F, G(\mathbf{x}, \cdot) \rangle,$$

applied to different choices of F .

Proof of Proposition A.1. — To determine \mathcal{N} , we consider separately the case of horizontal and vertical forcing. By symmetry, x_1 and x_2 can be interchanged.

Horizontal force: $\mathcal{L}\mathbf{v} = \mathbf{e}_1\delta$. — We choose $h_2 = 0$, so that boundary conditions become

$$\begin{cases} (1 - 2\nu)(\partial_3 h_1 + \partial_1 h_3) - x_1 \partial_{13}^2 h_1 - \partial_{13}^2 \beta = 0, \\ (1 - 2\nu)\partial_2 h_3 - x_1 \partial_{23}^2 h_1 - \partial_{23}^2 \beta = 0, & \text{for } x_3 = 0. \\ 2\nu \partial_1 h_1 + 2(1 - \nu) \partial_3 h_3 - x_1 \partial_{33}^2 h_1 - \partial_{33}^2 \beta = 0, \end{cases}$$

Differentiating the first equation with respect to x_1 , the second with respect to x_2 and taking the difference, we obtain

$$0 = (1 - 2\nu)\partial_{23}^2 h_1 + \partial_{23}^2 h_1 = 2(1 - \nu)\partial_{23}^2 h_1 \quad \text{for } x_3 = 0,$$

which suggests, after integration with respect to x_2 , the choice $F := \partial_3 h_1$. Applying (A.4),

$$\partial_3 h_1 = -\partial_{y_3} G \Big|_{\mathbf{y}=-\mathbf{e}_3} = -\partial_3(\phi + \tilde{\phi}) \quad \text{for } x_3 < 0,$$

and thus $h_1 = -(\phi + \tilde{\phi})$.

Being $\partial_3 h_1$ null for $x_3 = 0$, integration of the second boundary condition encourages the choice $F := (1 - 2\nu)h_3 - \partial_3 \beta$ which is zero for $x_3 = 0$. Hence, since $\Delta F = 0$, we deduce

$$(A.5) \quad (1 - 2\nu)h_3 - \partial_3 \beta = 0 \quad \text{for } x_3 < 0.$$

Concerning the third boundary condition, we observe that

$$\begin{aligned} \partial_1 h_1 &= x_1(\phi^3 + \tilde{\phi}^3) = 2x_1 \tilde{\phi}^3 = -2\partial_1 \tilde{\phi} \\ x_1 \partial_{33}^2 h_1 &= x_1(\phi^3 + \tilde{\phi}^3 - 3\phi^5 - 3\tilde{\phi}^5) = 2x_1(\tilde{\phi}^3 - 3\tilde{\phi}^5) = -2(\partial_1 \tilde{\phi} - \partial_{13}^2 \tilde{\phi}) \end{aligned} \quad \text{for } x_3 = 0,$$

since ϕ and $\tilde{\phi}$ coincide when $x_3 = 0$. Substituting in the third boundary condition, we obtain

$$F := 2(1 - \nu)\partial_3 h_3 - \partial_{33}^2 \beta + 2(1 - 2\nu)\partial_1 \tilde{\phi} - 2\partial_{13}^2 \tilde{\phi} = 0 \quad \text{for } x_3 = 0.$$

Since $\Delta F = 0$, we infer

$$2(1 - \nu)\partial_3 h_3 - \partial_{33}^2 \beta + 2(1 - 2\nu)\partial_1 \tilde{\phi} - 2\partial_{13}^2 \tilde{\phi} = 0 \quad \text{for } x_3 < 0,$$

and thus, being $\partial_1 \tilde{\phi} = -x_1 \tilde{\phi}^3 = -\partial_3(x_1 \tilde{\phi} \tilde{\psi})$,

$$2(1 - \nu)h_3 - \partial_3 \beta = -2x_1 \tilde{\phi}^3 + 2(1 - 2\nu)x_1 \tilde{\phi} \tilde{\psi} \quad \text{for } x_3 < 0.$$

Coupling with (A.5), we deduce

$$\begin{cases} h_3 = -2x_1\tilde{\phi}^3 + 2(1-2\nu)x_1\tilde{\phi}\tilde{\psi} \\ \partial_3\beta = -2(1-2\nu)x_1\tilde{\phi}^3 + 2(1-2\nu)^2x_1\tilde{\phi}\tilde{\psi} \end{cases} \quad \text{for } x_3 < 0.$$

Recalling that $\tilde{\phi}^3 = \partial_3(\tilde{\phi}\tilde{\psi})$ and $\tilde{\phi}\tilde{\psi} = \partial_3\tilde{\psi}$, by integration,

$$\beta = -2(1-2\nu)x_1\tilde{\phi}\tilde{\psi} + 2(1-2\nu)^2x_1\tilde{\psi} \quad \text{for } x_3 < 0.$$

Using the identity $(x_3-1)\tilde{\phi}\tilde{\psi} = \tilde{\psi} - \tilde{\phi}$, we infer

$$x_3h_3 + \beta = x_1\{-2(1-2\nu)\tilde{\phi} - 2x_3\tilde{\phi}^3 + c_\nu\tilde{\psi}\}.$$

Substituting in (A.2), we get the expressions for \mathcal{N}_{i1} given in (A.1).

Vertical force: $\mathcal{L}\mathbf{v} = \mathbf{e}_3\delta$. — Choosing $h_1 = h_2 = 0$, conditions (A.3) become

$$\begin{cases} (1-2\nu)\partial_\alpha h_3 - \partial_{\alpha 3}^2\beta = 0 & (\alpha = 1, 2), \\ 2(1-\nu)\partial_3 h_3 - \partial_{33}^2\beta = 0 \end{cases} \quad \text{for } x_3 = 0.$$

Integrating the first relation with respect to x_α , we obtain

$$\begin{cases} (1-2\nu)h_3 - \partial_3\beta = 0 \\ 2(1-\nu)\partial_3 h_3 - \partial_{33}^2\beta = 0 \end{cases} \quad \text{for } x_3 = 0.$$

Since $\Delta h_3 = \Delta\beta = \delta$, identity (A.4) with $F := (1-2\nu)h_3 - \partial_3\beta$ gives

$$(A.6) \quad \begin{aligned} (1-2\nu)h_3 - \partial_3\beta &= \{(1-2\nu)G + \partial_{y_3}G\}|_{\mathbf{y}=-\mathbf{e}_3} \\ &= (1-2\nu)(-\phi + \tilde{\phi}) - (x_3+1)\phi^3 - (x_3-1)\tilde{\phi}^3 \quad \text{for } x_3 < 0. \end{aligned}$$

Applying (A.4) to $F := 2(1-\nu)\partial_3 h_3 - \partial_{33}^2\beta$, we deduce

$$\begin{aligned} 2(1-\nu)\partial_3 h_3 - \partial_{33}^2\beta &= \{-2(1-\nu)\partial_{y_3}G - \partial_{y_3 y_3}^2 G\}|_{\mathbf{y}=-\mathbf{e}_3} \\ &= \partial_3\{-2(1-\nu)(\phi + \tilde{\phi}) + \partial_3(\phi - \tilde{\phi})\} \quad \text{for } x_3 < 0. \end{aligned}$$

Integrating with respect to x_3 , we infer

$$2(1-\nu)h_3 - \partial_3\beta = -2(1-\nu)(\phi + \tilde{\phi}) - (x_3+1)\phi^3 + (x_3-1)\tilde{\phi}^3 \quad \text{for } x_3 < 0.$$

Coupling with (A.6), we get explicit expressions for h_3 and $\partial_3\beta$, namely

$$\begin{cases} h_3 = -\phi - (3-4\nu)\tilde{\phi} + 2(x_3-1)\tilde{\phi}^3 \\ \partial_3\beta = (x_3+1)\phi^3 - c_\nu\tilde{\phi} + (3-4\nu)(x_3-1)\tilde{\phi}^3 \end{cases} \quad \text{for } x_3 < 0.$$

Differentiation of $\partial_3\beta$ with respect to x_α gives

$$\begin{aligned} \partial_{3\alpha}^2\beta &= -3x_\alpha(x_3+1)\phi^5 + c_\nu x_\alpha\tilde{\phi}^3 - 3(3-4\nu)x_\alpha(x_3-1)\tilde{\phi}^5 \\ &= \partial_3\{x_\alpha\phi^3 + c_\nu x_\alpha\tilde{\phi}\tilde{\psi} + (3-4\nu)x_\alpha\tilde{\phi}^3\} \end{aligned}$$

and thus

$$\partial_\alpha\beta = x_\alpha\{\phi^3 + c_\nu\tilde{\phi}\tilde{\psi} + (3-4\nu)\tilde{\phi}^3\} \quad \text{for } x_3 < 0.$$

Recalling identity (A.2), we deduce the corresponding expressions for \mathcal{N}_{i3} in (A.1). \square

The fundamental solution $\mathbf{N} = \mathbf{N}(\mathbf{x}, \mathbf{y})$ in the half-space $\{x_3 < 0\}$ is such that its columns $\mathbf{v}_1, \mathbf{v}_2$ and \mathbf{v}_3 solve $\mathcal{L}\mathbf{v}_i = \delta_{\mathbf{y}}\mathbf{e}_i$ where $\delta_{\mathbf{y}}$ is the Dirac delta concentrated at $\mathbf{y} = (y_1, y_2, y_3)$ with $y_3 < 0$. Thus, the Neumann function \mathbf{N} is given by

$$(A.7) \quad \mathbf{N}(\mathbf{x}, \mathbf{y}) = \frac{1}{|y_3|} \mathcal{N} \left(\frac{x_1 - y_1}{|y_3|}, \frac{x_2 - y_2}{|y_3|}, \frac{x_3}{|y_3|} \right)$$

as a result of the homogeneity of δ and the second order degree of \mathcal{L} .

Recalling the definitions of $\phi, \tilde{\phi}, \tilde{\psi}$ and computing at $(x_1 - y_1, x_2 - y_2, x_3)/|y_3|$, we obtain the identities

$$f := -\frac{\phi}{y_3} = \frac{1}{|\mathbf{x} - \mathbf{y}|}, \quad \tilde{f} := -\frac{\tilde{\phi}}{y_3} = \frac{1}{|\mathbf{x} - \tilde{\mathbf{y}}|}, \quad \tilde{g} := -\frac{\tilde{\psi}}{y_3} = \frac{1}{|\mathbf{x} - \tilde{\mathbf{y}}| - x_3 - y_3}.$$

where $\tilde{\mathbf{y}} = (y_1, y_2, -y_3)$. Hence, the components of $C_{\mu, \nu}^{-1}\mathbf{N}$ are given by

$$\begin{aligned} C_{\mu, \nu}^{-1}N_{\alpha\alpha} &= -(3 - 4\nu)f - (x_\alpha - y_\alpha)^2 f^3 - \tilde{f} - (3 - 4\nu)(x_\alpha - y_\alpha)^2 \tilde{f}^3 - c_\nu \tilde{g} \\ &\quad + c_\nu (x_\alpha - y_\alpha)^2 \tilde{f} \tilde{g}^2 - 2x_3 y_3 \tilde{f}^3 + 6(x_\alpha - y_\alpha)^2 x_3 y_3 \tilde{f}^5 \\ C_{\mu, \nu}^{-1}N_{\alpha\beta} &= (x_\alpha - y_\alpha)(x_\beta - y_\beta) \{-f^3 - (3 - 4\nu)\tilde{f}^3 + c_\nu \tilde{f} \tilde{g}^2 + 6x_3 y_3 \tilde{f}^5\} \\ C_{\mu, \nu}^{-1}N_{3\alpha} &= (x_\alpha - y_\alpha) \{-(x_3 - y_3)f^3 - (3 - 4\nu)(x_3 - y_3)\tilde{f}^3 + c_\nu \tilde{f} \tilde{g} \\ &\quad + 6x_3 y_3 (x_3 + y_3)\tilde{f}^5\} \\ C_{\mu, \nu}^{-1}N_{\alpha 3} &= (x_\alpha - y_\alpha) \{-(x_3 - y_3)f^3 - (3 - 4\nu)(x_3 - y_3)\tilde{f}^3 - c_\nu \tilde{f} \tilde{g} \\ &\quad - 6x_3 y_3 (x_3 + y_3)\tilde{f}^5\} \\ C_{\mu, \nu}^{-1}N_{33} &= -(3 - 4\nu)f - (x_3 - y_3)^2 f^3 - (1 + c_\nu)\tilde{f} - (3 - 4\nu)(x_3 + y_3)^2 \tilde{f}^3 \\ &\quad + 2x_3 y_3 \tilde{f}^3 - 6x_3 y_3 (x_3 + y_3)^2 \tilde{f}^5. \end{aligned}$$

Recollecting the expression for fundamental solution $\mathbf{\Gamma}$ in the whole space and using the relation $\tilde{f} = \tilde{g} - (x_3 + y_3)\tilde{f}\tilde{g}$, the above formulas can be rewritten as $\mathbf{N} = \mathbf{\Gamma} + \mathbf{R}$, where $\mathbf{\Gamma}$ is computed at $\mathbf{x} - \mathbf{y}$ and the component R_{ij} of \mathbf{R} are given by

$$\begin{aligned} R_{\alpha\alpha} &= C_{\mu, \nu} \{-(\tilde{f} + c_\nu \tilde{g}) - (3 - 4\nu)\eta_\alpha^2 \tilde{f}^3 + c_\nu \eta_\alpha^2 \tilde{f} \tilde{g}^2 - 2x_3 y_3 (\tilde{f}^3 - 3\eta_\alpha^2 \tilde{f}^5)\} \\ R_{\beta\alpha} &= C_{\mu, \nu} \eta_\alpha \eta_\beta \{-(3 - 4\nu)\tilde{f}^3 + c_\nu \tilde{f} \tilde{g}^2 + 6x_3 y_3 \tilde{f}^5\} \\ R_{3\alpha} &= C_{\mu, \nu} \eta_\alpha \{-(3 - 4\nu)(\eta_3 - 2y_3)\tilde{f}^3 + c_\nu \tilde{f} \tilde{g} + 6x_3 y_3 \eta_3 \tilde{f}^5\} \\ R_{\alpha 3} &= C_{\mu, \nu} \eta_\alpha \{-(3 - 4\nu)(\eta_3 - 2y_3)\tilde{f}^3 - c_\nu \tilde{f} \tilde{g} - 6x_3 y_3 \eta_3 \tilde{f}^5\} \\ R_{33} &= C_{\mu, \nu} \{-(\tilde{f} + c_\nu \tilde{g}) - (3 - 4\nu)\eta_3^2 \tilde{f}^3 + c_\nu \eta_3 \tilde{f} \tilde{g} + 2x_3 y_3 (\tilde{f}^3 - 3\eta_3^2 \tilde{f}^5)\}, \end{aligned}$$

where $\eta_\alpha = x_\alpha - y_\alpha$ for $\alpha = 1, 2$ and $\eta_3 = x_3 + y_3$, which can be recombined as

$$\begin{aligned} R_{ij} &= C_{\mu, \nu} \{-(\tilde{f} + c_\nu \tilde{g})\delta_{ij} - (3 - 4\nu)\eta_i \eta_j \tilde{f}^3 + 2(3 - 4\nu)y_3 [\delta_{3i}(1 - \delta_{3j})\eta_j + \delta_{3j}(1 - \delta_{3i})\eta_i] \tilde{f}^3 \\ &\quad + c_\nu [\delta_{i3}\eta_j - \delta_{3j}(1 - \delta_{3i})\eta_i] \tilde{f} \tilde{g} + c_\nu (1 - \delta_{3j})(1 - \delta_{3i})\eta_i \eta_j \tilde{f} \tilde{g}^2 \\ &\quad - 2(1 - 2\delta_{3j})x_3 y_3 (\delta_{ij} \tilde{f}^3 - 3\eta_i \eta_j \tilde{f}^5)\} \end{aligned}$$

for $i, j = 1, 2, 3$. Since $x_3 = \eta_3 - y_3$, we obtain the decomposition $R_{ij} := R_{ij}^1 + R_{ij}^2 + R_{ij}^3$, where

$$\begin{aligned} R_{ij}^1 &:= C_{\mu,\nu} \left\{ -(\tilde{f} + c_\nu \tilde{g}) \delta_{ij} - (3 - 4\nu) \eta_i \eta_j \tilde{f}^3 \right. \\ &\quad \left. + c_\nu [\delta_{3i} \eta_j - \delta_{3j} (1 - \delta_{3i}) \eta_i] \tilde{f} \tilde{g} + c_\nu (1 - \delta_{3j}) (1 - \delta_{3i}) \eta_i \eta_j \tilde{f} \tilde{g}^2 \right\} \\ R_{ij}^2 &:= 2C_{\mu,\nu} y_3 \left\{ (3 - 4\nu) [\delta_{3i} (1 - \delta_{3j}) \eta_j + \delta_{3j} (1 - \delta_{3i}) \eta_i] \tilde{f}^3 - (1 - 2\delta_{3j}) \delta_{ij} \eta_3 \tilde{f}^3 \right. \\ &\quad \left. + 3(1 - 2\delta_{3j}) \eta_i \eta_j \eta_3 \tilde{f}^5 \right\} \\ R_{ij}^3 &:= 2C_{\mu,\nu} (1 - 2\delta_{3j}) y_3^2 \left\{ \delta_{ij} \tilde{f}^3 - 3\eta_i \eta_j \tilde{f}^5 \right\}. \end{aligned}$$

The proof of Theorem 3.1 is complete.

REFERENCES

- [1] Y. A. ABRAMOVICH & C. D. ALIPRANTIS – *An invitation to operator theory*, Graduate Studies in Mathematics, vol. 50, American Mathematical Society, Providence, RI, 2002.
- [2] G. ALESSANDRINI, E. BERETTA, E. ROSSET & S. VESSELLA – “Optimal stability for inverse elliptic boundary value problems with unknown boundaries”, *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)* **29** (2000), no. 4, p. 755–806.
- [3] G. ALESSANDRINI, M. DI CRISTO, A. MORASSI & E. ROSSET – “Stable determination of an inclusion in an elastic body by boundary measurements”, *SIAM J. Math. Anal.* **46** (2014), no. 4, p. 2692–2729.
- [4] H. AMMARI, E. BRETIN, J. GARNIER, H. KANG, H. LEE & A. WAHAB – *Mathematical methods in elasticity imaging*, Princeton Series in Applied Mathematics, Princeton University Press, Princeton, NJ, 2015.
- [5] H. AMMARI & H. KANG – *Reconstruction of small inhomogeneities from boundary measurements*, Lect. Notes in Math., vol. 1846, Springer-Verlag, Berlin, 2004.
- [6] ———, *Polarization and moment tensors. with applications to inverse problems and effective medium theory*, Applied Mathematical Sciences, vol. 162, Springer, New York, 2007.
- [7] C. AMROUCHE, M. DAMBRINE & Y. RAUDIN – “An L^p theory of linear elasticity in the half-space”, *J. Differential Equations* **253** (2012), no. 3, p. 906–932.
- [8] A. ASPRI, E. BERETTA & C. MASCIÀ – “Asymptotic expansion for harmonic functions in the half-space with a pressurized cavity”, *Math. Methods Appl. Sci.* **39** (2016), no. 10, p. 2415–2430.
- [9] M. BATTAGLIA, J. GOTSMANN, D. CARBONE & J. FERNÁNDEZ – “4D volcano gravimetry”, *Geophysics* **73** (2008), no. 6, p. 3–18.
- [10] M. BATTAGLIA & D. P. HILL – “Analytical modeling of gravity changes and crustal deformation at volcanoes: the Long Valley caldera, California, case study”, *Tectonophysics* **471** (2009), p. 45–57.
- [11] D. J. CEDIO-FENGYA, S. MOSKOW & M. S. VOGELIUS – “Identification of conductivity imperfections of small diameter by boundary measurements. Continuous dependence and computational reconstruction”, *Inverse Problems* **14** (1998), no. 3, p. 553–595.
- [12] P. G. CIARLET – *Mathematical elasticity. Vol. I. Three-dimensional elasticity*, Studies in Mathematics and its Applications, vol. 20, North-Holland Publishing Co., Amsterdam, 1988.
- [13] G. CURRENTI, A. BONACCORSO, C. DEL NEGRO, D. SCANDURA & E. BOSCHI – “Elasto-plastic modeling of volcano ground deformation”, *Earth Planetary Science Letters* **296** (2010), p. 311–318.
- [14] G. CURRENTI, C. DEL NEGRO & G. GANCI – “Finite element modeling of ground deformation and gravity field at Mt. Etna”, *Annals Geophysics* **51** (2008), no. 1, p. 105–119.
- [15] B. E. J. DAHLBERG, C. E. KENIG & G. C. VERCHOTA – “Boundary value problems for the systems of elastostatics in Lipschitz domains”, *Duke Math. J.* **57** (1988), no. 3, p. 795–818.
- [16] P. M. DAVIS – “Surface deformation due to inflation of an arbitrarily oriented triaxial ellipsoidal cavity in an elastic half-space, with reference to Kilauea Volcano, Hawaii”, *J. Geophysical Research* **91** (1986), no. B7, p. 7429–7438.
- [17] G. DE NATALE & F. PINGUE – “Ground deformation modeling in volcanic areas”, in *Monitoring and Mitigation of Volcano Hazards* (R. Scarpa & R. I. Tilling, eds.), Springer, 1996, p. 365–388.

- [18] D. DZURISIN – *Volcano deformation. geodetic monitoring techniques*, Springer Praxis Books, Springer, 2006.
- [19] E. B. FABES, C. E. KENIG & G. C. VERCHOTA – “The Dirichlet problem for the Stokes system on Lipschitz domains”, *Duke Math. J.* **57** (1988), no. 3, p. 769–793.
- [20] Y. FIALKO, Y. KHAZAN & M. SIMONS – “Deformation due to a pressurized horizontal circular crack in an elastic half-space, with applications to volcano geodesy”, *Geophys. J. Int.* **146** (2001), no. 1, p. 181–190.
- [21] G. FICHERA – “Sull’esistenza e sul calcolo delle soluzioni dei problemi al contorno, relativi all’equilibrio di un corpo elastico”, *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (3)* **4** (1950), no. 248, p. 35–99.
- [22] A. FRIEDMAN & M. VOGELIUS – “Identification of small inhomogeneities of extreme conductivity by boundary measurements: a theorem on continuous dependence”, *Arch. Rational Mech. Anal.* **105** (1989), no. 4, p. 299–326.
- [23] M. E. GURTIN – *The linear theory of elasticity, encyclopedia of physics, Vol. VI a/2*, Springer-Verlag, 1972.
- [24] B. B. GUZINA & M. BONNET – “Topological derivative for the inverse scattering of elastic waves”, *Quart. J. Mech. Appl. Math.* **57** (2004), no. 2, p. 161–179.
- [25] T. KATO – *Perturbation theory for linear operators*, Springer-Verlag, Berlin, 1995.
- [26] R. KRESS – *Linear integral equations*, Applied Mathematical Sciences, vol. 82, Springer-Verlag, Berlin, 1989.
- [27] V. D. KUPRADZE – *Potential methods in the theory of elasticity*, Israel Program for Scientific Translations, Jerusalem; Daniel Davey & Co., Inc., New York, 1965.
- [28] M. LISOWSKI – “Analytical volcano deformation source models”, in *Volcano Deformation. Geodetic Monitoring Techniques* (D. Dzurisin, ed.), Springer Praxis Books, Springer, 2006, p. 279–304.
- [29] A. MANCONI, T. R. WALTER & F. AMELUNG – “Effects of mechanical layering on volcano deformation”, *Geophys. J. Int.* **170** (2007), no. 2, p. 952–958.
- [30] D. F. MCTIGUE – “Elastic stress and deformation near a finite spherical magma body: resolution of the point source paradox”, *Journal Geophysical Research* **92** (1987), no. 12, p. 931–940.
- [31] R. D. MINDLIN – “Force at a point in the interior of a semiinfinite solid”, *Journal of Applied Physics* **7** (1936), p. 195–202.
- [32] ———, “Force at a point in the interior of a semi-infinite solid”, in *Proceedings of The First Midwestern Conference on Solid Mechanics (University of Illinois, Urbana, Ill.)*, 1954.
- [33] K. MOGI – “Relations between the eruptions of various volcanoes and the deformation of the ground surfaces around them”, *Bull. Earthq. Res. Inst. U. Tokyo* **36** (1958), p. 99–134.
- [34] A. MORASSI & E. ROSSET – “Stable determination of cavities in elastic bodies”, *Inverse Problems* **20** (2004), no. 2, p. 453–480.
- [35] Y. OKADA – “Surface deformation due to shear and tensile faults in a half-space”, *Bull. Seismol. Soc. Am.* **75** (1985), p. 1135–1154.
- [36] P. SEGALL – *Earthquake and volcano deformation*, Princeton University Press, Princeton, NJ, 2010.
- [37] ———, “Volcano deformation and eruption forecasting”, *Geological Society Special Publication* **380** (2013), no. 1, p. 85–106.
- [38] D. VOLKOV – “A double layer surface traction free Green’s tensor”, *SIAM J. Appl. Math.* **69** (2009), no. 5, p. 1438–1456.
- [39] C. A. WILLIAMS & G. WADGE – “The effects of topography on magma chamber deformation models: application to Mt. Etna and radar interferometry”, *Geophysical Research Letters* **25** (1998), no. 10, p. 1549–1552.
- [40] X.-M. YANG & P. M. DAVIS – “Deformation from inflation of a dipping finite prolate spheroid in an elastic half-space as a model for volcanic stressing”, *Journal Geophysical Research* **93** (1988), no. B5, p. 4249–4257.
- [41] K. YOSIDA – *Functional analysis*, sixth ed., Grundlehren Math. Wiss., vol. 123, Springer-Verlag, Berlin-New York, 1980.

Manuscript received August 30, 2016
accepted January 26, 2017

ANDREA ASPRI, Dipartimento di Matematica “G. Castelnuovo”, Sapienza – Università di Roma,
P.le Aldo Moro, 2 - 00185 Roma, Italy
E-mail : aspri@mat.uniroma1.it

ELENA BERETTA, Dipartimento di Matematica, Politecnico di Milano,
Via Edoardo Bonardi 9 - 20133 Milano, Italy
E-mail : elena.beretta@polimi.it

CORRADO MASCIÀ, Dipartimento di Matematica “G. Castelnuovo”, Sapienza – Università di Roma,
P.le Aldo Moro, 2 - 00185 Roma, Italy
E-mail : mascia@mat.uniroma1.it