

Pathogenicity Factors in Group C and G Streptococci

CLAIRE E. TURNER,¹ LAURA BUBBA,^{2,3} and ANDROULLA EFSTRATIOU²

¹Department of Molecular Biology & Biotechnology, The Florey Institute, University of Sheffield, Sheffield, UK;

²Reference Microbiology Division, National Infection Service, Public Health England, London, United Kingdom;

³European Programme for Public Health Microbiology Training (EUPHEM), European Centre for Disease Prevention and Control (ECDC), Stockholm, Sweden

ABSTRACT Initially recognized zoonoses, streptococci belonging to Lancefield group C (GCS) and G (GGS) were subsequently recognised as human pathogens causing a diverse range of symptoms, from asymptomatic carriage to life threatening diseases. Their taxonomy has changed during the last decade. Asymptomatic carriage is <4% amongst the human population and invasive infections are often in association with chronic diseases such as diabetes, cardiovascular diseases or chronic skin infections. Other clinical manifestations include acute pharyngitis, pneumonia, endocarditis, bacteraemia and toxic-shock syndrome. Post streptococcal sequelae such as rheumatic fever and acute glomerulonephritis have also been described but mainly in developed countries and amongst specific populations. Putative virulence determinants for these organisms include adhesins, toxins, and other factors that are essential for dissemination in human tissues and for interference with the host immune responses. High nucleotide similarities among virulence genes and their association with mobile genetic elements supports the hypothesis of extensive horizontal gene transfer events between the various pyogenic streptococcal species belonging to Lancefield groups A, C and G. A better understanding of the mechanisms of pathogenesis should be apparent by whole-genome sequencing, and this would result in more effective clinical strategies for the pyogenic group in general.

INTRODUCTION

The pyogenic streptococci of Lancefield groups C and G were initially recognized as a cause of animal infections long before they were even considered as agents of human disease, even though they are widely distributed in animals and humans. They comprise a heterogeneous complex of streptococcal species that act as causative agents of a spectrum of diseases ranging from mild pharyngitis to skin infection to life-threatening systemic infections associated with high mortality rates. In

this article we provide an overview of the various group C and group G streptococcal species, the diseases they cause, and the major pathogenicity factors that contribute to their virulence ([Table 1](#)).

TAXONOMY AND IDENTIFICATION

Taxonomic classification of group C streptococci (GCS) and group G streptococci (GGS) has always proven to be a complex issue. However, extensive taxonomic studies over the past few years have distinguished most of the veterinary pathogens belonging to Lancefield groups C and G from the human pathogens. Previously, GCS and GGS were divided into the following species: *Strepto-*

Received: 2 February 2018, **Accepted:** 15 November 2018,

Published: 17 May 2019

Editors: Vincent A. Fischetti, The Rockefeller University, New York, NY; Richard P. Novick, Skirball Institute for Molecular Medicine, NYU Medical Center, New York, NY; Joseph J. Ferretti, Department of Microbiology & Immunology, University of Oklahoma Health Science Center, Oklahoma City, OK; Daniel A. Portnoy, Department of Molecular and Cellular Microbiology, University of California, Berkeley, Berkeley, CA; Miriam Braunstein, Department of Microbiology and Immunology, University of North Carolina-Chapel Hill, Chapel Hill, NC, and Julian I. Rood, Infection and Immunity Program, Monash Biomedicine Discovery Institute, Monash University, Melbourne, Australia

Citation: Turner CE, Bubba L, Efstratiou A. 2019. Pathogenicity factors in group C and G Streptococci. *Microbiol Spectrum* 7(3): GPP3-0020-2018. doi:10.1128/microbiolspec.GPP3-0020-2018.

Correspondence: Androulla Efstratiou. Androulla.efstratiou@phe.gov.uk

This chapter is dedicated to the memory of Gursharan Singh Chhatwal, an inspiring "streptococcologist" who leaves us with wonderful memories of Lancefield 2002 held in Goa, India, and many others.

© 2019 American Society for Microbiology. All rights reserved.

TABLE 1 Pathogenicity factors of group C and G streptococci

Pathogenicity factors	Organism	Reference(s)
Fibronectin binding proteins		
FnbA	<i>S. dysgalactiae</i>	59
FnbB	<i>S. dysgalactiae</i>	60
GfbA	<i>S. dysgalactiae</i>	64
FNZ/FNE	<i>S. zooepidemicus</i> , <i>S. equi</i>	61 , 65
FNZ2/FNEB	<i>S. zooepidemicus</i> , <i>S. equi</i>	66 , 69
SFS	<i>S. zooepidemicus</i> , <i>S. equi</i>	70
M-like proteins		
	<i>S. dysgalactiae</i>	91 , 105
	<i>S. equisimilis</i>	
FOG	<i>S. dysgalactiae</i>	93 , 94
DemA	<i>S. dysgalactiae</i>	105
SzM/SeM	<i>S. zooepidemicus</i> , <i>S. equi</i>	76 , 95–97 , 101
SzPSe/SzP	<i>S. zooepidemicus</i> , <i>S. equi</i>	102
Se18.9	<i>S. equi</i>	76 , 100
ScM	<i>S. canis</i>	104
Others		
C5a peptidase	<i>S. dysgalactiae</i>	108
SeCEP/SzoCEP	<i>S. zooepidemicus</i> , <i>S. equi</i>	112
Immunoglobulin binding proteins		
Protein G	<i>S. dysgalactiae</i>	116
MIG	<i>S. dysgalactiae</i>	120
MAG	<i>S. dysgalactiae</i>	122
ZAG	<i>S. zooepidemicus</i>	123
IgG-endopeptidases		
IdeE/IdeE2	<i>S. equi</i> , <i>S. zooepidemicus</i>	129
IdeE2/IdeZ2	<i>S. equi</i> , <i>S. zooepidemicus</i>	130
IdeP	<i>S. phocae</i>	133
Toxins		
Streptokinase	<i>S. dysgalactiae</i> , <i>S. equisimilis</i>	134
Streptolysin O	<i>S. dysgalactiae</i> , <i>S. equisimilis</i>	145
Streptolysin S	<i>S. dysgalactiae</i> , <i>S. equisimilis</i>	152 , 154
	<i>S. equi</i> , <i>S. zooepidemicus</i>	
Superantigens		
SpeG ^{dys} /SpeGg/SpeG	<i>S. equisimilis</i> , <i>S. canis</i>	158 , 159
SpeH	<i>S. equi</i>	167
SpeI	<i>S. equi</i>	167
SpeK	<i>S. equi</i> , <i>S. zooepidemicus</i> , <i>S. equisimilis</i>	158 , 168 , 169
SpeL	<i>S. equi</i> , <i>S. zooepidemicus</i> , <i>S. equisimilis</i>	158 , 168 , 169
SpeM	<i>S. equisimilis</i>	173
ssa	<i>S. equisimilis</i>	173
SDM/SpeM	<i>S. dysgalactiae</i>	171
SpeA	<i>S. equisimilis</i>	170
SpeC	<i>S. equisimilis</i>	170
SzeN	<i>S. zooepidemicus</i>	170
SzeP	<i>S. zooepidemicus</i>	170
SzeF	<i>S. zooepidemicus</i>	170

coccus dysgalactiae subsp. *equisimilis* (only GCS, human pathogen), *Streptococcus dysgalactiae* subsp. *dysgalactiae* (GCS, animal pathogen), *Streptococcus equi* (animal GCS), *Streptococcus zooepidemicus* (animal and human GCS), *Streptococcus canis* (animal GGS), the *Streptococcus anginosus* group, and *Streptococcus phocae* ([1](#)). GCS and GGS of human origin are now considered to constitute a single subspecies, *S. dysgalactiae* subsp. *equisimilis*. Therefore, the current taxonomy characterizes the

species as follows: *S. dysgalactiae* is divided into the subspecies *S. dysgalactiae* subsp. *equisimilis* and *S. dysgalactiae* subsp. *dysgalactiae* (hereafter referred to in this article as *S. equisimilis* and *S. dysgalactiae*). *S. equi* is divided into the subspecies *S. equi* subsp. *equi*, *S. equi* subsp. *zooepidemicus*, and *S. equi* subsp. *ruminatorum* (hereafter referred to as *S. equi*, *S. zooepidemicus*, and *S. ruminatorum*). However, it has been suggested that *S. equi* may simply be regarded as a subclone of *S. zooepidemicus* ([2](#)). On the basis of genetic evidence, *S. dysgalactiae*, *S. equi*, and *S. canis* are more closely related to each other than to the *S. anginosus* group and constitute species with the “large-colony” colony phenotype. *S. phocae* is a new species expressing the group C antigen thus far only isolated from seals ([3](#), [4](#)). Some of these species may also contain strains which express the Lancefield group A or group F antigens ([Table 2](#)).

HUMAN DISEASE AND DIAGNOSIS

GCS and GGS can cause a wide range of diseases from mild to severe, among both human and animal populations. GCS and GGS generally colonize the human respiratory, gastrointestinal, and genitourinary tracts, with an estimated <4% asymptomatic pharyngeal carriage rate in adults ([5–7](#)). Human invasive infections caused by GCS and GGS have been associated with underlying conditions, such as diabetes, cardiovascular diseases, and chronic skin conditions ([8–10](#)). While *S. dysgalactiae* is considered an animal pathogen, *S. equisimilis* is almost exclusively a human pathogen, with increasing prevalence and overlaps in clinical manifestations with group A *Streptococcus* (GAS). *S. equisimilis* infections include acute pharyngitis, pneumonia, endocarditis, cellulitis, peritonitis, septic arthritis, bacteremia, and toxic shock syndrome ([11–23](#)). Like GAS, *S. equisimilis* has also been linked to the poststreptococcal sequelae rheumatic heart disease, and in high endemic areas of rheu-

TABLE 2 Species of Lancefield group C and G streptococci^a

Species	Known sources	Lancefield group
<i>S. dysgalactiae</i> subsp. <i>dysgalactiae</i>	Animals	C
<i>S. dysgalactiae</i> subsp. <i>equisimilis</i>	Humans, animals (rare)	C, G
<i>S. equi</i> subsp. <i>equi</i>	Animals	C
<i>S. canis</i>	Animals, humans (rare)	G
<i>S. equi</i> subsp. <i>zooepidemicus</i>	Animals, humans	C
<i>S. phocae</i>	Animals (seal)	C
<i>S. anginosus</i> group	Humans	A, C, F, G

^aFrom reference [1](#).

matic fever, carriage rates of GCS/GGS have been found to be higher than those of GAS (24, 25).

Infections with *S. zooepidemicus* and *S. canis*, normally considered zoonotic species, have also been reported in humans (26–29), including severe infective endocarditis (30).

Generally, person to person transmission of GCS/GGS occurs via respiratory droplets or skin contacts, but other zoonotic vehicles of transmission such as unpasteurized milk products are also possible (26, 27, 31).

S. dysgalactiae predominantly resides in domestic animals such as cattle, sheep, cats, and dogs that are either healthy carriers of the bacterium or go on to develop diseases such as pneumonia, arthritis, septicemia, and abscesses, particularly bovine mastitis. *S. equi*, a pathogen primarily restricted to horses and donkeys, causes strangles, which is a highly contagious disease characterized by purulent discharges from the respiratory tract and the development of abscesses (32). This organism is not considered to be part of the normal flora, because of its close association with disease. In contrast, *S. zooepidemicus* is an opportunistic commensal that colonizes mucosal surfaces and causes rhinopharyngitis, pneumonia (33), endometritis, neonatal septicemia, and wound infections in horses (34) as well as disease in other domestic animals such as cattle, sheep, pigs, and chickens. Although uncommon, *S. zooepidemicus* is also responsible for a range of zoonotic infections in humans (26, 27).

The small-colony phenotype of group C and G streptococci is expressed by the *S. anginosus* group, formerly known as *Streptococcus milleri*. The group contains three species, *S. anginosus*, *Streptococcus constellatus*, and *Streptococcus intermedius* (1), and forms a distinct subclade by phylogenomic analysis (35).

Strains belonging to the *S. anginosus* group express group antigens F, C, A, and G or no antigen. These strains are also likely to be non-beta-hemolytic. Members of these species are recognized as common commensal organisms of the human oral cavity, gastrointestinal tract, and genitourinary tract. They are also associated with abscess formation in the mouth and other body sites (36, 37) as well as pharyngitis (38) and endocarditis (39, 40). It has also been reported that *S. anginosus*, along with *Streptococcus mitis* and *Treponema denticola*, can be isolated from esophageal cancer tissue and by initiating inflammation in the cancerous tissue has a role in the development or progression of these cancers (41, 42).

Several assays are available to detect, classify, and genetically describe GCS and GGS. Routine microbiological diagnosis follows the same guidelines that are in use for the identification of the other beta-hemolytic

streptococci (43). The beta-hemolytic streptococci isolates are divided into large and small colonies forming groups based on the growth on sheep blood agar: the large-colony-forming group is “pyogenic.” The small-colony-forming species make up the anginosus group and are not referred to as GCS or GGS, even though they may cross-react with C or G sera. Lancefield agglutination tests are still used to group beta-hemolytic streptococci into the Lancefield groups. Fermentation tests are used to classify GCS as *S. equisimilis* or *S. zooepidemicus* and are based on a spectrum of biochemical tests which are now available commercially (44). More recently, matrix-assisted laser desorption ionization–time of flight mass spectrometry has been described as a rapid alternative for the identification of streptococci (45). In addition, multilocus sequence analysis of seven housekeeping genes (46) and other sequence-based assays are performed to analyze the streptococcal phylogenies. In particular, sequencing of the *emm* gene, which encodes the hyper-variable surface protein M, and 16s rRNA genes is undertaken for phylogenetic analysis of these streptococci based on sequence similarities (47, 48).

Next-generation sequencing is increasingly being used for genetic characterization, phylogenetic analyses, virulence and pathogenicity profiles, and antibiotic resistance analyses (22, 49). *S. equisimilis* and *Streptococcus pyogenes* (or GAS) share a high genetic similarity (~72%) and similar virulence genes, suggesting a common evolutionary origin and genetic exchange (22, 25, 50).

MECHANISMS FOR ADHERENCE

S. equisimilis causes a spectrum of disease in humans similar to that caused by *S. pyogenes*, and molecular studies have demonstrated virulence determinants that are almost identical. Putative virulence determinants of *S. equisimilis* include adhesins, toxins, and factors that are essential for dissemination in human tissues and for interference with the host immune responses.

Adhesion of microorganisms to host tissues represents a critical phase in the development of infection. It is therefore unsurprising that microorganisms have evolved dedicated mechanisms for attachment and adherence to extracellular matrix components of the host (51, 52). Of these components, the high-molecular-weight glycoprotein fibronectin appears to be the major attachment target of Gram-positive cocci, including GGS and GCS. Fibronectin itself is responsible for substrate adhesion of eukaryotic cells via specific cell surface factors of the integrin family. It also specifically interacts with other matrix components, such as collagen, fibrin, and sulfated

glycosaminoglycans, demonstrating that this molecule fulfills multifunctional roles within the extracellular network (53). Being present in the extracellular matrix of most tissues, as well as in plasma and other body fluids in its soluble form, fibronectin represents an exquisite target for bacteria to exploit the cell attachment properties of this molecule by linking the pathogen to specific target cells. Epithelial cells of the human upper respiratory tract are bathed in secretions containing fibronectin in its soluble form. Once bound to the bacterial surface, it enables the pathogen to attach and subsequently colonize the primary site of infection.

Fibronectin binding proteins (FBPs) were first identified in *Staphylococcus aureus* and *S. pyogenes* (54), and approximately 11 have been defined in *S. pyogenes*, highlighting their importance for this pathogen (55). Binding by these bacteria to eukaryotic cells via fibronectin is also an important preliminary event prior to the invasion of these cells (56–58). The FBPs from streptococci and staphylococci share a common architecture, with a putative signal sequence at the N terminus and a wall- and membrane-spanning region. The major fibronectin binding domains are located within the C-terminal part of the proteins and are composed of 3 to 5 repetitive units that consist of 35 to 37 amino acid residues and bind to the 29-kDa N-terminal fragment of fibronectin. Further binding studies of PrtF/SfbI from *S. pyogenes* and FnBPA in *S. aureus* also indicate the presence of a secondary fibronectin binding site upstream of the repeat regions (54).

The first FBPs to be identified in group C and G streptococci were FnB_A and FnB_B from *S. dysgalactiae* (59). FnB_A can opacify serum and bind fibrinogen, like the serum opacity factor of *S. pyogenes* (60). Both FnB_A and FnB_B bind fibronectin through C-terminal repeat regions (51).

Three FBPs have been defined in *S. zooepidemicus*: FNZ, FNZ2, and SFS. Although the modular organization of FNZ, an *S. zooepidemicus* FBP, is similar to those from other species, homology at the amino acid level is weak (61). Fibronectin binding is mediated through a repeat region and a second upstream region containing the amino acid motif LAGESGET. This motif is also present in the secondary binding domain of SfbI (62), where it acts independently of the repeat region in binding to fibronectin (63). This domain is also present in GfbA, the homologue of SfbI in *S. dysgalactiae* (64). *fne* is the homologous gene to *fnz* in *S. equi* (65). However, FNE is unique in that a single nucleotide deletion in the *fne* gene has resulted in the truncation of the protein, so with the loss of the cell-wall binding motif, it is secreted into the surrounding environment. The truncated FNE

also lacks the classic fibronectin binding repeats found in the C-terminal region of FNZ but is able to bind fibronectin through the amino-half domain, also found in FNZ. Consequently, fibronectin binding at the bacterial surface is much lower for *S. equi* than *S. zooepidemicus*, although both have an additional cell wall-anchored FBP: FNEB and FNZ2 (66). FNE also binds to collagen, and the protein affects the interstitial fluid pressure *in vivo* (67). The molecular structure of the interaction between FNE and fibronectin has been reported (68). FNZ2 also has both collagen and fibronectin binding properties (69). SFS of both *S. equi* and *S. zooepidemicus* strains contains a signal peptide but no cell wall binding motifs or traditional fibronectin binding motifs. SFS inhibits binding between fibronectin and collagen. If SFS were to bind fibronectin attached to the bacterial surface through FNEB/FNZ/FNZ2, this might inhibit binding of fibronectin to collagen, which could have several physiological consequences (70). The molecular interaction between SFS and fibronectin has been reported (71).

Some of the FBPs and collagen binding proteins identified in *S. pyogenes* are chromosomally located within the fibronectin and collagen binding proteins and T antigen encoding (FCT) locus. There are at least nine types of FCT regions in *S. pyogenes* that vary by gene content and order, but within this region are the genes required to make the pili. The pilus has been shown to contribute to the formation of biofilm and mediate adherence to host cells (72–75). Whole-genome sequence analysis of group C and G streptococci has also identified potential FCT regions. One putative pilin locus, FimI, exists within the genome of *S. equi* strain 4047 (76). Although no pili structures have been identified for *S. equi*, the proteins of FimI are expressed during growth and have the potential to affect adherence *in vivo* (77). The gene encoding the collagen binding protein CNE of *S. equi* (78) is present in this FimI locus. A homologous locus was also found in *S. zooepidemicus* isolates, as well as an additional one or two potential pili loci (76, 79). Two FCT regions were also found in *S. equisimilis*, which share high levels of genetic identity to FCT regions from different *S. pyogenes* genotypes, suggesting multiple horizontal gene transfer events (22, 80).

An alternative way for group C and G streptococci to adhere to host cells is via binding to other extracellular matrix molecules, including fibrinogen, vitronectin, laminin, collagen, and plasminogen (81, 82). The M-protein-like fibrinogen binding protein of G streptococci (FOG) can act as an adhesin by binding collagen IV (83) and has been shown to also bind collagen I fibrils in dermis *in vivo* (84).

Vitronectin is a multifunctional serum protein that affects the humoral immune system by binding to and inhibiting the complement membrane attack complex (85) and is also a major matrix-associated adhesive glycoprotein that regulates blood coagulation. The ability of group C and G streptococci specifically to interact with vitronectin (86) and mediate the adherence to both epithelial and endothelial cells (87, 88) was demonstrated some time ago. However, a specific vitronectin binding protein has never been identified in streptococci.

ANTIPHAGOCYtic FACTORS

A major requirement of pathogenic streptococci is to be able to resist phagocytosis. The streptococcal M protein, first identified and characterized in *S. pyogenes*, is the major antiphagocytic factor (89). It is a multidomain surface-exposed molecule that forms a coiled-coil secondary structure with significant irregularities that in the B-repeat region are essential for the fibrinogen binding properties (90). Binding of complement factors, fibrinogen, and inhibition of C3b deposition on the bacterial surface are mechanisms by which the M-protein can inhibit opsonization of the organism by the alternative complement pathway, thus evading the host's nonspecific immune defense mechanism.

Several M-like proteins have been identified in group C and G streptococci. Protein MG1, the first group G streptococcal M-like protein characterized on the molecular level (91), exhibits typical structural and biological features of M proteins, such as coiled-coil structure and the ability to generate type-specific opsonizing antibodies. Protein MG1 shares highly homologous sequences with the C-terminal repeat region of class I M proteins, which are frequently associated with rheumatic fever. M proteins of GGS are also responsible for conferring resistance to phagocytosis (92).

In human strains of GGS, FOG is critical for bacterial pathogenesis through its antiphagocytic action of binding fibrinogen (93). It also has the ability to bind IgG subclasses IgG1 and IgG2, although FOG-bound IgG1 can trigger the complement cascade through C1/C1q activation. It is unclear whether this is detrimental to the bacteria, because FOG remains protective against phagocytosis (94).

The antiphagocytic protein SeM (FgBP) appears to be the predominant M-like protein on the surface of *S. equi*, capable of binding fibrinogen and IgG4 and IgG7 subclasses (95–97). SeM confers resistance to phagocytosis, but this also requires the presence of the hyaluronic acid capsule (98). Like the M protein of *S. pyogenes*, SeM

is variable at the N-terminal region, suggesting selective pressure (99). Interestingly, this does not appear to be the case for two other M-like proteins expressed by *S. equi*: Se18.9 and SzPSe (99). The antiphagocytic protein Se18.9 is commonly found in strains of *S. equi*, but rarely in strains of *S. zooepidemicus* (76), and acts by binding to fibrinogen and the complement regulatory protein, factor H (100). SzP proteins are also antiphagocytic and are expressed by both *S. equi* and *S. zooepidemicus*. Whole-genome sequence analysis of *S. zooepidemicus* strain H70 identified a potential M-protein homologue in SzM that resembles SeM in its secondary structure, which consists of a predicted C-terminal coiled-coil (76, 101). The N-terminal of SeM is unique to *S. equi* and binds fibrinogen (97, 102). SzM can also bind fibrinogen but cannot bind IgG (103). The M protein of *S. canis* specifically binds the Fc region of IgG in a nonopsonic manner (104).

DemA is an *S. dysgalactiae* protein with homology to FgBP identified by screening of a phagemid expression library (105). The mature DemA protein is 54 kDa in size, contains a signal sequence and a cell wall binding domain, and is predicted to have a coiled-coil secondary structure. DemA shows greatest homology to the FgBP at the C-terminal end in a region that does not participate in fibrinogen binding. The amino acid motif VSKDLADKL present with the repeat units of both DemA and FgBP suggests that the sequence may have an important biological function. DemA is able to bind IgG from various animal sources, reminiscent of type IIa Ig-binding proteins of *S. pyogenes* in a domain distinct from the fibrinogen binding domain. Nucleotide sequencing of the *demA* locus identified an open reading frame upstream of *demA* that is homologous to *mga*, a positive regulator of M-protein expression in *S. pyogenes* (105).

C5a peptidase of *S. pyogenes* (ScpA) is well known to specifically cleave and inactivate the chemoattractant C5a, but it has also recently been shown to cleave the complement factor C3 and the chemoattractant C3a (106), expanding its impact on immune evasion. ScpA has also been identified as an adhesion factor enabling bacteria to adhere to endothelial and epithelial cells (106). Homologues of ScpA have been identified in group B streptococci (ScpB) and group C and G streptococci isolated from humans but not from GGS from animals (107, 108). Both *S. equi* and *S. zooepidemicus* also carry homologous C5a peptidase genes (*scpE* and *scpZ*, respectively). However, the C5a peptidase gene in *S. equi* is reported to be a pseudogene (76). Unlike in *S. pyogenes*, where *scpA* is located in the *mga* regulon that includes the *emm*-gene, the *scp* genes of group B, C, and G streptococci are associated with a transposon, suggesting

that lateral genes transfer between these species (107, 109).

The *S. pyogenes* cell wall envelope proteinase, SpyCEP, is another protease that contributes to the prevention of phagocytosis through the specific cleavage of interleukin-8 and other chemokines to prevent the activation and migration of neutrophils (110, 111). Similar enzymes have been identified in *S. equi* (SeCEP) and *S. zooepidemicus* (SzoCEP) which share 98% identity to each other and 59% identity to SpyCEP (108). SeCEP has been shown to cleave both human and equine IL-8, and vaccination with a recombinant portion of SpyCEP prevented bacterial dissemination in a murine model of *S. equi* invasive infection, suggesting an important contribution to disease (112).

IMMUNOGLOBULIN BINDING AND INACTIVATING PROTEINS

Streptococcal protein G is a surface molecule associated with the majority of group C and G streptococcal isolates of human origin. Protein G interactions with immunoglobulins and other host proteins have been the subject of detailed reviews (113–115). Protein G is defined as a type III IgG Fc receptor and interacts with a wide species range of immunoglobulins, as well as human serum albumin, kininogen, and α_2 -macroglobulin. Protein G exhibits a modular structure in which the binding sites for IgG are located in the C-terminal repeat region (114–116). The central A/B-repeat region constitutes the binding domain for serum albumin, and the N-terminal E region is responsible for interacting with the native form of α_2 -macroglobulin (82). In contrast to human pathogenic strains of GGS that exclusively bind to the native (slow) form of α_2 -macroglobulin via protein G, animal-derived isolates of bovine and equine origin bind the proteinase-complexed (fast) form of the molecule. The B1 domain of protein G involved in immunoglobulin binding consists of an α -helix and four β -strand sheets. This domain has been used as a model structure in numerous biochemical studies examining protein folding, protein interaction, and synthetic protein design (117–119).

Two protein G-related proteins, MIG and MAG, from two mastitis-causing *S. dysgalactiae* strains (120) and MAG (121, 122) and ZAG (123) from *S. zooepidemicus* have been characterized on the molecular level. Like protein G, MAG and ZAG exhibit serum albumin and type III Fc receptor activity, whereas the protein MIG lacks albumin-binding activity. However, MAG is able to bind to immunoglobulins from a greater number of animal sources than protein G. MAG, ZAG, and MIG

also bind to the fast form of α_2 -macroglobulin. The α_2 -macroglobulin binding region in MIG is not homologous to those of protein G, MAG, ZAG, and GRAB. In phagocytosis assays a MIG isogenic mutant strain of *S. dysgalactiae* was not as resistant to opsonization by bovine neutrophils as the parental strain (124). MIG has also been shown to bind to bovine immunoglobulin A (125) and can inhibit bacterial internalization into host cells (126).

As well as IgG binding proteins, group C and G streptococci can express variants of the *S. pyogenes* IdeS and EndoS, which are IgG-degrading enzymes. While IdeS cleaves the hinge region of IgG (127), EndoS removes core IgG glycans (128). Both *S. equi* and *S. zooepidemicus* express two forms of IgG-endopeptidases, IdeE/IdeE2 and IdeZ/IdeZ2, respectively, that are capable of cleaving IgG from several mammalian species (129, 130), although IdeE2/IdeZ2 cleave horse IgG with greater efficiency than IdeE/IdeZ (130). EndoS homologues have also been identified in *S. equi* (EndoSe) and *S. zooepidemicus* (EndoSz), and they share 86 to 89% sequence identity to each other and 70% identity to EndoS of *S. pyogenes* (131). EndoSd has also been found in *S. equisimilis* with IgG hydrolyzing activity (132), and IdeP was identified in the group C *S. phocae* subsp. *phocae* that affects marine animals, although its function has not been confirmed (133).

ENZYMES AND TOXINS

After colonization, adherence, and evasion of host immune responses, the dissemination of pathogenic streptococci in tissues is regarded to be an important step for the onset and development of an invasive disease. One of the factors involved in this process is streptokinase, a protein found in groups C, G, and A streptococci (134). The formation of a streptokinase/plasminogen complex results in the exposure of the plasminogen active site, which then catalyzes the conversion of other plasminogen molecules into plasmin (135). Plasmin is a key serine protease in the fibrinolytic system that is able to break down tissue barriers, thereby enabling the dissemination of streptococci. M proteins coordinately interact with the secreted streptokinase by binding either fibrinogen (136) or plasminogen (137). A high level of variation exists within the streptokinase gene, *ska*, and alleles of *ska* have been associated with tissue tropisms and differing levels of plasminogen activation (138). There is some overlap of alleles between human strains of group C and G streptococci and GAS (139). Streptokinases have been isolated from both human and animal group C and G isolates and have specificity for the plasminogens of their respective hosts (140). Thus, streptokinases from human

group A, C, and G streptococci isolates have greater homology to each other than to animal isolates of group C and G streptococci. A study by Caballero et al. (58) found the amino acid homology between streptokinases from human and animal *S. equisimilis* isolates to be only 35%. Homology between the streptokinase from *S. equisimilis* of equine and porcine origins was only 21%. The ability of streptokinases to cleave plasminogen from specific species may therefore be a critical factor in determining the host range of individual streptococcal strains (141). Invasive human strains of *S. equisimilis* have been found to express higher levels of streptokinase, and this can drive virulence in a murine model of invasive necrotizing fasciitis, suggesting an important role for streptokinase in disease (142, 143).

Streptolysin O (SLO) is the prototype of a family of thiol-activated cytolysins produced by the genus *Streptococcus* and by other Gram-positive bacteria, including *Bacillus*, *Clostridium*, and *Listeria* species (144). The genes coding for SLO of group C and G streptococci (*S. dysgalactiae*) are almost identical to that of *S. pyogenes* (145). SLO homologues have not been described in *S. equi*. SLO is able to disrupt the cytoplasmic membrane of several eukaryotic cell types that include erythrocytes, leukocytes, macrophages, platelets, and epithelial cells. Separate from its pore-forming activity, SLO also acts to translocate NADase into host cells (146), which contributes to cytotoxicity by depleting energy stores. Other functions of these two streptococcal toxins include limiting neutrophil responsiveness and potentiating bacterial survival, replication, and persistence inside keratinocytes, epithelial cells, and macrophages, which may protect GAS from the immune response and antimicrobial therapy (147–151).

Streptolysin S (SLS) is another cytolysin secreted by groups A, C, and G streptococci, including the animal-pathogenic *S. equi* species (152). SLS belongs to a distinct group of hemolytic toxins that are characterized by their resistance to oxidation and sensitivity to trypan blue. SLS activity results in damage to membranes of various cell types as well as subcellular organelles (153). In contrast to the 57-kDa SLO, SLS is a small, 57-amino acid protein. The gene encoding SLS, *sagA*, is part of a locus containing nine open reading frames which contain significant homology with genes from bacteriocin loci (154, 155). In *S. equisimilis*, SLS expression is under the control of both the *covRS* and *fasCAX* two-component regulatory systems, which in *S. pyogenes* have been shown to be involved in the regulation of multiple virulence factors (156). In a mouse infection model, *S. equisimilis* expressing SLS proliferates and induces necrotic lesions

at the site of infection, whereas SLS-negative strains do not, suggesting that SLS is an important factor in the development of necrotizing fasciitis (154).

Streptococcal pyogenic toxins (spe) are superantigen genes capable of cross-linking the major histocompatibility complex class II on antigen-presenting cells to the T cell receptor, leading to proliferation of T cells and substantial release of inflammatory cytokines. Eleven superantigen genes have been identified in *S. pyogenes*, and they are thought to drive the development of scarlet fever and toxic shock syndrome, the latter contributing to high mortality rates following necrotizing fasciitis. Recent work has also identified a role for superantigens in upper respiratory tract infection (157). Homologues of the GAS superantigens have been identified in group C and G streptococci (158). The most commonly found superantigen in *S. equisimilis* is *speG* (*spegg* or *speG^{dys}*) (159, 160), and genomic analysis indicates that *S. pyogenes* and *S. equisimilis* *speG* genes are orthologues that are direct descendants from a common ancestor gene (161). The gene for *speG* can also be found in *S. canis* (158). The role of *S. equisimilis* *speG* in humans is unclear because the presence of this gene does not correlate with disease severity and it does not confer mitogenic activity toward human mononuclear cells, although it can stimulate bovine T cells (160, 162, 163). The streptococcal superantigens *speA*, *speC*, *speJ*, *speK*, *speH*, *speL*, *speM*, and *ssa* are found infrequently in *S. equisimilis* and other human group C and G streptococci (160, 164–166).

Homologues of *S. pyogenes* superantigens *speH* and *speI* can be carried by strains of *S. equi* (167), and homologues of *speK* and *speL* can be carried by both *S. equi* and *S. zooepidemicus*, although because of confusion with the nomenclature, *speK* homologues were originally termed *speL_{se}/seeL/szeL*, and *speL* homologues were termed *speM_{se}/seem/szeM* (160, 167–169). Three additional superantigen genes, termed *szeN*, *szeP*, and *szeF*, have been identified in *S. zooepidemicus* only and are capable of stimulating equine peripheral blood mononuclear cells (170). *S. dysgalactiae*-derived mitogen shows homology to *speM* of *S. pyogenes* and can stimulate human mononuclear cells (160, 171). The majority of the superantigens found in *S. pyogenes* and group C and G streptococci are associated with prophages which may drive the lateral transfer of these virulence factors between streptococcal species (76, 172, 173).

CONCLUSIONS

Many recent studies have highlighted the increasing number of systemic infections caused by *S. equisimilis*, par-

ticularly among immunocompromised individuals and specific populations, and this therefore suggests that this species will gain even more clinical importance in the future. High nucleotide similarities among virulence genes and their association with mobile genetic elements support the hypothesis of extensive horizontal gene transfer events between streptococcal species of the pyogenic group. A better understanding of the mechanisms of pathogenesis will hopefully be revealed by whole-genome sequencing, and this may result in more effective clinical strategies for the pyogenic group of streptococci in generally.

REFERENCES

- Facklam R. 2002. What happened to the streptococci: overview of taxonomic and nomenclature changes. *Clin Microbiol Rev* 15:613–630 <http://dx.doi.org/10.1128/CMR.15.4.613-630.2002>.
- Chanter N, Collin N, Holmes N, Binns M, Mumford J. 1997. Characterization of the Lancefield group C *Streptococcus* 16S-23S RNA gene intergenic spacer and its potential for identification and sub-specific typing. *Epidemiol Infect* 118:125–135 <http://dx.doi.org/10.1017/S0950268896007285>.
- Vossen A, Abdulmawjood A, Lämmle C, Weiss R, Siebert U. 2004. Streptococci isolated from harbor seals (*Phoca vitulina*) and grey seals (*Halichoerus grypus*) of the German North and Baltic Seas. *J Clin Microbiol* 42:469–473 <http://dx.doi.org/10.1128/JCM.42.1.469-473.2004>.
- Skaar I, Gaustad P, Tønjum T, Holm B, Stenwig H. 1994. *Streptococcus phocae* sp. nov., a new species isolated from clinical specimens from seals. *Int J Syst Bacteriol* 44:646–650 <http://dx.doi.org/10.1099/00207713-44-4-646>.
- Gunnarsson RK, Holm SE, Söderström M. 1997. The prevalence of beta-haemolytic streptococci in throat specimens from healthy children and adults. Implications for the clinical value of throat cultures. *Scand J Prim Health Care* 15:149–155 <http://dx.doi.org/10.3109/02813439709018506>.
- Siljander T, Karppelin M, Vähäkuopus S, Syrjänen J, Toropainen M, Kere J, Vuento R, Jussila T, Vuopio-Varkila J. 2008. Acute bacterial, nonnecrotizing cellulitis in Finland: microbiological findings. *Clin Infect Dis* 46:855–861 <http://dx.doi.org/10.1086/527388>.
- Bélar S, Toepfner N, Arnold B, Alabi AS, Berner R. 2015. β -Hemolytic streptococcal throat carriage and tonsillopharyngitis: a cross-sectional prevalence study in Gabon, Central Africa. *Infection* 43:177–183 <http://dx.doi.org/10.1007/s15010-014-0709-y>.
- Broyles LN, Van Beneden C, Beall B, Facklam R, Shewmaker PL, Malpiedi P, Daily P, Reingold A, Farley MM. 2009. Population-based study of invasive disease due to beta-hemolytic streptococci of groups other than A and B. *Clin Infect Dis* 48:706–712 <http://dx.doi.org/10.1086/597035>.
- Rantala S, Vuopio-Varkila J, Vuento R, Huhtala H, Syrjänen J. 2009. Clinical presentations and epidemiology of beta-haemolytic streptococcal bacteraemia: a population-based study. *Clin Microbiol Infect* 15:286–288 <http://dx.doi.org/10.1111/j.1469-0691.2008.02672.x>.
- Ekelund K, Skinhøj P, Madsen J, Konradsen HB. 2005. Invasive group A, B, C and G streptococcal infections in Denmark 1999–2002: epidemiological and clinical aspects. *Clin Microbiol Infect* 11:569–576 <http://dx.doi.org/10.1111/j.1469-0691.2005.01169.x>.
- Bradley SF, Gordon JJ, Baumgartner DD, Marasco WA, Kauffman CA. 1991. Group C streptococcal bacteremia: analysis of 88 cases. *Rev Infect Dis* 13:270–280 <http://dx.doi.org/10.1093/clinids/13.2.270>.
- Brandt CM, Spellerberg B. 2009. Human infections due to *Streptococcus dysgalactiae* subspecies *equisimilis*. *Clin Infect Dis* 49:766–772 <http://dx.doi.org/10.1086/605085>.
- Bruun T, Oppegaard O, Kittang BR, Mylvaganam H, Langeland N, Skrede S. 2015. Etiology of cellulitis and clinical prediction of streptococcal disease: a prospective study. *Open Forum Infect Dis* 3:ofv181 <http://dx.doi.org/10.1093/ofid/ofv181>.
- Carmeli Y, Ruoff KL. 1995. Report of cases of and taxonomic considerations for large-colony-forming Lancefield group C streptococcal bacteremia. *J Clin Microbiol* 33:2114–2117.
- Dubost JJ, Soubrier M, De Champs C, Ristori JM, Sauvezie B. 2004. Streptococcal septic arthritis in adults. A study of 55 cases with a literature review. *Joint Bone Spine* 71:303–311 [http://dx.doi.org/10.1016/S1297-319X\(03\)00122-2](http://dx.doi.org/10.1016/S1297-319X(03)00122-2).
- Freitas DM. 2017. Group G *Streptococcus dysgalactiae* subspecies *equisimilis*, the clinical significance of a rare infection: endocarditis, polyarthritis, septic bursitis and pneumonia with complicated parapneumonic effusion. *BMJ Case Rep* 2017:2017.
- González Terán B, Roiz MP, Ruiz Jimeno T, Rosas J, Calvo-Alén J. 2001. Acute bacterial arthritis caused by group C streptococci. *Semin Arthritis Rheum* 31:43–51 <http://dx.doi.org/10.1053/sarh.2001.21405>.
- Keiser P, Campbell W. 1992. 'Toxic strep syndrome' associated with group C *Streptococcus*. *Arch Intern Med* 152:882, 884 <http://dx.doi.org/10.1001/archinte.1992.00400160162042>.
- Korman TM, Boers A, Gooding TM, Curtis N, Visvanathan K. 2004. Fatal case of toxic shock-like syndrome due to group C *Streptococcus* associated with superantigen exotoxin. *J Clin Microbiol* 42:2866–2869 <http://dx.doi.org/10.1128/JCM.42.6.2866-2869.2004>.
- Lothar SA, Jassal DS, Lagacé-Wiens P, Keynan Y. 2017. Emerging group C and group G streptococcal endocarditis: a Canadian perspective. *Int J Infect Dis* 65:128–132 <http://dx.doi.org/10.1016/j.ijid.2017.10.018>.
- Naik TB, Nadagir SD, Biradar A. 2016. Prevalence of beta-hemolytic streptococci groups A, C, and G in patients with acute pharyngitis. *J Lab Physicians* 8:45–49 <http://dx.doi.org/10.4103/0974-2727.176235>.
- Shimomura Y, Okumura K, Murayama SY, Yagi J, Ubukata K, Kirikae T, Miyoshi-Akiyama T. 2011. Complete genome sequencing and analysis of a Lancefield group G *Streptococcus dysgalactiae* subsp. *equisimilis* strain causing streptococcal toxic shock syndrome (STSS). *BMC Genomics* 12:17 <http://dx.doi.org/10.1186/1471-2164-12-17>.
- Zaoutis T, Attia M, Gross R, Klein J. 2004. The role of group C and group G streptococci in acute pharyngitis in children. *Clin Microbiol Infect* 10:37–40 <http://dx.doi.org/10.1111/j.1469-0691.2004.00732.x>.
- Haidan A, Talay SR, Rohde M, Sriprakash KS, Currie BJ, Chhatwal GS. 2000. Pharyngeal carriage of group C and group G streptococci and acute rheumatic fever in an Aboriginal population. *Lancet* 356:1167–1169 [http://dx.doi.org/10.1016/S0140-6736\(00\)02765-3](http://dx.doi.org/10.1016/S0140-6736(00)02765-3).
- Bramhachari PV, Kaul SY, McMillan DJ, Shaila MS, Karmarkar MG, Sriprakash KS. 2010. Disease burden due to *Streptococcus dysgalactiae* subsp. *equisimilis* (group G and C streptococcus) is higher than that due to *Streptococcus pyogenes* among Mumbai school children. *J Med Microbiol* 59:220–223 <http://dx.doi.org/10.1099/jmm.0.015644-0>.
- Kittang BR, Pettersen VK, Oppegaard O, Skutlaberg DH, Dale H, Wiker HG, Skrede S. 2017. Zoonotic necrotizing myositis caused by *Streptococcus equi* subsp. *zooepidemicus* in a farmer. *BMC Infect Dis* 17:147 <http://dx.doi.org/10.1186/s12879-017-2262-7>.
- Pelkonen S, Lindahl SB, Suomala P, Karhukorpi J, Vuorinen S, Koivula I, Väisänen T, Pentikäinen J, Autio T, Tuuminen T. 2013. Transmission of *Streptococcus equi* subspecies *zooepidemicus* infection from horses to humans. *Emerg Infect Dis* 19:1041–1048 <http://dx.doi.org/10.3201/eid1907.121365>.
- Galpérine T, Cazorla C, Blanchard E, Boineau F, Ragnaud JM, Neau D. 2007. *Streptococcus canis* infections in humans: retrospective study of 54 patients. *J Infect* 55:23–26 <http://dx.doi.org/10.1016/j.jinf.2006.12.013>.
- Takeda N, Kikuchi K, Asano R, Harada T, Totsuka K, Sumiyoshi T, Uchiyama T, Hosoda S, Norihiko Takeda, Ken Kikuchi, Ryuta. 2001. Recurrent septicemia caused by *Streptococcus canis* after a dog bite. *Scand J Infect Dis* 33:927–928 <http://dx.doi.org/10.1080/00365540110076903>.

30. Lacave G, Coutard A, Troché G, Augusto S, Pons S, Zuber B, Laurent V, Amara M, Couzon B, Bédos JP, Pangon B, Grimaldi D. 2016. Endocarditis caused by *Streptococcus canis*: an emerging zoonosis? *Infection* 44:111–114 <http://dx.doi.org/10.1007/s15010-015-0809-3>.
31. Bordes-Benitez A, Sánchez-Oñoro M, Suárez-Bordón P, García-Rojas AJ, Saéz-Nieto JA, González-García A, Alamo-Antúnez I, Sánchez-Maroto A, Bolaños-Rivero M. 2006. Outbreak of *Streptococcus equi* subsp. *zooepidemicus* infections on the island of Gran Canaria associated with the consumption of inadequately pasteurized cheese. *Eur J Clin Microbiol Infect Dis* 25:242–246 <http://dx.doi.org/10.1007/s10096-006-0119-x>.
32. Harrington DJ, Sutcliffe IC, Chanter N. 2002. The molecular basis of *Streptococcus equi* infection and disease. *Microbes Infect* 4:501–510 [http://dx.doi.org/10.1016/S1286-4579\(02\)01565-4](http://dx.doi.org/10.1016/S1286-4579(02)01565-4).
33. Yoshikawa H, Yasu T, Ueki H, Oyama T, Oishi H, Anzai T, Oikawa M, Yoshikawa T. 2003. Pneumonia in horses induced by intrapulmonary inoculation of *Streptococcus equi* subsp. *zooepidemicus*. *J Vet Med Sci* 65:787–792 <http://dx.doi.org/10.1292/jvms.65.787>.
34. Welsh RD. 1984. The significance of *Streptococcus zooepidemicus* in the horse. *Equine Pract* 6:6–16.
35. Patel S, Gupta RS. 2018. Robust demarcation of fourteen different species groups within the genus *Streptococcus* based on genome-based phylogenies and molecular signatures. *Infect Genet Evol* 66:130–151 <http://dx.doi.org/10.1016/j.meegid.2018.09.020>.
36. Claridge JE III, Attorri S, Musher DM, Hebert J, Dunbar S. 2001. *Streptococcus intermedius*, *Streptococcus constellatus*, and *Streptococcus anginosus* (“*Streptococcus milleri* group”) are of different clinical importance and are not equally associated with abscess. *Clin Infect Dis* 32:1511–1515 <http://dx.doi.org/10.1086/320163>.
37. Ruoff KL. 1988. *Streptococcus anginosus* (“*Streptococcus milleri*”): the unrecognized pathogen. *Clin Microbiol Rev* 1:102–108 <http://dx.doi.org/10.1128/CMR.1.1.102>.
38. Whiley RA, Hall LM, Hardie JM, Beighton D. 1999. A study of small-colony, beta-haemolytic, Lancefield group C streptococci within the anginosus group: description of *Streptococcus constellatus* subsp. *pharyngis* subsp. nov., associated with the human throat and pharyngitis. *Int J Syst Bacteriol* 49:1443–1449 <http://dx.doi.org/10.1099/00207713-49-4-1443>.
39. Kitada K, Inoue M, Kitano M. 1997. Infective endocarditis-inducing abilities of “*Streptococcus milleri*” group. *Adv Exp Med Biol* 418:161–163 http://dx.doi.org/10.1007/978-1-4899-1825-3_39.
40. Kitada K, Inoue M, Kitano M. 1997. Experimental endocarditis induction and platelet aggregation by *Streptococcus anginosus*, *Streptococcus constellatus* and *Streptococcus intermedius*. *FEMS Immunol Med Microbiol* 19:25–32 <http://dx.doi.org/10.1111/j.1574-695X.1997.tb01069.x>.
41. Morita E, Narikiyo M, Yano A, Nishimura E, Igaki H, Sasaki H, Terada M, Hanada N, Kawabe R. 2003. Different frequencies of *Streptococcus anginosus* infection in oral cancer and esophageal cancer. *Cancer Sci* 94:492–496 <http://dx.doi.org/10.1111/j.1349-7006.2003.tb01471.x>.
42. Narikiyo M, Tanabe C, Yamada Y, Igaki H, Tachimori Y, Kato H, Muto M, Montesano R, Sakamoto H, Nakajima Y, Sasaki H. 2004. Frequent and preferential infection of *Treponema denticola*, *Streptococcus mitis*, and *Streptococcus anginosus* in esophageal cancers. *Cancer Sci* 95:569–574 <http://dx.doi.org/10.1111/j.1349-7006.2004.tb02488.x>.
43. Public Health England. 2013. Group C and group G *Streptococcus*: guidance, data and analysis. <https://www.gov.uk/government/collections/group-c-and-group-g-streptococcus-guidance-data-and-analysis#diagnosis-and-treatment>.
44. Efstratiou A. 1997. Pyogenic streptococci of Lancefield groups C and G as pathogens in man. *Soc Appl Bacteriol Symp Ser* 26(S1):72S–79S <http://dx.doi.org/10.1046/j.1365-2672.83.s1.8.x>.
45. Schulthess B, Brodner K, Bloemberg GV, Zbinden R, Böttger EC, Hombach M. 2013. Identification of Gram-positive cocci by use of matrix-assisted laser desorption ionization-time of flight mass spectrometry: comparison of different preparation methods and implementation of a practical algorithm for routine diagnostics. *J Clin Microbiol* 51:1834–1840 <http://dx.doi.org/10.1128/JCM.02654-12>.
46. Bishop CJ, Aanensen DM, Jordan GE, Kilian M, Hanage WP, Spratt BG. 2009. Assigning strains to bacterial species via the Internet. *BMC Biol* 7:3 <http://dx.doi.org/10.1186/1741-7007-7-3>.
47. Lal D, Verma M, Lal R. 2011. Exploring internal features of 16S rRNA gene for identification of clinically relevant species of the genus *Streptococcus*. *Ann Clin Microbiol Antimicrob* 10:28 <http://dx.doi.org/10.1186/1476-0711-10-28>.
48. Centre disease and control (CDC). 2008. Introduction to emm typing: M protein gene (emm) typing *Streptococcus pyogenes*. <https://www.cdc.gov/streplab/groupa-strep/emm-background.html>.
49. Wang X, Zhang X, Zong Z. 2016. Genome sequence and virulence factors of a group G *Streptococcus dysgalactiae* subsp. *equisimilis* strain with a new element carrying *erm(B)*. *Sci Rep* 6:20389 <http://dx.doi.org/10.1038/srep20389>.
50. Davies MR, McMillan DJ, Beiko RG, Barroso V, Geffers R, Sriprakash KS, Chhatwal GS. 2007. Virulence profiling of *Streptococcus dysgalactiae* subspecies *equisimilis* isolated from infected humans reveals 2 distinct genetic lineages that do not segregate with their phenotypes or propensity to cause diseases. *Clin Infect Dis* 44:1442–1454 <http://dx.doi.org/10.1086/516780>.
51. Joh D, Speziale P, Gurusiddappa S, Manor J, Höök M. 1998. Multiple specificities of the staphylococcal and streptococcal fibronectin-binding microbial surface components recognizing adhesive matrix molecules. *Eur J Biochem* 258:897–905 <http://dx.doi.org/10.1046/j.1432-1327.1998.2580897.x>.
52. Patti JM, Allen BL, McGavin MJ, Höök M. 1994. MSCRAMM-mediated adherence of microorganisms to host tissues. *Annu Rev Microbiol* 48:585–617 <http://dx.doi.org/10.1146/annurev.mi.48.100194.003101>.
53. Hynes RO, Yamada KM. 1982. Fibronectins: multifunctional modular glycoproteins. *J Cell Biol* 95:369–377 <http://dx.doi.org/10.1083/jcb.95.2.369>.
54. Schwarz-Linek U, Höök M, Potts JR. 2004. The molecular basis of fibronectin-mediated bacterial adherence to host cells. *Mol Microbiol* 52:631–641 <http://dx.doi.org/10.1111/j.1365-2958.2004.04027.x>.
55. Ryan PA, Juncosa B. 2016. Group A streptococcal adherence. In Ferretti JJ, Stevens DL, Fischetti VA (ed), *Streptococcus pyogenes: Basic Biology to Clinical Manifestations*. University of Oklahoma Health Sciences Center, Oklahoma City, OK.
56. Cue D, Dombek PE, Lam H, Cleary PP. 1998. *Streptococcus pyogenes* serotype M1 encodes multiple pathways for entry into human epithelial cells. *Infect Immun* 66:4593–4601.
57. Molinari G, Talay SR, Valentin-Weigand P, Rohde M, Chhatwal GS. 1997. The fibronectin-binding protein of *Streptococcus pyogenes*, SfbI, is involved in the internalization of group A streptococci by epithelial cells. *Infect Immun* 65:1357–1363.
58. Caballero AR, Lottenberg R, Johnston KH. 1999. Cloning, expression, sequence analysis, and characterization of streptokinases secreted by porcine and equine isolates of *Streptococcus equisimilis*. *Infect Immun* 67:6478–6486.
59. Lindgren PE, McGavin MJ, Signäs C, Guss B, Gurusiddappa S, Höök M, Lindberg M. 1993. Two different genes coding for fibronectin-binding proteins from *Streptococcus dysgalactiae*. The complete nucleotide sequences and characterization of the binding domains. *Eur J Biochem* 214:819–827 <http://dx.doi.org/10.1111/j.1432-1033.1993.tb17985.x>.
60. Courtney HS, Hasty DL, Li Y, Chiang HC, Thacker JL, Dale JB. 1999. Serum opacity factor is a major fibronectin-binding protein and a virulence determinant of M type 2 *Streptococcus pyogenes*. *Mol Microbiol* 32:89–98 <http://dx.doi.org/10.1046/j.1365-2958.1999.01328.x>.
61. Lindmark H, Jacobsson K, Frykberg L, Guss B. 1996. Fibronectin-binding protein of *Streptococcus equi* subsp. *zooepidemicus*. *Infect Immun* 64:3993–3999.
62. Talay SR, Valentin-Weigand P, Jerlström PG, Timmis KN, Chhatwal GS. 1992. Fibronectin-binding protein of *Streptococcus pyogenes*:

sequence of the binding domain involved in adherence of streptococci to epithelial cells. *Infect Immun* 60:3837–3844.

63. Sela S, Aviv A, Tovi A, Burstein I, Caparon MG, Hanski E. 1993. Protein F: an adhesin of *Streptococcus pyogenes* binds fibronectin via two distinct domains. *Mol Microbiol* 10:1049–1055 <http://dx.doi.org/10.1111/j.1365-2958.1993.tb00975.x>.
64. Kline JB, Xu S, Bisno AL, Collins CM. 1996. Identification of a fibronectin-binding protein (GfbA) in pathogenic group G streptococci. *Infect Immun* 64:2122–2129.
65. Lindmark H, Nilsson M, Guss B. 2001. Comparison of the fibronectin-binding protein FNE from *Streptococcus equi* subspecies *equi* with FNZ from *S. equi* subspecies *zooepidemicus* reveals a major and conserved difference. *Infect Immun* 69:3159–3163 <http://dx.doi.org/10.1128/IAI.69.5.3159-3163.2001>.
66. Lannergård J, Flock M, Johansson S, Flock JI, Guss B. 2005. Studies of fibronectin-binding proteins of *Streptococcus equi*. *Infect Immun* 73:7243–7251 <http://dx.doi.org/10.1128/IAI.73.11.7243-7251.2005>.
67. Lidén A, van Wieringen T, Lannergård J, Kassner A, Heinegård D, Reed RK, Guss B, Rubin K. 2008. A secreted collagen- and fibronectin-binding streptococcal protein modulates cell-mediated collagen gel contraction and interstitial fluid pressure. *J Biol Chem* 283:1234–1242 <http://dx.doi.org/10.1074/jbc.M704827200>.
68. Tiouajni M, Durand D, Blondeau K, Graille M, Urvoas A, Valerio-Lepiniec M, Guellouz A, Aumont-Nicaise M, Minard P, van Tilbeurgh H. 2014. Structural and functional analysis of the fibronectin-binding protein FNE from *Streptococcus equi* spp. *FEBS J* 281:5513–5531 <http://dx.doi.org/10.1111/febs.13092>.
69. Hong K. 2005. Identification and characterization of a novel fibronectin-binding protein gene from *Streptococcus equi* subspecies *zooepidemicus* strain VTU211. *FEMS Immunol Med Microbiol* 45:231–237 <http://dx.doi.org/10.1016/j.femsim.2005.04.006>.
70. Lindmark H, Guss B. 1999. SFS, a novel fibronectin-binding protein from *Streptococcus equi*, inhibits the binding between fibronectin and collagen. *Infect Immun* 67:2383–2388.
71. Ma W, Ma H, Fogerty FJ, Mosher DF. 2015. Bivalent ligation of the collagen-binding modules of fibronectin by SFS, a non-anchored bacterial protein of *Streptococcus equi*. *J Biol Chem* 290:4866–4876 <http://dx.doi.org/10.1074/jbc.M114.612259>.
72. Abbot EL, Smith WD, Siou GP, Chiriboga C, Smith RJ, Wilson JA, Hirst BH, Kehoe MA. 2007. Pili mediate specific adhesion of *Streptococcus pyogenes* to human tonsil and skin. *Cell Microbiol* 9:1822–1833 <http://dx.doi.org/10.1111/j.1462-5822.2007.00918.x>.
73. Crotty Alexander LE, Maisey HC, Timmer AM, Rooijackers SH, Gallo RL, von Kockritz-Blickwede M, Nizet V. 2010. M1T1 group A streptococcal pili promote epithelial colonization but diminish systemic virulence through neutrophil extracellular entrapment. *J Mol Med (Berl)* 88:371–381 <http://dx.doi.org/10.1007/s00109-009-0566-9>.
74. Manetti AG, Zingaretti C, Falugi F, Capo S, Bombaci M, Bagnoli F, Gambellini G, Bensi G, Mora M, Edwards AM, Musser JM, Graviss EA, Telford JL, Grandi G, Margarit I. 2007. *Streptococcus pyogenes* pili promote pharyngeal cell adhesion and biofilm formation. *Mol Microbiol* 64:968–983 <http://dx.doi.org/10.1111/j.1365-2958.2007.05704.x>.
75. Smith WD, Pointon JA, Abbot E, Kang HJ, Baker EN, Hirst BH, Wilson JA, Banfield MJ, Kehoe MA. 2010. Roles of minor pilin subunits Spy0125 and Spy0130 in the serotype M1 *Streptococcus pyogenes* strain SF370. *J Bacteriol* 192:4651–4659 <http://dx.doi.org/10.1128/JB.00071-10>.
76. Holden MT, Heather Z, Paillot R, Steward KF, Webb K, Ainslie F, Jourdan T, Bason NC, Holroyd NE, Mungall K, Quail MA, Sanders M, Simmonds M, Willey D, Brooks K, Aanensen DM, Spratt BG, Jolley KA, Maiden MC, Kehoe M, Chanter N, Bentley SD, Robinson C, Maskell DJ, Parkhill J, Waller AS. 2009. Genomic evidence for the evolution of *Streptococcus equi*: host restriction, increased virulence, and genetic exchange with human pathogens. *PLoS Pathog* 5:e1000346 <http://dx.doi.org/10.1371/journal.ppat.1000346>.
77. Steward KF, Robinson C, Maskell DJ, Nenci C, Waller AS. 2017. Investigation of the Fim1 putative pilus locus of *Streptococcus equi* subspecies *equi*. *Microbiology* 10.1099/mic.0.000506.
78. Lannergård J, Frykberg L, Guss B. 2003. CNE, a collagen-binding protein of *Streptococcus equi*. *FEMS Microbiol Lett* 222:69–74 [http://dx.doi.org/10.1016/S0378-1097\(03\)00222-2](http://dx.doi.org/10.1016/S0378-1097(03)00222-2).
79. Beres SB, Sesso R, Pinto SW, Hoe NP, Porcella SF, Deleo FR, Musser JM. 2008. Genome sequence of a Lancefield group C *Streptococcus zooepidemicus* strain causing epidemic nephritis: new information about an old disease. *PLoS One* 3:e3026 <http://dx.doi.org/10.1371/journal.pone.0003026>.
80. Oppegaard O, Mylvaganam H, Skrede S, Lindemann PC, Kittang BR. 2017. Emergence of a *Streptococcus dysgalactiae* subspecies *equisimilis* stG62647-lineage associated with severe clinical manifestations. *Sci Rep* 7:7589 <http://dx.doi.org/10.1038/s41598-017-08162-z>.
81. Allen BL, Höök M. 2002. Isolation of a putative laminin binding protein from *Streptococcus anginosus*. *Microb Pathog* 33:23–31 <http://dx.doi.org/10.1006/mpat.2002.0510>.
82. Müller HP, Rantamäki LK. 1995. Binding of native alpha 2-macroglobulin to human group G streptococci. *Infect Immun* 63:2833–2839.
83. Dinkla K, Nitsche-Schmitz DP, Barroso V, Reissmann S, Johansson HM, Frick IM, Rohde M, Chhatwal GS. 2007. Identification of a streptococcal octapeptide motif involved in acute rheumatic fever. *J Biol Chem* 282:18686–18693 <http://dx.doi.org/10.1074/jbc.M701047200>.
84. Nitsche DP, Johansson HM, Frick IM, Mörgelin M. 2006. Streptococcal protein FOG, a novel matrix adhesin interacting with collagen I *in vivo*. *J Biol Chem* 281:1670–1679 <http://dx.doi.org/10.1074/jbc.M506776200>.
85. Preissner KT. 1991. Structure and biological role of vitronectin. *Annu Rev Cell Biol* 7:275–310 <http://dx.doi.org/10.1146/annurev.cb.07.110191.001423>.
86. Chhatwal GS, Preissner KT, Müller-Berghaus G, Blobel H. 1987. Specific binding of the human S protein (vitronectin) to streptococci, *Staphylococcus aureus*, and *Escherichia coli*. *Infect Immun* 55:1878–1883.
87. Filippesen LF, Valentin-Weigand P, Blobel H, Preissner KT, Chhatwal GS. 1990. Role of complement S protein (vitronectin) in adherence of *Streptococcus dysgalactiae* to bovine epithelial cells. *Am J Vet Res* 51:861–865.
88. Valentin-Weigand P, Grulich-Henn J, Chhatwal GS, Müller-Berghaus G, Blobel H, Preissner KT. 1988. Mediation of adherence of streptococci to human endothelial cells by complement S protein (vitronectin). *Infect Immun* 56:2851–2855.
89. Fischetti VA. 1991. Streptococcal M protein. *Sci Am* 264:58–65 <http://dx.doi.org/10.1038/scientificamerican0691-58>.
90. McNamara C, Zinkernagel AS, Macheboeuf P, Cunningham MW, Nizet V, Ghosh P. 2008. Coiled-coil irregularities and instabilities in group A *Streptococcus* M1 are required for virulence. *Science* 319:1405–1408 <http://dx.doi.org/10.1126/science.1154470>.
91. Collins CM, Kimura A, Bisno AL. 1992. Group G streptococcal M protein exhibits structural features analogous to those of class I M protein of group A streptococci. *Infect Immun* 60:3689–3696.
92. Campo RE, Schultz DR, Bisno AL. 1995. M proteins of group G streptococci: mechanisms of resistance to phagocytosis. *J Infect Dis* 171:601–606 <http://dx.doi.org/10.1093/infdis/171.3.601>.
93. Johansson HM, Mörgelin M, Frick IM. 2004. Protein FOG: a streptococcal inhibitor of neutrophil function. *Microbiology* 150:4211–4221 <http://dx.doi.org/10.1099/mic.0.27269-0>.
94. Nitsche-Schmitz DP, Johansson HM, Sastalla I, Reissmann S, Frick IM, Chhatwal GS. 2007. Group G streptococcal IgG binding molecules FOG and protein G have different impacts on opsonization by C1q. *J Biol Chem* 282:17530–17536 <http://dx.doi.org/10.1074/jbc.M702612200>.
95. Lewis MJ, Meehan M, Owen P, Woof JM. 2008. A common theme in interaction of bacterial immunoglobulin-binding proteins with immuno-

- globulins illustrated in the equine system. *J Biol Chem* 283:17615–17623 <http://dx.doi.org/10.1074/jbc.M709844200>.
96. Meehan M, Lynagh Y, Woods C, Owen P. 2001. The fibrinogen-binding protein (FgBP) of *Streptococcus equi* subsp. *equi* additionally binds IgG and contributes to virulence in a mouse model. *Microbiology* 147:3311–3322 <http://dx.doi.org/10.1099/00221287-147-12-3311>.
97. Meehan M, Muldowney DA, Watkins NJ, Owen P. 2000. Localization and characterization of the ligand-binding domain of the fibrinogen-binding protein (FgBP) of *Streptococcus equi* subsp. *equi*. *Microbiology* 146:1187–1194 <http://dx.doi.org/10.1099/00221287-146-5-1187>.
98. Timoney JF, Suther P, Velineni S, Artiushin SC. 2014. The antiphagocytic activity of SeM of *Streptococcus equi* requires capsule. *J Equine Sci* 25:53–56 <http://dx.doi.org/10.1294/jes.25.53>.
99. Ijaz M, Velineni S, Timoney JF. 2011. Selective pressure for allelic diversity in SeM of *Streptococcus equi* does not affect immunoreactive proteins SzPse or Se18.9. *Infect Genet Evol* 11:1159–1163 <http://dx.doi.org/10.1016/j.meegid.2011.01.011>.
100. Tiwari R, Qin A, Artiushin S, Timoney JF. 2007. Se18.9, an antiphagocytic factor H binding protein of *Streptococcus equi*. *Vet Microbiol* 121:105–115 <http://dx.doi.org/10.1016/j.vetmic.2006.11.023>.
101. Kelly C, Bugg M, Robinson C, Mitchell Z, Davis-Poynter N, Newton JR, Jolley KA, Maiden MC, Waller AS. 2006. Sequence variation of the SeM gene of *Streptococcus equi* allows discrimination of the source of strangles outbreaks. *J Clin Microbiol* 44:480–486 <http://dx.doi.org/10.1128/JCM.44.2.480-486.2006>.
102. Timoney JF, Artiushin SC, Boschwitz JS. 1997. Comparison of the sequences and functions of *Streptococcus equi* M-like proteins SeM and SzPse. *Infect Immun* 65:3600–3605.
103. Velineni S, Timoney JF. 2013. Characterization and protective immunogenicity of the SzM protein of *Streptococcus zooepidemicus* NC78 from a clonal outbreak of equine respiratory disease. *Clin Vaccine Immunol* 20:1181–1188 <http://dx.doi.org/10.1128/CVI.00069-13>.
104. Bergmann S, Eichhorn I, Kohler TP, Hammerschmidt S, Goldmann O, Rohde M, Fulde M. 2017. SCM, the M protein of *Streptococcus canis* binds immunoglobulin G. *Front Cell Infect Microbiol* 7:80 <http://dx.doi.org/10.3389/fcimb.2017.00080>.
105. Vasi J, Frykberg L, Carlsson LE, Lindberg M, Guss B. 2000. M-like proteins of *Streptococcus dysgalactiae*. *Infect Immun* 68:294–302 <http://dx.doi.org/10.1128/IAI.68.1.294-302.2000>.
106. Lynskey NN, Reglinski M, Calay D, Siggins MK, Mason JC, Botto M, Sriskandan S. 2017. Multi-functional mechanisms of immune evasion by the streptococcal complement inhibitor C5a peptidase. *PLoS Pathog* 13:e1006493 <http://dx.doi.org/10.1371/journal.ppat.1006493>.
107. Franken C, Haase G, Brandt C, Weber-Heynemann J, Martin S, Lämmler C, Podbielski A, Lütticken R, Spellerberg B. 2001. Horizontal gene transfer and host specificity of beta-haemolytic streptococci: the role of a putative composite transposon containing scpB and lmb. *Mol Microbiol* 41:925–935 <http://dx.doi.org/10.1046/j.1365-2958.2001.02563.x>.
108. Cleary PP, Peterson J, Chen C, Nelson C. 1991. Virulent human strains of group G streptococci express a C5a peptidase enzyme similar to that produced by group A streptococci. *Infect Immun* 59:2305–2310.
109. Sriprakash KS, Hartas J. 1996. Lateral genetic transfers between group A and G streptococci for M-like genes are ongoing. *Microb Pathog* 20:275–285 <http://dx.doi.org/10.1006/mpat.1996.0026>.
110. Kurupati P, Turner CE, Tziona I, Lawrenson RA, Alam FM, Nohadani M, Stamp GW, Zinkernagel AS, Nizet V, Edwards RJ, Sriskandan S. 2010. Chemokine-cleaving *Streptococcus pyogenes* protease SpyCEP is necessary and sufficient for bacterial dissemination within soft tissues and the respiratory tract. *Mol Microbiol* 76:1387–1397 <http://dx.doi.org/10.1111/j.1365-2958.2010.07065.x>.
111. Edwards RJ, Taylor GW, Ferguson M, Murray S, Rendell N, Wrigley A, Bai Z, Boyle J, Finney SJ, Jones A, Russell HH, Turner C, Cohen J, Faulkner L, Sriskandan S. 2005. Specific C-terminal cleavage and inactivation of interleukin-8 by invasive disease isolates of *Streptococcus pyogenes*. *J Infect Dis* 192:783–790 <http://dx.doi.org/10.1086/432485>.
112. Turner CE, Kurupati P, Wiles S, Edwards RJ, Sriskandan S. 2009. Impact of immunization against SpyCEP during invasive disease with two streptococcal species: *Streptococcus pyogenes* and *Streptococcus equi*. *Vaccine* 27:4923–4929 <http://dx.doi.org/10.1016/j.vaccine.2009.06.042>.
113. Malke H. 2000. Genetics and pathogenicity factors of group C and group G streptococci, p 163–176. In Fischetti VA, Novick RP, Ferretti JJ, Portnoy DA, Rood JI (ed), *Gram-Positive Pathogens*. ASM Press, Washington, DC.
114. Navarre WW, Schneewind O. 1999. Surface proteins of Gram-positive bacteria and mechanisms of their targeting to the cell wall envelope. *Microbiol Mol Biol Rev* 63:174–229.
115. Sjöbring U, Björck L, Kastern W. 1991. Streptococcal protein G. Gene structure and protein binding properties. *J Biol Chem* 266:399–405.
116. Guss B, Eliasson M, Olsson A, Uhlén M, Frej AK, Jörnvall H, Flock JI, Lindberg M. 1986. Structure of the IgG-binding regions of streptococcal protein G. *EMBO J* 5:1567–1575 <http://dx.doi.org/10.1002/j.1460-2075.1986.tb04398.x>.
117. Byeon IJ, Louis JM, Gronenborn AM. 2003. A protein contortionist: core mutations of GB1 that induce dimerization and domain swapping. *J Mol Biol* 333:141–152 [http://dx.doi.org/10.1016/S0022-2836\(03\)00928-8](http://dx.doi.org/10.1016/S0022-2836(03)00928-8).
118. Ding K, Louis JM, Gronenborn AM. 2004. Insights into conformation and dynamics of protein GB1 during folding and unfolding by NMR. *J Mol Biol* 335:1299–1307 <http://dx.doi.org/10.1016/j.jmb.2003.11.042>.
119. Gronenborn AM, Filpula DR, Essig NZ, Achari A, Whitlow M, Wingfield PT, Clore GM. 1991. A novel, highly stable fold of the immunoglobulin binding domain of streptococcal protein G. *Science* 253:657–661 <http://dx.doi.org/10.1126/science.1871600>.
120. Jonsson H, Müller HP. 1994. The type-III Fc receptor from *Streptococcus dysgalactiae* is also an alpha 2-macroglobulin receptor. *Eur J Biochem* 220:819–826 <http://dx.doi.org/10.1111/j.1432-1033.1994.tb18684.x>.
121. Jonsson H, Burtsoff-Asp C, Guss B. 1995. Streptococcal protein MAG: a protein with broad albumin binding specificity. *Biochim Biophys Acta* 1249:65–71 [http://dx.doi.org/10.1016/0167-4838\(95\)00065-3](http://dx.doi.org/10.1016/0167-4838(95)00065-3).
122. Jonsson H, Frykberg L, Rantamäki L, Guss B. 1994. MAG, a novel plasma protein receptor from *Streptococcus dysgalactiae*. *Gene* 143:85–89 [http://dx.doi.org/10.1016/0378-1119\(94\)90609-2](http://dx.doi.org/10.1016/0378-1119(94)90609-2).
123. Jonsson H, Lindmark H, Guss B. 1995. A protein G-related cell surface protein in *Streptococcus zooepidemicus*. *Infect Immun* 63:2968–2975.
124. Song XM, Perez-Casal J, Bolton A, Potter AA. 2001. Surface-expressed Mig protein protects *Streptococcus dysgalactiae* against phagocytosis by bovine neutrophils. *Infect Immun* 69:6030–6037 <http://dx.doi.org/10.1128/IAI.69.10.6030-6037.2001>.
125. Song XM, Perez-Casal J, Fontaine MC, Potter AA. 2002. Bovine immunoglobulin A (IgA)-binding activities of the surface-expressed Mig protein of *Streptococcus dysgalactiae*. *Microbiology* 148:2055–2064 <http://dx.doi.org/10.1099/00221287-148-7-2055>.
126. Song XM, Perez-Casal J, Potter AA. 2004. The Mig protein of *Streptococcus dysgalactiae* inhibits bacterial internalization into bovine mammary gland epithelial cells. *FEMS Microbiol Lett* 231:33–38 [http://dx.doi.org/10.1016/S0378-1097\(03\)00923-6](http://dx.doi.org/10.1016/S0378-1097(03)00923-6).
127. von Pawel-Rammingen U, Johansson BP, Björck L. 2002. IdeS, a novel streptococcal cysteine proteinase with unique specificity for immunoglobulin G. *EMBO J* 21:1607–1615 <http://dx.doi.org/10.1093/emboj/21.7.1607>.
128. Collin M, Olsén A. 2001. EndoS, a novel secreted protein from *Streptococcus pyogenes* with endoglycosidase activity on human IgG. *EMBO J* 20:3046–3055 <http://dx.doi.org/10.1093/emboj/20.12.3046>.
129. Lannergård J, Guss B. 2006. IdeE, an IgG-endopeptidase of *Streptococcus equi* ssp. *equi*. *FEMS Microbiol Lett* 262:230–235 <http://dx.doi.org/10.1111/j.1574-6968.2006.00404.x>.

130. Hulting G, Flock M, Frykberg L, Lannergård J, Flock JI, Guss B. 2009. Two novel IgG endopeptidases of *Streptococcus equi*. *FEMS Microbiol Lett* 298:44–50 <http://dx.doi.org/10.1111/j.1574-6968.2009.01698.x>.
131. Flock M, Frykberg L, Sköld M, Guss B, Flock JI. 2012. Antiphagocytic function of an IgG glycosyl hydrolase from *Streptococcus equi* subsp. *equi* and its use as a vaccine component. *Infect Immun* 80:2914–2919 <http://dx.doi.org/10.1128/IAI.06083-11>.
132. Shadnezhad A, Naegeli A, Sjögren J, Adamczyk B, Leo F, Allhorn M, Karlsson NG, Jensen A, Collin M. 2016. EndoSd: an IgG glycan hydrolyzing enzyme in *Streptococcus dysgalactiae* subspecies *dysgalactiae*. *Future Microbiol* 11:721–736 <http://dx.doi.org/10.2217/fmb.16.14>.
133. Rungelrath V, Wohlsein JC, Siebert U, Stott J, Prenger-Berninghoff E, von Pawel-Rammingen U, Valentin-Weigand P, Baums CG, Seele J. 2017. Identification of a novel host-specific IgG protease in *Streptococcus phocae* subsp. *phocae*. *Vet Microbiol* 201:42–48 <http://dx.doi.org/10.1016/j.vetmic.2017.01.009>.
134. Ben Nasr A, Wistedt A, Ringdahl U, Sjöbring U. 1994. Streptokinase activates plasminogen bound to human group C and G streptococci through M-like proteins. *Eur J Biochem* 222:267–276 <http://dx.doi.org/10.1111/j.1432-1033.1994.tb18865.x>.
135. Lottenberg R, Minning-Wenz D, Boyle MD. 1994. Capturing host plasmin(ogen): a common mechanism for invasive pathogens? *Trends Microbiol* 2:20–24 [http://dx.doi.org/10.1016/0966-842X\(94\)90340-9](http://dx.doi.org/10.1016/0966-842X(94)90340-9).
136. Wang H, Lottenberg R, Boyle MD. 1995. A role for fibrinogen in the streptokinase-dependent acquisition of plasmin(ogen) by group A streptococci. *J Infect Dis* 171:85–92 <http://dx.doi.org/10.1093/infdis/171.1.85>.
137. Tewodros W, Karlsson I, Kronvall G. 1996. Allelic variation of the streptokinase gene in beta-hemolytic streptococci group C and G isolates of human origin. *FEMS Immunol Med Microbiol* 13:29–34.
138. McArthur JD, McKay FC, Ramachandran V, Shyam P, Cork AJ, Sanderson-Smith ML, Cole JN, Ringdahl U, Sjöbring U, Ranson M, Walker MJ. 2008. Allelic variants of streptokinase from *Streptococcus pyogenes* display functional differences in plasminogen activation. *FASEB J* 22:3146–3153 <http://dx.doi.org/10.1096/fj.08-109348>.
139. Keramati M, Roohvand F, Eslaminejad Z, Mirzaie A, Nikbin VS, Aslani MM. 2012. PCR/RFLP-based allelic variants of streptokinase and their plasminogen activation potencies. *FEMS Microbiol Lett* 335:79–85 <http://dx.doi.org/10.1111/j.1574-6968.2012.02640.x>.
140. McCoy HE, Broder CC, Lottenberg R. 1991. Streptokinases produced by pathogenic group C streptococci demonstrate species-specific plasminogen activation. *J Infect Dis* 164:515–521 <http://dx.doi.org/10.1093/infdis/164.3.515>.
141. Schroeder B, Boyle MD, Sheerin BR, Asbury AC, Lottenberg R. 1999. Species specificity of plasminogen activation and acquisition of surface-associated proteolytic activity by group C streptococci grown in plasma. *Infect Immun* 67:6487–6495.
142. Andreoni F, Ugolini F, Keller N, Neff A, Nizet V, Hollands A, Marques Maggio E, Zinkernagel AS, Schuepbach RA. 2017. Immunoglobulin attenuates streptokinase-mediated virulence in *Streptococcus dysgalactiae* subsp. *equisimilis* necrotizing fasciitis. *J Infect Dis* 217:270–279.
143. Siemens N, Kittang BR, Chakrakodi B, Oppegaard O, Johansson L, Bruun T, Mylvaganam H, Svensson M, Skrede S, Norrby-Teglund A, INFECT Study Group. 2015. Increased cytotoxicity and streptolysin O activity in group G streptococcal strains causing invasive tissue infections. *Sci Rep* 5:16945 <http://dx.doi.org/10.1038/srep16945>.
144. Billington SJ, Jost BH, Songer JG. 2000. Thiol-activated cytolysins: structure, function and role in pathogenesis. *FEMS Microbiol Lett* 182:197–205 [http://dx.doi.org/10.1016/S0378-1097\(99\)00536-4](http://dx.doi.org/10.1016/S0378-1097(99)00536-4).
145. Okumura K, Hara A, Tanaka T, Nishiguchi I, Minamide W, Igarashi H, Yutsudo T. 1994. Cloning and sequencing the streptolysin O genes of group C and group G streptococci. *DNA Seq* 4:325–328 <http://dx.doi.org/10.3109/10425179409020859>.
146. Magassa N, Chandrasekaran S, Caparon MG. 2010. *Streptococcus pyogenes* cytolysin-mediated translocation does not require pore formation by streptolysin O. *EMBO Rep* 11:400–405 <http://dx.doi.org/10.1038/embor.2010.37>.
147. Barnett TC, Liebl D, Seymour LM, Gillen CM, Lim JY, Larock CN, Davies MR, Schulz BL, Nizet V, Teasdale RD, Walker MJ. 2013. The globally disseminated M1T1 clone of group A *Streptococcus* evades autophagy for intracellular replication. *Cell Host Microbe* 14:675–682 <http://dx.doi.org/10.1016/j.chom.2013.11.003>.
148. O'Neill AM, Thurston TL, Holden DW. 2016. Cytosolic replication of group A *Streptococcus* in human macrophages. *MBio* 7:e00020-16.
149. O'Seaghda M, Wessels MR. 2013. Streptolysin O and its co-toxin NAD-glycohydrolase protect group A *Streptococcus* from xenophagic killing. *PLoS Pathog* 9:e1003394 <http://dx.doi.org/10.1371/journal.ppat.1003394>.
150. Sharma O, O'Seaghda M, Velarde JJ, Wessels MR. 2016. NAD⁺-glycohydrolase promotes intracellular survival of group A *Streptococcus*. *PLoS Pathog* 12:e1005468 <http://dx.doi.org/10.1371/journal.ppat.1005468>.
151. Uchiyama S, Döhrmann S, Timmer AM, Dixit N, Ghojani M, Bhandari T, Timmer JC, Sprague K, Bubeck-Wardenburg J, Simon SI, Nizet V. 2015. Streptolysin O rapidly impairs neutrophil oxidative burst and antibacterial responses to group A *Streptococcus*. *Front Immunol* 6:581 <http://dx.doi.org/10.3389/fimmu.2015.00581>.
152. Flanagan J, Collin N, Timoney J, Mitchell T, Mumford JA, Chanter N. 1998. Characterization of the haemolytic activity of *Streptococcus equi*. *Microb Pathog* 24:211–221 <http://dx.doi.org/10.1006/mpat.1997.0190>.
153. Nizet V. 2002. Streptococcal beta-hemolysins: genetics and role in disease pathogenesis. *Trends Microbiol* 10:575–580 [http://dx.doi.org/10.1016/S0966-842X\(02\)02473-3](http://dx.doi.org/10.1016/S0966-842X(02)02473-3).
154. Humar D, Datta V, Bast DJ, Beall B, De Azavedo JC, Nizet V. 2002. Streptolysin S and necrotising infections produced by group G *streptococcus*. *Lancet* 359:124–129 [http://dx.doi.org/10.1016/S0140-6736\(02\)07371-3](http://dx.doi.org/10.1016/S0140-6736(02)07371-3).
155. Nizet V, Beall B, Bast DJ, Datta V, Kilburn L, Low DE, De Azavedo JC. 2000. Genetic locus for streptolysin S production by group A *streptococcus*. *Infect Immun* 68:4245–4254 <http://dx.doi.org/10.1128/IAI.68.7.4245-4254.2000>.
156. Steiner K, Malke H. 2002. Dual control of streptokinase and streptolysin S production by the *covRS* and *fasCAX* two-component regulators in *Streptococcus dysgalactiae* subsp. *equisimilis*. *Infect Immun* 70:3627–3636 <http://dx.doi.org/10.1128/IAI.70.7.3627-3636.2002>.
157. Zeppa JJ, Kasper KJ, Mohorovic I, Mazzuca DM, Haeryfar SMM, McCormick JK. 2017. Nasopharyngeal infection by *Streptococcus pyogenes* requires superantigen-responsive V β -specific T cells. *Proc Natl Acad Sci U S A* 114:10226–10231.
158. Commons RJ, Smeesters PR, Proft T, Fraser JD, Robins-Browne R, Curtis N. 2014. Streptococcal superantigens: categorization and clinical associations. *Trends Mol Med* 20:48–62 <http://dx.doi.org/10.1016/j.molmed.2013.10.004>.
159. Sachse S, Seidel P, Gerlach D, Günther E, Rödel J, Straube E, Schmidt KH. 2002. Superantigen-like gene(s) in human pathogenic *Streptococcus dysgalactiae*, subsp. *equisimilis*: genomic localisation of the gene encoding streptococcal pyrogenic exotoxin G (*speG(dys)*). *FEMS Immunol Med Microbiol* 34:159–167 <http://dx.doi.org/10.1111/j.1574-695X.2002.tb00618.x>.
160. Brandt CM, Schweizer KG, Holland R, Lütticken R, Freyaldenhoven BS. 2005. Lack of mitogenic activity of *speG*- and *speG(dys)*-positive *Streptococcus dysgalactiae* subspecies *equisimilis* isolates from patients with invasive infections. *Int J Med Microbiol* 295:539–546 <http://dx.doi.org/10.1016/j.ijmm.2005.07.013>.
161. Okumura K, Shimomura Y, Murayama SY, Yagi J, Ubukata K, Kirikae T, Miyoshi-Akiyama T. 2012. Evolutionary paths of streptococ-

- cal and staphylococcal superantigens. *BMC Genomics* 13:404 <http://dx.doi.org/10.1186/1471-2164-13-404>.
162. Korem M, Hidalgo-Grass C, Michael-Gayego A, Nir-Paz R, Salameh S, Moses AE. 2014. Streptococcal pyrogenic exotoxin G gene in blood and pharyngeal isolates of *Streptococcus dysgalactiae* subspecies *equisimilis* has a limited role in pathogenesis. *J Microbiol Immunol Infect* 47:292–296 <http://dx.doi.org/10.1016/j.jmii.2012.12.003>.
163. Zhao J, Hayashi T, Saarinen S, Papageorgiou AC, Kato H, Imanishi K, Kirikae T, Abe R, Uchiyama T, Miyoshi-Akiyama T. 2007. Cloning, expression, and characterization of the superantigen streptococcal pyrogenic exotoxin G from *Streptococcus dysgalactiae*. *Infect Immun* 75:1721–1729 <http://dx.doi.org/10.1128/IAI.01183-06>.
164. Kalia A, Bessen DE. 2003. Presence of streptococcal pyrogenic exotoxin A and C genes in human isolates of group G streptococci. *FEMS Microbiol Lett* 219:291–295 [http://dx.doi.org/10.1016/S0378-1097\(03\)00022-3](http://dx.doi.org/10.1016/S0378-1097(03)00022-3).
165. Traverso F, Blanco A, Villalón P, Beratz N, Sáez Nieto JA, Lopardo H, National Collaborative Group for the Study of Streptococci and Related Bacteria. 2016. Molecular characterization of invasive *Streptococcus dysgalactiae* subsp. *equisimilis*. Multicenter study: Argentina 2011–2012. *Rev Argent Microbiol* 48:279–289 <http://dx.doi.org/10.1016/j.ram.2016.07.001>.
166. Anand TD, Rajesh T, Rajendhran J, Gunasekaran P. 2012. Superantigen profiles of *emm* and *emm*-like typeable and nontypeable pharyngeal streptococcal isolates of South India. *Ann Clin Microbiol Antimicrob* 11:3 <http://dx.doi.org/10.1186/1476-0711-11-3>.
167. Artiushin SC, Timoney JF, Sheoran AS, Muthupalani SK. 2002. Characterization and immunogenicity of pyrogenic mitogens SePE-H and SePE-I of *Streptococcus equi*. *Microb Pathog* 32:71–85 <http://dx.doi.org/10.1006/mpat.2001.0482>.
168. Alber J, El-Sayed A, Estoepangestie S, Lämmler C, Zschöck M. 2005. Dissemination of the superantigen encoding genes *seeL*, *seeM*, *szeL* and *szeM* in *Streptococcus equi* subsp. *equi* and *Streptococcus equi* subsp. *zooepidemicus*. *Vet Microbiol* 109:135–141 <http://dx.doi.org/10.1016/j.vetmic.2005.05.001>.
169. Profit T, Webb PD, Handley V, Fraser JD. 2003. Two novel superantigens found in both group A and group C *Streptococcus*. *Infect Immun* 71:1361–1369 <http://dx.doi.org/10.1128/IAI.71.3.1361-1369.2003>.
170. Paillot R, Darby AC, Robinson C, Wright NL, Steward KF, Anderson E, Webb K, Holden MT, Efstratiou A, Broughton K, Jolley KA, Priestnall SL, Marotti Campi MC, Hughes MA, Radford A, Erles K, Waller AS. 2010. Identification of three novel superantigen-encoding genes in *Streptococcus equi* subsp. *zooepidemicus*, *szeF*, *szeN*, and *szeP*. *Infect Immun* 78:4817–4827 <http://dx.doi.org/10.1128/IAI.00751-10>.
171. Miyoshi-Akiyama T, Zhao J, Kato H, Kikuchi K, Totsuka K, Kataoka Y, Katsumi M, Uchiyama T. 2003. *Streptococcus dysgalactiae*-derived mitogen (SDM), a novel bacterial superantigen: characterization of its biological activity and predicted tertiary structure. *Mol Microbiol* 47:1589–1599 <http://dx.doi.org/10.1046/j.1365-2958.2003.03411.x>.
172. Lefébure T, Richards VP, Lang P, Pavinski-Bitar P, Stanhope MJ. 2012. Gene repertoire evolution of *Streptococcus pyogenes* inferred from phylogenomic analysis with *Streptococcus canis* and *Streptococcus dysgalactiae*. *PLoS One* 7:e37607 <http://dx.doi.org/10.1371/journal.pone.0037607>.
173. Igwe EI, Shewmaker PL, Facklam RR, Farley MM, van Beneden C, Beall B. 2003. Identification of superantigen genes *speM*, *ssa*, and *smeZ* in invasive strains of beta-hemolytic group C and G streptococci recovered from humans. *FEMS Microbiol Lett* 229:259–264 [http://dx.doi.org/10.1016/S0378-1097\(03\)00842-5](http://dx.doi.org/10.1016/S0378-1097(03)00842-5).