

Mixed QCD \otimes QED corrections to on-shell Z boson production at the LHC

Maximilian Delto,^a Matthieu Jaquier,^a Kirill Melnikov^a and Raoul Röntsch^b

^a*Institute for Theoretical Particle Physics,
KIT, Karlsruhe, Germany*

^b*Theoretical Physics Department, CERN,
1211 Geneva 23, Switzerland*

E-mail: maximilian.delto@kit.edu, matthieu.jaquier@kit.edu,
kirill.melnikov@kit.edu, raoul.rontsch@cern.ch

ABSTRACT: We compute the mixed QCD \otimes QED corrections to the production of on-shell Z bosons at the LHC at a fully-exclusive level. We also include the factorised NLO QCD correction to Z boson production and NLO QED correction to Z boson decay into two leptons. We make use of an abelianised version of the nested soft-collinear subtraction formalism to perform this computation. We study the phenomenological impact of the mixed QCD \otimes QED corrections for a number of observables relevant for LHC phenomenology.

KEYWORDS: QCD Phenomenology

ARXIV EPRINT: [1909.08428](https://arxiv.org/abs/1909.08428)

Contents

1	Introduction	1
2	Technical aspects of the calculation	3
2.1	Preliminary remarks	4
2.2	The mapping of colour factors	4
2.3	Implementation of Z boson decay	7
2.4	Checks of the computation	8
3	Results	9
3.1	Inclusive cross sections	9
3.2	Fiducial cross sections	10
3.3	Kinematic distributions	12
4	Conclusion and outlook	15

1 Introduction

The recent years have seen a transition of the Large Hadron Collider (LHC) from a discovery machine to a precision machine. The reason for this is the absence of a direct observation of New Physics which, even after the discovery of the Higgs boson [1, 2], is needed to clarify questions left open by the Standard Model. Forthcoming searches for physics Beyond the Standard Model (BSM) will focus on systematic studies of possible small deviations from Standard Model predictions in precision observables. A reliable theoretical description of such observables within the Standard Model is an important prerequisite for the success of this research program.

A case in point is the hadronic production of charged leptons via a virtual photon and/or the Z boson, the celebrated Drell-Yan (DY) process [3] (see [4] for a review). Its high production rate and distinct signature make it extremely useful for luminosity monitoring [5–7] and detector calibration [8]. Being theoretically well-understood, this process is also suited for electroweak (EW) precision physics, such as the measurement of the weak mixing angle [8, 9]. Moreover, DY production is used in parton distribution function (PDF) fits [10–13] and for searches for New Physics at high energies [14]. In such analyses, the rapidity distribution of the Z boson and the dilepton invariant mass, respectively, are of particular interest.

The inclusive next-to-leading order (NLO) QCD corrections to Drell-Yan production were first computed four decades ago [15]. Inclusive results at next-to-next-to-leading order (NNLO) in QCD have also been known for many years [16–18]. Arbitrary infrared safe

kinematic distributions are also available through NNLO QCD accuracy [19–24]. In addition, threshold effects at next-to-next-to-next-to leading order (N³LO) have been studied in refs. [25, 26]. EW corrections to $pp \rightarrow \ell^+\ell^-$ were computed in refs. [27–32].

Recently, an important milestone in the quest for high precision theoretical predictions for LHC processes has been reached with the calculation of Higgs boson production in hadronic collisions at N³LO QCD [33]. Since techniques developed in the course of that calculation put the N³LO QCD corrections to the Drell-Yan process within reach, it becomes important to know the mixed QCD-EW $\mathcal{O}(\alpha_s\alpha)$ corrections as well since, based on the sizes of strong and EW coupling constants, one expects both contributions to be comparable in magnitude. The computation of mixed QCD-EW $\mathcal{O}(\alpha_s\alpha)$ corrections requires the evaluation of complicated two-loop diagrams with up to two massive propagators, as well as the respective real-virtual and double-real contributions where a photon and/or a parton is emitted in the final state. All these contributions contain intertwined QCD and QED singularities, which need to be extracted and cancelled properly.

Several ingredients required for the calculation of $\mathcal{O}(\alpha_s\alpha)$ corrections to DY production have already appeared in the literature. In ref. [34] integrated double-real contributions to the production of a single on-shell gauge boson have been computed using the method of reverse unitarity [35]. Furthermore, the two-loop master integrals needed for the double-virtual contributions were recently presented in ref. [36]. Nevertheless, up to now, the various ingredients have not been combined in a way that allows one to compute physical observables.

Given the absence of the full calculation, different approximations have been used in the past to estimate mixed QCD \otimes EW corrections. In ref. [37] NNLO QCD corrections have been combined additively with the NLO EW ones. Ref. [38] has included the matching of the NLO QCD and EW corrections to QCD parton showers and multiple photon emissions.

Although the generic Drell-Yan process $pp \rightarrow \ell^+\ell^-$ is the target of many experimental analyses, the theoretical description of the *on-shell* production of Z bosons $pp \rightarrow Z \rightarrow \ell^+\ell^-$ offers significant simplifications. Indeed, for an on-shell Z boson, virtual and real contributions that connect incoming partons and outgoing leptons are suppressed by the ratio of the Z boson width to its mass, Γ_Z/M_Z . The EW corrections to the production of a lepton pair via an on-shell Z boson can thus be separated into gauge-invariant subsets according to whether the correction is associated with the production of the Z boson (initial) or with its decay (final). Similarly, mixed QCD \otimes EW corrections can be divided into an initial-initial and an initial-final contribution. It was shown in ref. [39] that these results follow from the systematic expansion of the production cross section around the resonance pole. Such an expansion was used in ref. [39] to compute the first subleading non-factorisable contribution, which was found to be negligible. In the same reference, the fact that the NLO EW corrections originating from the initial state are smaller than those arising from the final state was used to argue that the mixed QCD-EW initial-initial correction should be smaller than the initial-final one. The latter contribution was computed in ref. [40] in the resonance region.

Recently, mixed QCD \otimes QED corrections to the inclusive production of an on-shell Z boson in hadronic collisions have been computed in ref. [41]. These corrections provide

a gauge-invariant subset of the initial-initial QCD \otimes EW corrections; they can be obtained from the known NNLO QCD corrections to on-shell Z boson production through an abelianisation procedure.¹ The mixed QCD \otimes QED corrections computed in ref. [41] turned out to be quite significant at the LHC, being smaller than the NNLO QCD corrections by only a factor of three. This rather modest suppression of the initial-initial QCD \otimes QED corrections to the inclusive cross section relative to NNLO QCD corrections makes it interesting to study the mixed corrections to more exclusive observables.

In addition, the computation of mixed QCD \otimes QED corrections to Z boson production is an important step towards the calculation of the such corrections to the production of an on-shell W boson at the LHC, which is of high relevance for the W -mass determination. Indeed, while interactions of W bosons with photons introduce additional subtleties in the computation of such corrections compared to the Z boson case, understanding the infrared structure of mixed corrections in $pp \rightarrow Z$ is a prerequisite for the analysis of mixed corrections to $pp \rightarrow W^\pm$.

The goal of this paper is to present the calculation of the fully-differential mixed QCD \otimes QED corrections to the production of an on-shell Z boson in hadronic collisions (the *initial-initial* corrections). This contribution features the most complex structure of infrared singularities and, for this reason, represents an important step towards the computation of full QCD \otimes EW corrections to Z boson production. In addition to mixed initial-initial corrections, we also compute *initial-final* corrections that arise through an interplay of QCD corrections to Z production and QED corrections to its decay. Comparisons of the two contributions for various observables will allow us to quantify the degree of dominance of the initial-final corrections over the initial-initial ones.

The calculation is performed by extending the nested soft-collinear subtraction scheme presented in ref. [43] for NNLO QCD computations through the abelianisation procedure of ref. [41]. We make use of the NNPDF3.1luxQED PDF set [44–46], whose evolution is correct through $\mathcal{O}(\alpha_s\alpha)$; this enables us to remove collinear singularities from initial state radiation in a consistent way. We use the resulting code to study the impact of the QCD \otimes QED corrections on several distributions of phenomenological interest, including the transverse momentum and the rapidity spectra of the Z boson, the transverse momentum distributions of leptons and distributions in one of the so-called Collins-Soper angles θ^* [47].

The rest of this paper is organised as follows. In section 2 we briefly summarise technical details of the calculation and explain how a transition from NNLO QCD to mixed QCD \otimes QED corrections is accomplished. In section 3 we present phenomenological results. We conclude in section 4.

2 Technical aspects of the calculation

Our goal is to compute mixed QCD \otimes QED corrections starting from the existing implementation of NNLO QCD corrections to the on-shell Z boson production at a fully-differential level [48]. We describe the relevant technical aspects of this calculation in this section.

¹Such a procedure has already been considered earlier, see e.g. ref. [42].

2.1 Preliminary remarks

As we mentioned in the introduction, the mixed QCD \otimes QED corrections to the production cross section of an on-shell Z boson and its subsequent leptonic decay can be divided into *initial-initial* and *initial-final* contributions, while the interference between production and decay sub-processes is suppressed by the ratio of the Z boson width to its mass, Γ_Z/M_Z [39]. The required initial-final matrix elements can be constructed from the $\mathcal{O}(\alpha_s)$ and $\mathcal{O}(\alpha)$ helicity amplitudes for the production and decay sub-processes, respectively. The infrared singularities arising from these corrections can be handled using standard NLO techniques; we employ the Frixione-Kunszt-Signer subtraction scheme [49, 50] to deal with these. The only subtlety is that spin correlations between production and decay processes caused by the spin-one nature of the intermediate Z boson need to be properly accounted for.

The initial-initial corrections pose a greater challenge. The main ingredients required for the computation of these corrections are:

- the tree-level matrix elements for the *parton-initiated* processes $q\bar{q} \rightarrow Z + \gamma + g$, $q\bar{q} \rightarrow Z + q + \bar{q}$, $qq \rightarrow Z + q + q$ and $qg \rightarrow Z + q + \gamma$;
- the tree-level matrix elements for the *photon-initiated* processes $q\gamma \rightarrow Z + q + g$ and $g\gamma \rightarrow Z + q + \bar{q}$;
- the matrix elements for the one-loop QED correction to the *parton-initiated* processes $q\bar{q} \rightarrow Z + g$ and $qg \rightarrow Z + q$;
- the matrix elements for the one-loop QCD correction to the *parton-initiated* process $q\bar{q} \rightarrow Z + \gamma$;
- the matrix elements for the one-loop QCD correction to the *photon-initiated* process $q\gamma \rightarrow Z + q$;
- the matrix elements for the two-loop mixed QCD \otimes QED correction to $q\bar{q} \rightarrow Z$ production.

All these matrix elements contain infrared singularities due to soft and/or collinear emissions of gluons, photons and quark-antiquark pairs. These singularities have to be regularised and removed in an appropriate subtraction scheme, yielding a fully-differential description of on-shell Z boson production suitable for numerical integration. We achieve this goal by abelianising the NNLO QCD calculation of Z boson production performed within the nested soft-collinear subtraction scheme [43, 48, 51–53]. The abelianisation procedure was recently described in ref. [41]; in essence this is a set of rules that allows one to replace the SU(3) colour factors in the NNLO QCD formulas in such a way that the computation of mixed QCD \otimes QED corrections becomes possible. We discuss these replacement rules in the next section.

2.2 The mapping of colour factors

There are three SU(3) colour factors, C_F^2 , $C_A C_F$ and $C_F T_R$, which appear in NNLO QCD corrections to Z boson production. For the purpose of turning a NNLO QCD computa-



Figure 1. Contributions to the colour factors $C_F T_R$ (left) and $C_A C_F$ (right).

tion into a computation of QED \otimes QCD corrections, these colour factors require different modifications. We discuss them in turn.

We first consider the NNLO QCD computation of Z boson production from either a quark-antiquark or a quark-quark initial state. The colour factor $C_F T_R$ appears in diagrams with two disjoint quark lines, see e.g. figure 1(a). When such diagrams are squared and sums over colours of initial- and final-state particles are computed, two independent colour traces appear. These colour traces are of the form $\text{Tr}(T^a T^b) \text{Tr}(T^b T^a) = C_F T_R$. When a gluon is replaced by a photon in these diagrams the colour traces become $\text{Tr}(T^a) \text{Tr}(T^a)$ and vanish. Similarly, partonic processes $q_i q_j \rightarrow Z + q_i + q_j$ for $i \neq j$ that contribute at NNLO QCD become irrelevant for $\mathcal{O}(\alpha\alpha_s)$ corrections. We note that the consequence of that is the absence of terms proportional to products of two different electric quark charges in mixed QCD \otimes QED contributions. We conclude that NNLO QCD contributions proportional to the colour factor $C_F T_R$ have no counter-parts in the computation of QED \otimes QCD corrections and need to be removed. We achieve this by setting T_R to zero in the expressions for NNLO QCD corrections provided in ref. [48].

The case of the colour factor $C_A C_F$ is very similar. The colour factors C_A originate either from diagrams with three-gluon vertices (see figure 1(b)) or from the non-commutative nature of generators of SU(3) colour algebra in the fundamental representation. Neither of these issues apply to the case of mixed QCD \otimes QED corrections. The corresponding contributions can be eliminated by setting C_A to zero in the NNLO QCD computation.

Finally, we need to understand how C_F^2 colour factors should be modified for the purpose of computing mixed QCD \otimes QED corrections. We consider a collision of a quark q with an anti-quark \bar{q} , assume that the electric charge of the quark is e_q , and discuss a few illustrative examples.

Consider the double-virtual corrections, shown on the top line of figure 2. Upon setting C_A to zero, colour traces that appear in both planar and non-planar diagrams provide a colour factor C_F^2 . Since any of the two gluon lines can be replaced by a photon line in any of these diagrams, the required modification of the colour factor is $C_F^2 \rightarrow 2C_F e_q^2$. It is easy to see that the same holds for real-virtual contributions, shown on the second line of figure 2, and for interferences that arise between double-real contributions (see the final line figure 2).

A distinct situation arises in cases when two gluons appear in the final state, shown in figure 3. In this case, two diagrams in the QCD case are mapped onto two diagrams in the QCD \otimes QED case; hence, it appears at first sight that for these diagrams the correct




Figure 2. Contributions to the colour factor C_F^2 and their abelianised counterparts.




Figure 3. Contributions to the colour factor C_F^2 and their abelianised counterparts.

replacement rule is $C_F^2 \rightarrow C_F e_q^2$, so that the factor of two is missing. However, this is not the case because contributions of diagrams with two gluons to the cross section are multiplied by a factor $1/2!$ to account for the symmetric final state. Clearly, there is no such factor in case of the $g + \gamma$ final state. This mismatch is accounted for if the colour factor C_F^2 in the $q\bar{q} \rightarrow Z + g + g$ contribution is again replaced by $2C_F e_q^2$, in accord with what is needed for double-virtual and real-virtual contributions.

Hence, after an examination of all the cases, we conclude that for processes with an incoming quark-antiquark pair or interference-like contributions with two identical quarks, we replace

$$C_F^2 \rightarrow 2C_F e_q^2, \quad T_R \rightarrow 0, \quad C_A \rightarrow 0, \quad (2.1)$$

in the formulas that describe NNLO QCD corrections to Z boson production and obtain results for mixed QCD \otimes QED corrections. In eq. (2.1), e_q is the electric charge of the incoming quark.

For processes with an incoming (anti)quark and a gluon, there is no symmetry factor, and the two possible ways to replace a gluon by a photon amount to the two distinct processes $qg \rightarrow Z + q + \gamma$ and $q\gamma \rightarrow Z + q + g$. The replacement rules become

$$C_F^2 \rightarrow C_F e_q^2, \quad C_A \rightarrow 0. \quad (2.2)$$

Similarly, for processes induced by two gluons, replacing a gluon by a photon leads to the

processes $g\gamma \rightarrow Z + q\bar{q}$ and $\gamma g \rightarrow Z + q\bar{q}$. We then have the replacement rule

$$C_F^2 \rightarrow C_F e_q^2, \quad C_A \rightarrow 0. \quad (2.3)$$

We also note that, if photon-induced contributions are obtained from gluon-induced processes, the averaging over colour charges of the incoming partons has to be changed as well.

Making use of the procedure described above, we abelianised the fully-differential description of the on-shell Z boson production given in ref. [48], including regulated double-real and real-virtual contributions, integrated subtraction terms and the virtual contributions. This gives us an opportunity to compute mixed QCD \otimes QED corrections to any infrared-safe observable in the production of an on-shell Z boson.

2.3 Implementation of Z boson decay

We now discuss the treatment of the decay of the Z boson into massless leptons, $Z \rightarrow \ell^+\ell^-$, in QCD and QED perturbative expansions. At leading order, the cross section for the production of an on-shell Z boson is computed from the tree-level process $q\bar{q} \rightarrow Z \rightarrow \ell^+\ell^-$, where the square of the propagator of an intermediate Z boson is replaced by its narrow width limit

$$\frac{1}{(Q^2 - M_Z^2)^2 + M_Z^2 \Gamma_Z^2} \rightarrow \frac{\pi}{M_Z \Gamma_Z} \delta(Q^2 - M_Z^2). \quad (2.4)$$

In eq. (2.4) Q is the four-momentum of the Z boson and Γ_Z is its width. In principle, the width of the Z boson in eq. (2.4) receives perturbative corrections. These corrections should, partially, cancel QED corrections to $Z \rightarrow \ell^+\ell^-$ that are included in our calculation. In order to account for that, we rewrite the cross section for $pp \rightarrow Z \rightarrow \ell^+\ell^-$ as follows

$$d\sigma_{pp \rightarrow Z \rightarrow \ell^+\ell^-} = \frac{d\sigma_{pp \rightarrow Z} d\Gamma_{Z \rightarrow \ell^+\ell^-}}{\Gamma_Z} = \text{Br}_{Z \rightarrow \ell^+\ell^-} \times d\sigma_{pp \rightarrow Z} \times \frac{d\Gamma_{Z \rightarrow \ell^+\ell^-}}{\Gamma_{Z \rightarrow \ell^+\ell^-}}. \quad (2.5)$$

In eq. (2.5) we introduced the branching ratio of the Z boson decay to a massless $\ell^+\ell^-$ pair $\text{Br}_{Z \rightarrow \ell^+\ell^-}$; we will treat it as an experimental input parameter and will not expand it in α_s and α . However, all other terms in eq. (2.5) will be treated within QCD/QED perturbation theory. In particular, the ratio $d\Gamma_{Z \rightarrow \ell^+\ell^-} / \Gamma_{Z \rightarrow \ell^+\ell^-}$ must be expanded to first order in α . We write

$$\begin{aligned} \Gamma_{Z \rightarrow \ell^+\ell^-} &= \Gamma_{Z \rightarrow \ell^+\ell^-}^{(0)} + \Gamma_{Z \rightarrow \ell^+\ell^-}^{(1)} + \mathcal{O}(\alpha^2), \\ d\Gamma_{Z \rightarrow \ell^+\ell^-} &= d\Gamma_{Z \rightarrow \ell^+\ell^-}^{(0)} + d\Gamma_{Z \rightarrow \ell^+\ell^-}^{(1)} + \mathcal{O}(\alpha^2), \end{aligned} \quad (2.6)$$

and expand the ratio

$$\frac{d\Gamma_{Z \rightarrow \ell^+\ell^-}}{\Gamma_{Z \rightarrow \ell^+\ell^-}} = \frac{d\Gamma_{Z \rightarrow \ell^+\ell^-}^{(0)}}{\Gamma_{Z \rightarrow \ell^+\ell^-}^{(0)}} + \left[-\frac{d\Gamma_{Z \rightarrow \ell^+\ell^-}^{(0)} \Gamma_{Z \rightarrow \ell^+\ell^-}^{(1)}}{(\Gamma_{Z \rightarrow \ell^+\ell^-}^{(0)})^2} + \frac{d\Gamma_{Z \rightarrow \ell^+\ell^-}^{(1)}}{\Gamma_{Z \rightarrow \ell^+\ell^-}^{(0)}} \right] + \mathcal{O}(\alpha^2). \quad (2.7)$$

By construction, the above expression integrates to one over the unrestricted decay phase-space, so that terms in the square brackets integrate to zero. The expansion coefficients of

the leptonic Z width read

$$\Gamma_{Z \rightarrow \ell^+ \ell^-}^{(0)} = \frac{G_F M_Z^3}{6\sqrt{2}\pi} \left(\frac{1}{4} + \left(\frac{1}{2} - 2\sin^2 \theta_W \right)^2 \right), \quad (2.8)$$

$$\Gamma_{Z \rightarrow \ell^+ \ell^-}^{(1)} = \Gamma_{Z \rightarrow e^+ e^-}^{(0)} \times \frac{3\alpha}{4\pi}. \quad (2.9)$$

We now go back to eq. (2.5) and expand all relevant ingredients in series in the strong and electromagnetic coupling constants. To present the results of such an expansion, we denote the $\mathcal{O}(\alpha_s^n \alpha^m)$ contribution to the Z production cross section as $\sigma_{pp \rightarrow Z}^{(n,m)}$, and find the following results for $pp \rightarrow Z \rightarrow \ell^+ \ell^-$ cross sections²

$$d\sigma_{\text{LO}} = \text{Br}_{Z \rightarrow \ell^+ \ell^-} \times d\sigma_{pp \rightarrow Z}^{(0,0)} \times \frac{d\Gamma^{(0)}}{\Gamma^{(0)}}, \quad (2.10)$$

$$d\sigma_{\text{NLO}}^{(\alpha_s)} = \text{Br}_{Z \rightarrow \ell^+ \ell^-} \times d\sigma_{pp \rightarrow Z}^{(1,0)} \times \frac{d\Gamma^{(0)}}{\Gamma^{(0)}}, \quad (2.11)$$

$$d\sigma_{\text{NLO}}^{(\alpha)} = \text{Br}_{Z \rightarrow \ell^+ \ell^-} \times \left[d\sigma_{pp \rightarrow Z}^{(0,1)} \times \frac{d\Gamma^{(0)}}{\Gamma^{(0)}} + d\sigma_{pp \rightarrow Z}^{(0,0)} \times \left(\frac{d\Gamma^{(1)}}{\Gamma^{(0)}} - \frac{d\Gamma^{(0)}}{\Gamma^{(0)}} \times \frac{3\alpha}{4\pi} \right) \right], \quad (2.12)$$

$$d\sigma_{\text{NNLO}}^{(\alpha_s^2)} = \text{Br}_{Z \rightarrow \ell^+ \ell^-} \times d\sigma_{pp \rightarrow Z}^{(2,0)} \times \frac{d\Gamma^{(0)}}{\Gamma^{(0)}}, \quad (2.13)$$

$$d\sigma_{\text{NNLO}}^{(\alpha_s \alpha)} = \text{Br}_{Z \rightarrow \ell^+ \ell^-} \times \left[d\sigma_{pp \rightarrow Z}^{(1,1)} \times \frac{d\Gamma^{(0)}}{\Gamma^{(0)}} + d\sigma_{pp \rightarrow Z}^{(1,0)} \times \left(\frac{d\Gamma^{(1)}}{\Gamma^{(0)}} - \frac{d\Gamma^{(0)}}{\Gamma^{(0)}} \times \frac{3\alpha}{4\pi} \right) \right]. \quad (2.14)$$

Note that terms in round brackets in eqs. (2.12) and (2.14) integrate to zero over an unrestricted phase-space, such that the inclusive cross section is given by a product of the branching fraction and the production cross section, as expected from eq. (2.5). We emphasise again that we consider massless leptons throughout this paper.

To present our results for QCD \otimes QED corrections, we define ratios of contributions to the cross section at different perturbative orders. We write

$$\Delta_\alpha = \frac{\sigma_{\text{NLO}}^{(\alpha)}}{\sigma_{\text{LO}} + \sigma_{\text{NLO}}^{(\alpha_s)}}, \quad \Delta_{\alpha_s^2} = \frac{\sigma_{\text{NNLO}}^{(\alpha_s^2)}}{\sigma_{\text{LO}} + \sigma_{\text{NLO}}^{(\alpha_s)}}, \quad \Delta_{\alpha_s \alpha} = \frac{\sigma_{\text{NNLO}}^{(\alpha_s \alpha)}}{\sigma_{\text{LO}} + \sigma_{\text{NLO}}^{(\alpha_s)}}. \quad (2.15)$$

To discuss kinematic distributions, we define differential bin-by-bin corrections in a similar fashion

$$d\Delta_\alpha = \frac{d\sigma_{\text{NLO}}^{(\alpha)}}{d\sigma_{\text{LO}} + d\sigma_{\text{NLO}}^{(\alpha_s)}}, \quad d\Delta_{\alpha_s^2} = \frac{d\sigma_{\text{NNLO}}^{(\alpha_s^2)}}{d\sigma_{\text{LO}} + d\sigma_{\text{NLO}}^{(\alpha_s)}}, \quad d\Delta_{\alpha_s \alpha} = \frac{d\sigma_{\text{NNLO}}^{(\alpha_s \alpha)}}{d\sigma_{\text{LO}} + d\sigma_{\text{NLO}}^{(\alpha_s)}}. \quad (2.16)$$

2.4 Checks of the computation

Although the abelianisation procedure is, in principle, straightforward, its implementation in a fully-differential NNLO QCD computation is tedious. For this reason, it is important to check the implementation. We did this in the following way. In addition to abelianising the fully-differential NNLO QCD computation in ref. [48], we also abelianised the analytic

²From now on we drop the subscripts in $\Gamma_{Z \rightarrow \ell^+ \ell^-}$ and $d\Gamma_{Z \rightarrow \ell^+ \ell^-}$.

NNLO QCD coefficients for inclusive Z boson production given in ref. [16] and compared these to analytic expressions for mixed QCD \otimes QED corrections given in appendix B of ref. [41]. We then used our abelianised analytic expressions to compute the inclusive on-shell production cross section of $pp \rightarrow Z$ and check that it agrees with the cross section obtained using the fully-differential implementation of mixed QCD \otimes QED corrections.

3 Results

In this section we discuss mixed QCD \otimes QED corrections to various observables in $pp \rightarrow Z \rightarrow \ell^+\ell^-$ and compare them to other corrections. We consider the LHC with 13 TeV center-of-mass collision energy. We use $M_Z = 91.1876$ GeV for the mass of the Z boson, and consider its decay to a single flavour of massless leptons with a branching ratio $\text{Br}_{Z \rightarrow \ell^+\ell^-} = 0.033632$. We compute the couplings of the Z boson to leptons and quarks using $G_F = 1.16639 \times 10^{-5}$ GeV $^{-2}$, $M_W = 80.398$ GeV and $\sin^2 \theta_W = 0.2226459$ as the input parameters. We use the NNPDF3.1luxQED set with five active flavours [46] as provided by the LHAPDF interface [54], and take all quarks to be massless. To compute the Δ -corrections defined in the previous section, we always use parton distributions at NNLO accuracy to calculate all the relevant contributions. We set the renormalisation and factorisation scales to $\mu_R = \mu_F = M_Z$. The strong coupling constant is taken to be $\alpha_S(M_Z) = 0.118$ which is compatible with values provided by the NNPDF set. We describe photon interactions with leptons and quarks by the fine structure constant evaluated at the renormalisation scale $\mu = M_Z$; numerically, it is equal to $\alpha = 1/128$.

3.1 Inclusive cross sections

We use the above setup to compute the inclusive cross section of Z boson production at various (NLO QED, NNLO QCD and mixed QCD \otimes QED) approximations. Using notations introduced in the previous section, we find³

$$\Delta_\alpha = 3.2 \cdot 10^{-3}, \quad \Delta_{\alpha_s^2} = -6.4 \cdot 10^{-3}, \quad \Delta_{\alpha_s \alpha} = 2.9 \cdot 10^{-4}. \quad (3.1)$$

We note that the Δ ratios in eq. (3.1) receive contributions from corrections to the production only, since corrections to the decay of the Z boson $Z \rightarrow \ell^+\ell^-$ cancel with corrections to the partial decay width $\Gamma_{Z \rightarrow \ell^+\ell^-}$, as explained in the previous section.

It follows from eq. (3.1) that the magnitude of mixed QCD \otimes QED corrections to the inclusive cross section $pp \rightarrow Z$ is consistent with expectations. Indeed, they are smaller than NLO QED (NNLO QCD) corrections by a factor ten (twenty), respectively. These suppression factors are in accord with our expectations, based on the relative magnitude of strong and electromagnetic coupling constants.

We note that our results for the corrections to the inclusive cross section are different from the results of ref. [41]. In particular, this reference reported a smaller suppression of the mixed QCD \otimes QED corrections relative to the NNLO QCD one. The reason for this

³We neglect contributions of top quarks to Z -boson production cross section including the top-bottom triangle correction to the axial current. Such contributions were shown to be small in ref. [55].

is that ref. [41] employs a four-flavour scheme to compute both NNLO QCD and mixed QCD \otimes QED corrections. Since both NNLO QCD and mixed QCD \otimes QED corrections exhibit a strong sensitivity to input parameters, thanks to a very strong cancellation between (large) corrections to $q\bar{q}$ and qg partonic channels at the LHC energies [41], even small changes in the input can lead to significant changes in final results for corrections. We have confirmed that if we use the same input, we agree with the results of ref. [41].⁴

Finally, we comment on the scale dependence of the total NNLO cross section that includes the $\Delta_{\alpha_s^2}$ and $\Delta_{\alpha_s\alpha}$ corrections. Although this dependence is small, $\mathcal{O}(0.5 - 0.7\%)$, it is entirely dominated by pure QCD effects and it is not possible to unambiguously identify the impact of QCD \otimes QED corrections on it. For this reason, we decided to avoid presenting results for the scale dependence of the NNLO cross section. Instead, to understand the dependence of the QCD \otimes QED corrections on the scale choice, we consider the $\Delta_{\alpha_s\alpha}$ correction where the cancellation between the μ -dependence of parton distribution functions and the explicit scale dependence of the NNLO contribution cannot be expected. Because of that, it is not surprising that these corrections appear to be rather sensitive to the choice of the factorisation and the renormalisation scales μ . Indeed, by choosing the scale $\mu = [M_Z/2, M_Z, 2M_Z]$, we obtain the corrections $\Delta_{\alpha_s\alpha} = [5.6, 2.9, -0.28] \cdot 10^{-4}$. Although a stronger sensitivity of the *correction* $\Delta_{\alpha_s\alpha}$ to the choice of the scale is expected, there is yet another reason for large variations in $\Delta_{\alpha_s\alpha}$. In fact, the enhanced dependence on μ can be also traced back to a strong cancellation between quark- and gluon-initiated contributions to $\Delta_{\alpha_s\alpha}$ which is affected by the change of the scale μ in a significant way.

As we discuss in the next section, the cancellation between $q\bar{q}$ and qg channels is also an important feature of mixed corrections to fiducial cross sections. We therefore expect that for the fiducial cases the scale variation of mixed corrections will exhibit similar behaviour. For this reason, we will not discuss the scale dependences of those corrections any further in the next section.

3.2 Fiducial cross sections

Fiducial cross sections are defined through kinematic selection criteria applied to physical objects in final states. We define the selection criteria for $pp \rightarrow Z \rightarrow \ell^+\ell^-$ as

$$p_{\perp,\ell_1} > 24 \text{ GeV}, \quad p_{\perp,\ell_2} > 16 \text{ GeV}, \quad |y_{\ell_i}| < 2.4, \quad 50 \text{ GeV} < m_{\ell\bar{\ell}} < 120 \text{ GeV}, \quad (3.2)$$

where $\ell_{1,2}$ denote leptons with leading and subleading transverse momenta, respectively.

Since we work with massless leptons, their transverse momenta are not collinear-safe observables; for this reason, we need to introduce an analog of QCD jets for leptons by combining leptons with collinear photons. Such recombination procedures are also used in experimental measurements to define “physical” electrons subject to selection cuts. We note that this procedure is valid only in the case of electrons; for muons, a calculation that retains the muon mass is required. For the purposes of the computation of mixed QCD \otimes QED corrections, we choose a simplified version of the standard recipe [56]. To this end, we begin by computing two quantities $R_{\ell\pm\gamma} = \sqrt{(y_{\ell\pm} - y_\gamma)^2 + (\varphi_{\ell\pm} - \varphi_\gamma)^2}$ where

⁴We thank D. de Florian for clarifications and help with this comparison.

$y_{\ell^\pm, \gamma}$ are the rapidities and $\varphi_{\ell^\pm, \gamma}$ the azimuthal angles of the lepton or antilepton, and the photon, respectively. If $R_{\ell^\pm, \gamma}$ for the photon and one of the leptons is smaller than some R_{\min} , the photon is recombined with the lepton by adding their momenta; the new object is treated as a lepton inasmuch as the selection cuts eq. (3.2) are concerned. In this paper, we use the standard value [56] $R_{\min} = 0.1$.

As we already mentioned, there are three distinct sources of QED corrections. To show them separately, we decompose the NLO QED and mixed QCD \otimes QED results according to whether the QED correction is associated with the production of the Z boson (P), its decay (D), or its decay width $\Gamma_{Z \rightarrow e^+e^-}$ (W). Using the definitions for Δ 's in the previous section, we find

$$\begin{aligned} \Delta_\alpha &= (3.0 \cdot 10^{-3})_P - (7.2 \cdot 10^{-3})_D - (1.6 \cdot 10^{-3})_W, \\ \Delta_{\alpha_s^2} &= -(1.2 \cdot 10^{-2}), \\ \Delta_{\alpha_s \alpha} &= -(1.5 \cdot 10^{-4})_{P \otimes P} - (4.9 \cdot 10^{-3})_{P \otimes D} - (0.3 \cdot 10^{-3})_{P \otimes W}. \end{aligned} \tag{3.3}$$

The many results in eq. (3.3) can be compared in different ways. First, we note that the QCD corrections are larger than in the inclusive case by almost a factor of two whereas the NLO QED corrections (initial) do not change significantly. The mixed QCD \otimes QED correction to the production is, on the other hand, smaller by a factor of two than in the inclusive case. Similar to the inclusive case, the relative magnitude of NLO QED, NNLO QCD and mixed QCD \otimes QED corrections to the fiducial Z boson production cross section is consistent with expectations based on the relative sizes of QCD and QED couplings, despite the somewhat larger relative magnitude of the NNLO QCD correction.

It is interesting to understand how the final result for QCD \otimes QED corrections to the production comes about. To this end, it is instructive to decompose $[\Delta_{\alpha_s \alpha}]_{P \otimes P}$ into contributions of particular partonic channels, see table 1. We observe a sizeable, almost an order-of-magnitude cancellation between $q\bar{q}$ and qg channels. In fact a similar cancellation reduces the magnitude of NNLO QCD corrections which could have been quite a bit bigger than what they are if this cancellation was not present.

It is seen from table 1 that contributions of photon-induced channels are very small, as expected. However, due to the aforementioned cancellation, they still contribute roughly twenty percent to the total result for mixed initial-initial corrections. It is also interesting that the photon-induced contributions are *larger* than those of qg channels. This implies that neglecting contributions with photons in the initial state is not a good approximation if QCD \otimes QED precision is desired.

A rather different situation arises if we look at contributions to eq. (3.3) that involve corrections to Z boson decays. They are described by $[\Delta_\alpha]_D$ and by $[\Delta_{\alpha_s \alpha}]_{P \otimes D}$ for the QED and QCD \otimes QED corrections, respectively. Inspecting eq. (3.3), we observe that both of these contributions are large and that $[\Delta_{\alpha_s \alpha}]_{P \otimes D}$ is smaller than $[\Delta_\alpha]_D$ by only thirty percent in spite of being suppressed by one power of α_s . This implies that for fiducial cross sections QED radiation in the decay is very strongly affected by QCD radiation in the production.

This result illustrates that the selection criteria shown in eq. (3.2) are strongly impacted by the non-vanishing transverse momentum of the Z boson that in the case of a fixed-order

Partonic Channel	$[\Delta_{\alpha_s\alpha}]_{P\otimes P} \cdot 10^4$
$q\bar{q}$	5.60
qq	0.13
$qg + gq$	-7.01
$q\gamma + \gamma q$	-0.32
γg	0.06
Total	-1.54

Table 1. Contributions of the different partonic channels to $[\Delta_{\alpha_s\alpha}]_{P\otimes P}$.

computation is provided by the initial state QCD radiation. In addition, $R_{\min} = 0.1$ is probably too small an isolation cone to allow a stable perturbative description of QED radiation off the outgoing leptons. To illustrate this remark, we point out that, with $R_{\min} = 0.1$ and an additional selection cut $p_{\perp,\gamma} > 5$ GeV, the rate of the Z boson decay to three QED jets becomes close to four percent and thus much larger than a naive expectation based on $\alpha/\pi \sim 2 \times 10^{-3}$ suppression of events with additional radiation. It is clear that a quasi-collinear fragmentation of a lepton to a photon, allowed by the selection cuts, may strongly change the observable cross section by reducing the transverse momentum of the lepton. Furthermore, we also observe that by rejecting events with two leptons and a photon and keeping events with only Born-like kinematics, the size of $[\Delta_{\alpha_s\alpha}]_{P\otimes D}$ gets reduced relative to $[\Delta_{\alpha}]_D$.⁵

3.3 Kinematic distributions

We continue with the discussion of the impact of QCD \otimes QED corrections on kinematic distributions for the production of two leptons via an on-shell Z boson at the LHC. Below we show the respective distribution at NLO QCD accuracy in upper panels and the relative corrections $\Delta_{\alpha_s^2}$, $[\Delta_{\alpha_s\alpha}]_{P\otimes P}$ and $[\Delta_{\alpha_s\alpha}]_{P\otimes D}$ in lower panels. We use the selection criteria shown in eq. (3.2) and $R_{\min} = 0.1$ throughout this section.

We begin with the discussion of the transverse momentum distribution of the two leptons, $p_{\perp,\ell\ell}$, shown in the left panels of figure 4. It is seen from the plot that both $\Delta_{\alpha_s^2}$ and $[\Delta_{\alpha_s\alpha}]_{P\otimes P}$ corrections become flat and positive above $p_{\perp,\ell\ell} \sim 20$ GeV. In that region, the NNLO QCD corrections amount to $\mathcal{O}(40)$ percent. Although this is quite a large correction, we note that in this kinematic region they can be considered as NLO QCD corrections to Z + jet production. The mixed $[\Delta_{\alpha_s\alpha}]_{P\otimes P}$ corrections can be thought of as QED corrections to the NLO QCD distribution; this would suggest a percent-level correction. In fact, initial-initial corrections in this case turn out to be even smaller, of the order of two permille for $p_{T,\ell\ell} > 20$ GeV. This is partially due to the cancellation between $q\bar{q}$ - and qg -induced contributions that we already discussed. For smaller values of the Z

⁵In the future, it may be interesting to combine $\mathcal{O}(\alpha_s^2)$ corrections to the production of the Z boson and $\mathcal{O}(\alpha)$ to its decay in order to investigate the size of the correction due to a second emission from the initial state; since the first QCD emission already provides some boost to the Z boson, it is conceivable that the corrections due to second gluon emission will be much more moderate.




Figure 4. Relative differential corrections Δ for the transverse momentum (left) and rapidity (right) of the dilepton system, $p_{\perp,\ell\ell}$ and $y_{\ell\ell}$. See text for further details.

transverse momentum, the corrections become large and negative; however, resummation may be needed to obtain a reliable prediction in this region.

The initial-final correction $[\Delta_{\alpha_s\alpha}]_{P\otimes D}$ to the $p_{\perp,\ell\ell}$ distribution of a lepton pair shows quite a different behaviour, with a maximum at $p_{\perp,\ell\ell} \simeq 30$ GeV. This feature appears because of an interplay of a few contributions with different kinematics features. On the one hand, processes without initial-state QCD radiation but with final-state photon emission yield a pair of leptons with a total transverse momentum smaller than $M_Z/2$. Moreover, the selection cuts we use further restrict it to $p_{\perp,\ell\ell} \lesssim 31$ GeV. On the other hand, processes with initial state radiation boost the Z boson, leading to a tail which extends beyond this kinematic limit. This behaviour can already be observed in the initial and final contributions at NLO QED accuracy.

It is also interesting to point out that, although initial-final $[\Delta_{\alpha_s\alpha}]_{P\otimes D}$ corrections are indeed larger than initial-initial corrections $[\Delta_{\alpha_s\alpha}]_{P\otimes P}$ for almost all values of the transverse momentum $p_{\perp,\ell\ell}$, it is not the case for $p_{\perp,\ell\ell} \sim 30$ GeV where the initial-final correction passes through zero(s). For those values of $p_{\perp,\ell\ell}$, the initial-final and initial-initial QCD \otimes QED contributions become comparable.

The rapidity distribution of the dilepton system at NLO QCD and the different corrections to it are displayed in the right panels of figure 4. The shape of the rapidity distribution is determined by the selection cuts which flatten the distribution for values $|y_{\ell\ell}| < 1$. Inside this region, which represents the bulk of the cross section, the NNLO QCD correction is flat and negative and amounts to a decrease of the NLO QCD result by $\mathcal{O}(-2)$ percent. The NNLO QCD correction then crosses zero at rapidities $|y_{\ell\ell}| \simeq 1$ and increase to $\mathcal{O}(3)$ percent at large rapidities. The mixed $[\Delta_{\alpha_s\alpha}]_{P\otimes P}$ correction has a very




Figure 5. Relative differential corrections Δ for the transverse momentum of the leading (left) and subleading (right) lepton, p_{\perp,ℓ_1} and p_{\perp,ℓ_2} . See text for further details.

similar shape; it decreases the NLO QCD distribution by $\mathcal{O}(-0.4)$ permille in the central region and increases it to $\mathcal{O}(+0.4)$ permille outside. In contrast to the previous two corrections, the initial-final $[\Delta_{\alpha_s\alpha}]_{P\otimes D}$ correction is negative for all rapidities and amounts to a decrease by about $\mathcal{O}(-7)$ permille in the central region. For rapidities $|y_{\ell\ell}| > 1$, the initial-final correction becomes smaller but it remains a factor 5 larger than the initial-initial one. Hence, it follows that for the rapidity distribution of a lepton pair, the initial-final corrections always dominate over the initial-initial one.

We show the transverse momentum distributions of the leading and the subleading leptons in figure 5. For the leading lepton, the NNLO QCD corrections enhance the distribution at large p_{\perp,ℓ_1} , which is consistent with the additional boost that leptons get from the second initial-state emission. The feature at $p_{\perp,\ell_1} = M_Z/2$ is a Sudakov shoulder effect, which is known to appear close to kinematic boundaries. The NNLO QCD correction factor stabilises at $\mathcal{O}(40)$ percent for large values of p_{\perp,ℓ_1} . The initial-initial correction $[\Delta_{\alpha_s\alpha}]_{P\otimes P}$ shows a similar behaviour at high p_{\perp,ℓ_1} where it is about two permille. Interestingly, the initial-initial correction also enhances the distribution at smaller values of the transverse momentum p_{\perp,ℓ_1} . The initial-final correction $[\Delta_{\alpha_s\alpha}]_{P\otimes D}$ is negative at high and positive at small p_{\perp,ℓ_1} ; this is consistent with the picture of the final state leptons losing energy to QED radiation and decreasing their transverse momenta. The correction $[\Delta_{\alpha_s\alpha}]_{P\otimes D}$ changes from $\mathcal{O}(-3)$ percent for $p_{\perp,\ell_1} > M_Z/2$ to $\mathcal{O}(+3)$ percent for $p_{\perp,\ell_1} < M_Z/2$.

The transverse momentum distribution of the subleading lepton at NLO QCD accuracy has some features which impact the respective corrections. Indeed, in addition to the Sudakov shoulder at $p_{\perp,\ell_2} = M_Z/2$, the distribution also features a similar effect at $p_{\perp,\ell_2} = 24$ GeV, due to the fact that 24 GeV is a cut on the minimal transverse momentum of the

leading lepton. Since at leading order the transverse momenta of the two leptons must be equal to each other, the leading order p_{\perp,ℓ_2} distribution is truncated at this value *also for the subleading lepton* and the lower bins are only populated through higher order corrections.

In order to avoid displaying large fluctuations of radiative effects around the Sudakov shoulder at $p_{\perp,\ell_2} = M_Z/2$, we combined two bins between 40 GeV and 48 GeV into a single bin to present various corrections. Similar to the leading-lepton case, we observe large positive NNLO QCD and mixed $[\Delta_{\alpha_s\alpha}]_{P\otimes D}$ corrections for $p_{\perp,\ell_2} > M_Z/2$ that can be as large as $\mathcal{O}(50)$ percent for QCD and $\mathcal{O}(2)$ permille for mixed QCD \otimes QED. The initial-final correction $[\Delta_{\alpha_s\alpha}]_{P\otimes D}$ is negative and takes values between $\mathcal{O}(-2)$ and $\mathcal{O}(-4)$ percent.

Finally, we discuss distributions of $\cos\theta^*$, where θ^* is the angle between the three-momentum of one of the leptons and a unit vector constructed from the difference between three-momenta of the colliding protons in the rest frame of the dilepton system. Its cosine is given by the following formula [47]

$$\cos\theta^* = \frac{P_{\ell^-}^+ P_{\ell^+}^- - P_{\ell^-}^- P_{\ell^+}^+}{\sqrt{m_{\ell^+\ell^-}^2 (m_{\ell^+\ell^-}^2 + p_{\perp,\ell^+\ell^-})}} \frac{p_{z,\ell^+\ell^-}}{|p_{z,\ell^+\ell^-}|}, \quad (3.4)$$

where $P_i^\pm = (E_i \pm p_{z,i})$. The $\cos\theta^*$ -distribution is one of the few observables with strong sensitivity to the weak mixing angle $\sin^2\theta_{\text{eff}}^\ell$; it is used in experimental analyses for this purpose [9]. The relevant input for this measurement is provided by $\cos\theta^*$ distributions for restricted $m_{\ell\ell}$ and $|y_{\ell\ell}|$ intervals.

To illustrate how various effects modify the $\cos\theta^*$ distributions, in figure 6 we show them for $0.6 < |y_{\ell\ell}| < 1.2$ and $1.8 < |y_{\ell\ell}| < 2.4$ rapidity intervals. The panels on the left describe the rapidity interval $0.6 < |y_{\ell\ell}| < 1.2$ and feature the distribution at NLO QCD accuracy with two well-separated maxima in the upper panel and NNLO QCD and mixed QCD \otimes QED corrections in the lower panel. The NNLO QCD corrections are below a percent level at small values of $\cos\theta^*$, but become large and negative at the boundaries of the distribution. The mixed $[\Delta_{\alpha_s\alpha}]_{P\otimes P}$ corrections are relatively flat and amount to roughly 0.3 permille in the bulk of the distribution. They increase slightly at small values of $\cos\theta^*$, but become large and negative in the outermost non-vanishing bins. The initial-final $[\Delta_{\alpha_s\alpha}]_{P\otimes D}$ contribution is negative for all values of $\cos\theta^*$ and is relatively flat, taking values between one and two permille.

The $\cos\theta^*$ distribution for rapidities $1.8 < |y_{\ell\ell}| < 2.4$ is quite different. First, two maxima merge into one maximum located at small values of $\cos\theta^*$. For this reason, we only display results in the interval $-0.5 < \cos\theta^* < 0.5$. Second, all corrections in this case are rather flat. However, their magnitudes are comparable to those in the interval $0.6 < |y_{\ell\ell}| < 1.2$, with the NNLO QCD corrections being just few percent and the initial-initial correction $[\Delta_{\alpha_s\alpha}]_{P\otimes P}$ below a permille level. The initial-final $[\Delta_{\alpha_s\alpha}]_{P\otimes D}$ corrections are negative and are close to two permille.

4 Conclusion and outlook

In this article, we presented the calculation of mixed QCD \otimes QED corrections to the production of an on-shell Z boson at the LHC. We made use of the nested soft-collinear




Figure 6. Relative differential corrections Δ for the $\cos \theta^*$ distribution for $0.6 < |y_{\ell\ell}| < 1.2$ (left) and $1.8 < |y_{\ell\ell}| < 2.4$ (right). See text for further details.

subtraction scheme developed for NNLO QCD computations at a fully-differential level, and extended it to cover the mixed QCD \otimes QED corrections. We adopted the abelianisation procedure introduced in ref. [41] and included partonic channels with photons in the initial state. Since we considered production of an on-shell Z boson, interactions between initial state partons and decay products of the Z boson can be neglected.

As an illustration of the fully-exclusive nature of our computation, we calculated the mixed QCD \otimes QED corrections to a number of observables such as the transverse momentum distributions of di-leptons and of the leading and subleading leptons, as well as the rapidity of the dilepton system and one of the Collins-Soper angles θ^* . Initial-initial QCD \otimes QED corrections typically change these distributions at below a permille level whereas initial-final ones change them by a few permille. This is a factor hundred (ten) smaller than the effects of NNLO QCD corrections, respectively.

Finally, we note that the production of the on-shell Z boson is a relatively simple case. In the future, it may be interesting to compute the mixed QCD \otimes QED corrections to the generic off-shell Drell-Yan process and to the production of an on-shell W boson. Thanks to recent developments, both of these computations are now feasible. The case of W production will require an extension of the nested soft-collinear subtraction scheme to the case of a charged resonance. This is an interesting problem that we plan to address in the future.

Acknowledgments

We are grateful to A. Behring, F. Caola, D. de Florian, P. F. Monni, and G. Salam for useful conversations. The research of K.M. is supported by BMBF grant 05H18VKCC1 and by the

DFG Collaborative Research Center TRR 257 “Particle Physics Phenomenology after the Higgs Discovery”. M.D. and M.J. are supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under grant 396021762 - TRR 257.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] ATLAS collaboration, *Observation of a new particle in the search for the standard model Higgs boson with the ATLAS detector at the LHC*, *Phys. Lett. B* **716** (2012) 1 [[arXiv:1207.7214](https://arxiv.org/abs/1207.7214)] [[INSPIRE](#)].
- [2] CMS collaboration, *Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC*, *Phys. Lett. B* **716** (2012) 30 [[arXiv:1207.7235](https://arxiv.org/abs/1207.7235)] [[INSPIRE](#)].
- [3] S.D. Drell and T.-M. Yan, *Massive lepton pair production in hadron-hadron collisions at high-energies*, *Phys. Rev. Lett.* **25** (1970) 316 [Erratum *ibid.* **25** (1970) 902] [[INSPIRE](#)].
- [4] M.L. Mangano, *Production of electroweak bosons at hadron colliders: theoretical aspects*, *Adv. Ser. Direct. High Energy Phys.* **26** (2016) 231 [[arXiv:1512.00220](https://arxiv.org/abs/1512.00220)] [[INSPIRE](#)].
- [5] M. Dittmar, F. Pauss and D. Zurcher, *Towards a precise parton luminosity determination at the CERN LHC*, *Phys. Rev. D* **56** (1997) 7284 [[hep-ex/9705004](https://arxiv.org/abs/hep-ex/9705004)] [[INSPIRE](#)].
- [6] V.A. Khoze, A.D. Martin, R. Orava and M.G. Ryskin, *Luminosity monitors at the LHC*, *Eur. Phys. J. C* **19** (2001) 313 [[hep-ph/0010163](https://arxiv.org/abs/hep-ph/0010163)] [[INSPIRE](#)].
- [7] W.T. Giele and S.A. Keller, *Hard scattering based luminosity measurement at hadron colliders*, [hep-ph/0104053](https://arxiv.org/abs/hep-ph/0104053) [[INSPIRE](#)].
- [8] S. Haywood et al., *Electroweak physics*, in the proceedings of the *CERN Workshop on standard model physics (and more) at the LHC*, May 25–26, CERN, Geneva (1999), [hep-ph/0003275](https://arxiv.org/abs/hep-ph/0003275) [[INSPIRE](#)].
- [9] CMS collaboration, *Measurement of the weak mixing angle using the forward-backward asymmetry of Drell-Yan events in pp collisions at 8 TeV*, *Eur. Phys. J. C* **78** (2018) 701 [[arXiv:1806.00863](https://arxiv.org/abs/1806.00863)] [[INSPIRE](#)].
- [10] NNPDF collaboration, *Parton distributions from high-precision collider data*, *Eur. Phys. J. C* **77** (2017) 663 [[arXiv:1706.00428](https://arxiv.org/abs/1706.00428)] [[INSPIRE](#)].
- [11] L.A. Harland-Lang, A.D. Martin, P. Motylinski and R.S. Thorne, *Parton distributions in the LHC era: MMHT 2014 PDFs*, *Eur. Phys. J. C* **75** (2015) 204 [[arXiv:1412.3989](https://arxiv.org/abs/1412.3989)] [[INSPIRE](#)].
- [12] T.-J. Hou et al., *Progress in the CTEQ-TEA NNLO global QCD analysis*, [arXiv:1908.11394](https://arxiv.org/abs/1908.11394) [[INSPIRE](#)].
- [13] S. Alekhin, J. Bluemlein, S.-O. Moch and R. Placakyte, *The new ABMP16 PDF*, *PoS DIS2016* (2016) 016 [[arXiv:1609.03327](https://arxiv.org/abs/1609.03327)] [[INSPIRE](#)].
- [14] M. Farina et al., *Energy helps accuracy: electroweak precision tests at hadron colliders*, *Phys. Lett. B* **772** (2017) 210 [[arXiv:1609.08157](https://arxiv.org/abs/1609.08157)] [[INSPIRE](#)].

- [15] G. Altarelli, R.K. Ellis and G. Martinelli, *Large perturbative corrections to the Drell-Yan process in QCD*, *Nucl. Phys. B* **157** (1979) 461 [INSPIRE].
- [16] R. Hamberg, W.L. van Neerven and T. Matsuura, *A complete calculation of the order α_s^2 correction to the Drell-Yan K factor*, *Nucl. Phys. B* **359** (1991) 343 [Erratum *ibid.* **B 644** (2002) 403] [INSPIRE].
- [17] W.L. van Neerven and E.B. Zijlstra, *The $O(\alpha_s^2)$ corrected Drell-Yan K factor in the DIS and \overline{MS} scheme*, *Nucl. Phys. B* **382** (1992) 11 [Erratum *ibid.* **B 680** (2004) 513] [INSPIRE].
- [18] R.V. Harlander and W.B. Kilgore, *Next-to-next-to-leading order Higgs production at hadron colliders*, *Phys. Rev. Lett.* **88** (2002) 201801 [hep-ph/0201206] [INSPIRE].
- [19] S. Catani et al., *Vector boson production at hadron colliders: a fully exclusive QCD calculation at NNLO*, *Phys. Rev. Lett.* **103** (2009) 082001 [arXiv:0903.2120] [INSPIRE].
- [20] K. Melnikov and F. Petriello, *The W boson production cross section at the LHC through $O(\alpha_s^2)$* , *Phys. Rev. Lett.* **96** (2006) 231803 [hep-ph/0603182] [INSPIRE].
- [21] K. Melnikov and F. Petriello, *Electroweak gauge boson production at hadron colliders through $O(\alpha_s^2)$* , *Phys. Rev. D* **74** (2006) 114017 [hep-ph/0609070] [INSPIRE].
- [22] R. Gavin, Y. Li, F. Petriello and S. Quackenbush, *FEWZ 2.0: a code for hadronic Z production at next-to-next-to-leading order*, *Comput. Phys. Commun.* **182** (2011) 2388 [arXiv:1011.3540] [INSPIRE].
- [23] R. Gavin, Y. Li, F. Petriello and S. Quackenbush, *W physics at the LHC with FEWZ 2.1*, *Comput. Phys. Commun.* **184** (2013) 208 [arXiv:1201.5896] [INSPIRE].
- [24] R. Boughezal et al., *Color singlet production at NNLO in MCFM*, *Eur. Phys. J. C* **77** (2017) 7 [arXiv:1605.08011] [INSPIRE].
- [25] T. Ahmed, M. Mahakhud, N. Rana and V. Ravindran, *Drell-Yan production at threshold to third order in QCD*, *Phys. Rev. Lett.* **113** (2014) 112002 [arXiv:1404.0366] [INSPIRE].
- [26] S. Catani et al., *Threshold resummation at N^3LL accuracy and soft-virtual cross sections at N^3LO* , *Nucl. Phys. B* **888** (2014) 75 [arXiv:1405.4827] [INSPIRE].
- [27] U. Baur et al., *Electroweak radiative corrections to neutral current Drell-Yan processes at hadron colliders*, *Phys. Rev. D* **65** (2002) 033007 [hep-ph/0108274] [INSPIRE].
- [28] U. Baur, S. Keller and W.K. Sakumoto, *QED radiative corrections to Z boson production and the forward backward asymmetry at hadron colliders*, *Phys. Rev. D* **57** (1998) 199 [hep-ph/9707301] [INSPIRE].
- [29] C.M. Carloni Calame, G. Montagna, O. Nicrosini and M. Treccani, *Multiple photon corrections to the neutral-current Drell-Yan process*, *JHEP* **05** (2005) 019 [hep-ph/0502218] [INSPIRE].
- [30] A. Arbuzov et al., *One-loop corrections to the Drell-Yan process in SANC. (II). The Neutral current case*, *Eur. Phys. J. C* **54** (2008) 451 [arXiv:0711.0625] [INSPIRE].
- [31] S. Dittmaier and M. Huber, *Radiative corrections to the neutral-current Drell-Yan process in the Standard Model and its minimal supersymmetric extension*, *JHEP* **01** (2010) 060 [arXiv:0911.2329] [INSPIRE].
- [32] C. Bernaciak and D. Wackerroth, *Combining NLO QCD and electroweak radiative corrections to W boson production at hadron colliders in the POWHEG framework*, *Phys. Rev. D* **85** (2012) 093003 [arXiv:1201.4804] [INSPIRE].

- [33] B. Mistlberger, *Higgs boson production at hadron colliders at N^3LO in QCD*, *JHEP* **05** (2018) 028 [[arXiv:1802.00833](#)] [[INSPIRE](#)].
- [34] R. Bonciani, F. Buccioni, R. Mondini and A. Vicini, *Double-real corrections at $\mathcal{O}(\alpha\alpha_s)$ to single gauge boson production*, *Eur. Phys. J. C* **77** (2017) 187 [[arXiv:1611.00645](#)] [[INSPIRE](#)].
- [35] C. Anastasiou and K. Melnikov, *Higgs boson production at hadron colliders in NNLO QCD*, *Nucl. Phys. B* **646** (2002) 220 [[hep-ph/0207004](#)] [[INSPIRE](#)].
- [36] M. Heller, A. von Manteuffel and R.M. Schabinger, *Multiple polylogarithms with algebraic arguments and the two-loop EW-QCD Drell-Yan master integrals*, [arXiv:1907.00491](#) [[INSPIRE](#)].
- [37] Y. Li and F. Petriello, *Combining QCD and electroweak corrections to dilepton production in FEWZ*, *Phys. Rev. D* **86** (2012) 094034 [[arXiv:1208.5967](#)] [[INSPIRE](#)].
- [38] L. Barze et al., *Neutral current Drell-Yan with combined QCD and electroweak corrections in the POWHEG BOX*, *Eur. Phys. J. C* **73** (2013) 2474 [[arXiv:1302.4606](#)] [[INSPIRE](#)].
- [39] S. Dittmaier, A. Huss and C. Schwinn, *Mixed QCD-electroweak $\mathcal{O}(\alpha_s\alpha)$ corrections to Drell-Yan processes in the resonance region: pole approximation and non-factorizable corrections*, *Nucl. Phys. B* **885** (2014) 318 [[arXiv:1403.3216](#)] [[INSPIRE](#)].
- [40] S. Dittmaier, A. Huss and C. Schwinn, *Dominant mixed QCD-electroweak $\mathcal{O}(\alpha_s\alpha)$ corrections to Drell-Yan processes in the resonance region*, *Nucl. Phys. B* **904** (2016) 216 [[arXiv:1511.08016](#)] [[INSPIRE](#)].
- [41] D. de Florian, M. Der and I. Fabre, *QCD \oplus QED NNLO corrections to Drell-Yan production*, *Phys. Rev. D* **98** (2018) 094008 [[arXiv:1805.12214](#)] [[INSPIRE](#)].
- [42] A.L. Kataev, *Higher order $\mathcal{O}(\alpha^2)$ and $\mathcal{O}(\alpha\alpha_s)$ corrections to sigma total $e^+e^- \rightarrow$ hadron and Z boson decay rate*, *Phys. Lett. B* **287** (1992) 209 [[INSPIRE](#)].
- [43] F. Caola, K. Melnikov and R. Röntsch, *Nested soft-collinear subtractions in NNLO QCD computations*, *Eur. Phys. J. C* **77** (2017) 248 [[arXiv:1702.01352](#)] [[INSPIRE](#)].
- [44] A. Manohar, P. Nason, G.P. Salam and G. Zanderighi, *How bright is the proton? A precise determination of the photon parton distribution function*, *Phys. Rev. Lett.* **117** (2016) 242002 [[arXiv:1607.04266](#)] [[INSPIRE](#)].
- [45] A.V. Manohar, P. Nason, G.P. Salam and G. Zanderighi, *The photon content of the proton*, *JHEP* **12** (2017) 046 [[arXiv:1708.01256](#)] [[INSPIRE](#)].
- [46] NNPDF collaboration, *Illuminating the photon content of the proton within a global PDF analysis*, *SciPost Phys.* **5** (2018) 008 [[arXiv:1712.07053](#)] [[INSPIRE](#)].
- [47] J.C. Collins and D.E. Soper, *Angular distribution of dileptons in high-energy hadron collisions*, *Phys. Rev. D* **16** (1977) 2219 [[INSPIRE](#)].
- [48] F. Caola, K. Melnikov and R. Röntsch, *Analytic results for color-singlet production at NNLO QCD with the nested soft-collinear subtraction scheme*, *Eur. Phys. J. C* **79** (2019) 386 [[arXiv:1902.02081](#)].
- [49] S. Frixione, Z. Kunszt and A. Signer, *Three jet cross-sections to next-to-leading order*, *Nucl. Phys. B* **467** (1996) 399 [[hep-ph/9512328](#)] [[INSPIRE](#)].
- [50] S. Frixione, *A general approach to jet cross-sections in QCD*, *Nucl. Phys. B* **507** (1997) 295 [[hep-ph/9706545](#)] [[INSPIRE](#)].

- [51] F. Caola, M. Delto, H. Frellesvig and K. Melnikov, *The double-soft integral for an arbitrary angle between hard radiators*, *Eur. Phys. J. C* **78** (2018) 687 [[arXiv:1807.05835](#)] [[INSPIRE](#)].
- [52] M. Delto and K. Melnikov, *Integrated triple-collinear counter-terms for the nested soft-collinear subtraction scheme*, *JHEP* **05** (2019) 148 [[arXiv:1901.05213](#)] [[INSPIRE](#)].
- [53] F. Caola, K. Melnikov and R. Röntsch, *Analytic results for decays of color singlets to $g\bar{g}$ and $q\bar{q}$ final states at NNLO QCD with the nested soft-collinear subtraction scheme*, *Eur. Phys. J. C* **79** (2019) 1013 [[arXiv:1907.05398](#)] [[INSPIRE](#)].
- [54] A. Buckley et al., *LHAPDF6: parton density access in the LHC precision era*, *Eur. Phys. J. C* **75** (2015) 132 [[arXiv:1412.7420](#)] [[INSPIRE](#)].
- [55] D.A. Dicus and S.S.D. Willenbrock, *Radiative corrections to the ratio of Z and W boson production*, *Phys. Rev. D* **34** (1986) 148 [[INSPIRE](#)].
- [56] S. Alioli et al., *Precision studies of observables in $pp \rightarrow W \rightarrow l\nu_l$ and $pp \rightarrow \gamma, Z \rightarrow l^+l^-$ processes at the LHC*, *Eur. Phys. J. C* **77** (2017) 280 [[arXiv:1606.02330](#)] [[INSPIRE](#)].