


Unexpected Isomerization of Hexa-*tert*-butyl-octaphosphaneToni Grell^[a] and Evamarie Hey-Hawkins^{*[a]}

Dedication ((optional))

Abstract: Octaphosphane $\{cyclo-(P_4tBu_3)_2\}$ (**1**) undergoes an unexpected isomerization reaction to the constitutional isomer 2,2',2'',2''',3,3'-hexa-*tert*-butyl-bicyclo[3.3.0]octaphosphane (**2**) in the presence of Lewis acidic metal salts. The mechanism of this reaction is discussed and elucidated with DFT calculations. The results underline the fascinating similarity between phosphorus-rich and isolobal carbon compounds. The new bicyclic octaphosphane **2** shows a dynamic behavior in solution and reacts with $[AuCl(tht)]$ (tht = tetrahydrothiophene) to give a mono- ($[AuCl(2-\kappa P^3)]$, **3**) and a dinuclear complex ($[(AuCl)_2(2-\kappa^2 P^3, \kappa P^3)]$, **4**). With *cis*- $[PdCl_2(cod)]$ (cod = 1,5-cyclooctadiene), the chelate complex ($[PdCl_2(2-\kappa^2 P^2, P^2)]$, **5**) with a different coordination mode of the ligand was obtained.

The chemistry of phosphorus-rich compounds is an interesting field with a history of almost 150 years. Since the discovery of “Phosphabenzol”^[1] and the pioneering work of Baudler^[2] who reported a vast number of these molecules, several groups focus on this topic today because of its outstanding importance in the direct activation of white phosphorus (P_4)^[3] or the preparation of phosphorus-rich metal phosphides^[4,5]. Surely one of the most fascinating features of these compounds is their similarity to organic molecules as they can exhibit constitutional, configurational as well conformational isomerism and also multiple bonds^[6] and even aromaticity^[7]. This analogy can be rationalized with the isolobal concept and is the reason why phosphorus is often referred to as the carbon copy.^[8]




Scheme 1. Isomerization reaction between octaphosphane **1** and **2** as well as for the isolobal bi(cyclobutane) to bicyclo[3.3.0]octane (pentalane) and bicyclo[3.2.1]octane.

We here report a discovery that further substantiates this statement. Recently, we reported on the facile synthesis of

hexa-*tert*-butyl-octaphosphane (**1**) which reacted with $[AuCl(tht)]$ (tht = tetrahydrothiophene) yields mono-, di- and trinuclear complexes in which up to four phosphorus atoms of the ligand are involved in coordinating gold(I) atoms.^[9] During these studies we observed a limited coordination behavior of **1**. Instead of coordination, the constitutional isomer octaphosphane **2** was formed on reaction with a large variety of Lewis acidic metal salts and complexes such as MX_2 or $[MX_2(L)]$ ($M = Mg, Mn, Fe, Co, Zn, Cd, Sn; X = Cl, Br; L = 2,2'$ -bipyridine or N,N,N',N' -tetramethylethane-1,2-diamine) (Scheme 1).

Isomerization reactions of phosphorus-rich molecules are well known in the literature.^[4,10] One famous example is the Cope-like rearrangement of the P_7^{3-} anion.^[11] In addition to that, the heavier homolog of octaphosphane **1**, $\{cyclo-(As_4tBu_3)_2\}$ spontaneously isomerizes in solution at room temperature to the homolog of **2**.^[12] Furthermore, the isolobal octane, bi(cyclobutane), irreversibly isomerizes to bicyclo[3.3.0]octane (pentalane) if treated with the Lewis acid $AlBr_3$ with formation of bicyclo[3.2.1]octane as the final product (Scheme 1).^[13]

In the case of the octaphosphane **1**, the isomerization requires a Lewis acidic metal salt, which influences the reaction rate and temperature. The reaction of **1** with different equivalents of $ZnCl_2$ (in C_6D_6 , reflux) was monitored by $^{31}P\{^1H\}$ NMR spectroscopy. In each case, the conversion stopped at approximately 90% and no complete conversion of **1** to **2** was observed. The amount of the salt merely influenced the time until this state was reached (Figure S1, SI). Conversely, under the same conditions the conversion of octaphosphane **2** stops at 10%. These observations thus imply an equilibrium between octaphosphanes **1** and **2** in the presence of a Lewis acidic metal salts. Accordingly, octaphosphane **2** was also obtained as a side product in the synthesis of **1** which involves the two metal salts $SnCl_2$ and $MgCl_2$.



[a] Dr. T. Grell, Prof. Dr. E. Hey-Hawkins
Faculty of Chemistry and Mineralogy
Institute of Inorganic Chemistry
Johannisallee 29, 04103 Leipzig, Germany
E-mail: hey@uni-leipzig.de

COMMUNICATION

Scheme 2. Proposed mechanism of the isomerization reaction between octaphosphanes **1** and **2**. Reaction energies (calculated with Zn^{2+} as M^{2+} and two Cl^- coligands) are given in $\text{kJ}\cdot\text{mol}^{-1}$. Bond formation (dashed line) and cleavage (wavy lines) are indicated.

In analogy to the mechanism that was postulated for bi(cyclobutane)^[13] we propose the mechanism depicted in Scheme 2 for the isomerization of **1** and **2**. Firstly, a monometallic complex (**i1**) is formed in which the metal cation is coordinated by a bridge P atom and one of the *t*Bu carrying P atoms usually involved in coordination (β -P)^[9]. This leads to a proximity of the second bridge P atom to the other *t*Bu carrying P atom thus facilitating bond cleavage and formation to give the spirocyclic intermediate **i2**. In this intermediate, cleavage of the conceivable P–P bonds would either lead to a diastereomer with wrong configuration (bond I in Scheme 2) or is impaired, as the phosphanido group is sterically blocked by the metal cation (bond II in Scheme 2). Therefore, a rearrangement to intermediate **i3** is required followed by formation of intermediate **i4**. Liberation of the metal and inversion of the phosphorus atom $P^\#$ finally gives **2**. DFT calculations involving a $ZnCl_2$ fragment showed that the reaction energies (Scheme 2) for the mechanism are plausible, especially given the elevated temperatures required. While the inversion barriers of phosphanes are usually high, they can be much smaller in oligophosphanes, and thus the inversion of $P^\#$ is conceivable.^[14]





Figure 1. Experimental (top) and simulated (bottom) $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum of octaphosphane **2** (AA'A''A'''BB'CC' spin system with C_{2v} symmetry). Chemical shifts in ppm: $\delta(P_A) = +66.3$, $\delta(P_B) = +35.0$, $\delta(P_C) = +23.7$. Selected coupling constants in Hz: $^1J_{AB} = -292.30(3)$, $^1J_{AC} = -262.67(3)$, $^1J_{CC'} = -311.99(12)$, $^2J_{AA''} = +143.18(8)$.

The analogs of octaphosphane **2** with Me, Et and *i*Pr instead of *t*Bu substituents are known but could not be characterized by X-ray diffraction.^[15] The identity of **2** was unambiguously verified by single crystal XRD. The molecular structure (Figure S18, SI) confirms the proposed connectivity of the P_8 skeleton in a C_2 -symmetrical octaphosphane. All structural parameters including the P–P bond lengths between the bridgehead phosphorus atoms are in the expected range. The structure is furthermore analogous

to As_8tBu_6 with respect to configuration and conformation. The $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum of octaphosphane **2** shows three symmetrical multiplets which could be successfully simulated as an AA'A''A'''BB'CC' spin system assuming C_{2v} symmetry (Figure 1). The obtained parameters confirm the constitution as well as the configuration of the oligophosphane, as the $^1J_{PP}$ coupling constants range between -262.67 and -311.99 Hz which is in excellent agreement for similar compounds with *t*Bu substituents in a *trans* arrangement.^[16] Moreover, the $^2J_{PP}$ coupling constant ($J_{AA''}$) is large and positive indicating lone pairs of electrons directed towards each other^[17], thus corroborating the *endo* configuration of the molecule. Finally, the values are also similar to the respective ethyl-substituted oligophosphane.^[18] This is also the case for the chemical shifts. While $\delta(P_A)$ and $\delta(P_B)$ are similar to the analogs, the signal P_C appears at higher field. This matches the trend for Me, Et and *i*Pr where an increasing steric demand of the substituent causes a growing shielding of this signal.^[15]




Scheme 3. Libration of octaphosphane **2**. The transition state corresponds to the average structure in solution. The remaining structures (left and right) correspond to the enantiomeric structure found in the solid state.

Obviously, there is a molecular process which causes a higher effective symmetry in solution (C_{2v}) compared to the solid state (C_2). This was confirmed by VT NMR measurements which show that on cooling the signal of P_A in the $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum (Figure S6, SI) appears to lose its mirror symmetry resulting into two partially overlapping multiplets. The situation is much clearer in the $^1\text{H}\{^{31}\text{P}\}$ NMR spectra (Figure S7, SI) where a proper decoalescence of the singlet of the *t*Bu group attached to P_A ($+1.38$ ppm, $I_{\text{rel}} = 2$) is observed at -75 °C. The resulting pattern at lower temperatures is thus consistent with an AA'BB'CC'DD' spin system (C_2) matching the molecular structure in the solid state. The molecular motion responsible for the process is the libration around the bond between the bridgehead atoms (Scheme 3). We performed a relaxed surface scan of the respective dihedral angle (henceforth λ) using DFT calculations (PBE0-D3/def2-TZVP). The profile (Figure S24, SI) indeed shows that the molecular structure in the solid state ($\lambda = 40.6^\circ$) presents a minimum on the energy surface potential, while the C_{2v} -symmetrical transition state ($\lambda = 0^\circ$) matches the average structure in solution. Furthermore, the respective activation energy of $+32.6$ $\text{kJ}\cdot\text{mol}^{-1}$ agrees well with the experiment according to the temperature range in which the coalescence is observed.


COMMUNICATION

In view of the discovery of octaphosphane **2**, which was stated to be unstable^[19], the discovery of further octaphosphanes become more conceivable. We calculated the relative energies of all eight possible octanes C₈H₁₄, phosphanes P₈H₆ as well as organophosphanes P₈tBu₆ (Scheme 4, detailed in SI) and draw some key conclusions from these calculations. Going from the octanes to the phosphanes (formal exchange of CH₂ with PH) changes the order of relative energies which is due to phosphorus preferring smaller bond angles than carbon. Replacing the hydrogen atom with tBu groups changes the order once again for the organophosphanes. Structure **A** is usually considerably less stable than the other ones due to the higher strain of the four-membered ring. However, attractive dispersion interactions between the tBu groups stabilize this structure for the organophosphane **1**. For P₈tBu₆ thus type **A** ($\Delta E_{\text{rel}} = 28.2 \text{ kJ}\cdot\text{mol}^{-1}$) and **B** ($\Delta E_{\text{rel}} = 0 \text{ kJ}\cdot\text{mol}^{-1}$, octaphosphane **2**) are the most stable structures. Nevertheless, structures **D** and **F** are only slightly higher in energy ($\Delta E_{\text{rel}} = 41.1$ and $60.4 \text{ kJ}\cdot\text{mol}^{-1}$) and hence they are expected to be isolable compounds.



Scheme 4. Possible constitutional isomers of C₈H₁₄, P₈H₆ or P₈tBu₆ precluding multiple bonds.


As its constitutional isomer, octaphosphane **2** is able to coordinate gold(I). When **2** is reacted with one or two equivalents of [AuCl(tht)] a mononuclear complex [AuCl(**2**)] (**3**) or a dinuclear complex [(AuCl)₂(**2**)] (**4**) are obtained, respectively (Scheme 5). The molecular structures of both complexes (Figures S19 and S20, SI) show that the peripheral phosphorus atoms of the ligand coordinate gold(I) chloride ($\kappa^{\text{P}^3}/\kappa^{\text{P}^3}$ binding mode) resulting in a linear coordination environment. The structural parameters of the ligand in **3** and **4** are not significantly different from the free octaphosphane **2** besides a slightly larger dihedral angle λ which seems to be increasing with the number of AuCl fragments (44.8° for **3** and 50.69° for **4**). Mass spectrometry and elemental analysis confirm the composition of the proposed complexes.



Scheme 5. Formation and isomerization reaction of gold(I) complexes **3** and **4**.

However, ³¹P{¹H} NMR spectroscopy indicates a more complex situation in solution. Thus, for **3**, five multiplets with a relative intensity (*I*_{rel}) of 1:2:2:2:1 are observed whereas the spectrum of **4** shows three multiplets with *I*_{rel} of 4:2:2. This is consistent with the respective solid-state structure assuming the same libration motion as for the free octaphosphane equally causing a higher effective symmetry. Furthermore, in the ³¹P{¹H} NMR spectrum of **3** (Figure S9, SI) additional signals corresponding to the free octaphosphane **2** and the bimetallic complex **4** are observed. This indicates an equilibrium in solution (Eq. 1, Scheme 5) analogous to the one observed for the gold(I) complexes of octaphosphane **1**.^[9,20] In fact, when octaphosphane **2** is reacted with the dinuclear complex **4**, the resulting NMR spectrum is identical to the one of complex **3**. Furthermore, this is also supported by the crystal structure of **3**, which cocrystallized with 4% of complex **4** (details in SI). The situation is different for complex **4**. In the ³¹P{¹H} NMR spectrum (Figure S12, SI), in addition to the signals which match the solid state structure, a second signal set is observed for which we propose a constitutional isomer (Eq. 2, Scheme 5). As the experimental PXR diffractogram of **4** matches well the theoretical pattern calculated from its single crystal structure data, it appears that the second isomer is only formed in solution.

The reaction of octaphosphane **2** with an excess of [AuCl(tht)] yields an insoluble substance, presumably, a coordination polymer. Elemental analysis indicates the formation of a tetranuclear complex [(AuCl)₄(**2**)], and the IR spectrum shows the same characteristic vibrations of the tBu group that were observed for **2-5**.



COMMUNICATION

Scheme 6. Reaction of octaphosphane **2** with *cis*-[PdCl₂(cod)].

When octaphosphane **2** is reacted with an excess of *cis*-[PdCl₂(cod)] (cod = 1,5-cyclooctadiene), the chelate complex [PdCl₂(2-κ²P²,P²)] (**5**) is obtained. The ligand features a different binding mode compared to the gold(I) complexes, now involving the phosphorus atoms (P_A) adjacent to the bridgehead atoms in coordination (Figure 2). The coordination of the Pd atom (square-planar coordination environment) furthermore induces some changes in the structure of the ligand. These include a very large dihedral angle λ (54.1°) as well as a significantly shorter bond between the bridgehead phosphorus atoms. The pattern of the ³¹P{¹H} NMR spectrum of **5** (Figure S15, SI) agrees well with the expected AA'BB'CC'DD' spin system (C₂ symmetry). Remarkably, the change in the chemical shift induced by coordination is enormous. Thus, the coordinating atoms (P_A) are deshielded by 90.1 ppm whereas all other nuclei experience a high-field shift by 45.1 (P_B), 39.1 (P_C) and even 89.0 ppm (P_D) for the bridgehead atoms. All other analytical data, namely mass spectrometry, IR spectroscopy (characteristic tBu vibrations), elemental analysis as well as ¹H{³¹P} NMR (3 singlets) and ¹³C{¹H,³¹P} NMR spectroscopy (6 singlets) are in excellent agreement with the observed structure.




Figure 2. Molecular structure of complex **5** (H atoms and co-crystallized solvent molecule omitted for clarity). Selected bond lengths [Å] and angles [°]: Pd1–P1 2.2657(4), Pd1–P5 2.2568(4), P4–P8 2.1564(5), P1–Pd1–P5 89.06(1), P1–Pd1–Cl2 90.89(1), P5–Pd1–Cl1 91.722(14), λ 54.1(1).

In conclusion, octaphosphane **2** exhibits a variety of different coordination modes, as shown in complexes **3–5**. The peripheral P atoms (P_B and equivalent) seem to be slightly better donors. Chelation of a metal, however, requires the coordination of the P atoms adjacent to the bridgehead atoms (P_A and equivalent). Discovery of additional coordination modes in the future are very likely. Further insights might also be gained preparing complexes with NMR active metals such as Rh. These results once again show how complex the coordination chemistry of an apparently

simple phosphine molecule can be.^[9] They furthermore underline the similarity between phosphorus-rich compounds and they isolobal organic molecules.

Acknowledgements

Financial support from the Studienstiftung des deutschen Volkes (doctoral grant for T.G.) and the Graduate School BuildMoNa are gratefully acknowledged.

Keywords: Isomerization • Oligophosphanes • Gold • Palladium • Phosphorus

- [1] a) H. Köhler, A. Michaelis, *Ber. Dtsch. Chem. Ges.* **1877**, *10*, 807–814; b) G. Boeck, T. Peppel, D. Selent, A. Schulz, *Nachr. Chem.* **2017**, *65*, 1030–1033.
- [2] a) M. Baudler, K. Glinka, *Chem. Rev.* **1993**, *93*, 1623–1667; b) M. Baudler, K. Glinka, *Chem. Rev.* **1994**, *94*, 1273–1297.
- [3] a) M. Caporali, L. Gonsalvi, A. Rossin, M. Peruzzini, *Chemical reviews* **2010**, *110*, 4178–4235; b) M. Scheer, G. Balázs, A. Seitz, *Chemical reviews* **2010**, *110*, 4236–4256.
- [4] A. Kircali, R. Frank, S. Gómez-Ruiz, B. Kirchner, E. Hey-Hawkins, *ChemPlusChem* **2012**, *77*, 341–344.
- [5] a) A. Kircali, P. Lönnecke, E. Hey-Hawkins, *Z. Anorg. Allg. Chem.* **2014**, *640*, 271–274; b) D. M. Yufanyi, T. Grell, M.-B. Sárosi, P. Lönnecke, E. Hey-Hawkins, *Pure Appl. Chem.* **2019**, *91*, 785–796.
- [6] M. Yoshifuji, I. Shima, N. Inamoto, K. Hirotsu, T. Higuchi, *J. Am. Chem. Soc.* **1981**, *103*, 4587–4589.
- [7] M. Baudler, S. Akpapoglou, D. Ouzounis, F. Wasgestian, B. Meinigke, H. Budzikiewicz, H. Münster, *Angew. Chem.* **1988**, *100*, 288–289; *Angew. Chem. Int. Ed.*, **1988**, *27*, 280–281.
- [8] K. B. Dillon, F. Mathey, J. F. Nixon, *Phosphorus. The carbon copy; from organophosphorus to phospho-organic chemistry*, Wiley, Chichester, **1998**.
- [9] T. Grell, E. Hey-Hawkins, *Eur. J. Inorg. Chem.* **2019**; accepted.
- [10] a) A. Schisler, P. Lönnecke, E. Hey-Hawkins, *Phosphorus Sulfur Silicon Relat. Elem.* **2010**, *177*, 1447–1450; b) S. Gómez-Ruiz, E. Hey-Hawkins, *Coord. Chem. Rev.* **2011**, *255*, 1360–1386; c) S. Gómez-Ruiz, E. Hey-Hawkins, *New J. Chem.* **2010**, *34*, 1525–1532.
- [11] M. Baudler, T. Pontzen, J. Hahn, H. Ternberger, W. Faber, *Z. Naturforsch. B* **1980**, *35*, 517–521.
- [12] C. von Hänisch, D. Fenske, *Z. Anorg. Allg. Chem.* **1997**, *623*, 1040–1042.
- [13] N. W. Tun, I. Y. Shchapin, A. I. Nekhaev, *Pet. Chem.* **2013**, *53*, 335–340.
- [14] a) M. Baudler, J. Hahn, V. Arndt, B. Koll, K. Kazmierczak, E. Därr, *Z. Anorg. Allg. Chem.* **1986**, *538*, 7–20; b) M. Baudler, H. Jachow, *Z. Anorg. Allg. Chem.* **1990**, *580*, 27–35; c) M. Baudler, H. Jachow, K.-F. Tebbe, *Z. Anorg. Allg. Chem.* **1991**, *593*, 9–16.
- [15] M. Baudler, Y. Aktalay, *Z. Anorg. Allg. Chem.* **1983**, *496*, 29–39.
- [16] a) B. Riegel, A. Pfitzner, Y. G. Heckmann, H. Binder, E. Fluck, *Z. Anorg. Allg. Chem.* **1995**, *621*, 1365–1372; b) M. Baudler, H. Tschäbunin, *Z. Anorg. Allg. Chem.* **1984**, *511*, 77–83; c) R. Schmidt, S. A. Moya, D. Villagra, H. Binder, Y. G. Heckmann, *J. Chil. Chem. Soc.* **1996**, *41*,


- 371–375; d) G. Fritz, J. Härer, K. Stoll, *Z. Anorg. Allg. Chem.* **1983**, *504*, 47–54.
- [17] J. Hahn in *Methods in stereochemical analysis*, Vol. 8 (Hrsg.: J. G. Verkade), VCH, Weinheim, **1987**.
- [18] M. Bädler, E. Därr, G. Binsch, D. S. Stephenson, *Z. Naturforsch. B* **1984**, *39*, 1671–1675.
- [19] M. Baudler, J. Hellmann, P. Bachmann, K.-F. Tebbe, R. Fröhlich, M. Fehér, *Angew. Chem.* **1981**, *93*, 415–417; *Angew. Chem. Int. Ed. Engl.*, **1981**, *20*, 406–408.
- [20] a) D. M. Stefanescu, H. F. Yuen, D. S. Glück, J. A. Golen, A. L. Rheingold, *Angew. Chem.* **2003**, *115*, 1076–1078; *Angew. Chem. Int. Ed.* **2003**, *42*, 1046–1048; b) E. J. Fernández, M. C. Gimeno, P. G. Jones, A. Laguna, M. Laguna, J. M. López-de-Luzuriaga, *Angew. Chem.* **1994**, *106*, 96–98; *Angew. Chem. Int. Ed.*, **1994**, *33*, 87–88; c) H. Schmidbaur, T. Pollok, G. Reber, G. Müller, *Chem. Ber.* **1988**, *121*, 1345–1348.
-

Entry for the Table of Contents (Please choose one layout)

Layout 1:

COMMUNICATION

The octaphosphane $\{\text{cyclo}-(\text{P}_4\text{tBu}_3)\}_2$ unexpectedly isomerizes to a new constitutional isomer in the presence of Lewis acidic metal salts. The mechanism of this reaction is elucidated and furthermore compared with the isolobal organic compounds. This novel octaphosphane 2,2',2'',2''',3,3'-hexa-*tert*-butyl-bicyclo[3.3.0]octaphosphane can coordinate up to two gold(I) chloride fragments while a tricyclic complex is formed with palladium(II).



Toni Grell, Evamarie Hey-Hawkins*

Page No. – Page No.

**Unexpected Isomerization of
Hexa-*tert*-butyl-octaphosphane**