



Contents lists available at ScienceDirect

# Safety and Health at Work

journal homepage: [www.e-shaw.net](http://www.e-shaw.net)

## Letter

# Chemical Exposure, Risk of Multiple Chemical Sensitivity, and Occupational Safety



### Keywords:

Chemical intolerance  
Hyperosmia  
Metals  
Staff persons  
Workplace chemicals

The study by Pérez-Crespo et al [1] reports interesting results about the prevalence of multiple chemical sensitivity (MCS) among university researchers in Spain [1]. Surprisingly, they have found no difference in the prevalence of MCS between women and men [1]. This differs strikingly from that in all previous data. Consistent with published literature, women are particularly at risk from chemical sensitivity caused by chemicals. We would like to recommend few additions to their study.

First and foremost, the article by Pérez-Crespo [1] does not mention or refer to possible chemical agents (a wide variety of chemical agents) used among participants employed as researchers in an academic setting [1] (refer Table 1 of our article).

Because multiple chemical toxicants may exist among laboratory workers engaged in their research investigations, we expected that the article by Pérez-Crespo et al. [1] would contain a rough approximation of chemical substances used by laboratory staff in their departments at the university (i.e., organic solvents, acids, amines, aldehydes, alcohols, ethers, halogens, and metals/non-metals/metalloids).

As a related example, occupational exposure to organic solvents may lead to chemical odor intolerance, which is cacostmia, in 60% of

workers exposed to solvents, as reported in an occupational health study [2].

Second, the first sentence of the Background section of the abstract states “Multiple Chemical Sensitivity (MCS) is an acquired disease which etiology remains unknown.” [1].

Pesticide [3–9] and toxic metal [10–13] exposure has long been known to be associated with MCS. We have found [10–12] that exposure to metallic elements is an important etiologic factor in MCS in the general Italian population [10–12]. In fact, it has been suggested that MCS is being diagnosed in an increasing number of adults with well-documented exposure to toxic and/or transition metals [10–12].

Furthermore, our previous studies [10–12] in patients with MCS provided clear evidence that allergic sensitization resulted from exposure to transition metals [10–12]. More importantly, epidemiologic data have suggested a link between pesticide exposure and MCS in adults [3–9]. High blood levels of p,p'-dichlorodiphenyldichloroethylene (DDE, an organochlorine pesticide used in agriculture and pest control) were associated with MCS [14].

Baines et al. [14] found a significant difference between the patients who were diagnosed with MCS and those who were in the control group [14].

Finally, for comparison, in our cohort of 175 patients who were given a medical diagnosis of MCS [11], in contrast to that of Pérez-Crespo et al. [1], none of them were employed as university researchers or laboratory staff in a laboratory.

We believe that these important risk factors for the development of MCS should be communicated accurately to the medical and scientific community.

### Funding

This study did not receive funding.

### Conflicts of interest

The authors have no conflicts of interest connected with this work.

### References

- [1] Perez-Crespo J, Lobato-Canon R, Solanes-Puchol A. Multiple chemical sensitivity in chemical laboratory workers. *Saf Health Work* 2018;9:473–8.
- [2] Morrow LA, Ryan CM, Hodgson MJ, Robin N. Alterations in cognitive and psychological functioning after organic solvent exposure. *J Occup Med* 1990;32:444–50.

**Table 1**

Chemicals potentially linked to multiple chemical sensitivity (MCS) in humans.

Chemical agents
Alcohols
Ammonia
Amines
Aldehydes
Carboxylic acids
Halogens
Ketones
Metallic elements
Organic solvents
Polycyclic aromatic hydrocarbons (PHAs)

- [3] Bell IR, Schwartz GE, Peterson JM, Amend D. Self-reported illness from chemical odors in young adults without clinical syndromes or occupational exposures. *Arch Environ Health* 1993;48:6–13.
- [4] Brown AE. Developing a pesticide policy for individuals with multiple chemical sensitivity: considerations for institutions. *Toxicol Ind Health* 1999;15:432–7.
- [5] Lee HS, Hong SY, Hong ZR, Gil HO, Yang JO, Lee EY, et al. Pesticide-initiated idiopathic environmental intolerance in South Korean farmers. *Inhal Toxicol* 2007;19:577–85.
- [6] Pall ML. NMDA sensitization and stimulation by peroxynitrite, nitric oxide, and organic solvents as the mechanism of chemical sensitivity in multiple chemical sensitivity. *FASEB J* 2002;16:1407–17.
- [7] Reid S, Hotopf M, Hull L, Ismail K, Unwin C, Wessely S. Multiple chemical sensitivity and chronic fatigue syndrome in British Gulf War veterans. *Am J Epidemiol* 2001;153:604–9.
- [8] Caress SM, Steinemann AC. A review of a two-phase population study of multiple chemical sensitivities. *Environ Health Perspect* 2003;111:1490–7.
- [9] Miller CS, Mitzel HC. Chemical sensitivity attributed to pesticide exposure versus remodeling. *Arch Environ Health* 1995;50:119–29.
- [10] Pigatto PD, Minoia C, Ronchi A, Brambilla L, Ferrucci SM, Spadari F, et al. Allergological and toxicological aspects in a multiple chemical sensitivity cohort. *Oxid Med Cell Longev* 2013;2013:356235.
- [11] Pigatto PD, Ronchi A, Dolcetta D, Brambilla L, Ferrucci S, Passoni M, et al. Exposure to metals, multiple chemical sensitivity and neurogenic inflammation. *J Clin Toxicol* 2018;8:1.
- [12] Guzzi G, Ronchi A, Barbaro M, Spadari F, Bombeccari G, Brambilla L, et al. Multiple chemical sensitivity and toxic metals. *Toxicol Lett* 2016;258(Suppl. 1):s113.
- [13] Pigatto PD, Guzzi G. Prevalence and risk factors for multiple chemical sensitivity in Australia. *Prev Med Rep* 2019;14:100856.
- [14] Baines CJ, McKeown-Eyssen GE, Riley N, Cole DE, Marshall L, Loescher B, et al. Case-control study of multiple chemical sensitivity, comparing haematology,

biochemistry, vitamins and serum volatile organic compound measures. *Occup Med (Lond)* 2004;54:408–18.

Paolo D. Pigatto

*Clinical Dermatology, IRCCS Istituto Ortopedico Galeazzi, Department of Biomedical, Surgical and Dental Sciences, University of Milan, Milan, Italy*

Anna Ronchi

*Istituti Clinici Scientifici Maugeri IRCCS, Laboratory of Experimental and Clinical Toxicology, Toxicology Unit, Pavia Poison Control Center and National Toxicology Information Center, Pavia, Italy*

Gianpaolo Guzzi\*

*Italian Association for Metals and Biocompatibility Research – A.I.R.M.E.B., Milan, Italy*

\*Corresponding author. *Italian Association for Metals and Biocompatibility Research – A.I.R.M.E.B. (not-for-profit organization), Via A. Banfi, 4, 20122 Milan, Italy.*

*E-mail address: [gianpaolo\\_guzzi@fastwebnet.it](mailto:gianpaolo_guzzi@fastwebnet.it) (G. Guzzi)*

1 July 2019

Available online 12 June 2020