



ZINC-FINGER interactions mediate transcriptional regulation of hypocotyl growth in *Arabidopsis*

Giorgio Perrella^a, Mhairi L. H. Davidson^a, Liz O'Donnell^a, Ana-Marie Nastase^{a,b}, Pawel Herzyk^{a,b}, Ghislain Breton^{c,1}, Jose L. Pruneda-Paz^c, Steve A. Kay^{c,2}, Joanne Chory^{d,e}, and Eirini Kaiserli^{a,d,3}

^aInstitute of Molecular, Cell and Systems Biology, University of Glasgow, G128QQ Glasgow, United Kingdom; ^bGlasgow Polyomics, University of Glasgow, G611QH Bearsden, United Kingdom; ^cSection of Cell and Developmental Biology, Division of Biological Sciences, University of California, San Diego, La Jolla, CA 92093; ^dPlant Biology Laboratory, The Salk Institute for Biological Studies, La Jolla, CA 92037; and ^eHoward Hughes Medical Institute, The Salk Institute for Biological Studies, La Jolla, CA 92037

Edited by Christian Fankhauser, University of Lausanne, 1015 Lausanne, Switzerland, and accepted by Editorial Board Member Caroline Dean March 24, 2018 (received for review October 17, 2017)

Integration of environmental signals and interactions among photoreceptors and transcriptional regulators is key in shaping plant development. TANDEM ZINC-FINGER PLUS3 (TZP) is an integrator of light and photoperiodic signaling that promotes flowering in *Arabidopsis thaliana*. Here we elucidate the molecular role of TZP as a positive regulator of hypocotyl elongation. We identify an interacting partner for TZP, the transcription factor ZINC-FINGER HOMEODOMAIN 10 (ZFHD10), and characterize its function in coregulating the expression of blue-light-dependent transcriptional regulators and growth-promoting genes. By employing a genome-wide approach, we reveal that ZFHD10 and TZP coassociate with promoter targets enriched in light-regulated elements. Furthermore, using a targeted approach, we show that ZFHD10 recruits TZP to the promoters of key coregulated genes. Our findings not only unveil the mechanism of TZP action in promoting hypocotyl elongation at the transcriptional level but also assign a function to an uncharacterized member of the ZFHD transcription factor family in promoting plant growth.

light | transcription | *Arabidopsis* | zinc-finger | development

Detention and flowering provide great examples in which environmental and endogenous signal integration is required to mediate major transitions and morphological changes during early and late plant development. Light, temperature, photoperiod, clock, and hormone pathways share signaling components that coordinate physiological responses such as hypocotyl elongation and flowering (1–3). Genome-wide transcript analysis, proteomic, imaging, and genetic studies indicate that photoreceptors and circadian and hormone signaling components converge in the nucleus and act on gene expression at the level of transcription (1, 2, 4–11). Direct interactions among plant photoreceptors, key transcription factors (TFs) such as the bHLH family of PHYTOCHROME INTERACTING FACTORS (PIFs), BRASSINAZOLE RESISTANT 1 (BZR1), and TIMING OF CAB EXPRESSION (TOC1), as well as transcriptional regulators, such as HISTONE DEACETYLASES (HDA15, HDA19), TOPLESS (TPL), and scaffold proteins such as EARLY FLOWERING 3 (ELF3), have been shown to play important roles in controlling the expression of growth-promoting genes (2, 6–8, 12–17). Furthermore, recent reports have identified protein components that contribute to the repertoire of signal integrators of light and circadian regulation of hypocotyl elongation and photoperiodic flowering (7, 8, 18).

TANDEM ZINC-FINGER PLUS3 (TZP) was originally identified as a regulator of morning-specific hypocotyl growth based on a Quantitative Trait Locus study on recombinant-inbred lines of two natural *Arabidopsis* ecotypes (*Shahdara* and *Bay-0*) (19). Microarray analyses showed that TZP up-regulates the expression of growth-promoting and auxin-related genes in a blue-light and dawn-specific manner; however, the molecular mechanism of its action remains unknown (19). Recent studies have revealed a dual role for TZP in regulating plant development. In addition to its role in blue-light-induced photomorphogenesis, TZP was recently shown to integrate light and photoperiodic signaling by interacting

with the red-light receptor phytochrome B (phyB) in transcriptional nuclear photobodies (8). More specifically, ChIP assays and gene expression analyses revealed that TZP regulates the expression of the flower inducer *FLOWERING LOCUS T (FT)* to promote flowering initiation in long days (8). An independent study employing a proteomic approach showed that TZP is part of an in planta multiprotein complex consisting of known and novel light signaling and circadian components including phytochromes A-E, ELF3, and PHOTOPERIODIC CONTROL OF HYPOCOTYL1 (PCH1), placing TZP at the crossroads of light, clock, and photoperiodic pathways (7, 20, 21).

Here we uncover the molecular mechanism of TZP action in regulating blue-light-induced hypocotyl elongation. TZP acts at the transcriptional level by recruiting a member of the plant-specific

Significance

Light coordinates energy production, growth, and survival throughout plant development. In *Arabidopsis*, light stimulates transcriptional reprogramming during developmental transitions such as photomorphogenesis and flowering through the action of photoreceptors, transcription factors, and signaling components. Here we assign a function to a member of the zinc-finger homeodomain (ZFHD) transcription factor family in regulating light-induced development. Our findings reveal ZFHD10 to be a missing link in understanding how the recently discovered integrator of light and photoperiodic flowering, TANDEM ZINC-FINGER PLUS3 (TZP), controls the expression of growth-promoting transcriptional regulators via direct association with light-regulated promoter elements. Elucidating how such novel protein complexes coordinate gene expression will allow scientists and breeders to optimize plant growth and development in response to unfavorable environmental conditions.

Author contributions: G.P., M.L.H.D., G.B., J.L.P.-P., S.A.K., J.C., and E.K. designed research; G.P., M.L.H.D., L.O., G.B., J.L.P.-P., and E.K. performed research; P.H., G.B., J.L.P.-P., and S.A.K. contributed new reagents/analytic tools; G.P., M.L.H.D., A.-M.N., P.H., G.B., and E.K. analyzed data; and E.K. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission. C.F. is a guest editor invited by the Editorial Board.

This open access article is distributed under [Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 \(CC BY-NC-ND\)](https://creativecommons.org/licenses/by-nc-nd/4.0/).

Data deposition: The sequences reported in this paper have been deposited in the National Center for Biotechnology Information Gene Expression Omnibus database and Sequence Read Archive series (accession no. [GSE112889](https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE112889)).

¹Present address: Department of Integrative Biology and Pharmacology, McGovern Medical School, University of Texas Health Science Center at Houston, Houston, TX 77030.

²Present address: Department of Neurology, University of Southern California, Los Angeles, California.

³To whom correspondence should be addressed. Email: eirini.kaiserli@glasgow.ac.uk.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1718099115/-DCSupplemental.

Published online April 23, 2018.




Fig. 1. Identification of ZFHD10 as an interacting partner for TZP, using a genome-wide *Arabidopsis* TF library. (A) Yeast-two-hybrid analysis of pDEST32-TZP (GAL4BD-bait) and pDEST22-ZFHD10 (GAL4AD-prey) interactions by assessing growth on nonselective media (L⁻W⁻), selective media (L⁻W⁺ 100 mM 3-AT), or the x-galactosidase assay. pD22, pDEST22; pD32, pDEST32; 3-AT, 3-amino-1,2,4-triazole. (B) Schematic representation of TZP and ZFHD10 domain composition as a guide for the deletion constructs used for the interaction studies shown in Fig. 1C and *SI Appendix, Fig. S1 E and F*. (C) Deletion analysis of the interaction between TZP and ZFHD10. Yeast-two-hybrid analysis of pDEST32-TZP, pDEST32-TZP^{Nt}, pDEST32-TZP^{ZFPLUS}, pDEST32-TZP^{PLUS} (GAL4BD-bait), pDEST22-ZFHD10, pDEST22-ZFHD10^{ZF}, pDEST22-ZFHD10^{HD} (GAL4AD-prey) interaction, using the quantitative β -galactosidase assay. Data shown are representative of three independent experimental repeats.

zinc-finger homeodomain (ZFHD) TF family, ZFHD10. *Arabidopsis* contains 14 ZFHD family members with limited information on their molecular function in shaping plant growth, patterning, and embryo development, contrary to the well-established role of the closely related HD-ZIP TFs (22–25). Our genome-wide ChIP studies and quantitative transcript analysis reveal that TZP and ZFHD10 associate with common promoter elements of transcriptional regulators, light-responsive and growth-promoting genes required for hypocotyl elongation. Moreover, our study assigns a role for ZFHD TFs in light-regulated plant development via a direct interaction with TZP.

Results

TZP Interacts with a Member of the ZFHD Transcription Factor Family.

TZP contains two ZF domains and a PLUS3 domain, all of which have potential nucleic acid binding activity; however, there is no evidence suggesting that these domains confer transcriptional activation or repression activity (8, 19, 26–28). To investigate the molecular role of TZP in regulating transcriptional control of gene expression, a large-scale directed yeast-two-hybrid screen was performed using a gold standard TF ORFeome library (29) with TZP as the bait (*SI Appendix, Fig. S1B*). Nine putative TZP-interacting TFs were identified and verified, seven of which were members of the TEOSINTE BRANCHED 1/CYCLOIDEA/PROLIFERATING CELL NUCLEAR ANTIGEN FACTOR (TCP) TF family, one abscisic-acid-responsive NAC TF (ANAC), and one ZFHD TF (*SI Appendix, Fig. S1 B and C*). No autoactivation was observed for either the bait (TZP) or the positive prey TFs (*SI Appendix, Fig. S1 A and D*). The strongest interaction was observed between TZP and ZFHD10 (At5g39760), a member of the ZFHD TF family, which became the central focus of this study (Fig. 1A and *SI Appendix, Fig. S1 B and C*).

Deletion analysis revealed that the ZF of ZFHD10 is sufficient to associate with the ZFPLUS3 and PLUS3 domains of TZP (Fig.

1B and C and *SI Appendix, Fig. S1E*). ZFHD10 showed homodimerization properties through the ZF (CHCC3H2) motif consistent with a previous report (*SI Appendix, Fig. S1F*) (22). No interaction was observed between TZP and ZFHD9, the closest ZFHD family member to ZFHD10 or the homeodomain-leucine zipper (HD-ZIP), ATHB23, recently shown to interact with phyB (*SI Appendix, Fig. S1G*) (30).

To validate the interaction between TZP and ZFHD10 in planta, we used bimolecular fluorescence complementation (31, 32). C-terminal translational fusions of ZFHD10-nYFP (spyNe) and TZP-cYFP (spyCe) showed reconstitution of YFP fluorescence in the nucleus when coexpressed transiently in *Nicotiana benthamiana* epidermal cells, whereas no signal was detected for the expression of TZP-cYFP or ZFHD10-nYFP with the empty vector controls (spyNe and spyCe, respectively; Fig. 2A and *SI Appendix, Fig. S2C*). Colocalization studies also confirmed that TZP-mCherry and ZFHD10-GFP reside in the nucleus in light-grown plants (Fig. 2B). In planta coimmunoprecipitation (co-IP) was also performed to further verify the interaction between TZP-GFP and ZFHD10-RFP (Fig. 2C and *SI Appendix, Fig. S2D*). Overall, these data demonstrate that TZP and ZFHD10 interact in yeast and in planta.

TZP and ZFHD10 Promote Hypocotyl Elongation.

qRT-PCR showed that *ZFHD10* is abundant in seedlings primarily when grown in blue light, which correlates with the expression pattern of *TZP*, whereas no correlation was observed for *ZFHD9* (*SI Appendix, Fig. S3 D–G*). No significant diurnal regulation was observed for *ZFHD10* transcript or protein abundance in transgenic lines expressing 35S_{pro}ZFHD10-GFP/Col-0 (OXZFHD10) (*SI Appendix, Fig. S3C*) (33, 34). A closer look at seedlings revealed that TZP and ZFHD10 are present in the cotyledons as well as the hypocotyl, with an increase in abundance in the hypocotyl apex (*SI Appendix, Fig. S3I* and refs. 35 and 36). TZP is also highly expressed in roots (*SI Appendix, Fig. S3 F and H* and ref. 37), suggesting a potential role in other tissues.

To further explore the physiological significance of TZP-ZFHD10 interactions, we examined the photomorphogenic phenotypes of knockout and overexpressing lines for ZFHD10 and TZP (Fig. 3 and *SI Appendix, Figs. S3 A and B, S4, and S5 A and B*).




Fig. 2. TZP colocalizes and interacts with ZFHD10 in planta. (A) Bimolecular fluorescence complementation assay shows YFP reconstitution between TZP-spyCe and ZFHD10-spyNe when coexpressed transiently in *N. benthamiana* leaves. (Negative and positive controls are shown in *SI Appendix, Fig. S2C*.) (B) Representative images of *N. benthamiana* leaves coexpressing TZP-mCherry and ZFHD10-GFP. (C) Coimmunoprecipitation analysis of TZP-GFP and ZFHD10-RFP coexpressed transiently in *N. benthamiana*. Single infiltration of TZP-GFP or ZFHD10-RFP were used as negative controls. Plants were grown in white light before examination using confocal microscopy and coimmunoprecipitation. Data shown are representative of three independent experimental repeats. (Scale bars, 20 μ m.)




Fig. 3. ZFHD10 and TZIP promote hypocotyl elongation. (A) Hypocotyl measurements and representative images of homozygous transgenic *Arabidopsis* lines overexpressing ZFHD10 or TZIP in Col-0. Plants were grown for 7 d in blue light ($1 \mu\text{mol m}^{-2}\text{s}^{-1}$). (B) Hypocotyl measurements of the indicated genotypes. Plants were grown for 7 d in blue light ($1 \mu\text{mol m}^{-2}\text{s}^{-1}$). Col-0 was used as the wild-type control. Representative images are shown in *SI Appendix, Fig. S5A*. The data presented are mean \pm SE ($n = 15$ seedlings). (C) qRT-PCR analysis of *HFR1*, *ATHB2*, *ATXTH17*, and *PIF7* mRNA normalized to housekeeping gene *ISU1* of the indicated genotypes. Seedlings were grown in continuous blue light ($1 \mu\text{mol m}^{-2}\text{s}^{-1}$) for 7 d. Bars are means \pm SE ($n = 4$ technical replicates). Graphs are representative of three independent experimental repeats. Asterisks indicate difference to Col-0 at $P < 0.05$. An independent biological repeat is shown in *SI Appendix, Fig. S5C*.

It has previously been shown that overexpression of TZIP promotes hypocotyl elongation, whereas a premature stop codon within the *TZIP* locus in Bay-0 results in shorter hypocotyls primarily in response to blue light (19). Hypocotyl elongation measurements showed that OXZFHD10 phenocopies OXTZP in response to blue-light irradiation, whereas *tzp* and *zfh10* knockout/knockdown mutants exhibited shorter hypocotyls relative to the wild-type (Col-0), OXTZP, and OXZFHD10, primarily under low fluence rate blue light (Fig. 3A and B and *SI Appendix, Fig. S5A and B*) (19). We also generated and examined transgenic lines overexpressing TZIP in the *zfh10* background, and although they showed partial rescue of the mutant *zfh10* phenotype, they never reached the level of elongation demonstrated by OXTZP/Col-0 (Fig. 3B). Overexpression of ZFHD10 in the absence of functional TZIP (OXZFHD10/Bay-0) showed rescue of the short hypocotyl Bay-0 phenotype (*SI Appendix, Fig. S6*). These observations suggest that TZIP requires, at least in part, ZFHD10 for function and positions ZFHD10 downstream of TZIP action with regard to blue-light-regulated hypocotyl elongation.

ZFHD10 and TZIP Regulate Common Growth-Promoting Gene Targets.

TZIP is known to regulate the expression of growth-promoting genes based on microarray analysis (19). At this time, there is no report of transcriptional targets for ZFHD10. Because ZFHD10 interacts with TZIP and regulates hypocotyl elongation, we were interested in examining whether ZFHD10 controls the expression of TZIP-regulated growth-promoting genes (19). Indeed, ZFHD10 induces the expression of *ATHB2*, *LONG HYPOCOTYL IN FAR-RED (HFR1)*, *XYLOGLUCAN ENDOTRANSGLUCOSYLASE/HYDROLASE 17 (ATXTH17)*, and *PIF7*, as shown by qRT-PCR analysis performed on 7-day-old seedlings irradiated with low blue light (Fig. 3C and *SI Appendix, Fig. S5C*). *ATHB2* is an HD-ZIP TF that promotes hypocotyl growth under low-light conditions and in response to auxin (38–40). *HFR1* is a non-DNA-binding bHLH TF that associates with PIF TFs to regulate stem elongation in response to far-red, blue, and shade (38, 41–43). *ATXTH17* is a light-induced

plant cell wall enzyme essential for growth promotion in response to light and hormones (44, 45). *PIF7* is known to accumulate in response to light and interacts with phyB to promote plant growth in response to shade (4, 46, 47). An increase in the expression of these genes and the function of the proteins they encode has been associated with light-regulated hypocotyl elongation (19).

OXTZP and OXZFHD10 show increased transcript levels of the aforementioned genes, whereas *tzp* and *zfh10* mutants show partial reduction in their expression (Fig. 3C and *SI Appendix, Fig. S5C*). Collectively, TZIP and ZFHD10 show a similar trend in the regulation of the expression of well-characterized genes, which directly correlate with the hypocotyl phenotypes described earlier.

To test whether TZIP and ZFHD10 could directly control transcription by associating with the promoter regions of the genes they coregulate, we performed ChIP assays followed by qPCR. TZIP associates preferentially with the transcriptional start site (TSS) of *ATHB2*, *ATXTH17*, and *PIF7* and with the G-box (CACGTG), a well-characterized light-regulated element, of *HFR1* (Fig. 4 and *SI Appendix, Fig. S5D*). No significant enrichment was observed for the 3' untranslated region, the *IAA1* promoter, or Col-0 (Fig. 4). ZFHD10 showed a similar pattern of preferential binding to the TSS of *ATHB2*, *PIF7*, as well as the first G-box and TSS of *HFR1* (Fig. 4A, B, and D and *SI Appendix, Fig. S5D*). ZFHD10 showed no enrichment on two canonical ZFHD binding sites on the *ATHB2* promoter, whereas binding was observed on the HUD element (Hormone Up at Dawn) of *ATXTH17_{pro}* (Fig. 4B and C and *SI Appendix, Fig. S5D*) (19, 22, 33, 48).

We previously showed that TZIP^{PLUS3} is sufficient for binding nonspecific ssDNA in vitro, so we were interested in investigating whether the interaction of TZIP with ZFHD10 confers sequence specificity and guides TZIP to specific promoter sequences (8). ChIP-qPCR on transgenic plants expressing equal protein levels of TZIP-GFP in Col-0 and *zfh10* showed a considerable decrease in the recruitment of TZIP on *HFR1*, *ATHB2*, and *ATXTH17* promoters (Fig. 5 and *SI Appendix, Fig. S5E*). These results




Fig. 4. TZP and ZFHD10 associate with common genomic regions of growth-promoting genes. (A–D) Relative enrichment of TZP and ZFHD10 on *HFR1*, *ATHB2*, *ATXTH17*, and *PIF7* loci. Col-0, a region in the 3' untranslated region of each locus, and the *IAA1* promoter were used as negative controls. Seedlings were grown for 7 d under blue light ($1 \mu\text{mol m}^{-2}\text{s}^{-1}$). Bars are means \pm SE ($n = 4$ technical replicates). Graphs shown are representative of three independent experimental repeats. An independent experimental repeat is shown in *SI Appendix, Fig. S5D*. G, G-box promoter element; HUD, Hormone Up at Dawn promoter element (CACATG); ZfHD, Zinc finger Homeo-Domain binding site TAAATTG.

suggest that ZFHD10 is important for recruiting TZP on specific promoter sequences.

TZP and ZFHD10 Associate with Common Chromatin Regions Enriched in Light-Responsive Elements. In addition to examining whether TZP and ZFHD10 regulate and bind to promoters of genes

known to be involved in light-regulated hypocotyl elongation, an unbiased genome-wide approach was also performed using ChIP coupled to next-generation sequencing on 7-day-old GFP-tagged transgenic lines irradiated with low blue light. Bioinformatics analysis on data obtained by ChIP sequencing revealed that TZP and ZFHD10 preferentially associate with promoter regions, as




Fig. 5. ZFHD10 is important for the recruitment of TZP on light-regulated promoters. (A) Western blot analysis of TZP protein levels in OX TZP/Col-0 and OX TZP/*zfhd10-1* transgenic lines. Col-0 was used as a negative control for the anti-GFP antibody, and UGPase was used as a loading control. Relative enrichment of TZP on *HFR1* (B), *ATHB2* (C), *PIF7* (D), and *ATXTH17* (E) loci when expressed in Col-0 or *zfhd10* mutant background. Col-0 and the 3' untranslated region were used as negative controls. Seedlings were grown for 7 d under blue light ($1 \mu\text{mol m}^{-2}\text{s}^{-1}$). Bars are means \pm SE ($n = 4$ technical replicates). Graphs shown are representative of two independent experimental repeats. An independent experimental repeat is shown in *SI Appendix, Fig. S5E*. G, G-box.




Fig. 6. TZIP and ZFHD10 bind to common genomic regions. (A) Relative binding-peak distribution of TZIP and ZFHD10 across genomic regions. Values were normalized to those of Col-0. (B) Venn diagram depicting the overlap between TZIP and ZFHD10 targets (promoters and TSS) determined by whole-genome ChIP sequencing analysis. (C) Enriched motifs identified within 1 kb of the surrounding peak summits for overlapping TZIP and ZFHD10 ChIP-seq peaks.

80% and 50% of ZFHD10 and TZIP genomic binding sites, respectively, were within 1 kb of the TSS of gene loci (Fig. 6A). Data analysis using stringent parameters revealed a significant amount of promoter regions corresponding to 1,137 genes bound by both TZIP and ZFHD10 (Fig. 6B and Dataset S2). The number of targets bound exclusively by ZFHD10 (5,587) was higher than the number identified for TZIP (1,439), possibly because ZFHD10 has a canonical DNA-binding domain (HD), whereas TZIP may associate with specific promoters indirectly via binding to TFs (Fig. 6B).

De novo analysis of consensus motifs enriched within the promoter-binding peaks of TZIP and ZFHD10 identified well-characterized light-, temperature-, and circadian-regulated elements such as the G-box (CACGTG), which is a variation of the E-box, and the SORLIP2 (Sequence Over-represented in Light-Induced Promoters; TGGGCC) (Fig. 6C) (3, 49–51). Furthermore, peaks specifically bound by ZFHD10 showed over-representation of the transcriptional initiator TATA-box (TATATATA). Recent studies have highlighted the significance of interactions between the TATA-box and cis-regulatory light-responsive promoter elements in activating gene expression in plants (52).

To verify the results obtained by ChIP-seq (Fig. 7A, D, and G), we performed ChIP-qPCR on selected genomic targets that showed binding by both TZIP and ZFHD10 (Fig. 7B, E, and H). ChIP of seedlings expressing TZIP-GFP and ZFHD10-GFP exposed to low blue light showed enrichment in the promoter regions of *ARABIDOPSIS AUXIN RESPONSE FACTOR* (*ARF6*), the histone-lysine *N*-methyltransferase *SU(VAR)3-9* (*SUVRI*), and *EARLY FLOWERING 4-LIKE* (*EFL4*; Fig. 7B, E, and H).

In *Arabidopsis*, *SUVRI* promotes transcriptional gene silencing by forming a complex with SNF2-related chromatin-remodeling proteins that modify nucleosome positioning at RNA-directed DNA methylation-dependent and independent loci (53). *EFL4* has 44% sequence identity to *ELF4*, a protein that regulates phyB-mediated seedling deetiolation in response to red light as well as diurnal regulation and flowering time (6, 54). *ARF6* is involved in regulating hypocotyl elongation and flower development (55–57). Recent studies have shown that *ARF6* interacts with *PIF4* and *BZR1* and coregulates genes that promote hypocotyl elongation in response to light, hormones, and temperature (55). Overexpression of TZIP leads to a higher induction of *ARF6*, *SUVRI*, and *EFL4* transcripts than *OXZFHD10*, even though

ZFHD10 showed a greater enrichment on these loci (Fig. 7B, C, E, F, H, and I). ChIP experiments on TZIP/*zfh10* lines showed a reduction in TZIP enrichment on *SUVRI* and *EFL4* promoters (SI Appendix, Fig. S8B and C). However, no difference in TZIP binding is observed for *ARF6* (SI Appendix, Fig. S8A), suggesting this target could be bound by separate TZIP and ZFHD10 protein complexes. Furthermore, *tzp* and *zfh10* mutants showed down-regulation of *SUVRI* and, to a lesser extent, *EFL4* and *ARF6* transcripts (Fig. 7C, F, and I). These observations could be a result of additional TZIP-interacting proteins and photoreceptor pathways regulating the expression of these genes.

Gene ontology analysis of common promoter targets of TZIP and ZFHD10 showed enrichment in TFs (LFY, CCA1, members of the AGAMOUS-like, Dof ZF, WRKY, TCP, ZFHD, and HD-ZIP TF families), transcriptional regulators (ORIGIN OF REPLICATION COMPLEX 1B, NUCLEAR RNA POLYMERASE 1), chromatin remodeling enzymes [HISTONE DEACETYLASE 2 and 19 (HDA2, HDA19), HISTONE MONOUBIQUITINATION 1 (HUB1)], F-box and RNA-binding proteins (splicing factor, poly-A binding proteins, RAD5, MEDIATOR 7), and hormone response factors (ETHYLENE RESPONSE FACTOR 10, EIN3-BINDING F-BOX), suggesting an active role in the transcriptional regulation of gene expression in response to environmental, biotic, abiotic, and endogenous stimuli (SI Appendix, Fig. S9A and Dataset S3).

Promoter regions exclusively associated by TZIP show a huge over-representation in proteins involved in transcription such as C2H2-ZF, WRKY, bHLH, and MYB TFs, as well as TOPLESS-related, brassinosteroid (*BZR1* and *BZR*-like homolog), clock (*PRR9* and *PRR5*), flowering (*VIN3*-like, *FLD*, *SPATULA*), and light signaling components (*PIF4*, *PIL6*, *PAP2*, *phyE*, *cry2*, *CIB1*, *SPA4*), reflecting its involvement in regulating gene expression in response to environmental stimuli such as light and temperature (SI Appendix, Fig. S9B). On the contrary, ZFHD10-specific targets were enriched in RNA processing factors as well as proteins involved in salt stress and ABA signaling, in addition to embryo development (SI Appendix, Fig. S9C). Overall, our findings show that TZIP and ZFHD10 coassociate with and regulate the expression of not only light-regulated loci but also transcriptional regulators to shape plant development in response to environmental stimuli.

Discussion

Plant growth and development are coordinated through the action of multiple light, hormone, and photoperiodic pathways. This study focuses on uncovering the molecular role of TZIP, a signal integrating component, in promoting hypocotyl elongation in response to low blue light.

To elucidate how TZIP promotes growth and regulates gene expression during the early stages of plant development, we screened for TF-interacting partners for TZIP, using a large-scale, directed Y2H approach. Nine proteins exhibited positive interactions of various strength with TZIP, seven of which were members of the TCP TF family (*TCP8*, *TCP9*, *TCP12*, *TCP15*, *TCP20*, *TCP22*, *TCP23*) and one ANAC TF (*ANAC87*; SI Appendix, Fig. S1B). TCP TFs regulate many aspects of plant development, including circadian rhythms and photoperiodic flowering (58–60). However, in this study, we focus on the characterization and functional significance of the strongest TZIP interactor, which is a member of the ZFHD family (Fig. 1).

Homeobox TFs have key functions in shaping animal and plant development (25). In *Arabidopsis*, there are more than 100 proteins that contain DNA-binding HDs. The HD-ZIP class of TFs is one of the most well-characterized classes of HD-containing proteins with major roles in hormone and light-regulated developmental responses (24, 39). A unique combination of ZF and HD modules has given rise to the plant-specific ZFHD class of TFs originally discovered in the C4 plant *Flaveria*




Fig. 7. ChIP sequencing identifies overlapping targets for TZP and ZFHD10. (A, D, and G) Visualization of ChIP-seq data in the genomic regions encompassing three representative genes, *ARF6*, *SUVR1*, and *EFL4*. The ChIP tracks show the pile-up distribution of the raw reads from three pooled biological replicates of ChIP-seq data. Gray bars underneath the peaks indicate differences identified by SICER between TZP vs. Col-0 and ZFHD10 vs. Col-0 (TZP peaks ≥ 1.5 and ZFHD10 \geq twofold change to Col-0). Gene regions are represented as arrow-shaped boxes, which shows transcriptional orientation. (B, E, and H) Relative enrichment of TZP and ZFHD10 on *ARF6*, *SUVR1*, and *EFL4* promoter regions identified by SICER. Col-0 was used as a negative control. Bars are means \pm SE ($n = 4$ technical replicates). (C, F, and I) qRT-PCR analysis of absolute *ARF6*, *SUVR1*, and *EFL4* mRNA levels normalized to housekeeping gene *ISU1* of the above-indicated genotypes. For all the experiments, seedlings were grown in continuous blue light ($1 \mu\text{mol m}^{-2}\text{s}^{-1}$) for 7 d. Bars are means \pm SE ($n = 4$ technical replicates). Graphs are representative of two independent experimental replicates.

(61). Seminal reports on specific *Arabidopsis* ZFHDs clearly support their action as transcriptional regulators of drought tolerance and flower development via direct DNA binding through the HD (22, 48). However, there is no current study characterizing the function of ZFHD10 or the role of ZFHDs in light signaling and photomorphogenesis.

ZF domains are highly conserved and act as sites for protein or nucleic acid association (28). In the case of ZFHD10 and TZP, deletion analyses showed association through ZFPLUS3 (TZP)–ZF(ZFHD10) interactions (Fig. 1 B and C). There is no evidence suggesting that ZFHDs bind DNA as dimers or monomers. It would be interesting to further investigate whether the interaction

of TZP with ZFHD10 interferes with the homodimerization status of ZFHD10 or its heterodimerization with other ZFHD TFs.

Our in planta studies (colocalization, co-IP, hypocotyl measurements) verify that the interaction between TZP and ZFHD10 is physiologically significant and reveal a function for ZFHD10 as a positive regulator of hypocotyl growth in response to low blue light (Figs. 2 and 3 A and B and *SI Appendix, Fig. S5B*). To dissect the molecular mechanism of TZP and ZFHD10 function, we performed gene expression analysis and discovered coregulation of key genes involved in light-induced hypocotyl elongation and showed that ZFHD10 guides TZP to light-regulated promoter sequences (Figs. 3C, 4, and 5 B–E and *SI Appendix,*

Fig. S5 D and E). Furthermore, targeted as well as genome-wide chromatin association studies clearly showed that TZP and ZFHD10 bind to common genomic loci enriched in promoter regions of light-, temperature-, and hormone-regulated growth-promoting genes (Figs. 4, 6, and 7 and Dataset S2).

Research on how transcriptional regulators and chromatin remodeling factors control plant development is currently flourishing; however, there is limited information on how the expression of chromatin-modifying enzymes is regulated by environmental stimuli. Our ChIP sequencing analysis shows that TZP and ZFHD10 associate with the promoters of histone deacetylases and methyltransferases and an E3 ubiquitin ligase that monoubiquitinates H2B. These findings will therefore open avenues in investigating a potential role for TZP and ZFHD10 in controlling the transcript levels of these enzymes and further understanding the role of chromatin remodeling in fine-tuning light-regulated plant growth and development. A putative EAR (Ethylene-responsive element binding factor-associated Amphiphilic Repression) (LxLxL) motif within the TZP PLUS3 domain indicates that TZP could also recruit and directly modulate the action of transcriptional corepressors and histone remodeling enzymes at the protein level (17, 62, 63).

Motif analysis of the TZP-ZFHD10 promoter targets revealed a clear over-representation of light-responsive elements such as the G-box: CACGTG and the SORLIP2 previously associated with cryptochrome and phytochrome regulated gene expression (Fig. 6C) (49–51).

The importance of signal integration among blue (cry1 and cry2) and red/far-red (phyA and phyB) light receptors in regulating plant development, growth, and photoprotection is well established (4, 43, 64–69). Our data show that there is a decrease in the blue-light-dependent induction of TZP in *cry1cry2*, *cry1*, and *phyA* compared with Col-0, suggesting that cry1 and phyA have a redundant role in regulating blue-light-induced TZP expression (SI Appendix, Fig. S7A). Furthermore, ChIP analysis revealed a reduction in the enrichment of TZP on the promoters of blue-light-regulated loci indicating that cry1, cry2, and phyA are important for recruiting TZP to light-regulated genomic regions (SI Appendix, Fig. S7C). Our ChIP and physiological analysis suggests that TZP regulates blue-light-induced hypocotyl elongation through the action of cry1, cry2, and phyA (SI Appendix, Fig. S7). Whether cry1, cry2, and/or phyA can positively or negatively regulate TZP or vice versa through direct interactions or via the action of other TFs requires further investigation.

At this time, we have no evidence suggesting a role for the interaction between TZP and phyB in regulating blue-light-mediated hypocotyl growth (7, 8, 20). ZFHD10 is recruited to nuclear bodies (NBs) only when coexpressed with TZP in *N. benthamiana* exposed to white light, possibly because of the phyB-induced recruitment of TZP (and its partners) to NBs. Neither TZP nor ZFHD10 form NBs in *Arabidopsis* or *N. benthamiana* leaves when exposed to blue light (8). These observations suggest that the blue-light regulation of gene expression conferred by TZP and ZFHD10 is independent of phyB-mediated signaling or red-light-induced NB formation. It would be interesting to investigate whether TZP and ZFHD10 regulate gene expression by interacting or interfering with the action of key bHLH PIF4 and PIF5 TFs in response to low levels of blue light (4, 43).

Future studies are necessary to establish whether TZP and ZFHD10 operate through a PIF-dependent or independent pathway. Employing imaging tools that have recently been developed to study cell-type-specific interactions among key transcriptional regulators of cell fate in *Arabidopsis* roots would provide invaluable insights in understanding how nuclear signal integration influences plant growth and development (70).

The significance of our findings is multivalent. First, we elucidate the molecular and physiological role of TZP as a positive regulator of hypocotyl growth by associating with the promoters of chromatin remodeling enzymes, transcriptional regulators, and




Fig. 8. The role of TZP and ZFHD10 in regulating hypocotyl elongation. TZP acts as a positive regulator of growth downstream of the clock and the photoreceptors. TZP and ZFHD10 directly interact through their ZFPLUS3 and ZF domains, respectively. ZFHD10 recruits TZP to light-regulated promoter elements (G-box, SORLIP), and they act in concert to control the expression of growth-promoting transcriptional regulators in response to blue light (LBL). Whether TZP and ZFHD10 act cooperatively or independently of the PIF TFs remains to be investigated.

well-characterized light-responsive genes. Second, we identify ZFHD10 as the TF that directly associates and acts in concert with TZP to regulate the expression of genes required for hypocotyl elongation in response to low blue light. Last and very important, we report a function for a member of the unique family of ZFHD TFs in light-induced plant growth. Our data position TZP and ZFHD10 downstream of the clock and the photoreceptors with direct links to light, temperature, and hormone signal integration, which is essential for coordinating optimal plant growth and development in response to a constantly changing environment (Fig. 8).

Materials and Methods

Plant material and growth conditions are described in SI Appendix, SI Materials and Methods. The ProQuest Two-Hybrid System (Invitrogen) was used to identify and verify direct interactions between TZP (bait, pDEST32) and a library of TFs (prey, pDEST22) by sequential transformation of the yeast strain MaV203. Construction of the pDEST22-TF library screened has been described previously (29). For more information, please see SI Appendix, SI Materials and Methods. Protein extraction and Western blot analysis are described in SI Appendix, SI Materials and Methods. Transient protein expression in *N. benthamiana* using Agrobacterium-mediated infiltration was performed as previously described (8). Coimmunoprecipitation was performed using the μ MACS GFP Tag Protein Isolation Kit (Miltenyi Biotech), using 500 μ g total protein extracted as described for *Arabidopsis* (6, 8). Confocal microscopy was performed with Leica SP2 and SP8 inverted microscopes, and image analysis was performed as described previously (8). Representative images from three independent biological repeats are shown in this study. Transient expression and imaging in *N. benthamiana* was performed as described previously (32). Cloning and genotyping strategies are described in SI Appendix, SI Materials and Methods. Quantitative RT-PCR and ChIP assays were performed as described previously, with minor modifications (8, 71). SI Appendix, SI Materials and Methods. Next-generation sequencing and analysis of the ChIP DNA was carried out in the Glasgow Polyomics Facility (University of Glasgow). Raw data, analysis, and gene ontology studies can be accessed in SI Appendix, SI Materials and Methods (Datasets S2 and S3). A complete list of primers used for genotyping, qRT-PCR, cloning, and ChIP-qPCR are listed in SI Appendix, SI Materials and Methods and Dataset S1. Accession numbers of *Arabidopsis* genes that are the main focus of this study are At5g43630 (TZP) and At5g39760 (ZFHD10).

ACKNOWLEDGMENTS. We are grateful to Prof. Olivier Loudet (French National Institute for Agricultural Research, Versailles) for sharing the original 41AV and Bay-0 seeds. Many thanks to Prof. John M. Christie for critically reading the manuscript. We are indebted to Dr. J. Stuart Sullivan for providing the RNA from basal and apical hypocotyl segments. E.K. is indebted to the John Grieve Bequest for supporting her lectureship in biochemistry and the UK Biotechnology and Biological Sciences Research

Council for New Investigator Grant Award BB/M023079/1. The early stages of this work were supported by NIH Grant R01-GM52413 and the Howard Hughes Medical Institute (to J.C.). J.C. is an investigator at the Howard Hughes Medical Institute. E.K. was supported by fellowships from the Human Frontier Science Program (LT000159), the Salk Institute (Pioneer Fund,

Kirby and Stern Foundations), and the University of Glasgow (Lord Kelvin and Adam Smith Fellowship). G.B., J.L.P.-P., and S.A.K. acknowledge support from the NIH (R01 GM050006 and GM067837 to S.A.K.). M.L.H.D. is funded by a College of Medical, Veterinary, and Life Sciences PhD studentship from the University of Glasgow.

- Seaton DD, et al. (2015) Linked circadian outputs control elongation growth and flowering in response to photoperiod and temperature. *Mol Syst Biol* 11:776.
- Zhu JY, Oh E, Wang T, Wang ZY (2016) TOC1-PIF4 interaction mediates the circadian gating of thermoresponsive growth in Arabidopsis. *Nat Commun* 7:13692.
- Ezer D, et al. (2017) The evening complex coordinates environmental and endogenous signals in Arabidopsis. *Nat Plants* 3:17087.
- Pedmale UV, et al. (2016) Cryptochromes interact directly with PIFs to control plant growth in limiting blue light. *Cell* 164:233–245.
- de Wit M, Galvão VC, Fankhauser C (2016) Light-mediated hormonal regulation of plant growth and development. *Annu Rev Plant Biol* 67:513–537.
- Nusinow DA, et al. (2011) The ELF4-ELF3-LUX complex links the circadian clock to diurnal control of hypocotyl growth. *Nature* 475:398–402.
- Huang H, et al. (2016) PCH1 integrates circadian and light-signaling pathways to control photoperiod-responsive growth in Arabidopsis. *eLife* 5:e13292.
- Kaiserli E, et al. (2015) Integration of light and photoperiodic signaling in transcriptional nuclear foci. *Dev Cell* 35:311–321.
- Van Buskirk EK, Decker PV, Chen M (2012) Photobodies in light signaling. *Plant Physiol* 158:52–60.
- Galvão RM, et al. (2012) Photoactivated phytochromes interact with HEMERA and promote its accumulation to establish photomorphogenesis in Arabidopsis. *Genes Dev* 26:1851–1863.
- Chen M, et al. (2010) Arabidopsis HEMERA/pTAC12 initiates photomorphogenesis by phytochromes. *Cell* 141:1230–1240.
- Huang H, Nusinow DA (2016) Into the evening: Complex interactions in the Arabidopsis circadian clock. *Trends Genet* 32:674–686.
- Sawa M, Nusinow DA, Kay SA, Imaizumi T (2007) FKF1 and GIGANTEA complex formation is required for day-length measurement in Arabidopsis. *Science* 318:261–265.
- Inigo S, Alvarez MJ, Strasser B, Califano A, Cerdan PD (2012) PFT1, the MED25 subunit of the plant mediator complex, promotes flowering through CONSTANS dependent and independent mechanisms in Arabidopsis. *Plant J* 69:601–612.
- Liu X, et al. (2013) Phytochrome interacting factor3 associates with the histone deacetylase HDA15 in repression of chlorophyll biosynthesis and photosynthesis in etiolated Arabidopsis seedlings. *Plant Cell* 25:1258–1273.
- Oh E, Zhu JY, Ryu H, Hwang I, Wang ZY (2014) TOPLESS mediates brassinosteroid-induced transcriptional repression through interaction with BZR1. *Nat Commun* 5:4140.
- Wang L, Kim J, Somers DE (2013) Transcriptional corepressor TOPLESS complexes with pseudoresponse regulator proteins and histone deacetylases to regulate circadian transcription. *Proc Natl Acad Sci USA* 110:761–766.
- Endo M, Tanigawa Y, Murakami T, Araki T, Nagatani A (2013) Phytochrome-dependent late-flowering accelerates flowering through physical interactions with phytochrome B and constans. *Proc Natl Acad Sci USA* 110:18017–18022.
- Loudet O, et al. (2008) A zinc knuckle protein that negatively controls morning-specific growth in Arabidopsis thaliana. *Proc Natl Acad Sci USA* 105:17193–17198.
- Huang H, et al. (2016) Identification of evening complex associated proteins in Arabidopsis by affinity purification and mass spectrometry. *Mol Cell Proteomics* 15:201–217.
- Huang H, Alvarez S, Nusinow DA (2016) Data on the identification of protein interactors with the evening complex and PCH1 in Arabidopsis using tandem affinity purification and mass spectrometry (TAP-MS). *Data Brief* 8:56–60.
- Tan QK, Irish VF (2006) The Arabidopsis zinc finger-homeodomain genes encode proteins with unique biochemical properties that are coordinately expressed during floral development. *Plant Physiol* 140:1095–1108.
- Perotti MF, Ribone PA, Chan RL (2017) Plant transcription factors from the homeodomain-leucine zipper family I. Role in development and stress responses. *IUBMB Life* 69:280–289.
- Turchi L, Baima S, Morelli G, Ruberti I (2015) Interplay of HD-Zip II and III transcription factors in auxin-regulated plant development. *J Exp Bot* 66:5043–5053.
- Hu W, dePamphilis CW, Ma H (2008) Phylogenetic analysis of the plant-specific zinc finger-homeobox and mini zinc finger gene families. *J Integr Plant Biol* 50:1031–1045.
- Wier AD, Mayekar MK, Héroux A, Arndt KM, VanDemark AP (2013) Structural basis for Spt5-mediated recruitment of the Paf1 complex to chromatin. *Proc Natl Acad Sci USA* 110:17290–17295.
- de Jong RN, et al. (2008) Structure and DNA binding of the human Rtf1 Plus3 domain. *Structure* 16:149–159.
- Ciftci-Yilmaz S, Mittler R (2008) The zinc finger network of plants. *Cell Mol Life Sci* 65:1150–1160.
- Pruneda-Paz JL, et al. (2014) A genome-scale resource for the functional characterization of Arabidopsis transcription factors. *Cell Rep* 8:622–632.
- Choi H, et al. (2014) The homeodomain-leucine zipper ATHB23, a phytochrome B-interacting protein, is important for phytochrome B-mediated red light signaling. *Physiol Plant* 150:308–320.
- Walter M, et al. (2004) Visualization of protein interactions in living plant cells using bimolecular fluorescence complementation. *Plant J* 40:428–438.
- Kaiserli E, Sullivan S, Jones MA, Feeny KA, Christie JM (2009) Domain swapping to assess the mechanistic basis of Arabidopsis phototropin 1 receptor kinase activation and endocytosis by blue light. *Plant Cell* 21:3226–3244.
- Michael TP, et al. (2008) A morning-specific phytohormone gene expression program underlying rhythmic plant growth. *PLoS Biol* 6:e225.
- Mockler TC, et al. (2007) The diurnal project: Diurnal and circadian expression profiling, model-based pattern matching, and promoter analysis. *Cold Spring Harb Symp Quant Biol* 72:353–363.
- Kohnen MV, et al. (2016) Neighbor detection induces organ-specific transcriptomes, revealing patterns underlying hypocotyl-specific growth. *Plant Cell* 28:2889–2904.
- Sullivan S, et al. (2016) Functional characterization of Arabidopsis phototropin 1 in the hypocotyl apex. *Plant J* 88:907–920.
- Wendrich JR, et al. (2017) Framework for gradual progression of cell ontogeny in the Arabidopsis root meristem. *Proc Natl Acad Sci USA* 114:E8922–E8929.
- Sessa G, et al. (2005) A dynamic balance between gene activation and repression regulates the shade avoidance response in Arabidopsis. *Genes Dev* 19:2811–2815.
- Sawa S, et al. (2002) The HAT2 gene, a member of the HD-Zip gene family, isolated as an auxin inducible gene by DNA microarray screening, affects auxin response in Arabidopsis. *Plant J* 32:1011–1022.
- Kunihiro A, et al. (2011) Phytochrome-interacting factor 4 and 5 (PIF4 and PIF5) activate the homeobox ATHB2 and auxin-inducible IAA29 genes in the coincidence mechanism underlying photoperiodic control of plant growth of Arabidopsis thaliana. *Plant Cell Physiol* 52:1315–1329.
- Duek PD, Fankhauser C (2003) HFR1, a putative bHLH transcription factor, mediates both phytochrome A and cryptochrome signalling. *Plant J* 34:827–836.
- Hornitschek P, Lorrain S, Zoete V, Michielin O, Fankhauser C (2009) Inhibition of the shade avoidance response by formation of non-DNA binding bHLH heterodimers. *EMBO J* 28:3893–3902.
- de Wit M, et al. (2016) Integration of phytochrome and cryptochrome signals determines plant growth during competition for light. *Curr Biol* 26:3320–3326.
- Sasidharan R, Pierik R (2010) Cell wall modification involving XTHs controls phytochrome-mediated petiole elongation in Arabidopsis thaliana. *Plant Signal Behav* 5:1491–1492.
- Vissenberg K, et al. (2005) Differential expression of AtXTH17, AtXTH18, AtXTH19 and AtXTH20 genes in Arabidopsis roots. Physiological roles in specification in cell wall construction. *Plant Cell Physiol* 46:192–200.
- Leivar P, Quail PH (2011) PIFs: Pivotal components in a cellular signaling hub. *Trends Plant Sci* 16:19–28.
- Li L, et al. (2012) Linking photoreceptor excitation to changes in plant architecture. *Genes Dev* 26:785–790.
- Tran LS, et al. (2007) Co-expression of the stress-inducible zinc finger homeodomain ZFHD1 and NAC transcription factors enhances expression of the ERD1 gene in Arabidopsis. *Plant J* 49:46–63.
- Giuliano G, et al. (1988) An evolutionarily conserved protein binding sequence upstream of a plant light-regulated gene. *Proc Natl Acad Sci USA* 85:7089–7093.
- Michael TP, McClung CR (2003) Enhancer trapping reveals widespread circadian clock transcriptional control in Arabidopsis. *Plant Physiol* 132:629–639.
- Hudson ME, Quail PH (2003) Identification of promoter motifs involved in the network of phytochrome A-regulated gene expression by combined analysis of genomic sequence and microarray data. *Plant Physiol* 133:1605–1616.
- Srivastava R, et al. (2014) Distinct role of core promoter architecture in regulation of light-mediated responses in plant genes. *Mol Plant* 7:626–641.
- Han YF, et al. (2014) SUV2 is involved in transcriptional gene silencing by associating with SNF2-related chromatin-remodeling proteins in Arabidopsis. *Cell Res* 24:1445–1465.
- Khanna R, Kikis EA, Quail PH (2003) Early flowering 4 functions in phytochrome B-regulated seedling de-etiolation. *Plant Physiol* 133:1530–1538.
- Oh E, et al. (2014) Cell elongation is regulated through a central circuit of interacting transcription factors in the Arabidopsis hypocotyl. *eLife* 3:e03031.
- Wu MF, Tian Q, Reed JW (2006) Arabidopsis microRNA167 controls patterns of ARF6 and ARF8 expression, and regulates both female and male reproduction. *Development* 133:4211–4218.
- Nagpal P, et al. (2005) Auxin response factors ARF6 and ARF8 promote jasmonic acid production and flower maturation. *Development* 132:4107–4118.
- Nicolas M, Cubas P (2016) TCP factors: New kids on the signaling block. *Curr Opin Plant Biol* 33:33–41.
- Cubas P, Lauter N, Doebley J, Coen E (1999) The TCP domain: A motif found in proteins regulating plant growth and development. *Plant J* 18:215–222.
- Pruneda-Paz JL, Breton G, Para A, Kay SA (2009) A functional genomics approach reveals CHE as a component of the Arabidopsis circadian clock. *Science* 323:1481–1485.
- Windhövel A, Hein I, Dabrowa R, Stockhaus J (2001) Characterization of a novel class of plant homeodomain proteins that bind to the C4 phosphoenolpyruvate carboxylase gene of Flaveria trinervia. *Plant Mol Biol* 45:201–214.
- Krogan NT, Hogan K, Long JA (2012) APETALA2 negatively regulates multiple floral organ identity genes in Arabidopsis by recruiting the co-repressor TOPLESS and the histone deacetylase HDA19. *Development* 139:4180–4190.

